

LOG HOMOGENEOUS COMPACTIFICATIONS OF SOME CLASSICAL GROUPS

MATHIEU HURUGUEN

Received: January 8, 2014

Communicated by Alexander Merkurjev

ABSTRACT. We generalize in positive characteristics some results of Bien and Brion on log homogeneous compactifications of a homogeneous space under the action of a connected reductive group. We also construct an explicit smooth log homogeneous compactification of the general linear group by successive blow-ups starting from a grassmannian. By taking fixed points of certain involutions on this compactification, we obtain smooth log homogeneous compactifications of the special orthogonal and the symplectic groups.

2010 Mathematics Subject Classification: 14L35, 14M12, 14M17, 14M25

INTRODUCTION

Let k be an algebraically closed field and G a connected reductive group defined over k . Given a homogeneous space Ω under the action of the group G it is natural to consider equivariant COMPACTIFICATIONS or partial equivariant compactifications of it. EMBEDDINGS are normal irreducible varieties equipped with an action of G and containing Ω as a dense orbit, and compactifications are complete embeddings. Compactifications have shown to be powerful tools to produce interesting representations of the group G or to solve enumerative problems. In the influent paper [21], Luna and Vust developed a classification theory of embeddings of the homogeneous space Ω assuming that the field k is of characteristic zero. Their theory can be made very explicit and extended to all characteristics, see for instance [15], in the SPHERICAL case, that is, when a Borel subgroup of G possesses a dense orbit in the homogeneous space Ω . In this case, the embeddings of Ω are classified by combinatorial objects called COLORED FANS. If the homogeneous space is a torus acting on itself by

multiplication then one recovers the classification of torus embeddings or toric varieties in terms of fans, see for instance [14].

In the first part of the paper we focus on a certain category of “good” compactifications of the homogeneous space Ω . For example, these compactifications are smooth and the boundaries are strict normal crossing divisors. There are several notions of “good” compactifications in the literature. Some of them are defined by geometric conditions, as for example the TOROIDAL compactifications of Mumford [14], the REGULAR compactifications of Bifet De Concini and Procesi [3], the LOG HOMOGENEOUS compactifications of Brion [5] and some of them are defined by conditions from the embedding theory of Luna and Vust, as for example the COLORLESS compactifications. As it was shown by Bien and Brion [5], if the base field k is of characteristic zero then the homogeneous space Ω admits a log homogeneous compactification if and only if it is spherical, and in that case the four different notions of “good” compactifications mentioned above coincide. We generalize their results in positive characteristics in Section 1. We prove that a homogeneous space admitting a log homogeneous compactification is necessarily SEPARABLY SPHERICAL in the sense of Proposition-Definition 1.7. In that case, we relate the log homogeneous compactifications to the regular and the colorless one, see Theorem 1.8 for a precise statement. We do not know whether the condition of being separably spherical is sufficient for a homogeneous space to have a log homogeneous compactification. Along the way we prove Theorem 1.4, which is of independent interest, on the local structure of colorless compactifications of spherical homogeneous spaces, generalizing a result of Brion, Luna and Vust, see [7].

In Section 2, we focus on the explicit construction of equivariant compactifications of a connected reductive group. That is, the homogeneous space Ω is a connected reductive group G acted upon by $G \times G$ by left and right translations. The construction of “good” compactifications of a reductive group is a very old problem, with roots in the 19th century in the work of Chasles, Schubert, who were motivated by questions from enumerative geometry. When the group G is semi-simple there is a particular compactification \overline{G} called CANONICAL which possesses interesting properties, making it particularly convenient to work with. For example, the boundary is a divisor whose irreducible components intersect properly and the closure of the $G \times G$ -orbits are exactly the partial intersections of these prime divisors. Also, there is a unique closed orbit of $G \times G$ in the canonical compactification of G . Moreover, every toroidal compactification of G has a dominant equivariant morphism to \overline{G} . If the canonical compactification \overline{G} is smooth, then it is WONDERFUL in sense of Luna [20]. When the group G is of adjoint type, its canonical compactification is smooth, and there are many known constructions of this wonderful compactification, see for example [29], [17], [18], [19], [30], [27], [26] for the case of the projective linear group $\mathrm{PGL}(n)$ and [8], [24], [4] for the general case. In general the canonical compactification is not smooth, as it can be seen for example when G is the special orthogonal group $\mathrm{SO}(2n)$.

One way to construct a compactification of G is by considering a linear representation V of G and taking the closure of G in the projective space $\mathbb{P}(\text{End}(V))$. The compactifications arising in this way are called linear. It was shown by De Concini and Procesi [8] that the linear compactifications of a semi-simple group of adjoint type are of particular interest. Recently, Timashev [28], Gandini and Ruzzi [11], found combinatorial criterions for certain linear compactifications to be normal, or smooth. In [10], Gandini classifies the linear compactifications of the odd special orthogonal group having one closed orbit. By a very new and elegant approach, Martens and Thaddeus [22] recently discovered a general construction of the toroidal compactifications of a connected reductive group G as the coarse moduli spaces of certain algebraic stacks parametrizing objects called “framed principal G -bundles over chain of lines”.

Our approach is much more classical. In Section 2, we construct a log homogeneous compactification \mathcal{G}_n of the general linear group $\text{GL}(n)$ by successive blow-ups, starting from a grassmannian. The compactification \mathcal{G}_n is defined over an arbitrary base scheme. We then identify the compactifications of the special orthogonal group or the symplectic group obtained by taking the fixed points of certain involutions on the compactification \mathcal{G}_n . This provides a new construction of the wonderful compactification of the odd orthogonal group $\text{SO}(2n + 1)$, which is of adjoint type, of the symplectic group $\text{Sp}(2n)$, which is not of adjoint type, and of a toroidal desingularization of the canonical compactification of the even orthogonal group $\text{SO}(2n)$ having only two closed orbits. This is the minimal number of closed orbits on a smooth log homogeneous compactification, as the canonical compactification of $\text{SO}(2n)$ is not smooth.

Our procedure is similar to that used by Vainsencher, see [30], to construct the wonderful compactification of the projective linear group $\text{PGL}(n)$ or that of Kausz, see [13], to construct his compactification of the general linear group $\text{GL}(n)$. However, unlike Kausz, we are not able to describe the functor of points of our compactification \mathcal{G}_n . In that direction, we obtained a partial result in [12], where we describe the set $\mathcal{G}_n(K)$ for every field K . We decided not to include this description in the present paper, as it is long and technical. The functor of points of the wonderful compactification of the projective linear group is described in [27] and that of the symplectic group is described in [1].

ACKNOWLEDGMENTS

This paper is part of my Ph.D thesis. I would like to thank my supervisor Michel Brion for his precious advices and his careful reading. I would also like to thank Antoine Chambert-Loir and Philippe Gille for helpful comments on a previous version of this paper.

1 LOG HOMOGENEOUS COMPACTIFICATIONS

First we fix some notations. Let k be an algebraically closed field of arbitrary characteristic p . By a variety over k we mean a separated integral k -scheme of finite type. If X is a variety over k and x is a point of X , we denote by $T_{X,x}$ the tangent space of X at x . If Y is a subvariety of X containing x , we denote by $N_{Y/X,x}$ the normal space to Y in X at x .

For an algebraic group $G, H, P \dots$ we denote by the corresponding gothic letter $\mathfrak{g}, \mathfrak{h}, \mathfrak{p} \dots$ its Lie algebra. Let G be a connected reductive group defined over k . A G -variety is a variety equipped with an action of G . Let X be a G -variety. For each point $x \in X$ we denote by G_x the isotropy group scheme of x . We also denote by orb_x the morphism

$$orb_x : G \rightarrow X, \quad g \mapsto g \cdot x.$$

The orbit of x under the action of G is called separable if the morphism orb_x is, that is, if its differential is surjective, or, equivalently, if the group scheme G_x is reduced.

We fix a homogeneous space Ω under the action of G . Let X be a smooth compactification of Ω , that is, a complete smooth G -variety containing Ω as an open dense orbit. We suppose that the complement D of Ω in X is a strict normal crossing divisor.

In [3], Bifet, De Concini and Procesi introduce and study the regular compactifications of a homogeneous space over an algebraically closed field of characteristic zero. We generalize their definition in two different ways :

DEFINITION 1.1. *The compactification X is REGULAR (resp. STRONGLY REGULAR) if the orbits of G in X are separable, the partial intersections of the irreducible components of D are precisely the closures of the G -orbits in X and, for each point $x \in X$, the isotropy group G_x possesses an open (resp. open and separable) orbit in the normal space $N_{Gx/X,x}$ to the orbit Gx in X at the point x .*

If the characteristic of the base field k is zero, then the notion of regular and strongly regular coincide with the original notion of [3]. This is no longer true in positive characteristic, as we shall see at the end of Section 1.2.

In [5], Brion defines the log homogeneous compactifications over an algebraically closed field - throughout his paper the base field is also of characteristic zero, but the definition makes sense in arbitrary characteristic. Recall that the logarithmic tangent bundle $T_X(-\log D)$ is the vector bundle over X whose sheaf of section is the subsheaf of the tangent sheaf of X consisting of the derivations that preserve the ideal sheaf $\mathcal{O}_X(-D)$ of D . As G acts on X and D is stable under the action of G , it is easily seen that the infinitesimal action of the Lie algebra \mathfrak{g} on X gives rise to a natural vector bundle morphism:

$$X \times \mathfrak{g} \rightarrow T_X(-\log D).$$

We refer the reader to [5] for further details.

DEFINITION 1.2. *The compactification X is called LOG HOMOGENEOUS if the morphism of vector bundles on X :*

$$X \times \mathfrak{g} \rightarrow T_X(-\log D)$$

is surjective.

Assuming that the characteristic of the base field is zero, Bien and Brion prove in [2] that the homogeneous space Ω possesses a log homogeneous compactification if and only if it is spherical. In this case, they also prove that it is equivalent for a smooth compactification X of Ω to be log homogeneous, regular or to have no color - as an embedding of a spherical homogeneous space, see [15]. Their proof relies heavily on a local structure theorem for spherical varieties in characteristic zero established by Brion, Luna and Vust in [7].

A generalization of the local structure theorem was obtained by Knop in [16]; essentially, one has to replace in the statement of that theorem an isomorphism by a finite surjective morphism. In Section 1.1 we shall prove that under a separability assumption, the finite surjective morphism in Knop's theorem is an isomorphism. Then, in Section 1.2 we prove that the smooth compactification X of Ω is regular if and only if the homogeneous space Ω is spherical, the embedding X has no color and each closed orbit of G in X is separable (Theorem 1.5). We also prove that the smooth compactification X of Ω is strongly regular if and only if it is log homogeneous (Theorem 1.6). Finally, we exhibit a class of spherical homogeneous spaces for which the notion of regular and strongly regular compactifications coincide. In Section 1.3 we show that log homogeneity is preserved under taking fixed points by an automorphism of finite order prime to the characteristic of the base field k . In Section 1.4 we recall the classification of Luna and Vust in the setting of compactification of reductive groups, as this will be useful in Section 2.

1.1 A LOCAL STRUCTURE THEOREM

Let X be a smooth G -variety. We assume that there is a unique closed orbit ω of G in X and that this orbit is complete and separable. We fix a point x on ω . The isotropy group G_x is a parabolic subgroup of G . We fix a Borel subgroup B of G such that BG_x is open in G . We fix a maximal torus T of G contained in G_x and B and we denote by P the opposite parabolic subgroup to G_x containing B . We also denote by L the Levi subgroup of P containing T and by $R_u(P)$ the unipotent radical of P . With these notations we have the following proposition, which relies on a result of Knop [16, Theorem 1.2].

PROPOSITION 1.3. *There exists an affine open subvariety X_s of X which is stable under the action of P and a closed subvariety Z of X_s stable under the action of T , containing x such that:*

(1) The variety Z is smooth at x and the vector space $T_{Z,x}$ endowed with the action of T is isomorphic to the vector space $N_{\omega/X,x}$ endowed with the action of T .

(2) The morphism:

$$\mu : R_u(P) \times Z \rightarrow X_s, \quad (p, z) \mapsto p \cdot z$$

is finite, surjective, étale at (e, x) , and the fiber $\mu^{-1}(x)$ is reduced to the single point $\{(e, x)\}$.

Proof. As the smooth G -variety X has a unique closed orbit, it is quasi-projective by a famous result of Sumihiro, see [25]. We fix a very ample line bundle \mathcal{L} on X . We fix a G -linearization of this line bundle. By [16, Theorem 2.10], there exists an integer N and a global section s of \mathcal{L}^N such that the nonzero locus X_s of s is an affine open subvariety containing the point x and the stabilizer of the line spanned by s in the vector space $H^0(X, \mathcal{L}^N)$ is P . The open subvariety X_s is therefore affine, contains the point x and is stable under the action of the parabolic subgroup P . Using the line bundle \mathcal{L}^N , we embed X into a projective space $\mathbb{P}(V)$ on which G acts linearly. We choose a T -stable complement S to $T_{\omega,x}$ in the tangent space $T_{\mathbb{P}(V),x}$, such that S is the direct sum of a T -stable complement of $T_{\omega,x}$ in $T_{X,x}$ and a T -stable complement of $T_{X,x}$ in $T_{\mathbb{P}(V),x}$. This is possible because T is a linearly reductive group.

We consider now the linear subspace S' of $\mathbb{P}(V)$ containing x and whose tangent space at x is S . It is a T -stable subvariety of $\mathbb{P}(V)$. By [16, Theorem 1.2], there is an irreducible component Z of $X_s \cap S'$ containing x and such that the morphisms

$$\begin{aligned} \mu : R_u(P) \times Z &\rightarrow X_s, & (p, z) &\mapsto p \cdot z \\ \nu : Z &\rightarrow X_s/R_u(P), & z &\mapsto zR_u(P) \end{aligned}$$

are finite and surjective. Moreover, the fiber $\mu^{-1}(x)$ is reduced to the single point (e, x) . We observe now that S' intersects X_s transversally at x . This implies that the subvariety Z is smooth at x . It is also T -stable, as an irreducible component of $X_s \cap S'$. By definition, the parabolic subgroup P contains the Borel subgroup B , therefore the orbit $Px = R_u(P)x$ is open in ω . Moreover, we have the direct sum decomposition $\mathfrak{g} = \mathfrak{g}_x \oplus \mathfrak{p}_u$, where \mathfrak{p}_u is the Lie algebra of the unipotent radical $R_u(P)$ of P . The morphism

$$d_e orb_x : \mathfrak{g} \rightarrow T_{\omega,x}$$

is surjective and identically zero on \mathfrak{g}_x . This proves that the restriction of this morphism to \mathfrak{p}_u is an isomorphism. The morphism

$$\mu : R_u(P) \times Z \rightarrow X_s, \quad (p, z) \mapsto p \cdot z$$

is therefore étale at (e, x) . Indeed, its differential at this point is:

$$\mathfrak{p}_u \times T_{Z,x} \rightarrow T_{X,x}, \quad (h, k) \mapsto d_e orb_x(h) + k.$$

We also see that the spaces $T_{Z,x}$ and $N_{\omega/X,x}$ endowed with their action of the torus T are isomorphic, completing the proof of the proposition. \square

We now suppose further that X is an embedding of the homogeneous space Ω . With this additional assumption we have:

THEOREM 1.4. *The following three properties are equivalent:*

- (1) *The homogeneous space Ω is spherical and the embedding X has no color.*
- (2) *The torus T possesses an open orbit in the normal space $N_{\omega/X,x}$. Moreover, the complement D of Ω in X is a strict normal crossing divisor and the partial intersections of the irreducible components of D are the closure of the G -orbits in X .*
- (3) *The set $X_0 = \{y \in X, x \in \overline{By}\}$ is an affine open subvariety of X which is stable by P . Moreover, there exists a closed subvariety Z of X_0 which is smooth, stable by L , on which the derived subgroup $[L, L]$ acts trivially and containing an open orbit of the torus $L/[L, L]$, such that the morphism:*

$$R_u(P) \times Z \rightarrow X_0, \quad (p, z) \mapsto p \cdot z$$

is an isomorphism. Finally, each orbit of G in X intersects Z along a unique orbit of T .

Proof. (3) \Rightarrow (1) As T possesses an open orbit in Z , we see that the Borel subgroup B has an open orbit in X , and the homogeneous space Ω is spherical. Moreover, let D be a B -stable prime divisor on X containing ω . Using the isomorphism in (3) we can write

$$D \cap X_0 = R_u(P) \times (D \cap Z).$$

As $D \cap Z$ is a closed irreducible T -stable subvariety of Z , it is the closure of a T -orbit in Z . As the T -orbits in Z corresponds bijectively to the G -orbits in X , we see that D is the closure of a G -orbit in X and is therefore stable under the action of G . This proves that the embedding X of Ω has no color.

(3) \Rightarrow (2) The isomorphism in (3) proves that the spaces $T_{Z,x}$ and $N_{\omega/X,x}$ endowed with their actions of the torus T are isomorphic. As T possesses an open dense orbit in the first one, it also has an open dense orbit in the latter. As Z is smooth toric variety, we see that the complement of the open orbit of T in Z is a strict normal crossing divisor whose associated strata are the T -orbits in Z . Using the isomorphism given by (3), we see that the complement of the open orbit of the parabolic subgroup P in X_0 is a strict normal crossing divisor whose associated strata are the products $R_u(P) \times \Omega'$, where Ω' runs over the set of T -orbits in Z . To complete the proof that property (2) is satisfied, we translate the open subvariety X_0 by elements of G and we use the fact that each G -orbit in X intersects Z along a unique T -orbit.

(1) \Rightarrow (3) We use the notations of Proposition 1.3. By [15, Lemma 6.5] the fact that the embedding X has no color implies that the parabolic subgroup P is the stabilizer of the open B -orbit Ω in X . Using this fact and [16, Theorem 2.8] we obtain that the derived subgroup of P , and therefore the derived group of L , acts trivially on $X_s/R_u(P)$. Moreover, as the homogeneous space Ω is spherical, the Levi subgroup L has an open orbit in $X_s/R_u(P)$. The torus T has therefore an open orbit in $X_s/R_u(P)$, as the derived group of L acts trivially. Using the finite surjective morphism ν appearing in the proof of Proposition 1.3, we see that T has an open orbit in Z . Z is therefore a smooth affine toric variety with a fixed point under the action of a quotient of T . Moreover, as the subvariety Z is left stable under the action of T and the derived group $[L, L]$ acts trivially on Z , we see that the Levi subgroup L leaves the subvariety Z invariant.

We observe now that the locus of points of $R_u(P) \times Z$ where μ is not étale is closed and stable under the actions of $R_u(P)$ and T . The unique closed orbit of $R_u(P) \rtimes T$ in $R_u(P) \times Z$ is $R_u(P)x$ and μ is étale at (e, x) , therefore we obtain that μ is an étale morphism. As the morphism μ is also finite of degree 1 (the fiber of $\{x\}$ being reduced to a single point), it is an isomorphism.

We prove now that each G -orbit in X intersects Z along a unique T -orbit. First, we observe that, as ω is the unique closed orbit of G in X , the open subvariety X_s intersects every G -orbit. We shall prove that the closures of the G -orbits in X corresponds bijectively to the closures of the T -orbits in Z . Let X' be the closure of a G -orbit in X . As X' is the closure of $X' \cap X_s$, it is also equal, using the isomorphism μ , to the closure of $R_u(P)(X' \cap Z)$. The closed subvariety $X' \cap Z$ of Z is therefore a closed irreducible T -stable subvariety. We can conclude that it is the closure of a T -orbit in Z . Conversely let Z' be the closure of a T -orbit in Z . As Z is a smooth toric variety, we can write

$$Z' = D'_1 \cap D'_2 \cap \dots \cap D'_r,$$

where the D'_i s are T -stable prime divisors on Z . We observe that the prime divisors

$$\overline{R_u(P)D'_1}, \dots, \overline{R_u(P)D'_r}$$

on X are stable under the action of P . Indeed, the orbits of P in X_s are exactly the orbits of $R_u(P) \rtimes T$ in X_s . As X has no color, the fact that these divisors contain the closed orbit ω proves that they are stable under the action of G . Their intersection $\overline{R_u(P)Z}$ is also G -stable. As it is irreducible, we can conclude that it is the closure of a G -orbit in X .

In order to complete the proof that (1) \Rightarrow (3), it remains to show that

$$X_s = \{y \in X, \quad x \in \overline{By}\}.$$

Let y be a point on X such that x belongs to \overline{By} . The intersection $X_s \cap \overline{By}$ is a non empty open subset of \overline{By} which is stable under the action of B . Therefore

it contains \overline{y} , that is, y belongs to X_s . Now let y be a point on X_s . The closed subvariety \overline{By} contains a closed B -orbit in X_s . As the unique closed orbit of B in X_s is the orbit of x , we see that x belongs to \overline{By} , completing the argument.

(2 \Rightarrow 3) We use the notations introduced in Proposition 1.3. By assumption, the torus T possesses an open orbit in the normal space $N_{\omega/X,x}$. Moreover, by Proposition 1.3, the spaces $T_{Z,x}$ and $N_{\omega/X,x}$ endowed with their actions of T are isomorphic. Therefore, the torus T possesses an open dense orbit in $T_{Z,x}$. It is then an easy exercise left to the reader to prove that the variety Z is a smooth toric variety for a quotient of T . The same arguments as above prove that the morphism μ is an isomorphism.

We prove now that each G -orbit in X intersects Z along a unique orbit of T . Let D be the complement of Ω in X . By assumption, it is a strict normal crossing divisor whose associated strata are the G -orbits in X . We denote by D_1, \dots, D_r the irreducible component of D . As there is a unique closed orbit of G on X each partial intersection $\bigcap_{i \in I} D_i$ is non empty and irreducible or, in other words, it is a stratum of D . The integer r is the codimension of the closed orbit ω in X , and there are exactly 2^r G -orbits in X . As the variety Z is a smooth affine toric variety of dimension r with a fixed point, we see that there are exactly 2^r orbits of T on Z . As each orbit of G in X intersect Z we see that the intersection of a G -orbit with Z is a single T -orbit.

Finally, we prove that the open subvariety X_s is equal to X_0 by the same argument as in the proof of (1) \Rightarrow (3), completing the proof of the theorem. \square

1.2 REGULAR, STRONGLY REGULAR AND LOG HOMOGENEOUS COMPACTIFICATIONS

In this section we use the following notation. Let X be a G -variety with a finite number of orbits (for example, a spherical variety). Let ω be an orbit of G in X . We denote by

$$X_{\omega,G} = \{y \in X, \omega \subseteq \overline{Gy}\}.$$

It is an open G -stable subvariety of X in which ω is the unique closed orbit.

THEOREM 1.5. *Let X be a smooth compactification of the homogeneous space Ω . The following two properties are equivalent:*

- (1) X is regular.
- (2) The homogeneous space Ω is spherical, the embedding X has no color and the orbits of G in X are separable.

Proof. Suppose that X is regular. Let D be the complement of Ω in X . It is a strict normal crossing divisor. Let ω be a closed, and therefore complete and separable, orbit of G in X . We use the notations introduced at the beginning of Section 1.4 with $X_{\omega,G}$ in place of X . The normal space $N_{\omega/X,x}$ is the normal space to a stratum of the divisor D and therefore possesses a natural direct

sum decomposition into a sum of lines, each of them being stable under the action of G_x (which is connected, as it is a parabolic subgroup of G). Therefore the representation of G_x in $N_{\omega/X,x}$ factors through the action of a torus. This proves that the derived group of L acts trivially in this space, proving that the torus T has a dense orbit in $N_{\omega/X,x}$. By Theorem 1.4 (applied to $X_{\omega,G}$) the homogeneous space Ω is spherical and the embedding $X_{\omega,G}$ has no color. As this is true for each closed orbit ω of G in X , we see that the embedding X has no color.

We assume now that Ω is spherical, X has no color and that each orbit of G in X is separable. By applying Theorem 1.4 to each open subvariety $X_{\omega,G}$, where ω runs over the set of closed orbits of X , we see that the complement D of Ω in X is a strict normal crossing divisor and that, for each point x in X , the isotropy group G_x has an open orbit in the normal space $N_{Gx/X,x}$. Moreover, by assumption, the G -orbits in X are separable. To complete the proof of the theorem, it remains to show that the partial intersections of the irreducible components of D are irreducible. But this is true on every colorless embedding of a spherical homogeneous space, due to the combinatorial description of these embeddings, see [15, Section 3]. \square

THEOREM 1.6. *Let X be a smooth compactification of Ω . The following two properties are equivalent:*

- (1) X is a log homogeneous compactification.
- (2) X is strongly regular.

Proof. We suppose first that the compactification X is log homogeneous. We denote by D the complement of Ω in X . It is a strict normal crossing divisor. Following the argument given in [5, Proposition 2.1.2] we prove that each stratum of the strict normal crossing divisor D is a single orbit under the action of G which is separable and that for each point $x \in X$, the isotropy group G_x possesses an open and separable orbit in the normal space $N_{Gx/X,x}$. In order to conclude, it remains to prove that the partial intersection of the irreducible components of D are irreducible. But the same argument as in the proof of Theorem 1.5 prove that Ω is spherical and X has no color, which is sufficient to complete the proof.

Conversely, if X is supposed to be strongly regular, the proof of [5, Proposition 2.1.2] adapts without change and shows that X is a log homogeneous compactification of Ω . \square

PROPOSITION-DEFINITION 1.7. *If the homogeneous space Ω possesses a log homogeneous compactification, then it satisfies the following equivalent conditions:*

- (1) *The homogeneous space Ω is spherical and there exists a Borel subgroup of G whose open orbit in Ω is separable.*

- (2) The homogeneous space Ω is spherical and the open orbit of each Borel subgroup of G in Ω is separable.
- (3) The homogeneous space Ω is separable under the action of G , and there exists a point x in X and a Borel subgroup B of G such that $\mathfrak{b} + \mathfrak{g}_x = \mathfrak{g}$.

A homogeneous space satisfying one of these properties is said to be SEPARABLY SPHERICAL.

Proof. We suppose first that the homogeneous space Ω possesses a log homogeneous compactification X and we prove that it satisfies the first condition. By Theorem 1.6 and 1.5, the homogeneous space Ω is spherical. Let ω be a closed, and therefore complete and separable, orbit of G in X . We apply Theorem 1.4 to the open subvariety $X_{\omega,G}$. We use the notations introduced for this theorem. As X is strongly regular, the maximal torus T has an open and separable orbit in $T_{Z,x} = N_{\omega/X,x}$. As this space endowed with its action of T is isomorphic to Z endowed with its action of T , because Z is an affine smooth toric variety with fixed point for a quotient of T , we see that the open orbit of T in Z is separable. Consequently, the open orbit of $R_u(P) \times T$ in $R_u(P) \times Z$ is separable, and the open orbit of B in Ω is separable.

We prove now that the three conditions in the statement of the proposition-definition are equivalent. As the Borel subgroups of G are conjugated, condition (1) and (2) are equivalent. Suppose now that condition (1) is satisfied. Let B be a Borel subgroup of G and x a point in the open and separable orbit of B in Ω . The linear map $d_e orb_x : \mathfrak{b} \rightarrow T_{Bx,x}$ is surjective. As the orbit Bx is open in Ω we see that the homogeneous space Ω is separable under the action of G and that

$$\mathfrak{b} + \mathfrak{g}_x = \mathfrak{g}.$$

Conversely, we suppose that condition (3) is satisfied. As the homogeneous space Ω is separable, the linear map

$$d_e orb_x : \mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{g}_x$$

is the natural projection. As we have $\mathfrak{b} + \mathfrak{g}_x = \mathfrak{g}$, we see that the linear map

$$d_e orb_x : \mathfrak{b} \rightarrow \mathfrak{g}/\mathfrak{g}_x$$

is surjective. This means precisely that the orbit Bx is open in Ω and separable. \square

Here are some example of separably spherical homogeneous spaces: separable quotients of tori, partial flag varieties, symmetric spaces in characteristic not 2 (Vust proves in [31] that symmetric spaces in characteristic zero are spherical; his proof extends to characteristic not 2 to show that symmetric spaces are separably spherical).

THEOREM 1.8. *We assume that the homogeneous space Ω is separably spherical. Let X be a smooth compactification of Ω . The following conditions are equivalent:*

- (1) X has no color and the closed orbits of G in X are separable.
- (2) X is regular.
- (3) X is strongly regular.
- (4) X is log homogeneous under the action of G .

Proof. In view of Theorem 1.5 and 1.6 it suffices to show that (1) \Rightarrow (3). We assume that condition (1) is satisfied. Let ω be a closed, and therefore separable orbit of G in X . We apply Theorem 1.4 to the open subvariety $X_{\omega,G}$ of X introduced in the proof of Theorem 1.5. We use the notations introduced for Theorem 1.4. As the open orbit of B in Ω is separable, we see that the quotient of T acting on Z is separable. As Z is a smooth affine toric variety with fixed point under this quotient, we see that the orbits of T in Z are all separable and that for each point $z \in Z$, the stabilizer T_z has an open and separable orbit in the normal space $N_{Tz/Z,z}$. From this we get readily that the embedding $X_{\omega,G}$ of Ω satisfies the conditions defining a strongly regular embedding. As this is true for each closed orbit ω , we see that X is a strongly regular compactification of Ω . \square

We end this section with an example of a regular compactification of a homogeneous space which is not strongly regular. We suppose that the base field k has characteristic 2. Let G be the group $\mathrm{SL}(2)$ acting on $X := \mathbb{P}^1 \times \mathbb{P}^1$. There are two orbits: the open orbit Ω of pairs of distinct points and the closed orbit ω , the diagonal, which has codimension one in X . These orbits are separable under the action of G . Moreover, the complement of the open orbit, that is, the closed orbit ω , is a strict normal crossing divisor and the partial intersections of its irreducible components are the closure of G -orbits in X . A quick computation shows that for each point on the closed orbit ω , the isotropy group has an open non separable orbit in the normal space to the closed orbit at that point. Therefore the compactification X of Ω is regular and not strongly regular. By Theorem 1.8 the homogeneous space cannot be separably spherical. This can be seen directly as follows. The homogeneous space Ω is the quotient of G by a maximal torus T . A Borel subgroup B of G has an open orbit in Ω if and only if it does not contain T . But in that case the intersection $B \cap T$ is the center of G , which is not reduced because the characteristic of the base field is 2.

1.3 LOG HOMOGENEOUS COMPACTIFICATIONS AND FIXED POINTS

Let X be a smooth variety over the field k and σ an automorphism of X which has finite order r prime to the characteristic p of k . Fogarty proves in [9] that the fixed point subscheme X^σ is smooth and that, for each fixed point x of σ in X , the tangent space to X^σ at x is $T_{X,x}^\sigma$.

We suppose now that X is a smooth log homogeneous compactification of the homogeneous space Ω . We also assume that the automorphism σ leaves Ω stable and is G -equivariant, in the sense that there exists an automorphism σ of the group G satisfying

$$\forall g \in G, \quad \forall x \in X, \quad \sigma(gx) = \sigma(g)\sigma(x).$$

By [23, Proposition 10.1.5], the neutral component G' of the group G^σ is a reductive group. Moreover, each connected component of the variety Ω^σ is a homogeneous space under the action of G' . We let Ω' be such a component and X' be the connected component of X^σ containing Ω' .

PROPOSITION 1.9. *X' is a log homogeneous compactification of Ω' under the action of G' .*

Proof. Let D be the complement of Ω in X . Let x be a point in X' . Let D_1, \dots, D_s be the irreducible components of D containing x . First we prove that the intersection $D' := D \cap X'$ is a strict normal crossing divisor. For each index i , the intersection $D'_i := D_i \cap X'$ is a divisor on X' . Indeed, X' is not contained in D_i as it contains Ω' . As x is fixed by the automorphism σ , we can assume that the components D_i s are ordered in such a way that

$$\begin{aligned} \sigma(D_2) = D_1 \quad \dots \quad \sigma(D_{i_1}) = D_{i_1-1}, \quad \sigma(D_1) = D_{i_1} \\ \dots \\ \sigma(D_{i_{t-1}+2}) = D_{i_{t-1}+1} \quad \dots \quad \sigma(D_{i_t}) = \sigma(D_s) = D_{i_t-1}, \quad \sigma(D_{i_{t-1}+1}) = D_{i_t}. \end{aligned}$$

By convention we define $i_0 = 0$. For each integer j from 1 to t , and each integer i from $i_{j-1} + 1$ to i_j we have $D'_i = D'_{i_j}$. Therefore we see that D'_{i_j} is the connected component of the smooth variety $(D_{i_{j-1}+1} \cap \dots \cap D_{i_j})^\sigma$ containing x . Consequently, it is smooth. For the moment, we have proved that D' is a divisor on X' whose irreducible components are smooth.

We prove now that the divisor $D'_{i_1}, \dots, D'_{i_t}$ intersect transversally at the point x . Let U_x be an open neighborhood of x in X which is stable by the automorphism σ and on which the equation of D is $u_1 \dots u_s = 0$, where $u_1 \dots u_s \in \mathcal{O}_X(U_x)$ are part of a regular local parameter system at x and satisfy:

$$\begin{aligned} \sigma(u_2) = u_1 \quad \dots \quad \sigma(u_{i_1}) = u_{i_1-1} \\ \dots \\ \sigma(u_{i_{t-1}+2}) = u_{i_{t-1}+1} \quad \dots \quad \sigma(u_{i_t}) = u_{i_t-1}. \end{aligned}$$

We aim to prove that the images of the differential $d_x u_{i_j}$ by the natural projection

$$(T_{X,x})^* \rightarrow (T_{X',x})^*$$

are linearly independent, where j run from 1 to t . As the point x is fixed by σ , σ acts by the differential on the tangent space $T_{X,x}$ and by the dual action

on $(T_{X,x})^*$. As the order of the automorphism σ is prime to the characteristic p , we have a direct sum decomposition:

$$(T_{X,x})^* = ((T_{X,x})^*)^\sigma \oplus \text{Ker}(id + \sigma + \cdots + \sigma^{r-1})$$

where the projection on the first factor is given by

$$l \mapsto \frac{1}{r}(l + \sigma(l) + \cdots + \sigma^{r-1}(l)).$$

Moreover, as $T_{X',x}$ is equal to $(T_{X,x})^\sigma$, the second factor in this decomposition is easily seen to be $(T_{X',x})^\perp$, so that the natural projection

$$(T_{X,x})^* \rightarrow (T_{X',x})^*$$

gives an isomorphism

$$((T_{X,x})^*)^\sigma \rightarrow (T_{X',x})^*.$$

Finally, the images of the differential $d_x u_{i_j}$ in $(T_{X',x})^*$ are linearly independent, because the differentials $d_x u_i$ are linearly independent in $(T_{X,x})^*$.

We have proved that the divisor D' is a strict normal crossing divisor. We leave it as an exercise to the reader to prove that there exists a natural exact sequence of vector bundle on X'

$$0 \rightarrow T_{X'}(-\log D') \rightarrow T_X(-\log D)|_{X'} \rightarrow N_{X'/X} \rightarrow 0,$$

and that the space $T_{X'}(-\log D')_x$ is the subspace of fixed point by σ in the space $T_X(-\log D)_x$. Now, the compactification X is log homogeneous, therefore the linear map

$$\mathfrak{g} \rightarrow T_X(-\log D)_x$$

is surjective. As r and p are relatively prime, this linear map is still surjective at the level of fixed points. That is, the linear map

$$\mathfrak{g}^\sigma \rightarrow T_X(-\log D)_x^\sigma = T_{X'}(-\log D')_x$$

is surjective. This complete the proof of the proposition. \square

1.4 THE EXAMPLE OF REDUCTIVE GROUPS

In this section the homogeneous space Ω is a connected reductive group G acted upon by the group $G \times G$ by the following formula:

$$\forall (g, h) \in G \times G, \quad \forall x \in G, \quad (g, h) \cdot x = gxh^{-1}$$

We would like to explain here the classification of smooth log homogeneous compactifications of G . Observe that the homogeneous space G under the action of $G \times G$ is actually separably spherical. By Theorem 1.8, its smooth log homogeneous compactifications are the smooth colorless compactifications

with separable closed orbits. The last condition is actually superfluous : by [6, Chapter 6], the closed orbits of $G \times G$ in a colorless compactification of G are isomorphic to $G/B \times G/B$, where B is a Borel subgroup of G . The log homogeneous compactifications of G are therefore the smooth colorless one.

We now recall the combinatorial description of the smooth colorless compactifications of G . Let T be a maximal torus of G and B a Borel subgroup of G containing T . We denote by V the \mathbb{Q} -vector space spanned by the one-parameter subgroups of T and by \mathcal{W} the Weyl chamber corresponding to B . Let X be a smooth colorless embedding of G . We let the torus T act “on the left” on X . For this action, the closure of T in X is a smooth complete toric variety. We associate to X the fan consisting of those cones in the fan of the toric variety \overline{T} which are included in $-\mathcal{W}$. This sets a map from the set of smooth colorless compactifications of G to the set of fans in V with support $-\mathcal{W}$ and which are smooth with respect to the lattice of one parameter subgroups in V . This map is actually a bijection, see for instance [6, Chapter 6].

2 EXPLICIT COMPACTIFICATIONS OF CLASSICAL GROUPS

We construct a log homogeneous compactification \mathcal{G}_n of the general linear group $\mathrm{GL}(n)$ by successive blow-ups, starting from a grassmannian. The precise procedure is explained in Section 2.1. The compactification \mathcal{G}_n is defined over an arbitrary base scheme. In Section 2.2 we study the local structure of the action of $\mathrm{GL}(n) \times \mathrm{GL}(n)$ on \mathcal{G}_n , still over an arbitrary base scheme. This enables us to compute the colored fan of \mathcal{G}_n over an algebraically closed field in Section 2.4. Using this computation, we are able to identify the compactifications of the special orthogonal group or the symplectic group obtained by taking the fixed points of certain involutions on the compactification \mathcal{G}_n . In the odd orthogonal and symplectic case we obtain the wonderful compactification. In the even orthogonal case we obtain a log homogeneous compactification with two closed orbits. This is the minimal number of closed orbits on a smooth log homogeneous compactification, as the canonical compactification of $\mathrm{SO}(2n)$ is not smooth.

2.1 THE COMPACTIFICATIONS \mathcal{G}_m

As we mentioned above our construction works over an arbitrary base scheme: until the end of Section 2.3 we work over a base scheme S . Let \mathcal{V}_1 and \mathcal{V}_2 be two free modules of constant finite rank n on S . We denote by \mathcal{V} the direct sum of \mathcal{V}_1 and \mathcal{V}_2 . We denote by p_1 and p_2 the projections respectively on the first and the second factor of this direct sum. We denote by G the group scheme $\mathrm{GL}(\mathcal{V}_1) \times \mathrm{GL}(\mathcal{V}_2)$ which is a subgroup scheme of $\mathrm{GL}(\mathcal{V})$.

DEFINITION 2.1. *We denote by $\Omega := \mathrm{Iso}(\mathcal{V}_2, \mathcal{V}_1)$ the scheme over S parametrizing the isomorphisms from \mathcal{V}_2 to \mathcal{V}_1 .*

There is a natural action of the group scheme G on Ω , via the following formulas

$$\forall (g_1, g_2) \in G, \quad \forall x \in \Omega, \quad (g_1, g_2) \cdot x = g_1 x g_2^{-1}$$

For this action, Ω is a homogeneous space under the action of G .

DEFINITION 2.2. *We denote by \mathcal{G} the grassmannian*

$$\pi : \mathcal{G}_{r_S}(n, \mathcal{V}) \rightarrow S$$

parametrizing the submodules of \mathcal{V} which are locally direct summands of rank n . We denote by \mathcal{T} the tautological module on \mathcal{G} .

The module \mathcal{T} is a submodule of $\pi^*\mathcal{V}$ which is locally a direct summand of finite constant rank n . There is a natural action of the group scheme $\mathrm{GL}(\mathcal{V})$, and therefore of the group scheme G , on the grassmannian \mathcal{G} . Moreover, Ω is contained in \mathcal{G} as a G -stable open subscheme via the graph

$$\Omega \rightarrow \mathcal{G}, \quad x \mapsto \mathrm{Graph}(x).$$

DEFINITION 2.3. *We denote by p the following morphism of modules on the grassmannian \mathcal{G} :*

$$p = \pi^*p_1 \oplus \pi^*p_2 : \mathcal{T}^{\oplus 2} \rightarrow \pi^*\mathcal{V}.$$

DEFINITION 2.4. *For $d \in \llbracket 0, n \rrbracket$, we denote by \mathcal{H}_d the locally free module*

$$\mathcal{H}om\left(\bigwedge^{n+d} (\mathcal{T}^{\oplus 2}), \bigwedge^{n+d} (\pi^*\mathcal{V})\right).$$

on the grassmannian \mathcal{G} .

DEFINITION 2.5. *For $d \in \llbracket 0, n \rrbracket$, the exterior power $\wedge^{n+d} p$ is a global section of \mathcal{H}_d . We denote by \mathcal{Z}_d the zero locus of $\wedge^{n+d} p$ on the grassmannian \mathcal{G} .*

We define in this way a sequence of G -stable closed subschemes on the grassmannian \mathcal{G}

$$\mathcal{Z}_0 \subset \mathcal{Z}_1 \subset \cdots \subset \mathcal{Z}_n \subset \mathcal{G}.$$

Observe that the closed subscheme \mathcal{Z}_0 is actually empty. Moreover, it is easy to prove that the open subscheme Ω is the complement of \mathcal{Z}_n in \mathcal{G} .

We will now define a sequence of blow-ups

$$\mathcal{G}_n \xrightarrow{b_n} \mathcal{G}_{n-1} \longrightarrow \cdots \longrightarrow \mathcal{G}_1 \xrightarrow{b_1} \mathcal{G}_0$$

and, for each integer m between 0 and n , a family of closed subschemes $\mathcal{Z}_{m,d}$ of \mathcal{G}_m , where d runs from m to n .

DEFINITION 2.6. *Let $m \in \llbracket 0, n \rrbracket$ and $d \in \llbracket m, n \rrbracket$. The definition is by induction:*

- For $m = 0$, we set $\mathcal{G}_0 := \mathcal{G}$ and $\mathcal{Z}_{0,d} := \mathcal{Z}_d$.
- Assuming that the scheme \mathcal{G}_{m-1} and its subschemes $\mathcal{Z}_{m-1,d}$ are defined, we define

$$b_m : \mathcal{G}_m \rightarrow \mathcal{G}_{m-1}$$

to be the blow-up centered at $\mathcal{Z}_{m-1,m}$ and, for each integer d from m to n , we define $\mathcal{Z}_{m,d}$ to be the strict transform of $\mathcal{Z}_{m-1,d}$ that is, the schematic closure of

$$b_m^{-1}(\mathcal{Z}_{m-1,d} \setminus \mathcal{Z}_{m-1,m})$$

in \mathcal{G}_m .

Moreover, we denote by $\mathcal{I}_{m,d}$ the ideal sheaf on \mathcal{G}_m defining $\mathcal{Z}_{m,d}$.

The group scheme G acts on the schemes \mathcal{G}_m and leaves the subschemes $\mathcal{Z}_{m,d}$ globally invariant. Modulo Proposition 2.17 below, we prove now:

THEOREM 2.7. *For each integer m from 0 to $n - 1$, the S -scheme \mathcal{G}_m is a smooth projective compactification of Ω .*

Proof. By Proposition 2.17 the scheme \mathcal{G}_m is covered by a collection of open subschemes isomorphic to affine spaces over S . In particular, the S -scheme \mathcal{G}_m is smooth. It is a classical fact that the grassmannian \mathcal{G} is projective over S . As the blow-up of a projective scheme over S along a closed subscheme is projective over S , we see that \mathcal{G}_m is projective over S . Finally, observe that the open subscheme Ω of \mathcal{G} is disjoint from the closed subscheme \mathcal{Z}_n and therefore from each of the closed subscheme \mathcal{Z}_d . As a consequence, Ω is an open subscheme of each of the \mathcal{G}_m . □

2.2 AN ATLAS OF AFFINE CHARTS FOR \mathcal{G}_m

Let V be the set $\llbracket 1, n \rrbracket \times \{1, 2\}$. We denote by V_1 the subset $\llbracket 1, n \rrbracket \times \{1\}$ and by V_2 the subset $\llbracket 1, n \rrbracket \times \{2\}$. We shall refer to elements of V_1 as elements of type 1 and elements of V_2 as elements of type 2. We fix a basis $v_i, i \in V$, of the free module \mathcal{V} . We suppose that $v_i, i \in V_1$ is a basis for \mathcal{V}_1 and $v_i, i \in V_2$ is a basis for \mathcal{V}_2 . Moreover, for each subset I of V , we denote by \mathcal{V}_I the free submodule of \mathcal{V} spanned by the $(v_i)_{i \in I}$. For every integer m from 1 to n , we denote by $V^{>m}$ the set $\llbracket m + 1, n \rrbracket \times \{1, 2\}$. We define the sets $V^{\geq m}, V^{<m}$ and $V^{\leq m}$ similarly. We also have, with obvious notations, the sets $V_1^{>m}, V_2^{>m}, V_1^{\geq m}, V_2^{\geq m} \dots$. If I is a subset of V we denote by I_1 the set $I \cap V_1$ and by I_2 the set $I \cap V_2$.

One word on terminology. If X is an S -scheme, by a point x of X we mean an S -scheme S' and a point x of the set $X(S')$. However, as it is usually unnecessary, we do not mention the S -scheme S' and simply write: let x be a point of X .

DEFINITION 2.8. We denote by R the set of permutations f of V such that, for each integer m from 1 to n , the elements $f(m, 1)$ and $f(m, 2)$ of V have different types.

DEFINITION 2.9. Let $f \in R$. We denote by \mathcal{U}_f the affine space

$$\mathrm{Spec}(\mathcal{O}_S[x_{i,j}, (i, j) \in f(V_1) \times f(V_2)])$$

over S . It is equipped with a structural morphism π_f to S . Denote by \mathcal{F}_f the closed subscheme

$$\mathrm{Spec}(\mathcal{O}_S[x_{i,j}, (i, j) \in (f(V_1)_1 \times f(V_2)_2) \sqcup (f(V_1)_2 \times f(V_2)_1)])$$

We think of a point x of \mathcal{U}_f as a matrix indexed by the set $f(V_1) \times f(V_2)$. For a subset I_1 of $f(V_1)$ and I_2 of $f(V_2)$, we denote by x_{I_1, I_2} the submatrix of x indexed by $I_1 \times I_2$. For example, the closed subscheme \mathcal{F}_f is defined by the vanishing of the two matrices $x_{f(V_1)_1 \times f(V_2)_1}$ and $x_{f(V_1)_2 \times f(V_2)_2}$.

PROPOSITION-DEFINITION 2.10. Let $f \in R$. There exists a unique morphism

$$\iota_f : \mathcal{U}_f \rightarrow \mathcal{G}$$

such that $\mathcal{T}_f := \iota_f^* \mathcal{T}$ is the submodule of $\pi_f^* \mathcal{V}$ spanned by

$$\pi_f^* v_j + \sum_{i \in f(V_1)} x_{i,j} \pi_f^* v_i$$

where j runs over the set $f(V_2)$. The morphism ι_f is an open immersion. We denote by \mathcal{G}_f the image of the open immersion ι_f . The open subschemes \mathcal{G}_f cover the grassmannian \mathcal{G} as f runs over the set R .

Proof. This is classical. The open subscheme \mathcal{G}_f of the grassmannian parametrizes the complementary submodules of $\mathcal{V}_{f(V_1)}$ in \mathcal{V} . \square

DEFINITION 2.11. Let $f \in R$. We denote by $P_{f,0}$ the subgroup scheme

$$\mathrm{Stab}_G(\mathcal{V}_{f(V_1)})$$

of G . It is a parabolic subgroup. We also denote by $L_{f,0}$ its Levi subgroup

$$L_{f,0} := \mathrm{Stab}_G(\mathcal{V}_{f(V_1)}, \mathcal{V}_{f(V_2)}) = \prod_{i,j \in \{1,2\}} \mathrm{GL}(\mathcal{V}_{f(V_i)_j})$$

In the next proposition we describe the local structure of the action of the group scheme G on \mathcal{G} . This is analogous to Proposition 1.3.

PROPOSITION 2.12. Let $f \in R$. The open subscheme \mathcal{G}_f of \mathcal{G} is left stable under the action of $P_{f,0}$. For the corresponding action of $P_{f,0}$ on \mathcal{U}_f through

the isomorphism ι_f , the closed subscheme \mathcal{F}_f is left stable under the action of $L_{f,0}$ and we have the following formulas

$$\forall g \in L_{f,0}, \quad \forall x \in \mathcal{F}_f,$$

$$x' = g \cdot x \text{ where } \begin{cases} x'_{f(V_1)_1, f(V_2)_2} = g_{f(V_1)_1, f(V_2)_2} g_{f(V_2)_2}^{-1} \\ x'_{f(V_1)_2, f(V_2)_1} = g_{f(V_1)_2, f(V_2)_1} g_{f(V_2)_1}^{-1} \end{cases}$$

Finally, the natural morphism

$$m_{f,0} : R_u(P_{f,0}) \times \mathcal{F}_f \rightarrow \mathcal{U}_f, \quad (r, x) \mapsto r \cdot x$$

is an isomorphism.

Proof. The open subscheme \mathcal{G}_f of the grassmannian parametrizes the complementary submodules of $\mathcal{V}_{f(V_1)}$ in \mathcal{V} . It follows that it is stable under the action of the stabilizer P of $\mathcal{V}_{f(V_1)}$ in $\text{GL}(\mathcal{V})$ and therefore under the action of its subgroup $P_{f,0}$.

Let x be a point of \mathcal{U}_f and g a point of P . By definition, the point $\iota_f(x)$ is the graph of x . Therefore, the point $g \cdot \iota_f(x)$ is the module consisting of elements of type

$$g(v + xv) = g_{f(V_2)}v + (g_{f(V_1), f(V_2)} + g_{f(V_1)}x)v, \quad v \in \mathcal{V}_{f(V_2)}.$$

It is thus equal to the point

$$\iota_f((g_{f(V_1), f(V_2)} + g_{f(V_1)}x)g_{f(V_2)}^{-1}).$$

In other words, the action of P on \mathcal{U}_f is given by

$$P \times \mathcal{U}_f \rightarrow \mathcal{U}_f, \quad (g, x) \mapsto (g_{f(V_1), f(V_2)} + g_{f(V_1)}x)g_{f(V_2)}^{-1}.$$

By specializing this action to the subgroup $P_{f,0}$ of P , we immediately see that \mathcal{F}_f is left stable under the action of $L_{f,0}$ we obtain the formulas in the statement of the proposition.

Moreover, still using the description of the action of P on \mathcal{U}_f found above, we see that if g is a point of $R_u(P_{f,0})$ and x a point of \mathcal{F}_f , then the point $x' = g \cdot x$ of \mathcal{U}_f is given by :

$$\begin{cases} x'_{f(V_1)_1, f(V_2)_1} = g_{f(V_1)_1, f(V_2)_1} \\ x'_{f(V_1)_1, f(V_2)_2} = x_{f(V_1)_1, f(V_2)_2} \\ x'_{f(V_1)_2, f(V_2)_1} = x_{f(V_1)_2, f(V_2)_1} \\ x'_{f(V_1)_2, f(V_2)_2} = g_{f(V_1)_2, f(V_2)_2} \end{cases}$$

This proves that the natural $P_{f,0}$ -equivariant morphism :

$$m_{f,0} : R_u(P_{f,0}) \times \mathcal{F}_f \rightarrow \mathcal{U}_f$$

is indeed an isomorphism. □

DEFINITION 2.13. Let $f \in R$ and $d \in \llbracket 0, n \rrbracket$. We denote by $\mathcal{I}_{f,0,d}$ the ideal sheaf on \mathcal{F}_f spanned by the minors of size d of the matrix

$$\begin{pmatrix} 0 & x_{f(V_1)_1, f(V_2)_2} \\ x_{f(V_1)_2, f(V_2)_1} & 0 \end{pmatrix}.$$

We denote by $\mathcal{Z}_{f,0,d}$ the closed subscheme of \mathcal{F}_f defined by the ideal sheaf $\mathcal{I}_{f,0,d}$.

PROPOSITION 2.14. Let $f \in R$ and $d \in \llbracket 0, n-1 \rrbracket$. Through the isomorphism

$$m_{f,0} : R_u(P_{f,0}) \times \mathcal{F}_f \rightarrow \mathcal{U}_f$$

of Proposition 2.12 the closed subscheme $\iota_f^{-1}(\mathcal{Z}_{0,d})$ is equal to $R_u(P_{f,0}) \times \mathcal{Z}_{f,0,d}$.

Proof. Due to the formula in the proof of Proposition 2.12, it suffices to show that the defining ideal of $\iota_f^{-1}(\mathcal{Z}_{0,d})$ on \mathcal{U}_f is spanned by the minors of size d of the matrix

$$\begin{pmatrix} 0 & x_{f(V_1)_1, f(V_2)_2} \\ x_{f(V_1)_2, f(V_2)_1} & 0 \end{pmatrix}.$$

To prove this, we express the matrix of the homomorphism

$$\iota_f^* p : \mathcal{T}_f^{\oplus 2} \rightarrow \pi_f^* \mathcal{V}$$

in appropriate basis. We choose the basis of \mathcal{T}_f described in Proposition-Definition 2.10. This basis is indexed by the set $f(V_2)$, which is the disjoint union of $f(V_2)_1$ and $f(V_2)_2$. We also choose the basis

$$\pi_f^*(v_i), \quad i \in f(V_1)_1, \quad \pi_f^*(v_j) + \sum_{i \in f(V_1)_1} x_{i,j} \pi_f^*(v_i), \quad j \in f(V_2)_1$$

for $\pi_f^* \mathcal{V}_1$ and

$$\pi_f^*(v_i), \quad i \in f(V_1)_2, \quad \pi_f^*(v_j) + \sum_{i \in f(V_1)_2} x_{i,j} \pi_f^*(v_i), \quad j \in f(V_2)_2$$

for $\pi_f^* \mathcal{V}_2$. The matrix of $\iota_f^* p$ in these basis can be expressed in blocks as follows:

$$\begin{pmatrix} 0 & x_{f(V_1)_1, f(V_2)_2} & 0 & 0 \\ Id & 0 & 0 & 0 \\ 0 & 0 & x_{f(V_1)_2, f(V_2)_1} & 0 \\ 0 & 0 & 0 & Id \end{pmatrix}.$$

By definition, the defining ideal of $\iota_f^{-1}(\mathcal{Z}_{0,d})$ is generated by the minors of size $n+d$ of this matrix. By reordering the vector in the basis, we get the block diagonal square matrix with blocks I_n and

$$\begin{pmatrix} 0 & x_{f(V_1)_1, f(V_2)_2} \\ x_{f(V_1)_2, f(V_2)_1} & 0 \end{pmatrix}.$$

We see therefore that the defining ideal of $\iota_f^{-1}(\mathcal{Z}_{0,d})$ is generated by the minors of size d of the last matrix, as we wanted. \square

DEFINITION 2.15. Let $f \in R$, $m \in \llbracket 0, n \rrbracket$ and $d \in \llbracket m, n \rrbracket$.

- We define a parabolic subgroup scheme $P_{f,m}$ of G by induction m . For m equals 0, we have already defined $P_{f,0}$. Then, assuming that $P_{f,m-1}$ has been defined, we set

$$P_{f,m} = \begin{cases} \text{Stab}_{P_{f,m-1}}(\mathcal{V}_{f(V_1^{>m}) \cap V_1}, \mathcal{V}_{\{f(m,2)\}}) \\ \text{if } f(m,1) \in V_1 \text{ and } f(m,2) \in V_2 \\ \text{Stab}_{P_{f,m-1}}(\mathcal{V}_{f(V_1^{>m}) \cap V_2}, \mathcal{V}_{\{f(m,2)\}}) \\ \text{if } f(m,1) \in V_2 \text{ and } f(m,2) \in V_1. \end{cases}$$

- We denote by $L_{f,m}$ the following Levi subgroup of $P_{f,m}$:

$$\prod_{i=1}^m (\text{GL}(\mathcal{V}_{f(i,1)}) \times \text{GL}(\mathcal{V}_{f(i,2)})) \times \prod_{i,j \in \{1,2\}} \text{GL}(\mathcal{V}_{f(V_i^{>m}) \cap V_j})$$

- We denote by $\mathcal{F}_{f,m}$ the affine space over S on the indeterminates $x_{i,j}$ where (i,j) runs over the union of the sets

$$\{(f(1,1), f(1,2)), \dots, (f(m,1), f(m,2))\}$$

and

$$((f(V_1^{>m})_1) \times (f(V_2^{>m})_2)) \cup ((f(V_1^{>m})_2) \times (f(V_2^{>m})_1)).$$

- We let the group scheme $L_{f,m}$ act on $\mathcal{F}_{f,m}$ by the following formulas

$$\begin{cases} x'_{f(1,1),f(1,2)} = g_{f(1,1)} g_{f(1,2)}^{-1} x_{f(1,1),f(1,2)} \\ x'_{f(i,1),f(i,2)} = g_{f(i,1)} g_{f(i-1,2)} g_{f(i-1,1)}^{-1} g_{f(i,2)}^{-1} x_{f(i,1),f(i,2)} \text{ for } i \in \llbracket 2, m \rrbracket \\ x'_{f(V_1^{>m})_1, f(V_2^{>m})_2} = g_{f(m,1)}^{-1} g_{f(m,2)} g_{f(V_1^{>m})_1} x_{f(V_1^{>m})_1, f(V_2^{>m})_2} g_{f(V_2^{>m})_2}^{-1} \\ x'_{f(V_1^{>m})_2, f(V_2^{>m})_1} = g_{f(m,1)}^{-1} g_{f(m,2)} g_{f(V_1^{>m})_2} x_{f(V_1^{>m})_2, f(V_2^{>m})_1} g_{f(V_2^{>m})_1}^{-1} \end{cases}$$

where g is a point of $L_{f,m}$, x a point of $\mathcal{F}_{f,m}$ and $x' := g \cdot x$.

- We denote by $\mathcal{U}_{f,m}$ the product

$$R_u(P_{f,m}) \times \mathcal{F}_{f,m}$$

acted upon by the group scheme $P_{f,m} = R_u(P_{f,m}) \rtimes L_{f,m}$ via the formula

$$\forall (r, l) \in P_{f,m}, \quad \forall (r', x) \in \mathcal{U}_{f,m}, \quad (r, l) \cdot (r', x) = (rlr'l^{-1}, l \cdot x).$$

- We denote by $\mathcal{I}_{f,m,d}$ the ideal sheaf on $\mathcal{F}_{f,m}$ spanned by the minors of size $d - m$ of the matrix

$$\begin{pmatrix} 0 & x_{f(V_1^{>m})_1, f(V_2^{>m})_2} \\ x_{f(V_1^{>m})_2, f(V_2^{>m})_1} & 0 \end{pmatrix}.$$

- We denote by $\mathcal{Z}'_{f,m,d}$ the closed subscheme of $\mathcal{F}_{f,m}$ defined by the ideal sheaf $\mathcal{I}_{f,m,d}$ and by $\mathcal{Z}_{f,m,d}$ the closed subscheme $R_u(P_{f,m,d}) \times \mathcal{Z}'_{f,m,d}$ of $\mathcal{U}_{f,m}$.
- We denote by $\mathcal{B}_{f,m}$ the blow-up of $\mathcal{U}_{f,m}$ along the closed subscheme $\mathcal{Z}_{f,m,m+1}$.

Let $f \in R$, $m \in \llbracket 1, n \rrbracket$ and $d \in \llbracket m, n \rrbracket$. The blow-up $\mathcal{B}_{f,m-1}$ is the closed subscheme of

$$\mathcal{U}_{f,m-1} \times \text{Proj}(\mathcal{O}_S[X_{i,j}, (i,j) \in (f(V_1^{\geq m})_1 \times f(V_2^{\geq m})_2) \sqcup (f(V_1^{\geq m})_2 \times f(V_2^{\geq m})_1)])$$

defined by the equations

$$\forall (i,j), (i',j') \in (f(V_1^{\geq m})_1 \times f(V_2^{\geq m})_2) \sqcup (f(V_1^{\geq m})_2 \times f(V_2^{\geq m})_1)$$

$$x_{i,j}X_{i',j'} - X_{i,j}x_{i',j'} = 0.$$

PROPOSITION 2.16. *With these notations, the open subscheme $\{X_{f(m,1),f(m,2)} \neq 0\}$ of $\mathcal{B}_{f,m-1}$ is left stable under the action of $P_{f,m}$ and is isomorphic, as a $P_{f,m}$ -scheme, to $\mathcal{U}_{f,m}$. Moreover, via this isomorphism, the strict transform of $\mathcal{Z}_{f,m-1,d}$ in $\mathcal{U}_{f,m}$ is $\mathcal{Z}_{f,m,d}$.*

Proof. We prove analogous statement for the the blow-up $\mathcal{B}'_{f,m-1}$ of $\mathcal{F}_{f,m-1}$ along the closed subscheme $\mathcal{Z}'_{f,m-1,m}$ from which the proposition is easily derived.

The scheme $\mathcal{B}'_{f,m-1}$ is the closed subscheme of

$$\mathcal{F}_{f,m-1} \times \text{Proj}(\mathcal{O}_S[X_{i,j}, (i,j) \in (f(V_1^{\geq m})_1 \times f(V_2^{\geq m})_2) \sqcup (f(V_1^{\geq m})_2 \times f(V_2^{\geq m})_1)])$$

defined by the equations

$$\forall (i,j), (i',j') \in (f(V_1^{\geq m})_1 \times f(V_2^{\geq m})_2) \sqcup (f(V_1^{\geq m})_2 \times f(V_2^{\geq m})_1)$$

$$x_{i,j}X_{i',j'} - X_{i,j}x_{i',j'} = 0.$$

Observe that the open subscheme $\mathcal{U}'_{f,m}$ of $\mathcal{B}'_{f,m-1}$ defined by $X_{f(m,1),f(m,2)} \neq 0$ is isomorphic, as a scheme over $\mathcal{F}_{f,m-1}$ to

$$b: \mathcal{F}_{f,m-1} \rightarrow \mathcal{F}_{f,m-1}, \quad x_{i,j} \mapsto \begin{cases} x_{f(m,1),f(m,2)}x_{i,j} & \text{if } (i,j) \in f(V_1^{\geq m}) \times f(V_2^{\geq m}) \\ & \text{and } i \text{ and } j \text{ have different types} \\ x_{i,j} & \text{otherwise.} \end{cases}$$

In the following we shall make use of this remark and use the coordinates $x_{i,j}$ to describe the points of $\mathcal{U}'_{f,m}$.

Observe that the center of the blow-up $\mathcal{B}'_{f,m-1} \rightarrow \mathcal{F}_{f,m-1}$ is stable under the action of $L_{f,m-1}$ and therefore the group scheme $L_{f,m-1}$ acts on $\mathcal{B}'_{f,m-1}$. This action can be described as follows.

$$\forall g \in L_{f,m-1}, \quad \forall (x, X) \in \mathcal{B}'_{f,m}, \quad (x', X') = g \cdot (x, X) \text{ where } x' = g \cdot x \text{ and}$$

$$\begin{cases} X'_{f(V_1^{\geq m})_1, f(V_2^{\geq m})_2} = g_{f(m-1,1)}^{-1} g_{f(m-1,2)} g_{f(V_1^{\geq m})_1} X_{f(V_1^{\geq m})_1, f(V_2^{\geq m})_2} g_{f(V_2^{\geq m})_2}^{-1} \\ X'_{f(V_1^{\geq m})_2, f(V_2^{\geq m})_1} = g_{f(m-1,1)}^{-1} g_{f(m-1,2)} g_{f(V_1^{\geq m})_2} X_{f(V_1^{\geq m})_2, f(V_2^{\geq m})_1} g_{f(V_2^{\geq m})_1}^{-1} \end{cases}$$

where we denote by X the matrix formed by the $X_{i,j}$. Observe now that the open subscheme $\mathcal{U}'_{f,m}$ is the locus of points (x, X) of $\mathcal{B}'_{f,m-1}$ such that $X_{f(m,1), f(m,2)}$ does not vanish. Therefore, it is left stable under the action of the parabolic subgroup of $L_{f,m-1}$

$$P = \begin{cases} \text{Stab}_{L_{f,m-1}}(\mathcal{V}_{f(V_1^{\geq m})}, \mathcal{V}_{f(m,2)}) \\ \quad \text{if } f(m,1) \text{ belongs to } V_1 \text{ and } f(m,2) \text{ to } V_2 \\ \text{Stab}_{L_{f,m-1}}(\mathcal{V}_{f(V_2^{\geq m})}, \mathcal{V}_{f(m,2)}) \\ \quad \text{if } f(m,1) \text{ belongs to } V_2 \text{ and } f(m,2) \text{ to } V_1 \end{cases}$$

The group scheme $L_{f,m}$ is a Levi subgroup scheme of P . Let g be a point of $L_{f,m}$ and x a point of $\mathcal{F}_{f,m}$. The point x corresponds to the couple $(b(x), X)$ in $\mathcal{B}'_{f,m-1}$, where

$$X_{i,j} = \begin{cases} x_{i,j} & \text{if } (i, j) \neq (f(m,1), f(m,2)) \\ 1 & \text{if } (i, j) = (f(m,1), f(m,2)). \end{cases}$$

Let $(x', X') = g \cdot (x, X)$. By a quick computation we get

$$\begin{cases} x'_{f(m,1), f(m,2)} = g_{f(m-1,1)}^{-1} g_{f(m-1,2)} g_{f(m,1)} g_{f(m,2)}^{-1} x_{f(m,1), f(m,2)} \\ X'_{f(m,1), f(m,2)} = g_{f(m-1,1)}^{-1} g_{f(m-1,2)} g_{f(m,1)} g_{f(m,2)}^{-1} \\ X'_{f(V_1^{\geq m})_1, f(V_2^{\geq m})_2} = g_{f(m-1,1)}^{-1} g_{f(m-1,2)} g_{f(V_1^{\geq m})_1} X_{f(V_1^{\geq m})_1, f(V_2^{\geq m})_2} g_{f(V_2^{\geq m})_2}^{-1} \\ X'_{f(V_1^{\geq m})_2, f(V_2^{\geq m})_1} = g_{f(m-1,1)}^{-1} g_{f(m-1,2)} g_{f(V_1^{\geq m})_2} X_{f(V_1^{\geq m})_2, f(V_2^{\geq m})_1} g_{f(V_2^{\geq m})_1}^{-1} \\ X'_{i,j} = 0 \text{ otherwise} \end{cases}$$

Therefore we see that the closed subscheme $\mathcal{F}_{f,m}$ of $\mathcal{U}'_{f,m}$ is left stable under the action of the Levi subgroup $L_{f,m}$ and, moreover, the action of $L_{f,m}$ on $\mathcal{F}_{f,m}$ is given by the formulas in Definition 2.15. In a similar way, we prove that the natural morphism

$$R_u(P) \times \mathcal{F}_{f,m} \rightarrow \mathcal{U}'_{f,m}, \quad (g, x) \mapsto g \cdot x = x'$$

is an isomorphism, given by the following formulas

$$\begin{cases} x'_{f(l,1),f(l,2)} = x_{f(l,1),f(l,2)} \text{ for all } l \in \llbracket 1, m \rrbracket \\ x'_{f(V_1^{>m})_1, f(m,2)} = g_{f(V_1^{>m})_1, f(m,1)} \\ x'_{f(m,1), f(V_2^{>m})_2} = -g_{f(m,2), f(V_2^{>m})_2} \\ x'_{f(V_1^{>m})_1, f(V_2^{>m})_2} = x_{f(V_1^{>m})_1, f(V_2^{>m})_2} - g_{f(V_1^{>m})_1, f(m,1)} g_{f(m,2), f(V_2^{>m})_2} \\ x'_{f(V_1^{>m})_2, f(V_2^{>m})_1} = x_{f(V_1^{>m})_2, f(V_2^{>m})_1} \end{cases}$$

Now we compute the strict transform of $\mathcal{Z}'_{f,m,d}$ in $\mathcal{U}'_{f,m}$. By definition, the strict transform of $\mathcal{Z}'_{f,m,d}$ is the schematic closure of

$$\mathcal{Z} := b^{-1}(\mathcal{Z}'_{f,m-1,d} \cap \{x_{f(m,1),f(m,2)} \neq 0\})$$

in $\mathcal{U}'_{f,m}$. Let x be a point of $\mathcal{U}'_{f,m}$. We denote (r, y) its components through the isomorphism

$$R_u(P) \times \mathcal{F}_{f,m} \rightarrow \mathcal{U}'_{f,m}.$$

By definition, the point x belongs to \mathcal{Z} if and only $x_{f(m,1),f(m,2)}$ is invertible and all the square $d - m + 1$ minors extracted from the following matrix

$$\begin{pmatrix} 0 & b(x_{f(V_1^{\geq m})_1, f(V_2^{\geq m})_2}) \\ b(x_{f(V_1^{\geq m})_2, f(V_2^{\geq m})_1}) & 0 \end{pmatrix}$$

are zero. By definition of the morphism b , each coefficient in this matrix is a multiple of $x_{f(m,1),f(m,2)}$ and the coefficient in place $(f(m,1), f(m,2))$ is exactly $x_{f(m,1),f(m,2)}$. As this indeterminate is invertible on the open subscheme $\{x_{f(m,1),f(m,2)} \neq 0\}$ we see that the point x belongs to \mathcal{Z} if and only $x_{f(m,1),f(m,2)}$ is invertible and all the minors of size $d - m + 1$ extracted from the matrix

$$\begin{pmatrix} 0 & x'_{f(V_1^{\geq m})_1, f(V_2^{\geq m})_2} \\ x'_{f(V_1^{\geq m})_2, f(V_2^{\geq m})_1} & 0 \end{pmatrix}$$

are zero, where $x'_{f(m,1),f(m,2)} = 1$ and $x'_{i,j} = x_{i,j}$ otherwise. By operating standard row and column operations on this matrix we see now that x belongs to \mathcal{Z} if and only if $x_{f(m,1),f(m,2)}$ is invertible and all the minors of size $d - m$ extracted from the matrix

$$\begin{pmatrix} 0 & x_{f(V_1^{>m})_1, f(V_2^{>m})_2} \\ x_{f(V_1^{>m})_2, f(V_2^{>m})_1} & 0 \end{pmatrix}$$

are zero. This proves that \mathcal{Z} is the intersection of $R_u(P) \times \mathcal{Z}'_{f,m,d}$ with the open subscheme $\{x_{f(m,1),f(m,2)} \neq 0\}$.

In order to complete the proof, we now show that the schematic closure of \mathcal{Z} in $\mathcal{U}'_{f,m}$ is equal to $R_u(P) \times \mathcal{Z}'_{f,m,d}$. We can check this Zariski locally on the

base S and therefore assume that S is affine. First of all, it is obvious that the closure of \mathcal{Z} is contained in $R_u(P) \times \mathcal{Z}'_{f,m,d}$. Conversely, let φ be a global function on $\mathcal{U}'_{f,m}$ which vanishes on \mathcal{Z} . This means that there exists an integer q such that $x_{f(m,1),f(m,2)}^q \varphi$ belongs to the ideal spanned by the minors of size $d - m$ of the following matrix:

$$\begin{pmatrix} 0 & x_{f(V_1^{\geq m})_1, f(V_2^{\geq m})_2} \\ x_{f(V_1^{\geq m})_2, f(V_2^{\geq m})_1} & 0 \end{pmatrix}.$$

As the indeterminate $x_{f(m,1),f(m,2)}$ does not appear in this matrix, we can conclude that φ itself belongs to this ideal, completing the proof of the proposition. \square

PROPOSITION 2.17. *Let $f \in R$. There exists a unique collection of open immersions $\iota_{f,m}$, for m from 1 to n such that each of the squares below are commutative :*

$$\begin{array}{ccccccc} \mathcal{U}_{f,0} & \xleftarrow{b_{f,1}} & \mathcal{U}_{f,1} & \xleftarrow{\dots} & \mathcal{U}_{f,n-1} & \xleftarrow{b_{f,n}} & \mathcal{U}_{f,n} \\ \downarrow \iota_{f,0} & & \downarrow \iota_{f,1} & & \downarrow \iota_{f,n-1} & & \downarrow \iota_{f,n} \\ \mathcal{G}_0 & \xleftarrow{b_1} & \mathcal{G}_1 & \xleftarrow{\dots} & \mathcal{G}_{n-1} & \xleftarrow{b_n} & \mathcal{G}_n \end{array}$$

The open immersion $\iota_{f,m}$ is equivariant for the action of $P_{f,m}$. We denote by $\mathcal{G}_{f,m}$ the image of the open immersion $\iota_{f,m}$. The open subschemes $\mathcal{G}_{f,m}$ cover \mathcal{G}_m as f run over R .

Proof. The existence and $P_{f,m}$ -equivariance of the $\iota_{f,m}$ follows directly from Proposition 2.16. The uniqueness comes from the fact that for each index m , the intersection $\Omega \cap \mathcal{G}_{f,m}$ is dense in $\mathcal{G}_{f,m}$. The last assertion is proved as follows. By Proposition 2.10, the open subschemes $\mathcal{G}_{f,0}$ cover the scheme \mathcal{G}_0 . Moreover, it follows from Proposition 2.16 that the blow-up $\mathcal{B}_{f,m-1}$ is covered by the open subschemes $\mathcal{U}_{f',m}$ where f' runs over the elements of R having the same restriction as f to $\llbracket 1, k \rrbracket \times \{1, 2\}$ and satisfying $f(V_1) = f'(V_1)$ and $f(V_2) = f'(V_2)$. \square

2.3 AN ALTERNATIVE CONSTRUCTION

In this section we provide an alternative construction for the schemes \mathcal{G}_m . Recall from Section 2.1 that the section $\wedge^{n+d} p$ of the locally free module \mathcal{H}_d does not vanish on Ω . Therefore it defines an invertible submodule of \mathcal{H}_d on Ω that is locally a direct summand. In other words, it defines a morphism from Ω to the projective bundle $\mathbb{P}(\mathcal{H}_d)$ over Ω .

DEFINITION 2.18. Let $d \in \llbracket 0, n \rrbracket$. We denote by φ_d the morphism

$$\varphi_d : \Omega \rightarrow \mathbb{P}(\mathcal{H}_d).$$

defined by the global section $\wedge^{n+d} p$ of \mathcal{H}_d .

PROPOSITION 2.19. Let $d \in \llbracket 0, n \rrbracket$ and $m \in \llbracket d, n \rrbracket$. The morphism φ_d extends to \mathcal{G}_m in a unique way.

Proof. We first observe that Ω is dense in each of the schemes \mathcal{G}_m . If the morphism φ_d extends to \mathcal{G}_m it is therefore in a unique way. Moreover, it suffices to show that φ_d extends to \mathcal{G}_d , because then the composite

$$\mathcal{G}_m \xrightarrow{b_m} \mathcal{G}_{m-1} \rightarrow \dots \rightarrow \mathcal{G}_d \xrightarrow{\varphi_d} \mathbb{P}(\mathcal{H}_d)$$

is an extension of φ_d to \mathcal{G}_m .

By the same density argument as above it suffices to show that the morphism φ_d extends from $\Omega \cap \mathcal{G}_{f,d}$ to $\mathcal{G}_{f,d}$ for each element f of the set R . We identify the scheme $\mathcal{G}_{f,d}$ with $\mathcal{U}_{f,d}$ via the isomorphism $\iota_{f,d}$. Observe that the scheme $\mathbb{P}(\mathcal{H}_d)$ is equipped with an action of the group scheme G and the morphism φ_d is equivariant with respect to this action. By using this remark, we see that it suffices to extend the morphism φ_d from $\Omega \cap \mathcal{F}_{f,d}$ to $\mathcal{F}_{f,d}$.

We denote by c the composite $b_{f,1} \circ \dots \circ b_{f,d}$. We use the trivialization of \mathcal{H}_d on $\mathcal{G}_{f,0}$ as in the proof of Proposition 2.14. For this trivialization, the coordinates are indexed by the product $(V_2 \sqcup V_2) \times V$. The morphism φ_d is given over $\Omega \cap \mathcal{F}_{f,d}$ in this coordinates by the $n + d$ minors of the matrix

$$\begin{pmatrix} 0 & c(x)_{f(V_1)_1, f(V_2)_2} & 0 & 0 \\ Id & 0 & 0 & 0 \\ 0 & 0 & c(x)_{f(V_1)_2, f(V_2)_1} & 0 \\ 0 & 0 & 0 & Id \end{pmatrix}.$$

It follows from the definition of c that all these minors are multiples of

$$x_{f(1,1), f(1,2)}^d x_{f(1,2), f(2,2)}^{d-1} \cdots x_{f(d,1), f(d,2)}$$

and that one of them, namely the one indexed by the product of

$$f(V_1)_2 \sqcup f(V_2^{\leq d})_2 \sqcup (f(V_1^{\leq d})_2) \sqcup f(V_2)_2$$

and

$$f(V_1^{\leq d})_1 \sqcup f(V_2)_1 \sqcup f(V_1^{\leq d})_2 \sqcup f(V_2)_2$$

is exactly equal to this product or its opposite. By dividing each coordinate by this product we therefore extend the morphism φ_d to $\mathcal{F}_{f,d}$. \square

PROPOSITION 2.20. Let $m \in \llbracket 0, n \rrbracket$. The morphism

$$\psi_m := \varphi_0 \times \varphi_1 \times \dots \times \varphi_m : \mathcal{G}_m \rightarrow \mathbb{P}(\mathcal{H}_0) \times_{\mathcal{G}} \mathbb{P}(\mathcal{H}_1) \times_{\mathcal{G}} \dots \times_{\mathcal{G}} \mathbb{P}(\mathcal{H}_m)$$

is a closed immersion.

Proof. We prove this by induction on m . For $m = 0$ the morphism φ_0 is defined by the nowhere vanishing section $\wedge^n p$ of \mathcal{H}_0 and is therefore a closed immersion. We suppose now that ψ_{m-1} is a closed immersion and prove that ψ_m is also a closed immersion. As this morphism is proper, it suffices to check that it is a monomorphism in order to prove that it is a closed immersion. Let q_1 and q_2 be two points of \mathcal{G}_m which are mapped to the same point by ψ_m . We want to show that they are equal.

First, we suppose that q_1 and q_2 are points of $\mathcal{G}_{f,m}$, where f is an element of R . We identify $\mathcal{G}_{f,m}$ and $\mathcal{U}_{f,m}$ via the isomorphism $\iota_{f,m}$. We use the notations introduced in the proof of Proposition 2.16. The scheme $\mathcal{U}_{f,m}$ is isomorphic to the product $R_u(P_{f,m-1}) \times \mathcal{U}'_{f,m}$. Using the induction hypothesis, we can assume that q_1 and q_2 are actually points of $\mathcal{U}'_{f,m}$. Viewed as points of $\mathcal{U}'_{f,m}$ (which is isomorphic to $\mathcal{F}_{f,m-1}$ via the morphism b) we denote the coordinates of q_1 by $(x_{i,j,1})$ and the coordinates of q_2 by $(x_{i,j,2})$. Still by the induction hypothesis, we have $x_{i,j,1} = x_{i,j,2}$ for (i, j) in the following set

$$\{(f(1, 1), f(1, 2)), \dots, (f(m, 1), f(m, 2))\}$$

as it follows from the definition of the morphism b . Observe now that the morphism φ_m is defined over $\mathcal{U}'_{f,m}$ by exactly the same process as explained in the proof of Proposition 2.19. By this we mean that the coordinates of φ_d are obtained by computing the $n + d$ minors of the matrix

$$\begin{pmatrix} 0 & c(x)_{f(V_1)_1, f(V_2)_2} & 0 & 0 \\ Id & 0 & 0 & 0 \\ 0 & 0 & c(x)_{f(V_1)_2, f(V_2)_1} & 0 \\ 0 & 0 & 0 & Id \end{pmatrix}$$

and dividing by the product

$$x_{f(1,1), f(1,2)}^m x_{f(1,2), f(2,2)}^{m-1} \cdots x_{f(m,1), f(m,2)}.$$

Indeed this process makes sense over $\mathcal{U}'_{f,m}$ and extend φ_d over $\Omega \cap \mathcal{U}'_{f,m}$. Moreover, such an extension is unique. Let (i, j) be an element of

$$(f(V_1^{\geq m})_1 \times f(V_2^{\geq m})_2) \sqcup (f(V_1^{\geq m})_2 \times f(V_2^{\geq m})_1)$$

different from $(f(m, 1), f(m, 2))$. We suppose that i is of type 1 and j of type 2, the other case being entirely similar. The coordinate of φ_m corresponding to the minor indexed by the product of

$$f(V_1)_2 \sqcup f(V_2^{<d})_2 \sqcup \{j\} \sqcup (f(V_1^{\leq d})_2) \sqcup f(V_2)_2$$

and

$$f(V_1^{\leq d})_1 \sqcup f(V_2)_1 \sqcup f(V_1^{<d})_2 \sqcup \{i\} \sqcup f(V_2)_2.$$

is $x_{i,j}$ or its opposite. We can therefore conclude that $x_{i,j,1} = x_{i,j,2}$. Finally we have proved that $q_1 = q_2$.

We go back to the general case. Let f be an element of R . We prove now that the open subschemes $q_1^{-1}(\mathcal{G}_{f,m})$ and $q_2^{-1}(\mathcal{G}_{f,m})$ are equal. This is sufficient to complete the proof of the proposition. By the induction hypothesis, we already know that the open subschemes $q_1^{-1}(\mathcal{G}_{f,m-1})$ and $q_2^{-1}(\mathcal{G}_{f,m-1})$ are equal. Observe now that the computations above actually prove the following: through φ_m , the non zero locus on $\mathcal{B}_{f,m}$ of the coordinate of $\mathbb{P}(\mathcal{H}_m)$ corresponding to the minor indexed by the product of

$$f(V_1)_2 \sqcup f(V_2^{\leq m})_2 \sqcup (f(V_1^{\leq m})_2) \sqcup f(V_2)_2$$

and

$$f(V_1^{\leq m})_1 \sqcup f(V_2)_1 \sqcup f(V_1^{\leq m})_2 \sqcup f(V_2)_2$$

is $\mathcal{U}_{f,m}$. This implies the result. □

2.4 THE COLORED FAN OF \mathcal{G}_n

In this section we assume that the base scheme S is the spectrum of an algebraically closed field k of arbitrary characteristic. The scheme \mathcal{G}_n is an equivariant compactification of the homogeneous space $\text{Iso}(\mathcal{V}_2, \mathcal{V}_1)$ under the action of the group $\text{GL}(\mathcal{V}_1) \times \text{GL}(\mathcal{V}_2)$. Through the fixed trivializations of the free modules \mathcal{V}_1 and \mathcal{V}_2 we see that \mathcal{G}_n is an equivariant compactification of the general linear group $\text{GL}(n)$ under the action of $\text{GL}(n) \times \text{GL}(n)$. The aim of this section is to compute the colored fan of this compactification, as explained in Section 1.4.

But first, we say a word about the blow-up procedure explained in Section 2.1 in this setting. By definition, the set $\mathcal{Z}_d(k)$ is the set of n -dimensional subspaces of $\mathcal{V}(k)$ such that the sum of the ranks of $p_1(k)$ and $p_2(k)$ is strictly less than $n + d$ at every point. Another way to state this is that $\mathcal{Z}_d(k)$ is the set

$$\{F \in \mathcal{G}(k), \quad \dim(F \cap \mathcal{V}_1(k)) + \dim(F \cap \mathcal{V}_2(k)) > n - d\}.$$

For example, the set $\mathcal{Z}_1(k)$ is the set of n -dimensional subspaces of $\mathcal{V}(k)$ which are direct sum of a subspace of $\mathcal{V}_1(k)$ and a subspace of $\mathcal{V}_2(k)$. Using this description, it is not difficult to prove that $\mathcal{Z}_1(k)$ is the union of the closed orbits of $G(k)$ in $\mathcal{G}(k)$. We leave it as an exercise to the reader to prove that, for d from 1 to $n - 1$, an orbit ω of $G(k)$ in $\mathcal{G}(k)$ is contained in $\mathcal{Z}_d(k)$ if and only if its closure is the union of ω and some orbits contained in $\mathcal{Z}_{d-1}(k)$.

We use the notations introduced in Section 1.4. We choose for T the diagonal torus in $\text{GL}(n)$ and for B the Borel subgroup of upper triangular matrices. The torus T is naturally isomorphic to the torus \mathbb{G}_m^n . The vector space V is therefore naturally isomorphic to \mathbb{Q}^n . The Weyl chamber \mathcal{W} corresponding to the chosen Borel subgroup B of $\text{GL}(n)$ is given by

$$\mathcal{W} = \{(a_1, \dots, a_n) \in V, \quad a_1 \geq a_2 \geq \dots \geq a_n\}.$$

PROPOSITION-DEFINITION 2.21. We denote by Q the set of permutations g of $\llbracket 1, n \rrbracket$ such that

$$\exists m \in \llbracket 0, n \rrbracket, \quad g_{|g^{-1}(\llbracket 1, m \rrbracket)} \text{ is decreasing and } g_{|g^{-1}(\llbracket m, n \rrbracket)} \text{ is increasing.}$$

If such an integer m exists, it is unique. We call it the integer associated to g and denote it by m_g . Also, we denote by ε_g the function

$$\varepsilon_g : \llbracket 1, n \rrbracket \rightarrow \{+1, -1\}, \quad x \mapsto \begin{cases} -1 & \text{if } g(x) \in \llbracket 1, m_g \rrbracket \\ 1 & \text{if } g(x) \in \llbracket m_g + 1, n \rrbracket \end{cases}$$

Finally, we denote by C_g the following cone in V :

$$C_g := \{(a_1, \dots, a_n) \in V, \quad 0 \leq \varepsilon_g(1)a_{g(1)} \leq \dots \leq \varepsilon_g(n)a_{g(n)}\}$$

PROPOSITION 2.22. The compactification \mathcal{G}_n of $\mathrm{GL}(n)$ is log homogeneous and its colored fan consists of the cones C_g and their faces, where g runs over the set Q .

Proof. Let us first prove that the compactification \mathcal{G}_n is log homogeneous. Let f be an element of R . We claim that the complement of Ω in $\mathcal{F}_{f,n}$ is the union of the coordinate hyperplanes $x_{f(d,1),f(d,2)} = 0$, where d runs from 1 to n . Indeed, a point x of $\mathcal{F}_{f,n}$ belongs to Ω if and only if the point $x' = (b_{f,n} \circ \dots \circ b_{f,1})(x)$ belongs to Ω . Moreover, we have

$$\begin{cases} x'_{f(d,1),f(d,2)} = x_{f(1,1),f(1,2)} \cdots x_{f(d,1),f(d,2)} \text{ for all } d \in \llbracket 1, n \rrbracket \\ x'_{i,j} = 0 \text{ if } (i, j) \notin \{(f(1,1), f(1,2)), \dots, (f(n,1), f(n,2))\}. \end{cases}$$

By Proposition 2.14, the point x' belongs to Ω if and only if the product

$$x'_{f(1,1),f(1,2)} \cdots x'_{f(n,1),f(n,2)} = x_{f(1,1),f(1,2)}^n x_{f(2,1),f(2,2)}^{n-1} \cdots x_{f(n,1),f(n,2)}$$

does not vanish, that is, if and only if each of the $x_{f(d,1),f(d,2)}$ does not vanish. This proves the claim. In particular, the complement of Ω in \mathcal{G}_n is a strict normal crossing divisor. By Definition 2.15, the variety $\mathcal{F}_{f,n}$ is a smooth toric variety for a quotient of the torus $T \times T = L_{f,n}$. From this we see that it is log homogeneous. It is now straightforward to check that the $P_{f,n}$ -variety $\mathcal{U}_{f,n}$ is log homogeneous. It readily follows that the G -variety \mathcal{G}_n is log homogeneous.

We let T acts on \mathcal{G}_n on the left. By Section 1.4, the closure \overline{T} of T in \mathcal{G}_n is a toric variety under the action of T and we can use the fan of this toric variety to compute the colored fan of \mathcal{G}_n . We shall now identify some of the cones in the fan of \overline{T} . Let g be an element of Q . We fix an element f of R such that, for each integer d from 1 to n , $f(d,1) = (g(d), 1)$ if $\varepsilon_g(d) = 1$ and $f(d,2) = (g(d), 1)$ if $\varepsilon_g(d) = -1$. The variety $\mathcal{F}_{f,n}$ is an open affine toric

subvariety of \overline{T} . It corresponds to a cone in the fan of \overline{T} , namely the cone spanned by the one-parameter subgroups having a limit in $\mathcal{F}_{f,n}$ at 0. Let

$$\lambda : \mathbb{G}_m \rightarrow T, \quad t \rightarrow (t^{a_1}, \dots, t^{a_n})$$

be a one-parameter subgroup of T . By the formulas in Definition 2.15, we see that the one-parameter subgroup λ has a limit in $\mathcal{F}_{f,n}$ at 0 if and only if

$$0 \leq \varepsilon_g(1)a_{g(1)} \leq \dots \leq \varepsilon_g(n)a_{g(n)}$$

that is, if and only if λ belongs to C_g . This proves that, for each element g of Q , the cone C_g belongs to the fan of the toric variety \overline{T} .

To complete the proof, we show now that the cone $-\mathcal{W}$ is equal to the union of the cones C_g , where g runs over the set Q . Let g be an element of Q and let (a_1, \dots, a_n) be a point in C_g . Let also i be an integer between 1 and $n-1$. If $i < m_g$, then there are two integers $j > j'$ such that $g(j) = i$ and $g(j') = i+1$. These integers satisfy $\varepsilon(j) = -1$ and $\varepsilon(j') = -1$. By definition of the cone C_g , we have $\varepsilon(j')a_{g(j')} \leq \varepsilon(j)a_{g(j)}$, that is, $a_i \leq a_{i+1}$. The same kind of argument prove that $a_i \leq a_{i+1}$ for $i = m_g$ and for $i > m_g$. This proves that the cone C_g is contained in the cone $-\mathcal{W}$. We consider now an element (a_1, \dots, a_n) of $-\mathcal{W}$. By definition it satisfies $a_1 \leq a_2 \leq \dots \leq a_n$. Let m be an integer such that $a_m \leq 0$ and $a_{m+1} \geq 0$. The rational numbers $-a_1, \dots, -a_m$ and a_{m+1}, \dots, a_n are nonnegative. By ordering them in increasing order, we construct an element g of Q such that the point (a_1, \dots, a_n) belongs to C_g . □

2.5 FIXED POINTS

In this section, the scheme S is the spectrum of an algebraically closed field k of characteristic not 2. We apply the results obtained in Section 1.3 for some involutions on the log homogeneous compactification \mathcal{G}_n of $\mathrm{GL}(n)$. We denote by J_r the antidiagonal square matrix of size r with all coefficients equal to one on the antidiagonal.

Let b be a nondegenerate symmetric or antisymmetric bilinear form on k^n . Via the fixed trivializations of \mathcal{V}_1 and \mathcal{V}_2 we obtain nondegenerate symmetric or antisymmetric bilinear forms b_1 and b_2 on the k -vector spaces $\mathcal{V}_1(k)$ and $\mathcal{V}_2(k)$. We equip the direct sum $\mathcal{V}(k)$ of $\mathcal{V}_1(k)$ and $\mathcal{V}_2(k)$ with the nondegenerate symmetric or antisymmetric bilinear form $b_1 \oplus b_2$. We let σ be the involution of \mathcal{G} mapping a n -dimensional k -vector subspace F of $\mathcal{V}(k)$ to its orthogonal. It is an easy exercise to check that

$$\dim(F^\perp \cap \mathcal{V}_1(k)) = \dim(F \cap \mathcal{V}_2(k)) \text{ and } \dim(F^\perp \cap \mathcal{V}_2(k)) = \dim(F \cap \mathcal{V}_1(k)).$$

By the description of \mathcal{Z}_d given in Section 2.4, we see that the involution σ leaves each of the closed subvarieties \mathcal{Z}_d of \mathcal{G} invariant. Therefore it extends to an involution, still denoted σ , of each of the varieties \mathcal{G}_m . To prove that we are

in the setting of Section 1.3, it remains to observe that there is an involution σ of $\mathrm{GL}(n)$, namely the one associated to b , such that

$$\forall g \in \mathrm{GL}(n) \times \mathrm{GL}(n), \quad \forall x \in \mathcal{G}_m, \quad \sigma((g_1, g_2) \cdot x) = \sigma(g_1) \cdot \sigma(x) \cdot \sigma(g_2)^{-1}.$$

As in Section 1.3, we denote by G' the neutral component of G^σ and by \mathcal{G}'_n the connected component of \mathcal{G}_n^σ containing G' .

THE ODD ORTHOGONAL CASE. We suppose that $n = 2r + 1$ is odd. We let b be the scalar product with respect to the matrix J_{2r+1} . We have $G' := \mathrm{SO}(2r+1)$. We let T' be the intersection of T with G' and B' the intersection of B with G' . The maximal torus T' is naturally isomorphic to the split torus \mathbb{G}_m^r via the following morphism

$$\mathbb{G}_m^r \rightarrow T', \quad (t_1, \dots, t_r) \mapsto \mathrm{diag}(t_1, \dots, t_r, 1, t_r^{-1}, \dots, t_1^{-1}).$$

The space V' is therefore naturally isomorphic to \mathbb{Q}^r . It is contained in V via the following linear map

$$V' \rightarrow V, \quad (a'_1, \dots, a'_r, 0, -a'_r, \dots, -a'_1).$$

The Weyl chamber with respect to B' is given by

$$\mathcal{W}' = \{(a'_1, \dots, a'_r) \in V, \quad a'_1 \geq a'_2 \geq \dots \geq a'_r \geq 0\}.$$

PROPOSITION 2.23. *The compactification \mathcal{G}'_n of G' is the wonderful compactification.*

Proof. First of all, by Proposition 1.9, the compactification \mathcal{G}'_n is log homogeneous. As explained in Section 1.4, we use the closure of T' in \mathcal{G}'_n to compute the colored fan of \mathcal{G}'_n . Observe that T' is a subtorus of T and therefore the fan of the toric variety $\overline{T'}$ is the trace on V' of the fan of the toric variety \overline{T} . By Proposition 2.22, the cone

$$\{(a_1, \dots, a_{2r+1}) \in V, \quad 0 \leq a_{r+1} \leq -a_r \leq a_{r+2} \leq \dots \leq -a_1 \leq a_{2r+1}\}$$

belongs to the fan of \overline{T} . The trace of this cone on V' is $-\mathcal{W}'$, proving that the cone $-\mathcal{W}'$ belongs to the colored fan of \mathcal{G}'_n . But the only fan in V' with support $-\mathcal{W}'$ containing $-\mathcal{W}'$ is the fan formed by $-\mathcal{W}'$ and its faces. This completes the proof. □

THE EVEN ORTHOGONAL CASE. We suppose that $n = 2r$ is even. We let b be the scalar product with respect to the matrix J_{2r} . We have $G' := \mathrm{SO}(2r)$. We let T' be the intersection of T with G' and B' the intersection of B with G' . The maximal torus T' is naturally isomorphic to the split torus \mathbb{G}_m^r via the following morphism

$$\mathbb{G}_m^r \rightarrow T', \quad (t_1, \dots, t_r) \mapsto \mathrm{diag}(t_1, \dots, t_r, t_r^{-1}, \dots, t_1^{-1}).$$

The space V' is therefore naturally isomorphic to \mathbb{Q}^r . It is contained in V via the following linear map

$$V' \rightarrow V, \quad (a'_1, \dots, a'_r, -a'_r, \dots, -a'_1).$$

The Weyl chamber with respect to B' is given by

$$\mathcal{W}' = \{(a'_1, \dots, a'_r) \in V, \quad a'_1 \geq a'_2 \geq \dots \geq a'_{r-1} \geq |a'_r|\}.$$

PROPOSITION 2.24. *The compactification \mathcal{G}'_n of G' is log homogeneous and its fan consists of the cones*

$$C_+ := \{(a'_1, \dots, a'_r) \in V, \quad a'_1 \leq a'_2 \leq \dots \leq a'_{r-1} \leq a'_r \leq 0\}$$

and

$$C_- := \{(a'_1, \dots, a'_r) \in V, \quad a'_1 \leq a'_2 \leq \dots \leq a'_{r-1} \leq -a'_r \leq 0\}$$

and their faces.

Proof. The proof is similar to that of Proposition 2.23. With the arguments given in this proof it suffices to observe that the trace of the following cone in V :

$$\{(a_1, \dots, a_{2r}) \in V, \quad 0 \leq -a_r \leq a_{r+1} \leq \dots \leq -a_1 \leq a_{2r}\}$$

on V' is C_+ , the trace of

$$\{(a_1, \dots, a_{2r}) \in V, \quad 0 \leq a_r \leq -a_{r+1} \leq -a_{r-1} \leq a_{r+2} \leq \dots \leq -a_1 \leq a_{2r}\}$$

is C_- and that $-\mathcal{W}'$ is the union of C_+ and C_- . \square

Observe that the Weyl chamber \mathcal{W}' is not smooth with respect to the lattice of one-parameter subgroups of T' . Therefore the canonical compactification of G' is not smooth, and the compactification \mathcal{G}'_n is a minimal log homogeneous compactification, in the sense that it has a minimal number of closed orbits.

THE SYMPLECTIC CASE. We suppose that $n = 2r$ is even. We let b be the scalar product with respect to the block antidiagonal matrix

$$\begin{pmatrix} 0 & -J_r \\ J_r & 0 \end{pmatrix}.$$

We have $G' := \mathrm{Sp}(2r)$. We let T' be the intersection of T with G' and B' the intersection of B with G' . The maximal torus T' is naturally isomorphic to the split torus \mathbb{G}_m^r via the following morphism

$$\mathbb{G}_m^r \rightarrow T', \quad (t_1, \dots, t_r) \mapsto \mathrm{diag}(t_1, \dots, t_r, t_r^{-1}, \dots, t_1^{-1}).$$

The space V' is therefore naturally isomorphic to \mathbb{Q}^r . It is contained in V via the following linear map

$$V' \rightarrow V, \quad (a'_1, \dots, a'_r, -a'_r, \dots, -a'_1).$$

The Weyl chamber with respect to B' is given by

$$\mathcal{W}' = \{(a'_1, \dots, a'_r) \in V, \quad a'_1 \geq a'_2 \geq \dots \geq a'_r \geq 0\}.$$

PROPOSITION 2.25. *The compactification \mathcal{G}'_n of G' is the wonderful compactification.*

Proof. The proof is similar to that of Proposition 2.23. With the arguments given in this proof it suffices to observe that the trace of the following cone in V :

$$\{(a_1, \dots, a_{2r}) \in V, \quad 0 \leq -a_r \leq a_{r+1} \cdots \leq -a_1 \leq a_{2r}\}$$

on V' is $-\mathcal{W}'$. □

REFERENCES

- [1] T. Abe. Compactification of the symplectic group via generalized symplectic isomorphisms. In *Higher dimensional algebraic varieties and vector bundles*, RIMS Kôkyûroku Bessatsu, B9, pages 1–50. Res. Inst. Math. Sci. (RIMS), Kyoto, 2008.
- [2] F. Bien and M. Brion. Automorphisms and local rigidity of regular varieties. *Compositio Math.*, 104(1):1–26, 1996.
- [3] E. Bifet, C. De Concini, and C. Procesi. Cohomology of regular embeddings. *Adv. Math.*, 82(1):1–34, 1990.
- [4] M. Brion. Group completions via Hilbert schemes. *J. Algebraic Geom.*, 12(4):605–626, 2003.
- [5] M. Brion. Log homogeneous varieties. In *Proceedings of the XVIth Latin American Algebra Colloquium (Spanish)*, Bibl. Rev. Mat. Iberoamericana, pages 1–39. Rev. Mat. Iberoamericana, Madrid, 2007.
- [6] M. Brion and S. Kumar. *Frobenius splitting methods in geometry and representation theory*, volume 231 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 2005.
- [7] M. Brion, D. Luna, and Th. Vust. Espaces homogènes sphériques. *Invent. Math.*, 84(3):617–632, 1986.
- [8] C. De Concini and C. Procesi. Complete symmetric varieties. In *Invariant theory (Montecatini, 1982)*, volume 996 of *Lecture Notes in Math.*, pages 1–44. Springer, Berlin, 1983.
- [9] J. Fogarty. Fixed point schemes. *Amer. J. Math.*, 95:35–51, 1973.
- [10] J. Gandini. Simple linear compactifications of odd orthogonal groups. *J. Algebra*, 375:258–279, 2013.
- [11] J. Gandini and A. Ruzzi. Normality and smoothness of simple linear group compactifications. *ArXiv e-prints*, March 2012.

- [12] M. Huruguen. Compactification of spherical homogeneous spaces over an arbitrary field. *Ph.D thesis, available at <http://tel.archives-ouvertes.fr/tel-00716402>*, 2011.
- [13] I. Kausz. A modular compactification of the general linear group. *Doc. Math.*, 5:553–594 (electronic), 2000.
- [14] G. Kempf, F.F. Knudsen, D. Mumford, and B. Saint-Donat. *Toroidal embeddings. I*. Lecture Notes in Mathematics, Vol. 339. Springer-Verlag, Berlin, 1973.
- [15] F. Knop. The Luna-Vust theory of spherical embeddings. In *Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989)*, pages 225–249, Madras, 1991. Manoj Prakashan.
- [16] F. Knop. Über Bewertungen, welche unter einer reductiven Gruppe invariant sind. *Math. Ann.*, 295(2):333–363, 1993.
- [17] D. Laksov. Notes on the evolution of complete correlations. In *Enumerative geometry and classical algebraic geometry (Nice, 1981)*, volume 24 of *Progr. Math.*, pages 107–132. Birkhäuser Boston, Boston, MA, 1982.
- [18] D. Laksov. Completed quadrics and linear maps. In *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, volume 46 of *Proc. Sympos. Pure Math.*, pages 371–387. Amer. Math. Soc., Providence, RI, 1987.
- [19] D. Laksov. The geometry of complete linear maps. *Ark. Mat.*, 26(2):231–263, 1988.
- [20] D. Luna. Toute variété magnifique est sphérique. *Transform. Groups*, 1(3):249–258, 1996.
- [21] D. Luna and Th. Vust. Plongements d’espaces homogènes. *Comment. Math. Helv.*, 58(2):186–245, 1983.
- [22] J. Martens and M. Thaddeus. Compactifications of reductive groups as moduli stacks of bundles. *ArXiv e-prints*, May 2011.
- [23] R. W. Richardson. On orbits of algebraic groups and Lie groups. *Bull. Austral. Math. Soc.*, 25(1):1–28, 1982.
- [24] E. Strickland. A vanishing theorem for group compactifications. *Math. Ann.*, 277(1):165–171, 1987.
- [25] H. Sumihiro. Equivariant completion. *J. Math. Kyoto Univ.*, 14:1–28, 1974.
- [26] M. Thaddeus. Complete collineations revisited. *Math. Ann.*, 315(3):469–495, 1999.

- [27] A. Thorup and S. Kleiman. Complete bilinear forms. In *Algebraic geometry (Sundance, UT, 1986)*, volume 1311 of *Lecture Notes in Math.*, pages 253–320. Springer, Berlin, 1988.
- [28] D. A. Timashëv. Equivariant compactifications of reductive groups. *Mat. Sb.*, 194(4):119–146, 2003.
- [29] J. A. Tyrell. Complete quadrics and collineations in S_n . *Mathematika*, pages 69–79, 1956.
- [30] I. Vainsencher. Complete collineations and blowing up determinantal ideals. *Math. Ann.*, 267(3):417–432, 1984.
- [31] T. Vust. Opération de groupes réductifs dans un type de cônes presque homogènes. *Bull. Soc. Math. France*, 102:317–333, 1974.

Mathieu Huruguen
Department of Mathematics
University of British Columbia
1984 Mathematics Road
Vancouver, BC, V6T 1Z2, Canada
huruguen@math.ubc.ca

