

ASYMPTOTIC FORMS OF WEAKLY INCREASING POSITIVE SOLUTIONS FOR QUASILINEAR ORDINARY DIFFERENTIAL EQUATIONS

KEN-ICHI KAMO, HIROYUKI USAMI

ABSTRACT. Asymptotic forms are determined for positive solutions which are called weakly increasing solutions to quasilinear ordinary differential equations.

1. INTRODUCTION

In this paper we consider the equation

$$(|u'|^{\alpha-1}u')' + p(t)|u|^{\beta-1}u = 0 \quad (1.1)$$

under the following conditions:

- (A1) α and β are positive constants satisfying $\alpha \neq \beta$;
- (A2) $p(t)$ is a C^1 -function defined near $+\infty$ satisfying the asymptotic condition $p(t) \sim t^{-\sigma}$ for some $\sigma \in \mathbb{R}$ as $t \rightarrow \infty$.

By condition (A2) equation (1.1) can be rewritten in the form

$$(|u'|^{\alpha-1}u')' + t^{-\sigma}(1 + \varepsilon(t))|u|^{\beta-1}u = 0, \quad (1.2)$$

where $\varepsilon(t) = t^\sigma p(t) - 1$ satisfies $\lim_{t \rightarrow \infty} \varepsilon(t) = 0$. Of course, here and in what follows the symbol “ $f(t) \sim g(t)$ as $t \rightarrow \infty$ ” means that $\lim_{t \rightarrow \infty} f(t)/g(t) = 1$. Some preparatory results for equation (1.1) are still valid for more general equations than (1.1); so it is convenient to consider the auxiliary equation

$$(|u'|^{\alpha-1}u')' + q(t)|u|^{\beta-1}u = 0, \quad (1.3)$$

where we assume that α and β satisfy condition (A1) and $q \in C([t_0, \infty); (0, \infty))$. A function u is defined to be a solution of equation (1.3) if $u \in C^1[t_1, \infty)$ and $|u'|^{\alpha-1}u' \in C^1[t_1, \infty)$ and it satisfies equation (1.3) on $[t_1, \infty)$ for sufficiently large t_1 .

It is easily seen that all positive solutions $u(t)$ of (1.3) are classified into the following three types according as their asymptotic behavior as $t \rightarrow \infty$:

- (I) Asymptotically linear solution: $u(t) \sim c_1 t$ for some constant $c_1 > 0$;
- (II) Weakly increasing solution: $u'(t) \downarrow 0$ and $u(t) \uparrow \infty$;
- (III) Asymptotically constant solution: $u(t) \sim c_1$ for some constant $c_1 > 0$.

2000 *Mathematics Subject Classification*. 34E10, 34C41.

Key words and phrases. Quasilinear ordinary differential equation; positive solution.

©2007 Texas State University - San Marcos.

Submitted May 24, 2007. Published September 28, 2007.

Concerning qualitative properties of positive solutions, the study of asymptotic behavior of asymptotically linear solutions and asymptotically constant solutions are rather easy, because their first approximations are given by definition. On the other hand, we can not easily know how the weakly increasing positive solutions behave except in the case $\alpha = 1$; see [1, 3].

In [3, Section 20], equation (1.1) with $\alpha = 1$ has been considered systematically, and asymptotic forms of weakly increasing positive solutions are given by means of the parameters β and σ . When $\alpha \neq 1$, as far as the authors are aware, there are no works in which asymptotic forms of weakly increasing positive solutions are studied systematically.

Motivated by these facts in this paper we make an attempt to find out asymptotic forms of weakly increasing positive solutions of (1.1) for the general case $\alpha > 0$. Furthermore we will also establish more than obtained in [3] in the case $\alpha > \beta$. In fact, some of our results are new even though $\alpha = 1$.

To gain an insight into our problem, we consider the typical equation

$$(|u'|^{\alpha-1}u')' + t^{-\sigma}|u|^{\beta-1}u = 0, \quad (1.4)$$

where $\sigma \in \mathbb{R}$ is a constant. Note that equation (1.1) can be regarded as a perturbation of (1.4). Equation (1.4) has a weakly increasing positive solution of the form ct^ρ , ($c > 0, 0 < \rho < 1$) if and only if $\min\{\alpha, \beta\} + 1 < \sigma < \max\{\alpha, \beta\} + 1$. This solution is uniquely given by

$$u_0(t) = \hat{C}t^k, \quad (1.5)$$

where

$$k = \frac{\alpha - \sigma + 1}{\alpha - \beta} \in (0, 1), \quad \hat{C} = \{\alpha(1 - k)k^\alpha\}^{\frac{1}{\beta - \alpha}}. \quad (1.6)$$

From this simple observation we can see that asymptotic forms of weakly increasing positive solutions of (1.3) may be strongly affected by that of the coefficient function $q(t)$. Furthermore we conjecture that weakly increasing positive solutions u of (1.1) behave like $u_0(t)$ given by (1.5) and (1.6) if $|\varepsilon(t)|$ is sufficiently small at ∞ .

We will show that the above conjecture is true in many cases. In fact, we can obtain the following theorems which are the main results of the paper:

Theorem 1.1. *Let $\alpha > \beta$.*

- (i) *Suppose that $\beta + 1 < \sigma < \alpha + 1$. Then, every weakly increasing positive solution u of (1.1) has the asymptotic form*

$$u(t) \sim u_0(t) \quad \text{as } t \rightarrow \infty,$$

where u_0 is given by (1.5) and (1.6).

- (ii) *Suppose that $\sigma = \alpha + 1$; namely $p(t) \sim t^{-\alpha-1}$ as $t \rightarrow \infty$. Then, every weakly increasing positive solution u of (1.1) has the asymptotic form*

$$u(t) \sim \alpha^{-\frac{1}{\alpha-\beta}} \left(\frac{\alpha}{\alpha-\beta}\right)^{-\frac{\alpha}{\alpha-\beta}} (\log t)^{\frac{\alpha}{\alpha-\beta}} \quad \text{as } t \rightarrow \infty.$$

Theorem 1.2. *Let $\alpha < \beta$. Suppose that $\alpha \leq 1$ and $1/2 < k < 1$ ($\Leftrightarrow (\alpha + \beta + 2)/2 < \sigma < \beta + 1$). Suppose furthermore that either*

$$\int_0^\infty \frac{\varepsilon(t)^2}{t} dt < \infty \quad (1.7)$$

or

$$\int_0^\infty |\varepsilon'(t)| dt < \infty \quad (1.8)$$

holds. Then, every weakly increasing positive solution u of (1.1) has the asymptotic form

$$u(t) \sim u_0(t) \quad \text{as } t \rightarrow \infty,$$

where u_0 is given by (1.5) and (1.6).

Theorem 1.3. *Let $\alpha < \beta$. Suppose that $\alpha \geq 1$ and $0 < k < 1/2 (\Leftrightarrow \alpha + 1 < \sigma < (\alpha + \beta + 2)/2)$. Suppose furthermore that either (1.7) or (1.8) holds. Then, the same conclusion as in Theorem 1.2 holds.*

Remark 1.4. (i) In Theorem 1.1 the differentiability of p is actually unnecessary. Similarly, in Theorems 1.2 and 1.3, the differentiability of p is unnecessary when (1.7) is assumed.

(ii) When $\alpha = 1$ and $\varepsilon(t) \equiv 0$, Theorems 1.1, 1.2 and 1.3 were obtained by [1] and [3, Corollaries 20.2, 20.3].

Remark 1.5. Let $\alpha < \beta$. We can not prove whether or not our conjecture is true in the following cases: (i) $\alpha \leq 1$ and $0 < k < 1/2$; (ii) $\alpha \geq 1$ and $1/2 < k < 1$; and (iii) $k = 1/2$.

We note that existence results of weakly increasing positive solutions to (1.3) and (1.1) are known for the case $\alpha > \beta$. In fact, equation (1.1) has a weakly increasing positive solution if and only if $\beta + 1 < \sigma \leq \alpha + 1$; see Remark 3.4 in Section 3. In contrast, it seems that there are not such useful results for the case $\alpha < \beta$. But we can show many concrete examples of those equations that have weakly increasing positive solutions.

The paper is organized as follows. In Section 2 we give preparatory lemmas employed later. In Section 3 we consider equation (1.1), as well as (1.3), under the sub-homogeneity condition $\alpha > \beta$. When $q(t)$ satisfies $0 < \liminf_{t \rightarrow \infty} q(t)/t^{-\sigma} \leq \limsup_{t \rightarrow \infty} q(t)/t^{-\sigma} < \infty$ for some $\sigma \in \mathbb{R}$, we can obtain a result which may be called as asymptotic equivalence theorem for equation (1.3); see Corollary 3.3. Theorem 1.1 is a direct consequence of this corollary. In Section 4 we consider only equation (1.1) under the super-homogeneity condition $\alpha < \beta$, and we prove Theorems 1.2 and 1.3 there. Other related results are found in [2, 4, 5, 6].

2. PREPARATORY LEMMAS

Lemma 2.1. *Let $w \in C^1[t_0, \infty)$, $w'(t) = O(1)$ as $t \rightarrow \infty$, and $w \in L^\lambda[t_0, \infty)$ for some $\lambda > 0$. Then, $\lim_{t \rightarrow \infty} w(t) = 0$.*

Proof. We have

$$\begin{aligned} |w(t)|^\lambda w(t) &= |w(t_0)|^\lambda w(t_0) + \int_{t_0}^t (|w(s)|^\lambda w(s))' ds \\ &= |w(t_0)|^\lambda w(t_0) + (\lambda + 1) \int_{t_0}^t |w(s)|^\lambda w'(s) ds. \end{aligned}$$

By our assumptions the above integral converges. Hence $\lim_{t \rightarrow \infty} |w(t)|^\lambda w(t)$ exists in \mathbb{R} . Since $w \in L^\lambda[t_0, \infty)$, the limit must be 0. The proof is complete. \square

Lemma 2.2. *Let $u \in C^1[t_0, \infty)$, $u(t) > 0$ and $u'(t) \downarrow 0$ as $t \rightarrow \infty$. Then, $tu'(t) \leq u(t)$ for sufficiently large t , and the function $u(t)/t$ is decreasing near ∞ .*

Proof. Since

$$u(t) = u(t_0) + \int_{t_0}^t u'(s) ds \geq u(t_0) + u'(t)(t - t_0),$$

we have

$$tu'(t) - u(t) \leq t_0u'(t) - u(t_0).$$

Noting the assumption $u'(\infty) = 0$ we find that $tu'(t) - u(t) < 0$ near ∞ . Since $(u(t)/t)' = (tu'(t) - u(t))/t^2$, the proof is completed. \square

3. SUB-HOMOGENEOUS CASE: $\alpha > \beta$

Throughout the section we assume that $\alpha > \beta$. As a first step we give the growth estimates for weakly increasing positive solutions of (1.3):

Lemma 3.1. *Let u be a weakly increasing positive solution of (1.3). Then the following estimates hold near ∞ :*

$$\begin{aligned} & \left(\frac{\alpha - \beta}{\alpha}\right)^{\frac{\alpha}{\alpha - \beta}} \left\{ \int_{t_1}^t \left(\int_s^\infty q(r) dr \right)^{1/\alpha} ds \right\}^{\frac{\alpha}{\alpha - \beta}} \\ & \leq u(t) \\ & \leq \left(\frac{\alpha - \beta}{\alpha}\right)^{\frac{\alpha}{\alpha - \beta}} \left\{ \int_{t_1}^t s^{-\beta/\alpha} \left(\int_s^\infty r^\beta q(r) dr \right)^{1/\alpha} ds \right\}^{\frac{\alpha}{\alpha - \beta}} \end{aligned} \quad (3.1)$$

where t_1 is a sufficiently large number.

Note that $\int^\infty s^\beta q(s) ds < \infty$ if equation (1.3) has a weakly increasing positive solution; see (ii) of Remark 3.4.

Proof of Lemma 3.1. We may assume that $u, u' > 0$ for $t \geq t_1$. Since u satisfies for large t ,

$$u'(t)^\alpha = \int_t^\infty q(s) u(s)^\beta ds, \quad (3.2)$$

and u is increasing, we have

$$u'(t)^\alpha \geq u(t)^\beta \int_t^\infty q(s) ds,$$

that is

$$u'(t) u(t)^{-\beta/\alpha} \geq \left(\int_t^\infty q(s) ds \right)^{1/\alpha}. \quad (3.3)$$

An integration of this inequality on the interval $[t_1, t]$ gives

$$\frac{\alpha}{\alpha - \beta} \left\{ u(t)^{\frac{\alpha - \beta}{\alpha}} - u(t_1)^{\frac{\alpha - \beta}{\alpha}} \right\} \geq \int_{t_1}^t \left(\int_s^\infty q(r) dr \right)^{1/\alpha} ds,$$

which proves the first inequality of (3.1). On the other hand, by the decreasing nature of $u(t)/t$ shown in Lemma 2.2, we find from (3.2) that

$$u'(t)^\alpha \leq \left(\frac{u(t)}{t}\right)^\beta \int_t^\infty s^\beta q(s) ds.$$

Accordingly,

$$u'(t) u(t)^{-\beta/\alpha} \leq t^{-\beta/\alpha} \left(\int_t^\infty s^\beta q(s) ds \right)^{1/\alpha}.$$

As before we can get the second inequality in (3.1). The proof is complete. \square

To give the main result in this section, let us consider the two equations, of the same type,

$$(|u'|^{\alpha-1}u')' + q_1(t)|u|^{\beta-1}u = 0, \tag{3.4}$$

$$(|u'|^{\alpha-1}u')' + q_2(t)|u|^{\beta-1}u = 0. \tag{3.5}$$

Here, we assume that $0 < \alpha < \beta$ and $q_1, q_2 \in C([t_0, \infty); (0, \infty))$.

Theorem 3.2. *Suppose that*

$$q_1(t) \sim q_2(t) \quad \text{as } t \rightarrow \infty, \tag{3.6}$$

$$C \int_{t_0}^t s^{-\beta/\alpha} \left(\int_s^\infty r^\beta q_1(r) dr \right)^{1/\alpha} ds \leq \int_{t_0}^t \left(\int_s^\infty q_1(r) dr \right)^{1/\alpha} ds \tag{3.7}$$

hold for some constant $C > 0$. If u_1 and u_2 are weakly increasing positive solutions of equations (3.4) and (3.5), respectively, then $u_1(t) \sim u_2(t)$ as $t \rightarrow \infty$.

Corollary 3.3. *Suppose that q_1 and q_2 satisfy (3.6) and*

$$0 < \liminf_{t \rightarrow \infty} q_1(t)/t^{-\sigma} \leq \limsup_{t \rightarrow \infty} q_1(t)/t^{-\sigma} < \infty$$

for some $\sigma \in (\beta + 1, \alpha + 1]$. If u_1 and u_2 are weakly increasing positive solutions of equations (3.4) and (3.5), respectively, then $u_1(t) \sim u_2(t)$ as $t \rightarrow \infty$.

Theorem 1.1 is an immediate consequence of Corollary 3.3. Indeed, to see (ii) of Theorem 1.1, it suffices to notice the fact that the equation

$$(|u'|^{\alpha-1}u')' + t^{-\alpha-1} \left(1 - \frac{\beta}{(\alpha - \beta) \log t} \right) |u|^{\beta-1}u = 0$$

has a weakly increasing positive solution given explicitly by

$$\alpha^{-\frac{1}{\alpha-\beta}} \left(\frac{\alpha}{\alpha - \beta} \right)^{-\frac{\alpha}{\alpha-\beta}} (\log t)^{\frac{\alpha}{\alpha-\beta}}.$$

Proof of Theorem 3.2. Put $z(t) = u_1(t)/u_2(t)$, $t \geq t_0$, t_0 being sufficiently large. Then, z satisfies the equation

$$z'' + \frac{2u_2'(t)}{u_2(t)} z' + \frac{u_2(t)^{\beta-1}}{\alpha} [(u_2(t)z' + u_2'(t)z)^{1-\alpha} q_1(t)z^\beta - q_2(t)u_2'(t)^{1-\alpha} z] = 0.$$

If $z'(T) = 0$ for some T , then

$$z''(T) = \alpha^{-1} q_1(T) u_2(T)^{\beta-1} u_2'(T)^{1-\alpha} z(T) \left(\frac{q_2(T)}{q_1(T)} - z(T)^{\beta-\alpha} \right).$$

Thus, if $z' = 0$ in the region $z > (q_1(t)/q_2(t))^{1/(\alpha-\beta)}$, then z attains a local minimum here; while if $z' = 0$ in the region $0 < z < (q_1(t)/q_2(t))^{1/(\alpha-\beta)}$, then z attains a local maximum here. Note that by our assumption $\lim_{t \rightarrow \infty} (q_1(t)/q_2(t))^{1/(\alpha-\beta)} = 1$. These simple observations are used below.

Since Lemma 3.1 and conditions (3.6) and (3.7) imply that $z(t)$ is bounded and bounded from 0, we can put $0 < \ell = \liminf_{t \rightarrow \infty} z(t) \leq \limsup_{t \rightarrow \infty} z(t) = L < \infty$. We claim that $\ell = L$; that is $\lim_{t \rightarrow \infty} z(t) \in (0, \infty)$ exists. Suppose for contradiction that $\ell \neq L$. We treat the following four cases separately: (a) $L \geq 1 > \ell$; (b) $L > 1 \geq \ell$; (c) $L > \ell \geq 1$; (d) $1 \geq L > \ell$.

Suppose that case (a) occurs. We can find two sequences $\{T_n\}$ and $\{t_n\}$ satisfying

$$\lim_{n \rightarrow \infty} T_n = \lim_{n \rightarrow \infty} t_n = \infty \tag{3.8}$$

and

$$\lim_{n \rightarrow \infty} z(T_n) = L, \quad \lim_{n \rightarrow \infty} z(t_n) = \ell, \quad t_n < T_n < t_{n+1} \quad \text{for } n = 1, 2, \dots \quad (3.9)$$

Since $\lim_{t \rightarrow \infty} (q_1(t)/q_2(t))^{1/(\alpha-\beta)} = 1$, we may assume that

$$z(t_n) < (q_1(t_n)/q_2(t_n))^{1/(\alpha-\beta)}.$$

For sufficiently large $n \in \mathbb{N}$ the minimum of $z(t)$ on the interval $[T_{n-1}, T_n]$ must be attained at an interior point, say $t_* \in (T_{n-1}, T_n)$. Obviously, $z'(t_*) = 0$ and $z''(t_*) \geq 0$. However, since $z(t_*) \leq z(t_n)$ for sufficiently large n , we get a contradiction to the above observation. Hence case (a) does not occur.

Next suppose that case (c) occurs. As in case (a) we can find two sequences $\{T_n\}$ and $\{t_n\}$ satisfying (3.8) and (3.9). For sufficiently large $n \in \mathbb{N}$ the maximum of $z(t)$ on the interval $[t_n, t_{n+1}]$ must be attained at an interior point, say $t^* \in (t_n, t_{n+1})$. Obviously, $z'(t^*) = 0$ and $z''(t^*) \leq 0$. Since $z(t^*) \geq z(T_n)$ and $z(T_n) > (q_1(T_n)/q_2(T_n))^{1/(\alpha-\beta)}$ for sufficiently large n , we get a contradiction as before. Hence case (c) does not occur. Similarly we can show that the other cases can not occur. Therefore $\lim_{t \rightarrow \infty} z(t) = \lim_{t \rightarrow \infty} u_1(t)/u_2(t) = m \in (0, \infty)$ exists. Finally, by L'Hospital's rule we have

$$\begin{aligned} m &= \lim_{t \rightarrow \infty} \frac{u_1(t)}{u_2(t)} = \left(\lim_{t \rightarrow \infty} \frac{u_1'(t)^\alpha}{u_2'(t)^\alpha} \right)^{1/\alpha} \\ &= \left(\lim_{t \rightarrow \infty} \frac{[u_1'(t)^\alpha]'}{[u_2'(t)^\alpha]'} \right)^{1/\alpha} \\ &= \left(\lim_{t \rightarrow \infty} \frac{-q_1(t)u_1(t)^\beta}{-q_2(t)u_2(t)^\beta} \right)^{1/\alpha} = m^{\beta/\alpha}, \end{aligned}$$

that is $m = m^{\beta/\alpha}$. Since $\alpha > \beta$, we have $m = 1$. This completes the proof. \square

Remark 3.4. Concerning the existence properties of weakly increasing positive solutions, we know the following results:

(i) If

$$\int_0^\infty t^\beta q(t) dt < \infty; \quad \text{and} \quad \int_0^\infty \left(\int_t^\infty q(s) ds \right)^{1/\alpha} dt = \infty, \quad (3.10)$$

Then, equation (1.3) has weakly increasing positive solutions [2, Example 1].

(ii) Conversely, if equation (1.3) has a weakly increasing positive solution, then we can show that

$$\int_0^\infty t^\beta q(t) dt < \infty; \quad \text{and} \quad \int_0^\infty t^{-\beta/\alpha} \left(\int_t^\infty s^\beta q(s) ds \right)^{1/\alpha} dt = \infty. \quad (3.11)$$

In fact, the first condition in (3.11) follows from [2, Example 2]; while the second one is an immediate consequence of the estimates in Lemma 3.1. In particular, when $\alpha = 1$, we find that conditions (3.10) and (3.11) are the same; that is, equation (1.3) (with $\alpha = 1$) has a weakly increasing positive solution if and only if (3.10) (with $\alpha = 1$) holds.

4. SUPER-HOMOGENEOUS CASE: $\alpha < \beta$

Throughout this section we assume that $\alpha < \beta$. In this case the situation seems to be more complicated than in the previous case. The main purpose of the section is to give the proofs of Theorems 1.2 and 1.3. To this end we need several lemmas.

Lemma 4.1. *Let $0 < \liminf_{t \rightarrow \infty} q(t)/t^{-\sigma} \leq \limsup_{t \rightarrow \infty} q(t)/t^{-\sigma} < \infty$ for some $\sigma \in (\alpha+1, \beta+1)$. Then every weakly increasing positive solution u of (1.1) satisfies $u(t) = O(u_0(t))$ and $u'(t) = O(u'_0(t))$ as $t \rightarrow \infty$, where u_0 is the exact solution of (1.4) given by (1.5) and (1.6).*

Proof. As in the proof of Lemma 3.1 we obtain (3.3). An integration of (3.3) on the interval $[t, \infty)$ for t large will give

$$u(t) \leq C_1 \left\{ \int_t^\infty \left(\int_s^\infty q(r) dr \right)^{1/\alpha} ds \right\}^{-\alpha/(\beta-\alpha)} \leq C_2 u_0(t),$$

where C_1 and C_2 are positive constant. Furthermore, by (3.2) we find that

$$u'(t) = \left(\int_t^\infty q(s) u(s)^\beta ds \right)^{1/\alpha} \leq C_3 \left(\int_t^\infty s^{-\sigma+k\beta} ds \right)^{1/\alpha} = C_4 t^{k-1} = O(u'_0(t))$$

as $t \rightarrow \infty$, where C_3 and C_4 are positive constants. This completes the proof. \square

Lemma 4.2. *Let $\sigma \in (\alpha+1, \beta+1)$, and u be a weakly increasing positive solution of equation (1.1). Put $s = \log u_0(t)$ and $v = u/u_0$. Then*

- (i) $v, \dot{v} = O(1)$ as $s \rightarrow \infty$, and $v + \dot{v} > 0$ near ∞ , where $\cdot = d/ds$;
- (ii) $v(s)$ satisfies near ∞ the equation

$$\ddot{v} + av - bv + b(\dot{v} + v)^{1-\alpha} v^\beta + b\delta(s)(\dot{v} + v)^{1-\alpha} v^\beta = 0, \quad (4.1)$$

where $a = 2 - \frac{1}{k} \neq 0$, $b = \frac{1-k}{k} > 0$, and $\delta(s) = \varepsilon(t)$.

Proof. We will prove only (i), because (ii) can be proved by direct computations. We note that $u, u' > 0$. Since $u = u_0 v$, the boundedness of v follows from Lemma 4.1. Noting $du/dt = \hat{C} k t^{k-1} (v + \dot{v})$, we have $v + \dot{v} > 0$. On the other hand, since $dt/ds = t/k$, we have

$$|\dot{v}| = \left| \frac{d}{dt} \left(\frac{u}{u_0} \right) \frac{dt}{ds} \right| = \left| \frac{u' u_0 - u'_0 u}{u_0^2} \right| \frac{t}{k} \leq C \frac{t^{k-1} t^{kt}}{t^{2k}} = O(1) \quad \text{as } s \rightarrow \infty.$$

This completes the proof. \square

Lemma 4.3. *Let the assumption either of Theorem 1.2 or Theorem 1.3 hold, and v be as in Lemma 4.2. Then $\dot{v} \in L^2[s_0, \infty)$ for sufficiently large s_0 .*

Proof. We note that conditions (1.7) and (1.8), respectively, are equivalent to

$$\int_{s_0}^\infty \delta(s)^2 ds < \infty \quad (4.2)$$

and

$$\int_{s_0}^\infty |\dot{\delta}(s)| ds < \infty. \quad (4.3)$$

We firstly consider the case where assumptions of Theorem 1.2 hold. In this case, the constant a appearing in (4.1) is positive. We multiply the both sides of (4.1) by \dot{v} . Since $\alpha \leq 1$, we have $(1 + \delta(s))(\dot{v} + v)^{1-\alpha} \dot{v} \geq (1 + \delta(s))v^{1-\alpha} \dot{v}$; and so we obtain

$$\ddot{v} + av^2 - bv\dot{v} + bv^{1-\alpha+\beta} \dot{v} + b\delta(s)v^{1-\alpha+\beta} \dot{v} \leq 0. \quad (4.4)$$

An integration gives

$$\frac{\dot{v}^2}{2} + a \int_{s_0}^s \dot{v}^2 dr - \frac{b}{2} v^2 + \frac{bv^{2-\alpha+\beta}}{2-\alpha+\beta} + b \int_{s_0}^s \delta(r)v^{1-\alpha+\beta} \dot{v} dr \leq \text{const}; \quad (4.5)$$

that is

$$a \int_{s_0}^s \dot{v}^2 dr + b \int_{s_0}^s \delta(r)v^{1-\alpha+\beta} \dot{v} dr \leq O(1) \quad \text{as } s \rightarrow \infty.$$

Here we have employed (i) of Lemma 4.2. Let the integral condition (1.7) hold; that is, let (4.2) hold. By the Schwarz inequality we have

$$a \int_{s_0}^s \dot{v}^2 dr - C_1 \left(\int_{s_0}^{\infty} \delta(r)^2 dr \right)^{1/2} \left(\int_{s_0}^s \dot{v}^2 dr \right)^{1/2} \leq O(1)$$

for some constant $C_1 > 0$. Therefore $\dot{v} \in L^2[s_0, \infty)$. Next let (1.8) hold. Using integral by parts, we obtain from (4.5)

$$\frac{\dot{v}^2}{2} + a \int_{s_0}^s \dot{v}^2 dr - \frac{b}{2} v^2 + \frac{b[1+\delta(r)]v^{2-\alpha+\beta}}{2-\alpha+\beta} - \frac{b}{2-\alpha+\beta} \int_{s_0}^s \delta(r)v^{2-\alpha+\beta} dr \leq \text{const}.$$

Noting (i) of Lemma 4.2, we find that $\dot{v} \in L^2[s_0, \infty)$.

Secondly we consider the case where assumptions of Theorem 1.3 hold. As above, we multiply both the sides of (4.1) by \dot{v} . Since $\alpha \geq 1$, we have $(1+\delta(s))(v+\dot{v})^{1-\alpha} \dot{v} \leq (1+\delta(s))v^{1-\alpha} \dot{v}$, and so we obtain

$$|a|\dot{v}^2 \leq \dot{v}\ddot{v} - b\dot{v} + b(1+\delta(s))v^{1-\alpha+\beta}\dot{v}.$$

An integration on the interval $[s_0, s]$ gives

$$|a| \int_{s_0}^s \dot{v}^2 dr \leq \frac{\dot{v}^2}{2} - \frac{bv^2}{2} + \frac{bv^{2-\alpha+\beta}}{2-\alpha+\beta} + b \int_{s_0}^s \delta(r)v^{1-\alpha+\beta} \dot{v} dr + \text{const}.$$

As before, we will obtain $\dot{v} \in L^2[s_0, \infty)$. This completes the proof. \square

Proof of Theorem 1.2. To this end it suffices to show that $\lim_{s \rightarrow \infty} v(s) = 1$, where $v(s)$ is the function introduced in Lemma 4.2. The proof is divided into three steps.

Step 1. We claim that $\liminf_{s \rightarrow \infty} v(s) > 0$; or equivalently $\liminf_{t \rightarrow \infty} u(t)/u_0(t) > 0$. The proof is done by contradiction. Suppose that $\liminf_{s \rightarrow \infty} v(s) = 0$. Firstly, we suppose that $v(s)$ decrease to 0 as $s \rightarrow \infty$. This means that $u(t)/u_0(t)$ decreases to 0 as $t \rightarrow \infty$. Accordingly we have

$$\begin{aligned} u'(t)^\alpha &= \int_t^\infty p(r)u(r)^\beta dr \\ &= \int_t^\infty p(r)u_0(r)^\beta \left(\frac{u(r)}{u_0(r)} \right)^\beta dr \\ &\leq \left(\frac{u(t)}{u_0(t)} \right)^\beta \int_t^\infty p(r)u_0(r)^\beta dr \\ &= C_1 t^{1-\sigma} u(t)^\beta, \end{aligned}$$

where $C_1 > 0$ is a constant. Consequently we obtain the differential inequality $u' \leq C_2 t^{(1-\sigma)/\alpha} u^{\beta/\alpha}$ for some constant $C_2 > 0$ near ∞ . But this differential inequality implies that $u(t)/u_0(t) \equiv v(s) \geq C_3 > 0$ for some constant $C_3 > 0$. This is an obvious contradiction. Next suppose that $\liminf_{s \rightarrow \infty} v(s) = 0$ and $\dot{v}(s)$ changes the sign in any neighborhood of ∞ . We notice that, if $\dot{v} = 0$ in

the region $0 < v < [1 + \delta(s)]^{-1/(\beta-\alpha)}$, then $\ddot{v} > 0$; while if $\dot{v} = 0$ in the region $v > [1 + \delta(s)]^{-1/(\beta-\alpha)}$, then $\ddot{v} < 0$. Therefore, in this case the curve $v = v(s)$ must cross the curve $v = [1 + \delta(s)]^{-1/(\beta-\alpha)}$ infinitely many times as $s \rightarrow \infty$. Therefore, we can find out two sequences $\{\xi_n\}$ and $\{\eta_n\}$ satisfying

$$\xi_n < \eta_n < \xi_{n+1}, \quad n = 1, 2, \dots; \quad \lim_{n \rightarrow \infty} \xi_n = \lim_{n \rightarrow \infty} \eta_n = \infty$$

and

$$v(\eta_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad v(\xi_n) = [1 + \delta(\xi_n)]^{-\frac{1}{\beta-\alpha}} \rightarrow 1 \quad \text{as } n \rightarrow \infty.$$

An integration of (4.4) on $[\xi_n, \eta_n]$ yields

$$\begin{aligned} & \frac{1}{2} \{ \dot{v}(\eta_n)^2 - \dot{v}(\xi_n)^2 \} + a \int_{\xi_n}^{\eta_n} \dot{v}^2 dr - \frac{b}{2} \{ v(\eta_n)^2 - v(\xi_n)^2 \} \\ & + \frac{b}{2 - \alpha + \beta} \{ v(\eta_n)^{2-\alpha+\beta} - v(\xi_n)^{2-\alpha+\beta} \} + b \int_{\xi_n}^{\eta_n} \delta(r) v^{1-\alpha+\beta} \dot{v} dr \leq 0. \end{aligned} \quad (4.6)$$

From equation (4.1) and (i) of Lemma 4.2 we know that $\ddot{v} = O(1)$ as $s \rightarrow \infty$. This fact and the fact that $\dot{v} \in L^2[s_0, \infty)$ imply that $\lim_{s \rightarrow \infty} \dot{v}(s) = 0$ by Lemma 2.1. Accordingly (4.6) is equivalent to

$$o(1) + o(1) - \frac{b}{2}(o(1) - 1) + \frac{b}{2 - \alpha + \beta}(o(1) - 1) + o(1) \leq 0 \quad \text{as } s \rightarrow \infty.$$

Letting $n \rightarrow \infty$, we get $b/2 - b/(2 - \alpha + \beta) \leq 0$ a contradiction to the assumption $\beta > \alpha$. Therefore, $\liminf_{s \rightarrow \infty} v(s) > 0$.

Step 2. We claim that there is a limit $\lim_{s \rightarrow \infty} v(s) \in (0, \infty)$. To see this, we integrate (4.1) multiplied by \dot{v} :

$$\begin{aligned} & \frac{\dot{v}^2}{2} + a \int_{s_0}^s \dot{v}^2 dr - \frac{b}{2} v^2 + b \int_{s_0}^s (\dot{v} + v)^{1-\alpha} v^\beta \dot{v} dr \\ & + b \int_{s_0}^s \delta(r) (\dot{v} + v)^{1-\alpha} v^\beta \dot{v} dr = \text{const.} \end{aligned} \quad (4.7)$$

Suppose that condition (1.7) holds; or equivalently (4.2) holds. Since $\dot{v} \in L^2[s_0, \infty)$, the first, and the third integrals in the left hand side of (4.7) converge as $s \rightarrow \infty$. The mean value theorem shows that

$$(v + \dot{v})^{1-\alpha} = \left(1 + \frac{\dot{v}}{v}\right)^{1-\alpha} v^{1-\alpha} = v^{1-\alpha} + (1 - \alpha)(v + \theta(r)\dot{v})^{-\alpha} \dot{v}, \quad (4.8)$$

where $\theta(r)$ is a quantity satisfying $0 < \theta(r) < 1$. Therefore,

$$\begin{aligned} \int_{s_0}^s (\dot{v} + v)^{1-\alpha} v^\beta \dot{v} dr &= \int_{s_0}^s \{ v^{1-\alpha+\beta} \dot{v} + (1 - \alpha)(v + \theta(r)\dot{v})^{-\alpha} v^\beta \dot{v}^2 \} dr \\ &= \frac{v(s)^{2+\beta-\alpha}}{2 + \beta - \alpha} + \int_{s_0}^s O(1) \dot{v}^2 dr + \text{const.} \end{aligned}$$

So we find that the function $-v^2/2 + v^{2+\beta-\alpha}/(2+\beta-\alpha)$ has a finite limit. This fact shows that $\lim_{s \rightarrow \infty} v(s) = m \in (0, \infty)$ exists. Suppose that (1.8) or equivalently

(4.3) holds. By (4.8), we have

$$\begin{aligned} & \int_{s_0}^s \delta(r)(\dot{v} + v)^{1-\alpha} v^\beta \dot{v} dr \\ &= \int_{s_0}^s \{\delta(r)v^{1-\alpha+\beta} \dot{v} + \delta(r)(1-\alpha)(v + \theta(r)\dot{v})^{-\alpha} v^\beta \dot{v}^2\} dr \\ &= \frac{\delta(s)v^{2+\beta-\alpha}}{2+\beta-\alpha} - \frac{1}{2+\beta-\alpha} \int_{s_0}^s \dot{\delta}(r)v^{2+\beta-\alpha} dr + \text{const} + \int_{s_0}^s O(1)\dot{v}^2 dr \end{aligned}$$

as $s \rightarrow \infty$. Hence, as before we know that the function $-v^2/2 + v^{2+\beta-\alpha}/(2+\beta-\alpha)$ has a finite limit. Therefore $m = \lim_{s \rightarrow \infty} v(s) \in (0, \infty)$ exists.

Step 3. Finally, we let $s \rightarrow \infty$ in equation (4.1). Then, we have $\lim_{s \rightarrow \infty} \ddot{v}(s) = b(m - m^{1+\beta-\alpha})$. Since $\dot{v} = o(1)$, we must have $\lim_{s \rightarrow \infty} \ddot{v}(s) = 0$, implying $m = 1$. The proof of Theorem 1.2 is complete. \square

Proof of Theorem 1.3. Firstly we show that $\liminf_{s \rightarrow \infty} v(s) > 0$. The proof is done by contradiction. Suppose that $\liminf_{s \rightarrow \infty} v(s) = 0$. We may assume that \dot{v} changes the sign in any neighborhood of ∞ , because the case in which v decreases to 0 can be treated as in the proof of Theorem 1.2 (Step 1). Since $v(s)$ takes local maxima in the region $v \geq (1 + \delta(s))^{-1/(\beta-\alpha)}$, there are the following sequences $\{\underline{s}_n\}$ and $\{\bar{s}_n\}$ satisfying

$$\begin{aligned} \underline{s}_n < \bar{s}_n < \underline{s}_{n+1}, \quad \lim_{n \rightarrow \infty} \underline{s}_n = \lim_{n \rightarrow \infty} \bar{s}_n = \infty, \\ \dot{v}(\underline{s}_n) = \dot{v}(\bar{s}_n) = 0, \quad \lim_{n \rightarrow \infty} v(\underline{s}_n) = 0, \quad v(\bar{s}_n) \geq (1 + \delta(\bar{s}_n))^{-1/(\beta-\alpha)}. \end{aligned}$$

Now, we decompose α in the form $\alpha = m - \rho$, where $m \in \mathbb{N}$ and $\rho > 0$. Although there are infinitely many such choices of decomposition for α , we fix one choice for a moment. We rewrite equation (4.1) as

$$\ddot{v} - |a|\dot{v} - bv + b(\dot{v} + v)^{-m+1+\rho} v^\beta + b\delta(s)(\dot{v} + v)^{-m+1+\rho} v^\beta = 0.$$

We multiply both the sides by $(v + \dot{v})^m \dot{v}$ and then integrate the resulting equation on the interval $[\underline{s}_n, \bar{s}_n]$ to obtain

$$\begin{aligned} & \int_{\underline{s}_n}^{\bar{s}_n} \ddot{v} \dot{v} (v + \dot{v})^m dr - |a| \int_{\underline{s}_n}^{\bar{s}_n} (v + \dot{v})^m \dot{v}^2 dr - b \int_{\underline{s}_n}^{\bar{s}_n} v \dot{v} (v + \dot{v})^m dr \\ & + b \int_{\underline{s}_n}^{\bar{s}_n} (v + \dot{v})^{1+\rho} \dot{v} v^\beta dr + b \int_{\underline{s}_n}^{\bar{s}_n} \delta(r) (v + \dot{v})^{1+\rho} \dot{v} v^\beta dr = 0. \end{aligned} \tag{4.9}$$

The binomial expansion implies

$$\begin{aligned} & \sum_{k=0}^m c_k \int_{\underline{s}_n}^{\bar{s}_n} \ddot{v} \dot{v}^{k+1} v^{m-k} dr - |a| \int_{\underline{s}_n}^{\bar{s}_n} (v + \dot{v})^m \dot{v}^2 dr - b \sum_{k=0}^m c_k \int_{\underline{s}_n}^{\bar{s}_n} v^{m-k+1} \dot{v}^{k+1} dr \\ & + b \int_{\underline{s}_n}^{\bar{s}_n} (v + \dot{v})^{1+\rho} \dot{v} v^\beta dr + b \int_{\underline{s}_n}^{\bar{s}_n} \delta(r) (v + \dot{v})^{1+\rho} \dot{v} v^\beta dr = 0, \end{aligned}$$

where $c_k = \binom{m}{k}$ are the binomial coefficients. Now, we evaluate each term in the left hand side. For $k \in \{0, 1, \dots, m - 1\}$ we obtain

$$\begin{aligned} \int_{\underline{s}_n}^{\bar{s}_n} \ddot{v} \dot{v}^{k+1} v^{m-k} dr &= \int_{\underline{s}_n}^{\bar{s}_n} \frac{d}{dr} \left(\frac{\dot{v}^{k+2}}{k+2} \right) v^{m-k} dr \\ &= -\frac{m-k}{k+2} \int_{\underline{s}_n}^{\bar{s}_n} \dot{v}^{k+3} v^{m-k-1} dr = o(1) \quad \text{as } n \rightarrow \infty. \end{aligned}$$

For $k = m$ obviously we have $\int_{\underline{s}_n}^{\bar{s}_n} \ddot{v} \dot{v}^{k+1} dr = 0$. Hence the first term of the left hand side of (4.9) tends to 0 as $n \rightarrow \infty$. The second term is dominated by $\text{const} \int_{\underline{s}_n}^{\bar{s}_n} \dot{v}^2 dr$, and hence it tends to zero as $n \rightarrow \infty$. Next, we compute the third term. For $k \in \{1, 2, \dots, m\}$ we have $|\int_{\underline{s}_n}^{\bar{s}_n} v^{m-k+1} \dot{v}^{k+1} dr| \leq \text{const} \int_{\underline{s}_n}^{\bar{s}_n} \dot{v}^2 dr$. For $k = 0$, we have

$$\int_{\underline{s}_n}^{\bar{s}_n} v^{m+1} \dot{v} dr = \frac{1}{m+2} (v(\bar{s}_n)^{m+2} - v(\underline{s}_n)^{m+2}) = \frac{v(\bar{s}_n)^{m+2}}{m+2} + o(1) \quad \text{as } n \rightarrow \infty.$$

Therefore, the third term is equal to

$$o(1) - \frac{bv(\bar{s}_n)^{m+2}}{m+2} \quad \text{as } n \rightarrow \infty.$$

To evaluate the fourth term we employ the mean value theorem to obtain

$$(v + \dot{v})^{1+\rho} = v^{1+\rho} + (1 + \rho)(v + \theta(r)\dot{v})^\rho \dot{v},$$

where $\theta(r)$ is a quantity between 0 and 1. Hence we can compute

$$\begin{aligned} &\int_{\underline{s}_n}^{\bar{s}_n} (v + \dot{v})^{1+\rho} \dot{v}^\beta dr \\ &= \int_{\underline{s}_n}^{\bar{s}_n} v^{1+\rho+\beta} \dot{v}^\beta dr + (1 + \rho) \int_{\underline{s}_n}^{\bar{s}_n} (v + \theta(r)\dot{v})^\rho \dot{v}^2 v^\beta dr \\ &= \frac{v(\bar{s}_n)^{2+\rho+\beta} - v(\underline{s}_n)^{2+\rho+\beta}}{2 + \rho + \beta} + (1 + \rho) \int_{\underline{s}_n}^{\bar{s}_n} O(1) \dot{v}^2 dr \\ &= \frac{v(\bar{s}_n)^{2+\rho+\beta}}{2 + \rho + \beta} + o(1) \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Finally by Schwarz's inequality we find that the last term is dominated by the quantity

$$\text{const} \left(\int_{\underline{s}_n}^{\bar{s}_n} \delta(r)^2 dr \right)^{1/2} \left(\int_{\underline{s}_n}^{\bar{s}_n} \dot{v}^2 dr \right)^{1/2} = o(1) \quad \text{as } n \rightarrow \infty.$$

(Note that (1.7) or equivalently (4.2) is assumed.) Consequently, from (4.9) we obtain the formula

$$o(1) - \frac{b}{m+2} v(\bar{s}_n)^{m+2} + \frac{b}{2 + \rho + \beta} v(\bar{s}_n)^{2+\rho+\beta} + o(1) = 0 \quad \text{as } n \rightarrow \infty.$$

This implies that $\lim_{n \rightarrow \infty} v(\bar{s}_n) = [(m + 2 + \beta - \alpha)/(m + 2)]^{1/(\beta - \alpha)}$. Since m can be moved arbitrarily, this is an obvious contradiction. Therefore $\liminf_{s \rightarrow \infty} v > 0$.

We are now in a position to show $\lim_{s \rightarrow \infty} v(s) = 1$. Since $\liminf_{s \rightarrow \infty} v(s) > 0$, we find that $\liminf_{t \rightarrow \infty} u(t)/u_0(t) > 0$. Integrating equation (3.2) (with q replaced by p), we further find that $\liminf_{t \rightarrow \infty} u'(t)/u'_0(t) > 0$. Since $v + \dot{v} = u'(t)/u'_0(t)$, we

obtain $\liminf_{s \rightarrow \infty} (v + \dot{v}) > 0$. Recalling equation (4.1), we find that $\ddot{v}(s) = O(1)$ as $s \rightarrow \infty$. Since we have already shown that $\dot{v} \in L^2[s_0, \infty)$, Lemma 2.1 shows that $\lim_{s \rightarrow \infty} \dot{v} = 0$. Consequently, as in the proof of Theorem 1.2 we can prove that $\lim_{s \rightarrow \infty} v(s) = 1$. This completes the proof of Theorem 1.3. \square

REFERENCES

- [1] R. Bellman, *Stability theory of differential equations*, McGraw-Hill, New York, 1953. (Reprint: Dover Publications, Inc., 1969)
- [2] Á. Elbert and T. Kusano, *Oscillation and non-oscillation theorems for a class of second order quasilinear differential equations*, Acta Math. Hung. 56 (1990), 325-336.
- [3] I. T. Kiguradze and T. A. Chanturia, *Asymptotic properties of solutions of nonautonomous ordinary differential equations*, Kluwer Academic Publisher, Dordrecht, 1992.
- [4] M. Kitano and T. Kusano, *On a class of second order quasilinear ordinary differential equations*, Hiroshima Math. J. 25 (1995), 321-355.
- [5] R. A. Moore and Z. Nehari, *Nonoscillation theorems for a class of nonlinear differential equations*, Trans. Amer. Math. Soc. 93 (1959), 30-52.
- [6] K. Nishihara, *Asymptotic behaviors of solutions of second order differential equations*, J. Math. Anal. Appl. 189 (1995), 424-441.

KEN-ICHI KAMO

DIVISION OF MATHEMATICS, SCHOOL OF MEDICINE, LIBERAL ARTS AND SCIENCES, SAPPORO MEDICAL UNIVERSITY, S1W17, CHUOKU, SAPPORO 060-8543, JAPAN

E-mail address: kamo@sapmed.ac.jp

HIROYUKI USAMI

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, HIROSHIMA UNIVERSITY, HIGASHI - HIROSHIMA, 739-8521, JAPAN

E-mail address: usami@mis.hiroshima-u.ac.jp