

## NON-LOCAL PROBLEMS WITH INTEGRAL GLUING CONDITION FOR LOADED MIXED TYPE EQUATIONS INVOLVING THE CAPUTO FRACTIONAL DERIVATIVE

OBIDJON KH. ABDULLAEV, KISHIN B. SADARANGANI

ABSTRACT. In this work, we study the existence and uniqueness of solutions to non-local boundary value problems with integral gluing condition. Mixed type equations (parabolic-hyperbolic) involving the Caputo fractional derivative have loaded parts in Riemann-Liouville integrals. Thus we use the method of integral energy to prove uniqueness, and the method of integral equations to prove existence.

### 1. INTRODUCTION AND FORMULATION OF A PROBLEM

The models of fractional-order derivatives are more adequate than the previously used integer-order models, because fractional-order derivatives and integrals enable the description of the memory and hereditary properties of different substances [23]. This is the most significant advantage of the fractional-order models in comparison with integer-order models, in which such effects are neglected.

Fractional differential equations have recently been proved to be valuable tools in the modeling of many phenomena in various fields of science and engineering. Indeed, we can find numerous applications in viscoelasticity, neurons, electrochemistry, control, porous media, electromagnetism, etc., (see [6, 7, 8, 15, 18, 19]). There has been a significant development in fractional differential equations in recent years; see the monographs of Kilbas, Srivastava, Trujillo [14], Miller and Ross [20], Podlubny [23], Samko, Kilbas, Marichev. [28] and the references therein.

Very recently, some basic theory for the initial boundary value problems of fractional differential equations involving a Riemann-Liouville differential operator of order  $0 < \alpha \leq 1$  has been discussed by Lakshmikantham and Vatsala [16, 17]. In a series of papers (see [4, 5]) the authors considered some classes of initial value problems for functional differential equations involving Riemann-Liouville and Caputo fractional derivatives of order  $0 < \alpha \leq 1$ : For more details concerning geometric and physical interpretation of fractional derivatives of Riemann-Liouville and Caputo types see [24]. Note that works [3, 12, 13, 22] are devoted to the studying of boundary value problems (BVP) for parabolic-hyperbolic equations, involving fractional derivatives. BVPs for the mixed type equations involving the Caputo and

---

2010 *Mathematics Subject Classification.* 34K37, 35M10.

*Key words and phrases.* Caputo fractional derivatives; loaded equation; non-local problem; integral gluing condition; existence; uniqueness.

©2016 Texas State University.

Submitted April 8, 2016. Published June 28, 2016.

the Riemann-Liouville fractional differential operators were investigated in works [10, 11].

For the first time it was given the most general definition of a loaded equations and various loaded equations are classified in detail by Nakhushiev [21]. Note that with intensive research on problem of optimal control of the agro-economical system, regulating the label of ground waters and soil moisture, it has become necessary to investigate BVPs for such class of partial differential equations. More results on the theory of BVP's for the loaded equations parabolic, parabolic-hyperbolic and elliptic-hyperbolic types were published in works [9, 1]. Integral boundary conditions have various applications in thermoelasticity, chemical engineering, population dynamics, etc. Gluing conditions of integral form were used in [2, 27].

We consider the equation:

$$\begin{aligned} u_{xx} - {}_C D_{0y}^\alpha u + p(x, y) \int_x^1 (t-x)^{\beta-1} u(t, 0) dt &= 0 \quad \text{for } y > 0 \\ u_{xx} - u_{yy} - q(x+y) \int_{x+y}^1 (t-x-y)^{\gamma-1} u(t, 0) dt &= 0 \quad \text{for } y < 0, \end{aligned} \quad (1.1)$$

with the operator

$${}_C D_{0y}^\alpha f = \frac{1}{\Gamma(1-\alpha)} \int_0^y (y-t)^{-\alpha} f'(t) dt, \quad (1.2)$$

where  $0 < \alpha, \beta, \gamma < 1$ ,  $p(x, y)$  and  $q(x+y)$  are given functions. Let  $\Omega$  be domain, bounded with segments:  $A_1 A_2 = \{(x, y) : x = 1, 0 < y < h\}$ ,  $B_1 B_2 = \{(x, y) : x = 0, 0 < y < h\}$ ,  $B_2 A_2 = \{(x, y) : y = h, 0 < x < 1\}$  at the  $y > 0$ , and characteristics :  $A_1 C : x - y = 1$ ;  $B_1 C : x + y = 0$  of the equation (1.1) at  $y < 0$ , where  $A_1(1; 0)$ ,  $A_2(1; h)$ ,  $B_1(0; 0)$ ,  $B_2(0; h)$ ,  $C(1/2; -1/2)$ .

Let us introduce:  $\theta(x) = \frac{x+1}{2} + i \frac{x-1}{2}$ ,  $i^2 = -1$ ,

$$D_{x\bar{a}}^{-\beta} f(x) = \frac{1}{\Gamma(\beta)} \int_x^a (t-x)^{\beta-1} f(t) dt, \quad 0 < \beta < 1. \quad (1.3)$$

$\Omega^+ = \Omega \cap (y > 0)$ ,  $\Omega^- = \Omega \cap (y < 0)$ ,  $I_1 = \{x : \frac{1}{2} < x < 1\}$ ,  $I_2 = \{y : 0 < y < h\}$ . In the domain  $\Omega$  we study the following problem.

**Problem I.** Find a solution  $u(x, y)$  of equation (1.1) from the class of functions:

$$W = \{u(x, y) : u(x, y) \in C(\bar{\Omega}) \cap C^2(\Omega^-), u_{xx} \in C(\Omega^+), {}_C D_{0y}^\alpha u \in C(\Omega^+)\},$$

that satisfies the boundary conditions:

$$u(x, y)|_{A_1 A_2} = \varphi(y), \quad 0 \leq y \leq h; \quad (1.4)$$

$$u(x, y)|_{B_1 B_2} = \psi(y), \quad 0 \leq y \leq h; \quad (1.5)$$

$$\frac{d}{dx} u(\theta(x)) = a(x) u_y(x, 0) + b(x) u_x(x, 0) + c(x) u(x, 0) + d(x), \quad x \in I_1, \quad (1.6)$$

and gluing condition

$$\lim_{y \rightarrow +0} y^{1-\alpha} u_y(x, y) = \lambda_1(x) u_y(x, -0) + \lambda_2(x) \int_x^1 r(t) u(t, 0) dt, \quad 0 < x < 1 \quad (1.7)$$

where  $\varphi(y)$ ,  $\psi(y)$ ,  $a(x)$ ,  $b(x)$ ,  $c(x)$ ,  $d(x)$ , and  $\lambda_j(x)$ , are given functions, such that  $\sum_{j=1}^2 \lambda_j^2(x) \neq 0$ .

2. UNIQUENESS OF SOLUTION FOR PROBLEM I

It is known that equation (1.1) on the characteristics coordinate  $\xi = x + y$  and  $\eta = x - y$  at  $y \leq 0$  has the form

$$u_{\xi\eta} = \frac{q(\xi)}{4} \int_{\xi}^1 (t - \xi)^{\gamma-1} u(t, 0) dt. \tag{2.1}$$

We introduce the notation:  $u(x, 0) = \tau(x)$ ,  $0 \leq x \leq 1$ ;  $u_y(x, -0) = \nu^-(x)$ ,  $0 < x < 1$ ;

$$\lim_{y \rightarrow +0} y^{1-\alpha} u_y(x, y) = \nu^+(x), \quad 0 < x < 1.$$

A solution of the Cauchy problem for the equation (1.1) in the domain  $\Omega^-$  can be represented as

$$\begin{aligned} u(x, y) &= \frac{\tau(x + y) + \tau(x - y)}{2} - \frac{1}{2} \int_{x+y}^{x-y} \nu^-(t) dt \\ &+ \frac{1}{4} \int_{x+y}^{x-y} q(\xi) d\xi \int_{\xi}^{x-y} d\eta \int_{\xi}^1 (t - \xi)^{\gamma-1} \tau(t) dt. \end{aligned} \tag{2.2}$$

After using condition (1.6) and taking (1.3) into account, from (2.2) we obtain

$$\begin{aligned} &(2a(x) - 1)\nu^-(x) \\ &= \frac{1-x}{2} \Gamma(\gamma) q(x) D_{x-1}^{-\gamma} \tau(x) + (1 - 2b(x))\tau'(x) - 2c(x)\tau(x) - 2d(x). \end{aligned} \tag{2.3}$$

Considering above notation and gluing condition (1.7) we have

$$\nu^+(x) = \lambda_1(x)\nu^-(x) + \lambda_2(x) \int_x^1 r(t)\tau(t) dt, \tag{2.4}$$

Further from (1.1) at  $y \rightarrow +0$  considering (1.2), (2.4) and

$$\lim_{y \rightarrow 0} D_{0y}^{\alpha-1} f(y) = \Gamma(\alpha) \lim_{y \rightarrow 0} y^{1-\alpha} f(y),$$

we obtain k1:

$$\tau''(x) - \Gamma(\alpha)\lambda_1(x)\nu^-(x) - \Gamma(\alpha)\lambda_2(x) \int_x^1 r(t)\tau(t) dt + \Gamma(\beta)p(x, 0)D_{x1}^{-\beta} \tau(x) = 0. \tag{2.5}$$

**Main results.**

**Theorem 2.1.** *If the given functions satisfy conditions*

$$\left(\frac{\lambda_2(x)}{r(x)}\right)' \geq 0, \quad \frac{\lambda_1(0)q(0)}{2a(0) - 1} \geq 0, \quad p(0, 0) \leq 0, \quad p'(x, 0) \leq 0, \quad \frac{\lambda_2(0)}{r(0)} \geq 0; \tag{2.6}$$

$$\left(\frac{(1-x)q(x)\lambda_1(x)}{(2a(x) - 1)}\right)' \geq 0, \quad \frac{c(x)\lambda_1(x)}{(2a(x) - 1)} \leq 0, \quad \left(\frac{(1 - 2b(x))\lambda_1(x)}{(2a(x) - 1)}\right)' \leq 0. \tag{2.7}$$

then, the solution  $u(x, y)$  of the Problem I is unique.

*Proof.* Known that if homogeneous problem has only trivial solution, then we can state that original problem has unique solution. For this aim, we assume that the Problem I has two solutions, then denoting difference of these as  $u(x, y)$ , we get

appropriate homogenous problem. Equation (2.5) we multiply to  $\tau(x)$  and integrate from 0 to 1:

$$\int_0^1 \tau''(x)\tau(x)dx - \Gamma(\alpha) \int_0^1 \lambda_1(x)\tau(x)\nu^-(x)dx \\ - \Gamma(\alpha) \int_0^1 \lambda_2(x)\tau(x)dx \int_x^1 r(t)\tau(t)dt + \Gamma(\beta) \int_0^1 \tau(x)p(x,0)D_{x1}^{-\beta}\tau(x)dx = 0.$$

We investigate the integral

$$I = \Gamma(\alpha) \int_0^1 \lambda_1(x)\tau(x)\nu^-(x)dx + \Gamma(\alpha) \int_0^1 \lambda_2(x)\tau(x)dx \int_x^1 r(t)\tau(t)dt \\ - \Gamma(\beta) \int_0^1 \tau(x)p(x,0)D_{x1}^{-\beta}\tau(x)dx.$$

Taking (2.3) into account at  $d(x) = 0$ , we obtain

$$I = \frac{\Gamma(\alpha)\Gamma(\gamma)}{2} \int_0^1 \frac{(1-x)q(x)}{2a(x)-1} \lambda_1(x)\tau(x)D_{x1}^{-\gamma}\tau(x)dx \\ + \Gamma(\alpha) \int_0^1 \frac{(1-2b(x))\lambda_1(x)}{2a(x)-1} \tau(x)\tau'(x)dx - 2\Gamma(\alpha) \int_0^1 \frac{\lambda_1(x)c(x)}{2a(x)-1} \tau^2(x)dx \\ + \Gamma(\alpha) \int_0^1 \lambda_2(x)\tau(x)dx \int_0^x r(t)\tau(t)dt - \Gamma(\beta) \int_0^1 \tau(x)p(x,0)D_{x1}^{-\beta}\tau(x)dx \\ = \frac{\Gamma(\alpha)}{2} \int_0^1 \frac{(1-x)q(x)}{2a(x)-1} \lambda_1(x)\tau(x)dx \int_x^1 (t-x)^{\gamma-1}\tau(t)dt \quad (2.8) \\ + \frac{\Gamma(\alpha)}{2} \int_0^1 \frac{1-2b(x)}{2a(x)-1} \lambda_1(x)d(\tau^2(x)) \\ - 2\Gamma(\alpha) \int_0^1 \frac{\lambda_1(x)c(x)}{2a(x)-1} \tau^2(x)dx - \frac{\Gamma(\alpha)}{2} \int_0^1 \frac{\lambda_2(x)}{r(x)} d\left(\int_x^1 r(t)\tau(t)dt\right)^2 \\ - \int_0^1 \tau(x)p(x,0)dx \int_x^1 (t-x)^{\beta-1}\tau(t)dt.$$

Considering  $\tau(1) = 0$ ,  $\tau(0) = 0$  (which deduced from the conditions (1.4), (1.5) in homogeneous case) and on a base of the formula [29]:

$$|x-t|^{-\gamma} = \frac{1}{\Gamma(\gamma)\cos\frac{\pi\gamma}{2}} \int_0^\infty z^{\gamma-1}\cos[z(x-t)]dz, \quad 0 < \gamma < 1$$

After some simplifications from (2.8) we obtain

$$I = \frac{\Gamma(\alpha)q(0)\lambda_1(0)}{4(2a(0)-1)\Gamma(1-\gamma)\sin\frac{\pi\gamma}{2}} \int_0^\infty z^{-\gamma}[M^2(0,z) + N^2(0,z)]dz \\ + \frac{\Gamma(\alpha)}{4\Gamma(1-\gamma)\sin\frac{\pi\gamma}{2}} \int_0^\infty z^{-\gamma}dz \int_0^1 [M^2(x,z) + N^2(x,z)]d\left[\lambda_1(x)\frac{(1-x)q(x)}{2a(x)-1}\right] \\ - \frac{\Gamma(\alpha)}{2} \int_0^1 \tau^2(x)\left(\lambda_1(x)\frac{1-2b(x)}{2a(x)-1}\right)' dx - 2\Gamma(\alpha) \int_0^1 \frac{\lambda_1(x)c(x)}{2a(x)-1} \tau^2(x)dx \\ + \frac{\Gamma(\alpha)}{2} \frac{\lambda_2(0)}{r(0)} \left(\int_0^1 r(t)\tau(t)dt\right)^2 + \frac{\Gamma(\alpha)}{2} \int_0^1 \left(\frac{\lambda_2(x)}{r(x)}\right)' \left(\int_x^1 r(t)\tau(t)dt\right)^2 dx$$

$$\begin{aligned}
 & - \frac{p(0, 0)}{2\Gamma(1 - \beta) \sin \frac{\pi\beta}{2}} \int_0^\infty z^{-\beta} [M^2(0, z) + N^2(0, z)] dz \\
 & - \frac{1}{2 \sin \frac{\pi\beta}{2} \Gamma(1 - \beta)} \int_0^\infty z^{-\beta} dz \int_0^1 \frac{\partial}{\partial x} [p(x, 0)] [M^2(x, z) + N^2(x, z)] dx \quad (2.9)
 \end{aligned}$$

where  $M(x, z) = \int_x^1 \tau(t) \cos zt dt$ ,  $N(x, z) = \int_x^1 \tau(t) \sin zt dt$ .

Thus, owing to (2.6),(2.7) from (2.9) it is concluded, that  $\tau(x) \equiv 0$ . Hence, based on the solution of the first boundary problem for the (1.1) [11, 25] owing to account (1.4) and (1.5) we obtain  $u(x, y) \equiv 0$  in  $\overline{\Omega}^+$ . Further, from functional relations (2.3), taking into account  $\tau(x) \equiv 0$  we obtain that  $\nu^-(x) \equiv 0$ . Consequently, based on the solution (2.2) we obtain  $u(x, y) \equiv 0$  in closed domain  $\overline{\Omega}^-$ .  $\square$

### 3. EXISTENCE OF SOLUTIONS FOR PROBLEM I

**Theorem 3.1.** *If conditions (2.6), (2.7) and*

$$p(x, y) \in C(\overline{\Omega}^+) \cap C^2(\Omega^+), \quad q(x + y) \in C(\overline{\Omega}^-) \cap C^2(\Omega^-); \quad (3.1)$$

$$\varphi(y), \psi(y) \in C(\overline{I_2}) \cap C^1(I_2); a(x), b(x), c(x), d(x) \in C^1(\overline{I_1}) \cap C^2(I_1) \quad (3.2)$$

hold, then the solution of the investigating problem exists.

*Proof.* Taking (2.3) into account from (2.5) we obtain

$$\tau''(x) - A(x)\tau'(x) = f(x) - B(x)\tau(x) \quad (3.3)$$

where

$$\begin{aligned}
 f(x) = & \frac{\Gamma(\alpha)\Gamma(\gamma)(1-x)\lambda_1(x)q(x)}{2(2a(x)-1)} D_{x1}^{-\gamma} \tau(x) - \Gamma(\beta)p(x, 0) D_{x1}^{-\beta} \tau(x) \\
 & + \Gamma(\alpha)\lambda_2(x) \int_x^1 r(t)\tau(t)dt - \frac{2\Gamma(\alpha)\lambda_1(x)d(x)}{2a(x)-1}
 \end{aligned} \quad (3.4)$$

$$A(x) = \frac{\Gamma(\alpha)\lambda_1(x)(1-2b(x))}{2a(x)-1}, \quad B(x) = \frac{2\Gamma(\alpha)\lambda_1(x)c(x)}{1-2a(x)} \quad (3.5)$$

The solution of (3.3) with conditions

$$\tau(0) = \psi(0), \quad \tau(1) = \varphi(0) \quad (3.6)$$

has the form

$$\begin{aligned}
 \tau(x) = & \psi(0) + A_1(x) \left( \int_x^1 (B(t)\tau(t) - f(t)) A'_1(t) dt + \frac{\varphi(0) - \psi(0)}{A_1(1)} \right) \\
 & - \frac{A_1(x)}{A_1(1)} \int_0^1 (B(t)\tau(t) - f(t)) \frac{A_1(t)}{A'_1(t)} dt \\
 & + \int_0^x (B(t)\tau(t) - f(t)) \frac{A_1(t)}{A'_1(t)} dt
 \end{aligned} \quad (3.7)$$

where

$$A_1(x) = \int_0^x \exp \left( \int_0^t A(z) dz \right) dt. \quad (3.8)$$

Further, considering (3.4) and using (1.3) from (3.7) we obtain

$$\tau(x)$$

$$\begin{aligned}
&= f_1(x) + A_1(x) \int_x^1 B(t)\tau(t)dA_1(t) \\
&\quad - \frac{\Gamma(\alpha)}{2} \int_x^1 \frac{(1-t)\lambda_1(t)q(t)}{2a(t)-1} dA_1(t) \int_t^1 \frac{\tau(s)ds}{(s-t)^{1-\gamma}} \\
&\quad + A_1(x) \int_x^1 A'_1(t)dt \int_t^1 [p(t,0)(s-t)^{\beta-1} - \Gamma(\alpha)\lambda_2(t)r(s)]\tau(s)ds \\
&\quad - \frac{A_1(x)}{A_1(1)} \int_0^1 \frac{A_1(t)}{A'_1(t)} B(t)\tau(t)dt + \int_0^x \frac{A_1(t)}{A'_1(t)} p(t,0)dt \int_t^1 (s-t)^{\beta-1}\tau(s)ds \\
&\quad + \Gamma(\alpha) \frac{A_1(x)}{A_1(1)} \int_0^1 \frac{A_1(t)}{A'_1(t)} dt \int_t^1 \left[ \frac{(1-t)\lambda_1(t)q(t)(s-t)^{\gamma-1}}{2(2a(t)-1)} - \lambda_2(t)r(s) \right] \tau(s)ds \\
&\quad - \frac{A_1(x)}{A_1(1)} \int_0^1 \frac{A_1(t)}{A'_1(t)} p(t,0)dt \int_t^1 (s-t)^{\beta-1}\tau(s)ds + \int_0^x \frac{A_1(t)}{A'_1(t)} B(t)\tau(t)dt \\
&\quad - \Gamma(\alpha) \int_0^x \frac{A_1(t)}{A'_1(t)} dt \int_t^1 \left[ \frac{(1-t)\lambda_1(t)q(t)}{2(2a(t)-1)} (s-t)^{\gamma-1} + \lambda_2(t)r(s) \right] \tau(s)ds \quad (3.9)
\end{aligned}$$

where

$$\begin{aligned}
&f_1(x) \\
&= \left(1 - \frac{A_1(x)}{A_1(1)}\right) \int_0^x \frac{2\Gamma(\alpha)d(t)A_1(t)\lambda_1(t)}{A'_1(t)(2a(t)-1)} dt \\
&\quad + 2\Gamma(\alpha)A_1(x) \int_x^1 \frac{d(t)A'_1(t)\lambda_1(t)}{2a(t)-1} dt \\
&\quad - \frac{A_1(x)}{A_1(1)} \int_x^1 \frac{2\Gamma(\alpha)d(t)A_1(t)\lambda_1(t)}{A'_1(t)(2a(t)-1)} dt - \frac{A_1(x)}{A_1(1)} (\psi(0) - \varphi(0)) + \psi(0). \quad (3.10)
\end{aligned}$$

After some simplifications we rewrite (3.9) in the form

$$\begin{aligned}
&\tau(x) \\
&= A_1(x) \int_x^1 \tau(s)ds \int_x^s [p(t,0)(s-t)^{\beta-1} - \Gamma(\alpha)\lambda_2(t)r(s)]A'_1(t)dt \\
&\quad - \Gamma(\alpha)A_1(x) \int_x^1 \tau(s)ds \int_x^s (s-t)^{\gamma-1} \frac{(1-t)\lambda(t)q(t)}{2(2a(t)-1)} A'_1(t)dt \\
&\quad - \Gamma(\alpha) \int_x^1 \tau(s)ds \int_0^x \left[ \frac{(1-t)\lambda_1(t)q(t)}{2(2a(t)-1)} (s-t)^{\gamma-1} + \lambda_2(t)r(s) \right] \frac{A_1(t)}{A'_1(t)} dt \\
&\quad + \int_x^1 \tau(s)ds \int_0^x (s-t)^{\beta-1} \frac{A_1(t)}{A'_1(t)} p(t,0)dt + A_1(x) \int_x^1 A'_1(t)B(t)\tau(t)dt \\
&\quad + \Gamma(\alpha) \frac{A_1(x)}{A_1(1)} \int_0^1 \tau(s)ds \int_0^s \left[ \frac{(1-t)\lambda_1(t)q(t)(s-t)^{\gamma-1}}{2(2a(t)-1)} - \lambda_2(t)r(s) \right] \frac{A_1(t)}{A'_1(t)} dt \\
&\quad - \frac{A_1(x)}{A_1(1)} \int_0^1 \tau(s)ds \int_0^s (s-t)^{\beta-1} \frac{A_1(t)}{A'_1(t)} p(t,0)dt - \frac{A_1(x)}{A_1(1)} \int_0^1 \frac{A_1(t)}{A'_1(t)} B(t)\tau(t)dt \\
&\quad + \int_0^x \tau(s)ds \int_0^s (s-t)^{\beta-1} \frac{A_1(t)}{A'_1(t)} p(t,0)dt + \int_0^x \frac{A_1(t)}{A'_1(t)} B(t)\tau(t)dt \\
&\quad - \Gamma(\alpha) \int_0^x \tau(s)ds \int_0^s \left[ \frac{(1-t)\lambda_1(t)q(t)}{2(2a(t)-1)} (s-t)^{\gamma-1} + \lambda_2(t)r(s) \right] \frac{A_1(t)}{A'_1(t)} dt + f_1(x)
\end{aligned}$$

i.e., we have the integral equation:

$$\tau(x) = \int_0^1 K(x, t)\tau(t)dt + f_1(x), \tag{3.11}$$

where

$$K(x, t) = \begin{cases} K_1(x, s), & 0 \leq t \leq x, \\ K_2(x, s), & x \leq t \leq 1. \end{cases} \tag{3.12}$$

$$\begin{aligned} K_1(x, s) = & \left(\frac{A_1(x)}{A_1(1)} - 1\right) \frac{\Gamma(\alpha)}{2} \int_0^s (s-t)^{\gamma-1} \frac{(1-t)A_1(t)\lambda(t)q(t)}{(2a(t)-1)A'_1(t)} dt \\ & - \Gamma(\alpha) \left(\frac{A_1(x)}{A_1(1)} + 1\right) r(s) \int_0^s \lambda_2(t) \frac{A_1(t)}{A'_1(t)} dt \\ & - \left(\frac{A_1(x)}{A_1(1)} - 1\right) \left[ \int_0^s (s-t)^{\beta-1} \frac{A_1(t)}{A'_1(t)} p(t, 0) dt + \frac{A_1(s)}{A'_1(s)} B(s) \right] \end{aligned} \tag{3.13}$$

$$\begin{aligned} & K_2(x, s) \\ = & A_1(x) \left( A'_1(s) B(s) - \frac{\Gamma(\alpha)}{2} \int_x^s (s-t)^{\gamma-1} \frac{(1-t)\lambda(t)q(t)}{2a(t)-1} A'_1(t) dt \right) \\ & - \Gamma(\alpha) A_1(x) \int_x^s \lambda_2(t) A'_1(t) dt + A_1(x) \int_x^s (s-t)^{\beta-1} A'_1(t) p(t, 0) dt \\ & - \Gamma(\alpha) r(s) \int_0^x \frac{A_1(t)\lambda_2(t)}{A'_1(t)} dt \\ & + \frac{\Gamma(\alpha)}{2} \frac{A_1(x)}{A_1(1)} \int_0^s (s-t)^{\gamma-1} \frac{(1-t)A_1(t)\lambda(t)q(t)}{(2a(t)-1)A'_1(t)} dt \\ & - \frac{A_1(x)}{A_1(1)} \frac{A_1(s)}{A'_1(s)} B(s) - \Gamma(\alpha) \frac{A_1(x)}{A_1(1)} r(s) \int_0^s \frac{A_1(t)\lambda_2(t)}{A'_1(t)} dt \\ & - \frac{\Gamma(\alpha)}{2} \int_0^x \frac{(1-t)A_1(t)\lambda(t)q(t)}{A'_1(t)(2a(t)-1)} (s-t)^{\gamma-1} dt \\ & + \left(1 - \frac{A_1(x)}{A_1(1)}\right) \int_0^x \frac{A_1(t)(s-t)^{\beta-1}}{A'_1(t)} p(t, 0) dt. \end{aligned} \tag{3.14}$$

Owing to class (3.1), (3.2) of the given functions and after some evaluations from (3.13), (3.14) and (3.10), (3.12) we conclude that

$$|K(x, t)| \leq \text{const}, \quad |f_1(x)| \leq \text{const}.$$

Since kernel  $K(x, t)$  is continuous and function in right-side  $F(x)$  is continuously differentiable, solution of integral equation (3.11) we can write via resolvent-kernel:

$$\tau(x) = f_1(x) - \int_0^1 \mathfrak{R}(x, t)f_1(t)dt, \tag{3.15}$$

where  $\mathfrak{R}(x, t)$  is the resolvent-kernel of  $K(x, t)$ . Unknown functions  $\nu^-(x)$  and  $\nu^+(x)$  we found accordingly from (2.3) and (2.4):

$$\begin{aligned} \nu^-(x) = & \frac{(x-1)q(x)}{2(2a(x)-1)} \int_x^1 (t-x)^{\gamma-1} dt \int_0^1 \mathfrak{R}(t, s)f_1(s)ds \\ & + \frac{(1-x)q(x)}{2(2a(x)-1)} \int_x^1 (t-x)^{\gamma-1} f_1(t)dt \end{aligned}$$

$$\begin{aligned}
& + \frac{1-2b(x)}{2a(x)-1} f_1'(x) - \frac{1-2b(x)}{2a(x)-1} \int_0^1 \frac{\partial \mathfrak{R}(x,t)}{\partial x} f_1(t) dt - \frac{2c(x)}{2a(x)-1} f_1(x) \\
& + \frac{2c(x)}{2a(x)-1} \int_0^1 \mathfrak{R}(x,t) f_1(t) dt - \frac{2d(x)}{2a(x)-1}
\end{aligned}$$

and  $\nu^+(x) = \lambda(x)\nu^-(x)$ .

The solution of Problem I in the domain  $\Omega^+$  can be written as [11],

$$\begin{aligned}
& u(x,y) \\
& = \int_0^y G_\xi(x,y,0,\eta)\psi(\eta)d\eta - \int_0^y G_\xi(x,y,1,\eta)\varphi(\eta)d\eta \\
& + \int_0^1 G_0(x-\xi,y)\tau(\xi)d\xi - \int_0^y \int_0^1 G(x,y,0,\eta)p(\xi)d\xi d\eta \int_\xi^1 (t-\xi)^{\beta-1}\tau(t)dt
\end{aligned}$$

where

$$\begin{aligned}
G_0(x-\xi,y) & = \frac{1}{\Gamma(1-\alpha)} \int_0^y \eta^{-\alpha} G(x,y,\xi,\eta) d\eta, \\
G(x,y,\xi,\eta) & = \frac{(y-\eta)^{\alpha/2-1}}{2} \sum_{n=-\infty}^{\infty} \left[ e_{1,\alpha/2}^{1,\alpha/2} \left( -\frac{|x-\xi+2n|}{(y-\eta)^{\alpha/2}} \right) \right. \\
& \quad \left. - e_{1,\alpha/2}^{1,\alpha/2} \left( -\frac{|x+\xi+2n|}{(y-\eta)^{\alpha/2}} \right) \right]
\end{aligned}$$

Is the Green's function of the first boundary problem (1.1) in the domain  $\Omega^+$  with the Riemann-Liouville fractional differential operator instead of the Caputo ones [26],

$$e_{1,\delta}^{1,\delta}(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!\Gamma(\delta - \delta n)}$$

is the Wright type function [26]. Solution of the Problem I in the domain  $\Omega^-$  will be found by the formula (2.2). Hence, the proof is complete.  $\square$

We remark that if  $a(x) = 1/2$ , then from (2.3) we find  $\tau(x)$  as a solution Volterra type integral equation. After that we can find  $\nu^+(x)$  from the first boundary value problem for the (1.1), and  $\nu^-(x)$  will be defined from the gluing condition (2.4).

#### REFERENCES

- [1] O. Kh. Abdullaev; About a method of research of the non-local problem for the loaded mixed type equation in double-connected domain, *Bulletin KRASEC. Phys. Math. Sci.*, 2014, vol. 9, no. 2, pp. 11-16. ISSN 2313-0156
- [2] Abdumavlen S. Berdyshev, Erkinjon T. Karimov, Nazgul S. Akhtaeva; On a boundary-value problem for the parabolic-hyperbolic equation with the fractional derivative and the sewing condition of the integral form. *AIP Proceedings*, Vol. 1611, 2014, pp.133-137.
- [3] E. Y. Arlanova; A problem with a shift for the mixed type equation with the generalized operators of fractional integration and differentiation in a boundary condition, *Vestnik Samarsk Gosudarstvennogo Universiteta.* (2008) vol. 6, no. 65, pp. 396-406.
- [4] A. Belarbi, M. Benchohra, A. Ouahab; Uniqueness results for fractional functional differential equations with infinite delay in Frechet spaces. *Appl. Anal.* 85(2006), No. 12, 1459-1470.
- [5] M. Benchohra, J. Henderson, S. K. Ntouyas, A. Ouahab; Existence results for fractional order functional differential equations with infinite delay. *J. Math. Anal. Appl.* 338(2008), No. 2, 1340-1350.

- [6] K. Diethelm, A. D. Freed; On the solution of nonlinear fractional order differential equations used in the modeling of viscoelasticity, in: F. Keil, W. Mackens, H. Voss, J. Werther (Eds.), *Scientific Computing in Chemical Engineering II Computational Fluid Dynamics, Reaction Engineering and Molecular Properties*, Springer-Verlag, Heidelberg.(1999), pp. 217–224.
- [7] W. G. Glockle, T. F. Nonnenmacher; A fractional calculus approach of self-similar protein dynamics, *Biophys. J.* 68 (1995) 46–53.
- [8] R. Hilfer; *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, (2000).
- [9] B. Islomov, U. Baltaeva; Boudanry-value problems for a third-order loaded parabolic-hyperbolic type equation with variable coefficients. *Electron. J. Differential Equ.*, Vol. 2015 (2015). No. 221. pp 1-10.
- [10] B. J. Kadirkulov; Boundary problems for mixed parabolic-hyperbolic equations with two lines of changing type and fractional derivative. *Electron. J. Differential Equ.*, Vol. 2014 (2014), No. 57, pp. 1–7.
- [11] E. T. Karimov, J. Akhatov; A boundary problem with integral gluing condition for a parabolic-hyperbolic equation involving the Caputo fractional derivative. *Electron. J. Differential Equ.*, Vol. 2014 (2014) No. 14. pp. 1–6.
- [12] A. A. Kilbas, O. A. Repin; Analogue of the Bitsadze-Samarskiy problem for an equation of mixed type with a fractional derivative; *Differentsialnye Uravneniya*. (2003) vol. 39, no. 5, pp. 638–719, *Translation in Journal of Difference Equations and Applications*. (2003). vol. 39, no. 5, pp. 674–680.
- [13] A. A. Kilbas, O. A. Repin; An analog of the Tricomi problem for a mixed type equation with a partial fractional derivative, *Fractional Calculus and Applied Analysis*. (2010) vol. 13, no. 1, pp. 69–84.
- [14] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo; *Theory and Applications of Fractional Differential Equations*, in: North-Holland Mathematics Studies, vol. 204, Elsevier Science B.V., Amsterdam. (2006).
- [15] J. W. Kirchner, X. Feng, C. Neal; Fractal streamchemistry and its implications for contaminant transport in catchments, *Nature* 403 (2000), 524–526.
- [16] V. Lakshmikantham, A. S. Vatsala; Basic theory of fractional differential equations. *Nonlinear Anal.* 69(2008), No. 8, 2677–2682.
- [17] V. Lakshmikantham, A. S. Vatsala; Theory of fractional differential inequalities and applications. *Commun. Appl. Anal.* 11(2007), No. 3-4, 395-402.
- [18] B. N. Lundstrom, M. H. Higgs, W. J. Spain, A. L. Fairhall; Fractional differentiation by neocortical pyramidal neurons, *Nat. Neurosci.* 11 (2008), 1335–1342.
- [19] F. Mainardi; Fractional calculus: some basic problems in continuum and statistical mechanics, in: A. Carpinteri, F. Mainardi (Eds.), *Fractals and Fractional Calculus in Continuum Mechanics*, Springer-Verlag, Wien.(1997), pp. 291-348.
- [20] K. S. Miller, B. Ross; *An Introduction to the Fractional Calculus and Differential Equations*, John Wiley, New York, (1993).
- [21] A. M. Nakhushiev; The loaded equations and their applications. *M. Nauka*, (2012). 232.
- [22] V. A. Nakhushieva; Boundary problems for mixed type heat equation, *Doklady AMAN* (2010) vol. 12, no. 2, pp. 3944. In Russian.
- [23] I. Podlubny; *Fractional Differential Equations*, Academic Press, New York, (1999).
- [24] I. Podlubny; Geometric and physical interpretation of fractional integration and fractional differentiation. *Dedicated to the 60th anniversary of Prof. Francesco Mainardi. Fract. Calc. Appl. Anal.* 5(2002), No. 4, 367-386.
- [25] A. V. Pskhu; Uravneniye v chasnykh proizvodnykh drobnogo poryadka. (in Russian) [Partial differential equation of fractional order] *Nauka*, Moscow, (2005). 200 pp.
- [26] A. V. Pskhu; Solution of boundary value problems fractional diffusion equation by the Green function method. *Differential equation*, 39(10) (2003), pp. 1509-1513
- [27] M. S. Salakhitdinov, E. T. Karimov; On a nonlocal problem with gluing condition of integral form for parabolic-hyperbolic equation with Caputo operator. *Reports of the Academy of Sciences of the Republic of Uzbekistan. (DAN RUz)*, No 4, (2014), 6-9.
- [28] S. G. Samko, A. A. Kilbas, O. I. Marichev; *Fractional Integral and Derivatives: Theory and Applications*, Gordon and Breach, Longhorne, PA, (1993).
- [29] M. M. Smirnov; *Mixed type equations*, M. Nauka. (2000).

OBIDJON KH. ABDULLAEV  
DEPARTMENT OF DIFFERENTIAL EQUATIONS AND MATHEMATICAL PHYSICS, NATIONAL UNIVERSITY  
OF UZBEKISTAN, 100114 UZBEKISTAN, TASHKENT, UZBEKISTAN  
*E-mail address:* `obidjon.mth@gmail.com`

KISHIN S. SADARANGANI  
DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LAS-PALMAS DE GRAN CANARIA, 35017 LAS PAL-  
MAS, SPAIN  
*E-mail address:* `ksadaran@dma.ulpgc.es`