

Processes with Inert Drift

David White
Dept. of Mathematics and Computer Science
Belmont University
Nashville TN 37212
white@u.washington.edu

Abstract

We construct a stochastic process whose drift is a function of the process's local time at a reflecting barrier. The process arose as a model of the interactions of a Brownian particle and an inert particle in [7]. We construct and give asymptotic results for two different arrangements of inert particles and Brownian particles, and construct the analogous process in \mathbb{R}^d .

Key words: Brownian motion; local time; Skorohod lemma.

AMS 2000 Subject Classification: Primary 60J6; Secondary: 60J55.

Submitted to EJP on April 5, 2007, final version accepted November 14, 2007.

1 Introduction

In this paper, we construct solutions (X, L) to the set of equations

$$\begin{aligned} X(t) &= B(t) + L(t) + \int_0^t \mu \circ L(s) \, ds \\ X(t) &\geq 0 \\ L(t) &= \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \int_0^t 1_{[0, \epsilon)} X(s) \, ds \end{aligned}$$

for certain monotone functions $\mu(l)$, where $B(t)$ is a standard Brownian motion with $B(0)=0$. The case when $\mu(l) = Kl$, $K > 0$ was studied by Frank Knight in [7], which was the starting point for this research.

The case when $\mu(l) = 0$ is the classical reflected Brownian motion. There are at least two approaches to constructing pathwise solutions in this case. The first is to let $X(t) = |B(t)|$, define $L(t)$ from $X(t)$ as above, and define a new Brownian motion $\tilde{B}(t) = X(t) - L(t)$. For the second approach we define

$$L(t) = -\inf_{s < t} B(s),$$

and take $X(t) = B(t) + L(t)$. The two approaches yield processes (X, L) with the same distribution. The second approach has the advantages of being easier to work with, and of retaining the original $B(t)$.

For other $\mu(l)$, $X(t)$ can be written as the difference of $B(t) + L(t)$ and $-\int_0^t \mu \circ L(s) \, ds$. These two pieces can be interpreted the path of a Brownian particle reflecting from the path $-\int_0^t \mu \circ L(s) \, ds$ of an inert particle. We call the second path inert because its derivative is constant when the two particles are apart. With this model in mind, we consider two other configurations of Brownian particles and inert particles. We also consider a generalization to \mathbb{R}^d .

In Section 2, we consider the one-dimensional case. We also obtain an explicit formula for the distribution of $\lim_{t \rightarrow \infty} \mu \circ L(t)$ and for the excursion measure of the process with drift.

In Section 3, we consider the case when a Brownian particle and an inert particle are confined to a one-dimensional ring. In other words, we have a process with drift confined to a closed interval, with reflection at both endpoints. The velocity of the inert particle turns out to be an interesting process. Indexed by local time, it is a piecewise linear process with Gaussian stationary distribution, though it is not a Gaussian process. Some results from this section have appeared in [3]. We show that under rescaling, the velocity process converges in distribution to an Ornstein-Uhlenbeck process.

Section 4 discusses the configuration consisting of two independent Brownian particles with an inert particle separating them. We would like to determine whether the process is recurrent or transient; that is, whether all three particles can meet in finite time, or whether the inert particle in the middle forces the distance between the Brownian particles to tend to infinity. In fact, this configuration is a critical case between the two behaviors. We show that under rescaling the distance between the two Brownian particles is a two-dimensional Bessel process. Dimension two is exactly the critical case for the family of Bessel processes.

Section 5 partially extends the results in Section 2 to domains in \mathbb{R}^d , $d \geq 2$. This sections uses results of Lions and Sznitman on the existence of reflected Brownian motion in domains in \mathbb{R}^d .

We show existence for the process with drift for the case when the velocity gained is proportional to the local time.

Some results similar to those in Section 2 have been found by S. Ramasubramanian in [9]. He in fact considers a slightly more general class of allowable μ , and extends the process to orthants in \mathbb{R}^d .

The author would like to thank the referees for their very helpful suggestions, and also Kavita Ramanan and one of the referees for the reference to [9].

This paper is partly based on a Ph.D. thesis written at the University of Washington under the guidance of Chris Burdzy.

2 Existence and Uniqueness in One Dimension

In this section, we prove the following version of the well-known Skorohod Lemma:

Theorem 2.1. *Let $f(t)$ be a continuous function with $f(0) \geq 0$, and let $\mu(l)$ be a continuous monotone function with*

$$\lambda(l) = \sup_{a < b < l} \frac{|\mu(b) - \mu(a)|}{b - a} < \infty \quad (1)$$

for every l . If $\mu(l) \rightarrow -\infty$, then we further require that

$$\sum_n (|\mu(n)| \vee 1)^{-1} = \infty. \quad (2)$$

There is a unique continuous $L(t)$ satisfying

1. $x(t) = f(t) + L(t) + \int_0^t \mu \circ L(s) ds \geq 0$,
2. $L(0) = 0$, $L(t)$ is nondecreasing,
3. $L(t)$ is flat off $\{t : x(t) = 0\}$; i.e., $\int_0^\infty 1_{\{x(s) > 0\}} dL(s) = 0$.

As a reminder to the reader, we quote the Skorohod Lemma. A proof can be found in, for example, [6].

Theorem 2.2 (The Skorohod Lemma). *Given a continuous function $f(t)$ with $f(0) \geq 0$, there is a unique continuous function $L(t)$ satisfying*

1. $x(t) \equiv f(t) + L(t) \geq 0$,
2. $L(0) = 0$, $L(t)$ is nondecreasing, and
3. $L(t)$ is flat off $\{t : x(t) = 0\}$; i.e., $\int_0^\infty 1_{\{x(s) > 0\}} dL(s) = 0$.

This function is given by

$$L(t) = \max \left[0, \max_{0 \leq s \leq t} (-f(s)) \right] \quad (3)$$

We will denote the unique $L(t)$ corresponding to $f(t)$ in equation (3) by $Lf(t)$.

A few lemmata and corollaries of the Skorohod equation will be used in the proof of Theorem 2.1. We begin with these.

Lemma 2.3. *If $f(t)$ and $g(t)$ are two continuous functions with $f(0) \geq 0$, and $f(t) \leq g(t)$ for $t \in [0, \infty)$, then $Lf(t) \geq Lg(t)$, $0 \leq t < \infty$. In particular, if $\phi(t) \geq 0$ then $L(f + \phi)(t) \leq Lf(t)$.*

Proof. This comes directly from equation (3). □

Lemma 2.4. *For continuous $f(t)$, $f(0) \geq 0$, and $S < T$, we have*

$$Lf(T) - Lf(S) \leq \max_{S \leq r \leq T} [f(S) - f(r)].$$

Proof. Using equation (3), we have

$$\begin{aligned} Lf(T) - Lf(S) &= \max_{S \leq r \leq T} [-f(r) - Lf(S)] \vee 0 \\ &\leq \max_{S \leq r \leq T} [f(S) - f(r)] \vee 0 \\ &= \max_{S \leq r \leq T} [f(S) - f(r)]. \end{aligned}$$

and the claim follows. □

Lemma 2.5. *If $f(t)$ and $g(t)$ are two continuous functions with $f(0) \geq 0$, and $f(t) = g(t)$ for $0 \leq t \leq T$, then $Lf(t) = Lg(t)$ for $0 \leq t \leq T$.*

Proof. This also comes directly from (3). □

Now we will prove Theorem 2.1.

Proof of Theorem 2.1. Uniqueness will be proved first. Assume that both $L(t)$ and $\tilde{L}(t)$ satisfy conditions 1–3 of Theorem 2.1. Let $Q = \inf\{t > 0 : L(t) \neq \tilde{L}(t)\}$, and suppose that $Q < \infty$. Define the function $M(t) = \lambda(L(t) \vee \tilde{L}(t))(t - Q)$. As the product of increasing functions, $M(t)$ is increasing, and $M(Q) = 0$. Choose $R > Q$ so that $M(R) < 1$.

By the continuity of $L(t)$ and $\tilde{L}(t)$, there exist T with $Q < T < R$, and $\delta > 0$, with the property that $|L(T) - \tilde{L}(T)| = \delta$, while $|L(t) - \tilde{L}(t)| < \delta$ for $t < T$. We will show that the case $L(T) - \tilde{L}(T) = \delta$ yields a contradiction; the same argument works for the other case, $\tilde{L}(T) - L(T) = \delta$.

Suppose that $L(T) - \tilde{L}(T) = \delta$ and $L(t) - \tilde{L}(t) < \delta$ for $t < T$. We have that

$$\begin{aligned} x(T) - \tilde{x}(T) &= (x(T) - x(Q)) - (\tilde{x}(T) - \tilde{x}(Q)) \\ &= [L(T) - L(Q) + \int_Q^T \mu \circ L(r) dr] \\ &\quad - [\tilde{L}(T) - \tilde{L}(Q) + \int_Q^T \mu \circ \tilde{L}(r) dr] \\ &= (L(T) - \tilde{L}(T)) + \int_Q^T (\mu \circ L(r) - \mu \circ \tilde{L}(r)) dr, \end{aligned}$$

and that

$$\begin{aligned} \left| \int_Q^T (\mu \circ L(r) - \mu \circ \tilde{L}(r)) dr \right| &\leq (T - Q) \lambda(L(T) \vee \tilde{L}(T)) \sup_{Q \leq t \leq T} (L(t) - \tilde{L}(t)) \\ &= M(T)(L(T) - \tilde{L}(T)). \end{aligned}$$

Combining these two equations we see that

$$x(T) - \tilde{x}(T) \geq (1 - M(T))(L(T) - \tilde{L}(T)) > 0.$$

Let $S = \sup\{t < T : x(t) = \tilde{x}(t)\}$. Then for $S < t \leq T$, $x(t) > \tilde{x}(t) \geq 0$, and by condition 3 of the theorem, $L(t) = L(S)$, and in particular, $L(T) = L(S)$. Then for $S < t \leq T$, $L(t) - \tilde{L}(t) = L(T) - \tilde{L}(t)$. Since $\tilde{L}(t)$ is nondecreasing by condition 2 of the theorem, $L(T) - \tilde{L}(t)$ is nonincreasing in t , and we see that $L(S) - \tilde{L}(S) \geq L(T) - \tilde{L}(T) = \delta$. This contradicts how T was chosen. Therefore, $L(t) = \tilde{L}(t)$ for all $t \geq 0$.

Next we will construct the $L(t)$ corresponding to $f(t)$ and $\mu(t)$. First, we may assume that $\mu(t) \geq 0$. Otherwise, set $m = \inf\{\mu(t), t \geq 0\}$, $\tilde{f}(t) = f(t) + mt$ and $\tilde{\mu}(t) = \mu(t) - m$, assuming for the moment that $m > -\infty$. It is easily checked that the $L(t)$ satisfying the theorem for $\tilde{f}(t)$ and $\tilde{\mu}(t)$ also works for $f(t)$ and $\mu(t)$.

For any $\varepsilon > 0$ we construct $L^\varepsilon(t)$ and $I^\varepsilon(t)$ so that $L^\varepsilon(t) = L(f + I^\varepsilon)(t)$, and $\frac{d}{dt}I^\varepsilon(t) = \mu(n\varepsilon)$, $n\varepsilon \leq L^\varepsilon(t) \leq (n+1)\varepsilon$. The construction is recursive. Let $I_0^\varepsilon = 0$, $L_0^\varepsilon(t) = Lf(t)$, and $T_0^\varepsilon = \inf\{t > 0 : L_0^\varepsilon(t) = \varepsilon\}$. Now for each positive integer n , define

$$\begin{aligned} I_{n+1}^\varepsilon(t) &= \begin{cases} I_n^\varepsilon(t), & t < T_n^\varepsilon, \\ I_n^\varepsilon(T_n^\varepsilon) + \mu(n\varepsilon)(t - T_n^\varepsilon), & T_n^\varepsilon \leq t \leq T_{n+1}^\varepsilon, \\ I_{n+1}^\varepsilon(T_{n+1}^\varepsilon), & T_{n+1}^\varepsilon < t, \end{cases} \quad (4) \\ L_{n+1}^\varepsilon(t) &= L(f + \sum_{j=1}^{n+1} I_j^\varepsilon)(t), \\ T_{n+1}^\varepsilon &= \inf\{t > 0 : L_n^\varepsilon(t) = (n+1)\varepsilon\}. \end{aligned}$$

For $m \geq n$, we have that $I_m^\varepsilon(t) \geq I_n^\varepsilon(t)$, so by Lemma 2.3, $L_m^\varepsilon(t) \leq L_n^\varepsilon(t)$, and from equation (3), $L_m^\varepsilon(t) = L_n^\varepsilon(t)$ for $t < T_{m \wedge n}^\varepsilon$. Let $L^\varepsilon(t) = \lim_{n \rightarrow \infty} L_n^\varepsilon(t)$, and $I^\varepsilon(t) = \lim_{n \rightarrow \infty} I_n^\varepsilon(t)$.

By Lemma 2.4, we see that

$$\begin{aligned} L^\varepsilon(t) - L^\varepsilon(s) &= L(f + I^\varepsilon)(t) - L(f + I^\varepsilon)(s) \\ &\leq \max_{s \leq r \leq t} \{(f + I^\varepsilon)(s) - (f + I^\varepsilon)(r)\} \\ &\leq \max_{s \leq r \leq t} \{f(s) - f(r)\}, \end{aligned}$$

that is, the family $\{L^\varepsilon(t)\}_{\varepsilon > 0}$ is equicontinuous. Since $I^\varepsilon(t) \geq 0$, by Lemma 2.3 $\{L^\varepsilon(t)\}_{\varepsilon > 0}$ is bounded for each t . Ascoli–Arzelà [10] then applies. For any sequence $\varepsilon(n) \rightarrow 0$, $L^{\varepsilon(n)}(t) \rightarrow L(t)$, uniformly on compact subsets of $[0, \infty)$. We will check that this $L(t)$ satisfies the conditions of the theorem. That $x(t) \geq 0$ follows from the definition of $L_n^\varepsilon(t)$ and equation (3), as does

the second condition. That $I^{\varepsilon_n}(t) \rightarrow \int_0^t \mu \circ L(s) ds$ follows from uniform convergence and the inequality

$$\int_0^t \mu \circ L^\varepsilon(t) ds - I^\varepsilon(t) \leq \varepsilon \lambda(t).$$

For condition 3, notice that $\{t : x(t) > \delta\} \subset \{t : x^{\varepsilon(n)}(t) > \delta/2\}$ for large enough n . The uniqueness of $L(t)$ proved above shows that $L(t)$ is independent of the $\varepsilon(n)$ chosen.

For the case where $m = \inf\{\mu(l), l \geq 0\} = -\infty$, repeat the above construction with $\mu^j(s) = \mu(s) \vee (-j)$, and denote the results as $L^j(t)$ and $x^j(t)$. By Lemma 2.5, $L^j(t)$ will agree with $L(t)$ for $0 \leq T^j$, where $T^j = \inf\{t : \mu \circ L(t) = -j\}$. To complete the proof, it is only necessary to show that $T^j \rightarrow \infty$. Suppose that $T^j \rightarrow T$. There are then $T_n \uparrow T$ so that $L(T_n) = n$ and $x(T_n) = 0$. We compute that

$$f(T_n) - f(T_{n+1}) = 1 + \int_{T_n}^{T_{n+1}} \mu \circ L(s) ds.$$

By the continuity of $f(t)$, the LHS approaches 0, so that

$$\left| \int_{T_n}^{T_{n+1}} \mu \circ L(s) ds \right| \rightarrow 1.$$

However,

$$\left| \int_{T_n}^{T_{n+1}} \mu \circ L(s) ds \right| \leq (T_{n+1} - T_n) \sup_{n \leq l \leq n+1} |\mu(l)|,$$

so that

$$T_{n+1} - T_n \geq \frac{0.9}{|\mu(n+1)|}$$

for sufficiently large n . Then $T = T_1 + \sum(T_{n+1} - T_n) \geq \infty$ by (2), a contradiction. Hence $T^j \rightarrow \infty$, and the proof is complete. \square

The following corollary restates Theorem 2.1 to be similar to Knight's original result [7, Theorem 1.1]. Theorem 2.1 yields a process with drift reflected at 0, this version yields a process reflected from a second curve $v(t)$.

Corollary 2.6. *Let $f(t)$ be a continuous function with $f(0) \geq 0$, and let $\mu(l)$ be a continuous increasing function with*

$$\lambda(l) = \sup_{a < b < l} \frac{|\mu(b) - \mu(a)|}{b - a} < \infty$$

for every $l \geq 0$. Then there is a unique continuous $L(t)$ satisfying

1. $\tilde{x}(t) = f(t) + L(t)$,
2. $L(0) = 0$, $L(t)$ is nondecreasing,

3. $L(t)$ is flat off $\{t : \tilde{x}(t) = v(t)\}$, and

4. $v(0) = 0$, $\frac{d}{dt}v(t) = -\mu \circ L(t)$.

Proof. This is just a restatement of Theorem 2.1. To see this, let

$$v(t) = - \int_0^t \mu \circ L(s) ds$$

and

$$\tilde{x}(t) = x(t) + v(t).$$

□




Figure 1: Two equivalent versions of the process, corresponding to Theorem 2.1 and Corollary 2.6, resp. Note that τ_∞ cannot be determined from the pictured portion of the graph, so the best candidate is what is labeled.

The remarks that follow demonstrate the necessity of the restrictions (1) and (2) on $\mu(t)$ in Theorem 2.1, at least for the deterministic version of the theorem.

Remark 1 (Non-uniqueness of $L(t)$). In the context of Theorem 2.1, if we let $f(t) = -t$, $\mu(l) = 1 - \sqrt{l}$, $L(t) = 0$, and $\tilde{L}(t) = t^2/4$, then $x(t) = \tilde{x}(t) = 0$, which shows that the theorem cannot be extended to general $f(t)$ and $\mu(l)$.

Remark 2 (Blow-up of $L(t)$). Again, in the context of Theorem 2.1, we let $f(t) = -t$, $\mu(l) = -l^2$, and $L(t) = \tan t$. Then

$$x(t) = -t + \tan t - \int_0^t \tan^2 s \, ds = 0,$$

so $L(t)$ satisfies the conditions of the theorem, but blows up at $t = \pi/2$.

The results so far do not rely on probability. We will now remedy that by applying the results to Brownian paths. For the rest of the section, we will need a standard Brownian motion $\{B_t, \mathcal{F}_t\}$, with $B_0 = 0$ a.s., and some fixed $\mu(t)$ satisfying (1). In the statements below, $L(t)$ will be a random function, $L(\omega, t) = LB(\omega, t)$.

Theorem 2.7. *If $f(t)$ in Theorem 2.1 is replaced with a Brownian motion B_t , then the corresponding $L(t)$ is the semimartingale local time at zero, $\Lambda_t(0)$, [6, p. 218] of $X_t \equiv B_t + L(t) + \int_0^t \mu \circ L(s) ds$, a.s.*

Further, we get that, a.s.,

$$2L(t) = \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} \int_0^t \mathbf{1}_{[0, \varepsilon)} \left(B_s + L(s) + \int_0^s \mu \circ L(r) dr \right) ds.$$

Proof. To see that $L(t) = \Lambda_t(0)$, use a version of Itô-Tanaka's formula (e.g. (7.4) on p. 218 of [6]), with the identifications $M_t = B_t$, $V_t = L(t) + \int_0^t \mu \circ L(s) ds$, and $f(x) = |x|$, and get that

$$\begin{aligned} |X_t| &= X_0 - \int_0^t \mathbf{1}_{\{0\}}(X_s) dB_s + \int_0^t \mathbf{1}_{(0, \infty)}(X_s) dB_s - \int_0^t \mathbf{1}_{\{0\}}(X_s) dL(s) \\ &\quad + \int_0^t \mu \circ L(s) ds + 2\Lambda_t(0) \\ &= B_t - L(t) + \int_0^t \mu \circ L(s) ds + 2\Lambda_t(0). \end{aligned}$$

On the other hand, if $f(t)$ is replaced by B_t in Theorem 2.1, we get since $X_t \geq 0$

$$|X_t| = B_t + L(t) + \int_0^t \mu \circ L(s) ds.$$

Therefore, $L(t) = \Lambda_t(0)$, a.s.

Having established the previous fact, a standard argument shows the other assertion. By [6, Theorem 7.1] we get

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} \int_0^t \mathbf{1}_{[0, \varepsilon)} \left(B_s + L(s) + \int_0^s \mu \circ L(r) dr \right) ds &= \lim_{\varepsilon \rightarrow 0} \frac{2}{\varepsilon} \int_0^\varepsilon \Lambda_t(a) da \\ &= 2\Lambda_t(0) = 2L(t). \end{aligned}$$

□

Corollary 2.8. For the case $\mu \equiv \lambda$, that is, Brownian motion with drift λ , reflected at 0, $X_t \equiv B_t + L(t) + \lambda t$,

$$L_X(t) \equiv L(t) = \max_{0 \leq s \leq t} (-B_t - \lambda t).$$

Proof. This follows directly from the previous theorem and equation (3) on page 1511. By the notation $L_X(t)$ we will mean the semimartingale local time at 0 of the reflected process $X(t)$ (not just for the specific case of constant drift). Note that in the stochastic case, unlike the deterministic case, the $L_X(t)$ can be recovered almost surely from $X(t)$. \square

We now fix λ , and define $X(t) \equiv B(t) + L_X(t) + \lambda t$ as above. Let ϕ be the right-continuous inverse of $L_X(t)$; that is,

$$\phi(b) = \inf\{t : L_X(t) = b\}.$$

By Corollary 2.8, we have that

$$\phi(b) \stackrel{d}{=} T_b,$$

where $T_b = \inf\{t : -B_t - \lambda t = b\}$. From [6, p. 196] we have that

$$P[T_b - T_a \in dt] = \frac{b-a}{\sqrt{2\pi t^3}} \exp\left[-\frac{(b-a)^2}{2t}\right] dt,$$

and a computation shows that for $\mu \equiv \lambda$,

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} P[T_{b+\varepsilon} - T_b \in dt] = \frac{dt}{\sqrt{2\pi t^3}} \exp\left[-\frac{\lambda^2 t}{2}\right].$$

Additionally, we have for $b > 0$ that

$$\begin{aligned} P\{T_b < \infty\} &= \begin{cases} 1, & \lambda \geq 0, \\ e^{-2|\lambda|b}, & \lambda < 0, \end{cases} \\ &= e^{-2|\lambda \wedge 0|b}. \end{aligned} \tag{5}$$

Theorem 2.9. Let $\tau_\infty = \inf\{s : \phi(s) = \infty\}$. Then

$$P\{\tau_\infty > \tau\} = \exp\left(-2 \int_0^\tau |\mu(s) \wedge 0| ds\right),$$

with $\mu(l)$ as in Theorem 2.1.

Proof. An equivalent definition of τ_∞ is $\tau_\infty = \sup\{L_X(t) : t \geq 0\}$. Recall from the proof of Theorem 2.1 the definition of $L^\varepsilon(t)$, and let $\tau_\infty^\varepsilon = \sup\{L^\varepsilon(t) : t \geq 0\}$. It is clear from the uniform convergence $L^\varepsilon(t)$ to $L_X(t)$ that $\tau_\infty = \lim_{\varepsilon \rightarrow 0} \tau_\infty^\varepsilon$.

For notational convenience, define $\underline{\tau} = \sup\{j\varepsilon : j\varepsilon \leq \tau, j \text{ an integer}\}$, where the value of ε should be apparent from the context. From (4) and (5) we see that

$$P(\tau_\infty^\varepsilon > \tau | \tau_\infty^\varepsilon > \underline{\tau}) = e^{-2|\mu(\underline{\tau}) \wedge 0|(\tau - \underline{\tau})}.$$

Using this, compute

$$\begin{aligned}
\frac{d}{d\tau}P(\tau_\infty^\varepsilon > \tau) &= \lim_{\delta \rightarrow 0} \delta^{-1} [P(\tau_\infty^\varepsilon > \tau + \delta) - P(\tau_\infty^\varepsilon > \tau)] \\
&= \lim_{\delta \rightarrow 0} \delta^{-1} [P(\tau_\infty^\varepsilon > \tau)P(\tau_\infty^\varepsilon > \tau + \delta | \tau_\infty^\varepsilon > \tau) - P(\tau_\infty^\varepsilon > \tau)] \\
&= \lim_{\delta \rightarrow 0} P(\tau_\infty^\varepsilon > \tau) \delta^{-1} (e^{-2|\mu(\underline{\tau}) \wedge 0| \delta} - 1) \\
&= P(\tau_\infty^\varepsilon > \tau) (-2|\mu(\underline{\tau}) \wedge 0|).
\end{aligned}$$

Solving this separable differential equation gives

$$P(\tau_\infty^\varepsilon > \tau) = \exp(-2 \int_0^\tau |\mu(\underline{s}) \wedge 0| ds),$$

and taking the limit as $\varepsilon \rightarrow 0$ gives the result. \square

For calculations, it is useful to consider the process we have constructed as a point process of excursions from the origin, indexed by local time. Here we derive some of the formulas that will be used in later sections.

Notation. For a process X_t and a subset C of $[0, \infty) \times [0, \infty]$, we define

$$\nu(X; C) = \# \{(\tau, \lambda) \in C : \phi(\tau) - \phi(\tau-) = \lambda\}.$$

We also define the measure

$$n(\cdot) = \text{E}\nu(X; \cdot).$$

Theorem 2.10. *The measure $n(\cdot)$ has density function*

$$\frac{1}{\sqrt{2\pi\lambda^3}} \exp\left(\frac{-\mu^2(\tau)\lambda}{2}\right) \exp\left(-2 \int_0^\tau |\mu(s) \wedge 0| ds\right),$$

with $\mu(l)$ as in Theorem 2.1.

Proof. For a fixed τ and λ , this decomposes as the product of the probability that an excursion of Brownian motion with drift $\mu(\tau)$ has time duration λ given that $\tau_\infty > \tau$, times the probability that $\tau_\infty > \tau$ (from Theorem 2.9). If $\tau_\infty < \tau$, no more excursions occur. \square

Finally, we calculate the probability that an excursion of Brownian motion with constant drift μ reaches height l . We will need this in the sections that follow.

Lemma 2.11. *For Brownian motion with constant drift μ , the intensity measure of excursions that reach level l before returning to the origin is given by*

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} P_\varepsilon^{(\mu)}(T_l < T_0) = \frac{\mu e^{\mu l}}{\sinh(\mu l)}.$$

Proof. Apply the Girsanov transform to get that

$$P_\varepsilon^{(\mu)}(T_l < T_0 \wedge t) = E[1_{\{T_l < T_0 \wedge t\}} Z_t],$$

where $Z_t = \exp(\mu(B_t - B_0) - \mu^2 t/2)$. By the Optional Sampling theorem, this is

$$\begin{aligned} &= E[1_{\{T_l < T_0 \wedge t\}} Z_{T_l \wedge t}] \\ &= E[1_{\{T_l < T_0 \wedge t\}} Z_{T_l}] \\ &= \exp(\mu(l - \varepsilon)) E_\varepsilon[\exp(-\mu^2 T_l/2) 1_{\{T_l < T_0\}}] \\ &= \exp(\mu(l - \varepsilon)) \frac{\sinh(\varepsilon\mu)}{\sinh(l\mu)}. \end{aligned}$$

The last formula comes from [6, 2.8.28]. □

3 A Process with Inert Drift in an Interval

The construction method in Section 2 for one inert particle and one Brownian particle can be extended to other configurations of particles. In this section we construct a process $X(t)$, which is a Brownian motion confined to the interval $[0, l]$, with drift $V(t) = K(L_0(t) - L_l(t))$, where $L_0(t)$ and $L_l(t)$ are the local times at 0 and l , resp., accumulated before time t .

Another way to look at this process is to allow the interval to move, that is, construct processes $Y_0(t) \leq X(t) \leq Y_l(t)$, where $Y_0(t)$ and $Y_l(t)$ describe the paths of inert particles with drift $V(t) = K(L_0(t) - L_l(t))$, with $Y_l(t) = Y_0(t) + l$, and with $X(t)$ the path of a Brownian particle reflected from $Y_0(t)$ and $Y_l(t)$. If we look at these processes modulo l , then $Y_0(t)$ and $Y_l(t)$ can be seen as two sides of a single inert particle, on the boundary of a disc. If we let $l = 2\pi$, then $\exp(iX(t))$ and $\exp(iY_0(t))$ trace out the paths of one Brownian particle and one inert particle on the boundary of the unit disc.

Theorem 3.1. *Given constants $l, K > 0, v \in \mathbb{R}$, and a Brownian motion $B(t)$ with $B(0) = 0$, there exist unique processes $L_0(t)$ and $L_l(t)$ satisfying*

1. $Y_0(t) \leq X(t) \leq Y_l(t)$, where

(a) $X(t) \equiv B(t) + L_0(t) - L_l(t)$

(b) $V(t) \equiv v - K(L_0(t) - L_l(t))$

(c) $Y_0(t) \equiv \int_0^t V(s) ds$

(d) $Y_l(t) \equiv Y_0(t) + l$

2. $L_0(t)$ and $L_l(t)$ are continuous, nondecreasing functions, with $L_0(0) = L_l(0) = 0$, and

3. $L_0(t)$ and $L_l(t)$ are flat off the sets $\{t : Y_0(t) = X(t)\}$ and $\{t : Y_l(t) = X(t)\}$, resp.

Proof. In Section 2 we constructed a similar process for just one Brownian and one inert particle. Because the two inert particles in this theorem are always distance l apart, we can carry out the construction piecewise; that is, do the construction for one inert particle and one Brownian particle as in the previous section until the distance between these two particles reaches l , then

continue the construction with the Brownian particle and the other inert particle, until the distance between these two reaches l , and continue switching between them.

Using Corollary 2.6, with the identifications $\mu(\tau) = K\tau$, $f(t) = B(t)$, we first construct unique processes $Y_0(t)$, $X(t)$, and $L_0(t)$ having the properties stated in the theorem, for $0 \leq t < T_0 \equiv \inf\{t : X(t) - Y_0(t) = l\}$. Applying Corollary 2.6 again, but changing the order of the Brownian and inert paths, we can extend $Y_0(t)$, $X(t)$, and $L_l(t)$ to the time interval $0 \leq t < T_1 \equiv \inf\{t > T_0 : Y_l(t) - X(t) = l\}$. Repeat this process to extend $Y_0(t)$ and $X(t)$ until $T_2 \equiv \inf\{t > T_1 : X(t) - Y_0(t) = l\}$. Then we are in the initial situation of the inert particle in contact with the upper inert particle $Y_l(t)$, and we can repeat the steps of the construction. \square




Figure 2: A process with drift in an interval

3.1 Density of the velocity process

The function $L(t) \equiv L_0(t) + L_l(t)$ is the total semimartingale local time that the Brownian particle $X(t)$ spends at the endpoints of the interval $[0, l]$. Since the velocity process $V(t)$ changes only at these endpoints, we will reparametrize V in terms of the local time. We define $\tilde{V}_\tau \equiv V(L^{-1}(\tau))$.

The process \tilde{V}_τ is a piecewise linear process consisting of segments with slope $\pm K$. The slope of the process at a particular time τ depends on which endpoint X has most recently visited. For this reason, \tilde{V}_τ is not Markov. We therefore introduce a second process J_τ , taking values 0 or 1, with value 0 indicating that X has most recently visited endpoint 0 and the velocity is increasing, and value 1 indicating that X has most recently visited endpoint l and the velocity is decreasing. This technique of introducing an additional state process is similar to that used in [1].

For convenience, we will define

$$\mathcal{D} = C^2(\mathbb{R} \times \{K, -K\}) \cap \{f : \lim_{|v| \rightarrow \infty} f(v, j) = 0\}, \quad (6)$$

that is, continuous, twice-differentiable functions on our state space that vanish at ∞ .

Lemma 3.2. *The infinitesimal generator, A , of the process (\tilde{V}_τ, J_τ) is given by*

$$\begin{aligned} Af(v, 1) &= -K \frac{\partial}{\partial v} f(v, 1) + \frac{ve^{-vl}}{\sinh(vl)} [f(v, 0) - f(v, 1)], \\ Af(v, 0) &= K \frac{\partial}{\partial v} f(v, 0) + \frac{ve^{vl}}{\sinh(vl)} [f(v, 1) - f(v, 0)]. \end{aligned}$$

Proof. We will prove the lemma using the definition of the generator (see [4]) and Lemma 2.11. Assume $f \in \mathcal{D}$.

$$Af(v, j) = \lim_{\tau \rightarrow 0} \frac{E^{(v, j)}[f(\tilde{V}_\tau, J_\tau) - f(v, j)]}{\tau}.$$

We will assume that $f \in C_b^2(\mathbb{R}) \cap \{f : \lim_{|x| \rightarrow \infty} f(x) = 0\}$.

By Lemma 2.11, excursions from 0 reach l with Poisson rate $ve^{vl}/\sinh(vl)$, and by symmetry, excursions from l that reach 0 occur with rate $ve^{-vl}/\sinh(vl)$. Let σ be the time of the first crossing. We can rewrite the previous equation for $j = 0$ as

$$\begin{aligned} Af(v, 0) &= \lim_{\tau \rightarrow 0} \left\{ \frac{1}{\tau} P^{(v, 0)}(\sigma > \tau) [f(v + K\tau, 0) - f(v, 0)] \right. \\ &\quad \left. + \frac{1}{\tau} P^{(v, 0)}(\sigma < \tau) [f(v + K\sigma - K(\tau - \sigma), 1) - f(v, 0)] \right\} \\ &= \lim_{\tau \rightarrow 0} \left\{ P^{(v, 0)}(\sigma > \tau) \frac{1}{\tau} [f(v + K\tau, 0) - f(v, 0)] \right. \\ &\quad \left. + \frac{1}{\tau} P^{(v, 0)}(\sigma < \tau) [f(v + K(2\sigma - \tau), 1) - f(v, 0)] \right\}. \end{aligned}$$

Because the limit of $P^{(v, 0)}(\sigma < \tau)/\tau$ is exactly the Poisson rate given above, $Af(v, 0)$ is as stated in the lemma. A similar calculation gives $Af(v, 1)$. \square

Lemma 3.3. *The (formal) adjoint, A^* of the generator A is given by*

$$\begin{aligned} A^*g(v, 1) &= K \frac{\partial}{\partial v} g(v, 1) + \frac{ve^{vl}}{\sinh(vl)} g(v, 0) - \frac{ve^{-vl}}{\sinh(vl)} g(v, 1), \\ A^*g(v, 0) &= -K \frac{\partial}{\partial v} g(v, 0) - \frac{ve^{vl}}{\sinh(vl)} g(v, 0) + \frac{ve^{-vl}}{\sinh(vl)} g(v, 1). \end{aligned}$$

Proof. The (formal) adjoint A^* of A is the operator satisfying, for all suitable f, g (including $f, g \in \mathcal{D}$),

$$\int (Af)g dv = \int f(A^*g) dv. \quad (7)$$

Let us assume that A is of the somewhat more general form

$$\begin{aligned} Af(v, 1) &= -K \frac{\partial}{\partial v} f(v, 1) + a(v)[f(v, 0) - f(v, 1)], \\ Af(v, 0) &= K \frac{\partial}{\partial v} f(v, 0) + b(v)[f(v, 1) - f(v, 0)]. \end{aligned}$$

Then we have that

$$\begin{aligned} \int (Af)g &= \int -K \frac{\partial}{\partial v} f(v, 1)g(v, 1) + \int a(v)[f(v, 0)g(v, 1) - f(v, 1)g(v, 1)] \\ &\quad + \int K \frac{\partial}{\partial v} f(v, 0)g(v, 0) + \int b(v)[f(v, 1)g(v, 0) - f(v, 0)g(v, 0)]. \end{aligned}$$

Integrating by parts,

$$\begin{aligned} \int (Af)g &= \int \{Kf(v, 1) \frac{\partial}{\partial v} g(v, 1) + f(v, 1)[b(v)g(v, 0) - a(v)g(v, 1)]\} \\ &\quad + \int \{-Kf(v, 0) \frac{\partial}{\partial v} g(v, 0) + f(v, 0)[-b(v)g(v, 0) + a(v)g(v, 1)]\}. \end{aligned}$$

Factoring out $f(v, j)$ leaves

$$\begin{aligned} A^*g(v, 1) &= K \frac{\partial}{\partial v} g(v, 1) + b(v)g(v, 0) - a(v)g(v, 1), \\ A^*g(v, 0) &= -K \frac{\partial}{\partial v} g(v, 0) - b(v)g(v, 0) + a(v)g(v, 1). \end{aligned}$$

□

Lemma 3.4. *The process (\tilde{V}_τ, J_τ) has stationary normal density*

$$g(v, j) = \frac{1}{2\sqrt{\pi K}} e^{-v^2/K} dv \quad (8)$$

Proof. The stationary distribution, μ , of a process is the probability measure satisfying

$$\int Af \, d\mu = 0 \quad (9)$$

for all f in the domain of A . If we assume that $d\mu$ is of the form $g(v, j)dv$, then this is equivalent to

$$\int f A^*g \, dv = 0, \quad (10)$$

so that it is sufficient to find $g(v, j)$ satisfying $A^*g(v, j) = 0$. By Lemma 3.3,

$$A^*g(v, 1) + A^*g(v, 0) = K \left(\frac{\partial}{\partial v} g(v, 1) - \frac{\partial}{\partial v} g(v, 0) \right),$$

so that $g(v, 0)$ and $g(v, 1)$ differ by a constant. Note that this does not depend on the jump intensities $a(v)$ and $b(v)$. Since $\int (g(v, 1) + g(v, 0))dv = 1$, $g(v, 0) = g(v, 1)$. Using this fact and Lemma 3.3, we get

$$\begin{aligned} A^*g(v, 1) &= K \frac{\partial}{\partial v} g(v, 1) + \frac{v(e^{vl} - e^{-vl})}{\sinh(vl)} g(v, 1) \\ &= K \frac{\partial}{\partial v} g(v, 1) + 2vg(v, 1). \end{aligned}$$

This gives the separable differential equation

$$0 = K \frac{\partial}{\partial v} g(v, 1) + 2vg(v, 1),$$

which has solutions of the form $g(v, 1) = C \exp(-v^2/K)$.

□

3.2 Behavior as $l \rightarrow 0$

In this section we will let $\tilde{V}^l(\tau)$ denote the process \tilde{V}_τ constructed in the previous section, with the constant $K = 1$. We will show that the sawtooth process $\tilde{V}^l(\tau)$ converges weakly to the Ornstein-Uhlenbeck process, under appropriate rescaling, as $l \rightarrow 0$. The proof uses results from section 11.2 of [11].

We first consider the sawtooth process $\tilde{V}^l(\tau)$ only at those times when it switches from decreasing to increasing, call them $\tau_0^l = 0, \tau_1^l, \tau_2^l, \dots$. We will also refer to those times when the process switches from increasing to decreasing, call them $\sigma_1^l, \sigma_2^l, \dots$. Let $p_n^l = \sigma_n^l - \sigma_{n-1}^l$, and $q_n^l = \tau_n^l - \sigma_n^l$.

Next, we construct the piecewise linear processes $\hat{V}^l(t)$ and $T^l(t)$. We define $\hat{V}^l(t)$ so that $\hat{V}^l(nl^2) = \tilde{V}^l(\tau_n^l)$, and $T^l(nl^2) = l\tau_n^l$.

As in the previous section, we will let $a(v, l) = ve^{vl}/\sinh(vl)$ and $b(v, l) = ve^{-vl}/\sinh(vl)$. We will use the following properties of a and b in our proof:

$$\begin{aligned} 0 &\leq \frac{\partial}{\partial v} a(v, l) \leq 2 \\ -2 &\leq \frac{\partial}{\partial v} b(v, l) \leq 0, \end{aligned}$$

so that $a(v, l)$ and $b(v, l)$ are monotone functions. We also use the property that $la(v, l) \rightarrow 1$ and $lb(v, l) \rightarrow 1$ as $l \rightarrow 0$:

$$\lim_{l \rightarrow 0} la(v, l) = \lim_{l \rightarrow 0} \frac{lve^{vl}}{\sinh vl} = \lim_{l \rightarrow 0} \frac{v(vl + 1)e^{vl}}{v \cosh vl} = 1.$$

Let $P_{v,t}^l$ be the probability measure corresponding to $\langle \hat{V}^l(t), T^l(t) \rangle$ starting from (v, t) . Let $P_{v,t}$ be the unique solution of the martingale problem (see [6] or [11]) for

$$L = \frac{\partial^2}{\partial v^2} - 2v \frac{\partial}{\partial v} + 2 \frac{\partial}{\partial T}$$

starting from (v, t) .

Lemma 3.5. $P_{v,t}^l \rightarrow P_{v,t}$ as $l \rightarrow 0$ uniformly on compact subsets of \mathbb{R}^2 .

The proof of Lemma 3.5 depends on and will follow a number of technical lemmas. We begin

with some definitions. The notation $P^{(v,t)}(dr, ds)$ denotes the transition density of (p^l, q^l) .

$$\begin{aligned}
E_l &= \{(u, d) | (u - d)^2 + l^2(u + d)^2 \leq 1\}, \\
a_l^{VV}(v, t) &= \frac{1}{l^2} \int_{E_l} (r - s)^2 P^{(v,t)}(p_1^l \in dr, q_1^l \in ds), \\
a_l^{VT}(v, t) &= \frac{1}{l^2} \int_{E_l} l(r + s)(r - s) P^{(v,t)}(p_1^l \in dr, q_1^l \in ds), \\
a_l^{TT}(v, t) &= \frac{1}{l^2} \int_{E_l} l^2(r + s)^2 P^{(v,t)}(p_1^l \in dr, q_1^l \in ds), \\
b_l^V(v, t) &= \frac{1}{l^2} \int_{E_l} (r - s) P^{(v,t)}(p_1^l \in dr, q_1^l \in ds), \\
b_l^T(v, t) &= \frac{1}{l^2} \int_{E_l} l(r + s) P^{(v,t)}(p_1^l \in dr, q_1^l \in ds), \\
\Delta_l^\varepsilon(x) &= \frac{1}{l^2} P^{(v,t)}((v_1^l, l\tau_1^l) > \varepsilon).
\end{aligned}$$

To prove the lemma, we need to show that the quantities $|a_l^{VV}(v, t) - 2|$, $|a_l^{VT}(v, t)|$, $|a_l^{TT}(v, t)|$, $|b_l^V(v, t) + 2v|$, $|b_l^T(v, t) - 2|$ and $|\Delta_l^\varepsilon(v, t)|$ converge to zero as $l \rightarrow 0$, uniformly for $|v| \leq R$. When this is done, Lemma 11.2.1 of [11] completes the proof.

Because the density $P^{(v,t)}(dr, ds)$ is a bit unwieldy for direct computation, we introduce \tilde{p}^l and \tilde{q}^l , which are exponential random variables with rate $a(v, l)$ and $b(v, l)$, respectively. Define $\tilde{a}_l^{VV}(v, t) \dots \tilde{b}_l^T(v, t)$ for \tilde{p}^l and \tilde{q}^l as above, using $\tilde{P}^{(v,t)}(dr, ds)$ for the transition density of $(\tilde{p}^l, \tilde{q}^l)$.

Remark 3. We will need to compare $P^{(v,t)}(dr, ds)$ and $\tilde{P}^{(v,t)}(dr, ds)$ in the calculations that follow. If the process $\hat{V}^l(t)$ stays between $v - \Delta$ and $v + \Delta$ for $0 \leq t \leq l^2$, then the rate of the process p^l is bounded by $a(v, l)$ and $a(v, l) + 2\Delta$, and the rate of the process q^l is bounded by $b(v, l) - 2\Delta$ and $b(v, l) + 2\Delta$. Then the densities satisfy

$$\begin{aligned}
a(v, l)(b(v, l) - 2\Delta)e^{(a(v,l)+2\Delta)r} e^{(b(v,l)+2\Delta)s} \\
\leq P^{(v,t)}(dr, ds) \leq (a(v, l) + 2\Delta)(b(v, l) + 2\Delta)e^{(a(v,l)-2\Delta)r} e^{(b(v,l)-2\Delta)s}.
\end{aligned}$$

Combining the RHS and LHS with the density for $\tilde{P}^{(v,t)}(dr, ds)$, we arrive at the inequality

$$\begin{aligned}
&|\tilde{P}^{(v,t)}(dr, ds) - P^{(v,t)}(dr, ds)| \\
&\leq \left(1 - e^{-2\Delta r} + \frac{2\Delta}{a(v, l)}\right) \left(e^{2\Delta s} - e^{-2\Delta s} + \frac{2\Delta(e^{2\Delta s} + e^{-2\Delta s})}{b(v, l)}\right) \tilde{P}^{(v,t)}(dr, ds).
\end{aligned}$$

Define $C(v, l, \Delta) =$

$$\sup_{r, s < \Delta} \left(1 - e^{-2\Delta r} + \frac{2\Delta}{a(v, l)}\right) \left(e^{2\Delta s} - e^{-2\Delta s} + \frac{2\Delta e^{2\Delta s} + 2\Delta e^{-2\Delta s}}{b(v, l)}\right) \quad (11)$$

$$\leq \left(1 - e^{-2\Delta^2} + \frac{2\Delta}{a(-R, l)}\right) \left(e^{2\Delta^2} - e^{-2\Delta^2} + \frac{2\Delta e^{2\Delta^2} + 2\Delta}{b(R, l)}\right). \quad (12)$$

It is clear that

$$\lim_{\Delta, l \rightarrow 0} \sup_{|v| \leq R} C(v, l, \Delta) \rightarrow 0.$$

Lemma 3.6.

$$|a_l^{VV}(v, t) - 2| \rightarrow 0.$$

Proof. We first need to show that

$$\frac{1}{l^2} \int_{E_l^c} (r-s)^2 \tilde{P}^{(v,t)}(dr, ds) \rightarrow 0. \quad (13)$$

Note that if $0 \leq r \leq 1/2$ and $0 \leq s \leq 1/2$, then $(r, s) \in E_l$ for $l < \sqrt{3}/2$. Then

$$\begin{aligned} \frac{1}{l^2} \int_{E_l^c} r^2 \tilde{P}^{(v,t)}(dr, ds) &\leq \frac{1}{l^2} \int_{r>1/2} r^2 \tilde{P}^{(v,t)}(dr, ds) + \frac{1}{l^2} \int_{s>1/2} r^2 \tilde{P}^{(v,t)}(dr, ds) \\ &= \frac{1}{l^2} \left(\int_{1/2}^{\infty} r^2 a(v, l) e^{-a(v, l)r} dr + E(\tilde{p}^l)^2 P(\tilde{q}^l > 1/2) \right) \\ &= \frac{1}{l^2} \left(\frac{2 + a(v, l) + a^2(v, l)/4}{a^2(v, l)} e^{-a(v, l)/2} + \frac{2}{a^2(v, l)} e^{-b(v, l)/2} \right) \\ &\leq \frac{1}{l^2} \left(\frac{2 + a(R, l) + a^2(R, l)/4}{a^2(-R, l)} e^{-a(-R, l)/2} + \frac{2}{a^2(-R, l)} e^{-b(R, l)/2} \right). \end{aligned}$$

Since $la(v, l) \rightarrow 1$ as $l \rightarrow 0$, we can show that this last term will converge to 0 if we can show that $a^2(R, l) \exp(-a(-R, l)/2) \rightarrow 0$:

$$a^2(R, l) e^{-a(-R, l)/2} \leq (1/l + 2R)^2 e^{-1/l + 2R},$$

and we know that $l^{-m} e^{-1/l} \rightarrow 0$ as $l \rightarrow 0$ for all m .

The same procedure shows that

$$\frac{1}{l^2} \int_{E_l^c} s^2 \tilde{P}^{(v,t)}(dr, ds) \rightarrow 0. \quad (14)$$

and these calculations (noting that $(r-s)^2 \leq r^2 + s^2$) give (13).

Since \tilde{p}^l and \tilde{q}^l are independent exponential random variables,

$$\begin{aligned} \tilde{a}_l^{VV}(v, t) &= \frac{1}{l^2} \int_{E_l} (r-s)^2 \tilde{P}^{(v,t)}(dr, ds) \\ &= \frac{1}{l^2} E(\tilde{p}^l - \tilde{q}^l)^2 - \frac{1}{l^2} \int_{E_l^c} (r-s)^2 \tilde{P}^{(v,t)}(dr, ds) \\ &= \frac{1}{l^2} \left(\frac{2}{a^2(v, l)} - \frac{2}{a(v, l)b(v, l)} + \frac{2}{b^2(v, l)} \right) - I. \end{aligned}$$

Since $a(v, l)$ and $b(v, l)$ are monotone, and $la(v, l)$ and $lb(v, l)$ converge to 1 for all v , and $I \rightarrow 0$, the RHS of the last equation converges to 2 uniformly for $|v| \leq R$. Combining this with equation (13) shows that

$$|\tilde{a}_l^{VV} - 2| \rightarrow 0.$$

Next we compare a_l^{VV} and \tilde{a}_l^{VV} .

$$\begin{aligned}
a_l^{VV} - \tilde{a}_l^{VV} &= \frac{1}{l^2} \int_{E_l} (r-s)^2 (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&= \frac{1}{l^2} \int_{E_l \cap \{r+s < \sqrt{l}\}} (r-s)^2 (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&\quad + \frac{1}{l^2} \int_{E_l \cap \{r+s > \sqrt{l}\}} (r-s)^2 (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&= II + III.
\end{aligned}$$

Since $(r-s)^2 \leq 1$ on E_l , the second term III can be bounded as follows:

$$\begin{aligned}
III &\leq \frac{1}{l^2} [P(p^l > \sqrt{l}/2) + P(q^l > \sqrt{l}/2, p^l < \sqrt{l}/2)] \\
&\quad + \frac{1}{l^2} [P(\tilde{p}^l > \sqrt{l}/2) + P(\tilde{q}^l > \sqrt{l}/2)]
\end{aligned} \tag{15}$$

$$\leq \frac{1}{l^2} [e^{-\sqrt{l}a(v,l)/2} + e^{-\sqrt{l}(b(v,l)-2K\sqrt{l})/2} + e^{-\sqrt{l}a(v,l)/2} + e^{-\sqrt{l}b(v,l)/2}] \tag{16}$$

since if $p^l < \sqrt{l}$, the rate of $q^l < b(v,l) + 2K\sqrt{l}$. We can compute that

$$\begin{aligned}
\frac{1}{l^2} e^{-\sqrt{l}a(v,l)/2} &< \frac{1}{l^2} e^{-\sqrt{l}(1/l-2R)} \\
&= \frac{1}{l^2} e^{-1/\sqrt{l}} e^{2R\sqrt{l}}.
\end{aligned}$$

The term $\exp(-1/\sqrt{l})/l^2 \rightarrow 0$ and $\exp(2R\sqrt{l}) \rightarrow 1$.

To bound the term II , we observe that on the set $E_l \cap \{r+s < \sqrt{l}\}$, \hat{V}^l must lie in the interval $(v - K\sqrt{l}, v + K\sqrt{l})$. Then

$$\begin{aligned}
II &\leq C(v, l, \sqrt{l}) \frac{1}{l^2} \int_{E_l \cap \{r+s < \sqrt{l}\}} (r-s)^2 \tilde{P}^{(v,t)}(dr, ds) \\
&\leq C(v, l, \sqrt{l}) \frac{1}{l^2} \int_{r,s > 0} (r-s)^2 \tilde{P}^{(v,t)}(dr, ds) \\
&= C(v, l, \sqrt{l}) \frac{1}{l^2} \left(\frac{2}{a^2(v, l)} - \frac{2}{a(v, l)b(v, l)} + \frac{2}{b^2(v, l)} \right).
\end{aligned}$$

As before, $(2/a^2 - 2/ab + 2/b)/l^2 \rightarrow 2$, and as shown above, $C(v, l, \sqrt{l}) \rightarrow 0$ uniformly on compact sets as $l \rightarrow 0$, so we conclude that $|a_l^{VV}(v, t) - \tilde{a}_l^{VV}(v, t)| \rightarrow 0$. Finally, we use the triangle inequality to conclude that $|a_l^{VV}(v, t) - 2| \rightarrow 0$. \square

Lemma 3.7.

$$|b_l^T(v, t) - 2| \rightarrow 0.$$

Proof. We will first show that

$$\frac{1}{l} \int_{E_l^c} (r+s) \tilde{P}^{(v,t)}(dr, ds) \rightarrow 0. \tag{17}$$

As in the previous case,

$$\begin{aligned}
\frac{1}{l} \int_{E_i^c} r \tilde{P}^{(v,t)}(dr, ds) &\leq \frac{1}{l} \int_{r>1/2} r \tilde{P}^{(v,t)}(dr, ds) + \frac{1}{l} \int_{s>1/2} r \tilde{P}^{(v,t)}(dr, ds) \\
&= \frac{1}{l} \left(\int_{1/2}^{\infty} r a(v, l) e^{-a(v,l)r} dr + E(\tilde{p}^l) P(\tilde{q}^l > 1/2) \right) \\
&= \frac{1}{l} \left(\frac{1 + a(v, l)/2}{a(v, l)} e^{-a(v,l)/2} + \frac{1}{a(v, l)} e^{-b(v,l)/2} \right) \\
&\leq \frac{1}{l} \left(\frac{1 + a(R, l)/2}{a(-R, l)} e^{-a(-R,l)/2} + \frac{1}{a(R, l)} e^{-b(R,l)/2} \right).
\end{aligned}$$

Since $la(v, l) \rightarrow 1$ as $l \rightarrow 0$, we can show that this last term will converge to 0 by showing that $a(R, l) \exp(-a(-R, l)/2) \rightarrow 0$:

$$a(R, l) e^{-a(-R,l)/2} \leq (1/l + 2R) e^{-1/l+2R},$$

and we know that $l^{-m} e^{-1/l} \rightarrow 0$ as $l \rightarrow 0$ for all m .

The same procedure shows that

$$\frac{1}{l} \int_{E_i^c} s \tilde{P}^{(v,t)}(dr, ds) \rightarrow 0.$$

and combining them gives (17).

Since \tilde{p}^l and \tilde{q}^l are independent exponential random variables,

$$\begin{aligned}
\tilde{b}_i^T(v, t) &= \frac{1}{l} \int_{E_i} (r + s) \tilde{P}^{(v,t)}(dr, ds) \\
&= \frac{1}{l} E(\tilde{p}^l + \tilde{q}^l) - \frac{1}{l} \int_{E_i^c} (r + s) \tilde{P}^{(v,t)}(dr, ds) \\
&= \frac{1}{l} \left(\frac{1}{a(v, l)} + \frac{1}{b(v, l)} \right) - I.
\end{aligned}$$

Since $a(v, l)$ and $b(v, l)$ are monotone, and $la(v, l)$ and $lb(v, l)$ converge to 1 for all v , and $I \rightarrow 0$, this quantity converges to 2 uniformly for $|v| \leq R$.

Next we compare b_i^T and \tilde{b}_i^T .

$$\begin{aligned}
b_i^T - \tilde{b}_i^T &= \frac{1}{l} \int_{E_i} (r + s) (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&= \frac{1}{l} \int_{E_i \cap \{r+s < \sqrt{l}\}} (r + s) (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&\quad + \frac{1}{l} \int_{E_i \cap \{r+s > \sqrt{l}\}} (r + s) (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&= II + III.
\end{aligned}$$

Since $r + s \leq 1/l$ on E_l , the second term III can be bounded as follows:

$$\begin{aligned} III &\leq \frac{1}{l^2} [P(p^l > \sqrt{l}/2) + P(q^l > \sqrt{l}/2, p^l < \sqrt{l}/2)] \\ &\quad + \frac{1}{l^2} [P(\tilde{p}^l > \sqrt{l}/2) + P(\tilde{q}^l > \sqrt{l}/2)] \end{aligned}$$

which we know converges to 0, since it is exactly the same as (15) on page 1526.

To bound the term II , we observe that on the set $E_l \cap \{r + s < \sqrt{l}\}$, \hat{V}^l must lie in the interval $(v - K\sqrt{l}, v + K\sqrt{l})$. Then

$$\begin{aligned} II &\leq C(v, l, \sqrt{l}) \frac{1}{l} \int_{E_l \cap \{r+s < \sqrt{l}\}} (r+s) \tilde{P}^{(v,t)}(dr, ds) \\ &\leq C(v, l, \sqrt{l}) \frac{1}{l} \int_{r,s > 0} (r+s) \tilde{P}^{(v,t)}(dr, ds) \\ &= C(v, l, \sqrt{l}) \frac{1}{l} \left(\frac{1}{a(v, l)} + \frac{1}{b(v, l)} \right). \end{aligned}$$

As before, $(1/a + 1/b)/l \rightarrow 2$, and as shown above, $C(v, l, \sqrt{l}) \rightarrow 0$ uniformly on compact sets as $l \rightarrow 0$, so we conclude that $|b_l^T(v, t) - \tilde{b}_l^T(v, t)| \rightarrow 0$. Finally, we use the triangle inequality to conclude that $|b_l^T(v, t) - 2| \rightarrow 0$. \square

Lemma 3.8.

$$|b_l^V(v, t) + 2v| \rightarrow 0$$

Proof. We first show that

$$\frac{1}{l^2} \int_{E_l^c} (r-s) \tilde{P}^{(v,t)}(dr, ds) \rightarrow 0. \quad (18)$$

As in the previous case,

$$\begin{aligned} \frac{1}{l^2} \int_{E_l^c} r \tilde{P}^{(v,t)}(dr, ds) &\leq \frac{1}{l^2} \int_{r > 1/2} r \tilde{P}^{(v,t)}(dr, ds) + \frac{1}{l^2} \int_{s > 1/2} r \tilde{P}^{(v,t)}(dr, ds) \\ &\leq \frac{1}{l^2} \left(\frac{1 + a(R, l)/2}{a(-R, l)} e^{-a(-R, l)/2} + \frac{1}{a(R, l)} e^{-b(R, l)/2} \right). \end{aligned}$$

Because $la(v, l) \rightarrow 1$ as $l \rightarrow 0$, this last term will converge to 0 if we can show that $(1/l)a(R, l) \exp(-a(-R, l)/2) \rightarrow 0$:

$$(1/l)a(R, l)e^{-a(-R, l)/2} \leq (1/l^2 + 2R/l)e^{-1/l+2R},$$

and we know that $l^{-m}e^{-1/l} \rightarrow 0$ as $l \rightarrow 0$ for all m .

The same procedure shows that

$$\frac{1}{l^2} \int_{E_l^c} s \tilde{P}^{(v,t)}(dr, ds) \rightarrow 0.$$

and combining them gives (18).

Since \tilde{p}^l and \tilde{q}^l are independent exponential random variables,

$$\begin{aligned}
\tilde{b}_l^V(v, t) &= \frac{1}{l^2} \int_{E_l} (r - s) \tilde{P}^{(v, t)}(dr, ds) \\
&= \frac{1}{l^2} E(\tilde{p}^l - \tilde{q}^l) - \frac{1}{l^2} \int_{E_l^c} (r - s) \tilde{P}^{(v, t)}(dr, ds) \\
&= \frac{1}{l^2} \left(\frac{1}{a(v, l)} - \frac{1}{b(v, l)} \right) - I \\
&= \frac{1}{l^2} \frac{1}{a(v, l)b(v, l)} (b(v, l) - a(v, l)) - I \\
&= \frac{1}{l^2} \frac{1}{a(v, l)b(v, l)} \frac{v(e^{-vl} - e^{vl})}{\sinh(vl)} - I \\
&= \frac{1}{l^2} \frac{1}{a(v, l)b(v, l)} (-2v) - I.
\end{aligned}$$

Since $a(v, l)$ and $b(v, l)$ are monotone, and $la(v, l)$ and $lb(v, l)$ converge to 1 for all v , and $I \rightarrow 0$, the last equation converges to $-2v$ uniformly for $|v| \leq R$.

Next we compare b_l^V and \tilde{b}_l^V .

$$\begin{aligned}
|b_l^V - \tilde{b}_l^V| &\leq \frac{1}{l^2} \int_{E_l} |r - s| (P^{(v, t)}(p^l \in dr, q^l \in ds) - P^{(v, t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&= \frac{1}{l^2} \int_{E_l \cap \{r+s < \sqrt{l}\}} |r - s| (P^{(v, t)}(p^l \in dr, q^l \in ds) - P^{(v, t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&\quad + \frac{1}{l^2} \int_{E_l \cap \{r+s > \sqrt{l}\}} |r - s| (P^{(v, t)}(p^l \in dr, q^l \in ds) - P^{(v, t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\
&= II + III.
\end{aligned}$$

Since $|r - s| \leq 1$ on E_l , the second term *III* can be bounded as follows:

$$\begin{aligned}
III &\leq \frac{1}{l^2} [P(p^l > \sqrt{l}/2) + P(q^l > \sqrt{l}/2, p^l < \sqrt{l}/2)] \\
&\quad + \frac{1}{l^2} [P(\tilde{p}^l > \sqrt{l}/2) + P(\tilde{q}^l > \sqrt{l}/2)]
\end{aligned}$$

which converges to 0, as (15) on page 1526.

To bound the term *II*, we observe that on the set $E_l \cap \{r + s < \sqrt{l}\}$, \hat{V}^l must lie in the interval $(v - K\sqrt{l}, v + K\sqrt{l})$. Then

$$\begin{aligned}
II &\leq \frac{1}{l} C(v, l, \sqrt{l}) \frac{1}{l} \int_{E_l \cap \{r+s < \sqrt{l}\}} (r + s) \tilde{P}^{(v, t)}(dr, ds) \\
&\leq \frac{1}{l} C(v, l, \sqrt{l}) \frac{1}{l} \int_{r, s > 0} (r + s) \tilde{P}^{(v, t)}(dr, ds) \\
&= \frac{1}{l} C(v, l, \sqrt{l}) \frac{1}{l} \left(\frac{1}{a(v, l)} + \frac{1}{b(v, l)} \right).
\end{aligned}$$

As before, $(1/a + 1/b)/l \rightarrow 2$. For this case, we need to show that $C(v, l, \sqrt{l})/l \rightarrow 0$ uniformly on compact sets as $l \rightarrow 0$. From (12), it is enough to show that

$$\frac{1}{l} \left(1 - e^{-2l} + \frac{2\sqrt{l}}{a(-R, l)} \right) \rightarrow 2$$

(in fact, any finite limit will do), and this is an application of l'Hôpital's rule. We conclude that $|b_l^V(v, t) - \tilde{b}_l^V(v, t)| \rightarrow 0$. Finally, we use the triangle inequality to conclude that $|b_l^V(v, t) + 2v| \rightarrow 0$. \square

Lemma 3.9.

$$|a_l^{TT}(v, t)| \rightarrow 0.$$

Proof. Since \tilde{p}^l and \tilde{q}^l are independent exponential random variables,

$$\begin{aligned} \tilde{a}_l^{TT}(v, t) &= \int_{E_l} (r+s)^2 \tilde{P}^{(v,t)}(dr, ds) \\ &\leq E(\tilde{p}^l + \tilde{q}^l)^2 \\ &= \frac{2}{a^2(v, l)} + \frac{2}{a(v, l)b(v, l)} + \frac{2}{b^2(v, l)}. \end{aligned}$$

Since $a(v, l)$ and $b(v, l)$ are monotone, and $a(v, l)$ and $b(v, l)$ converge to ∞ for all v , the last equation converges to 0 uniformly for $|v| \leq R$.

Next we compare a_l^{TT} and \tilde{a}_l^{TT} .

$$\begin{aligned} |a_l^{TT} - \tilde{a}_l^{TT}| &\leq \int_{E_l} (r+s)^2 (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\ &= \int_{E_l \cap \{r+s < \sqrt{l}\}} (r+s)^2 (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\ &\quad + \int_{E_l \cap \{r+s > \sqrt{l}\}} (r+s)^2 (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\ &= I + II. \end{aligned}$$

Since $(r+s)^2 \leq 1/l^2$ on E_l , the second term II can be bounded as follows:

$$\begin{aligned} II &\leq \frac{1}{l^2} [P(p^l > \sqrt{l}/2) + P(q^l > \sqrt{l}/2, p^l < \sqrt{l}/2)] \\ &\quad + \frac{1}{l^2} [P(\tilde{p}^l > \sqrt{l}/2) + P(\tilde{q}^l > \sqrt{l}/2)] \end{aligned}$$

which converges to 0, since it is the same as (15) on page 1526.

To bound the term I , we observe that on the set $E_l \cap \{r+s < \sqrt{l}\}$, \hat{V}^l must lie in the interval

$(v - K\sqrt{l}, v + K\sqrt{l})$. Then

$$\begin{aligned} I &\leq C(v, l, \sqrt{l}) \int_{E_l \cap \{r+s < \sqrt{l}\}} (r+s)^2 \tilde{P}^{(v,t)}(dr, ds) \\ &\leq C(v, l, \sqrt{l}) \int_{r,s > 0} (r+s)^2 \tilde{P}^{(v,t)}(dr, ds) \\ &= C(v, l, \sqrt{l}) \left(\frac{2}{a^2(v, l)} + \frac{2}{a(v, l)b(v, l)} + \frac{2}{b^2(v, l)} \right). \end{aligned}$$

In this case, both terms converge to 0 as $l \rightarrow 0$. We conclude that $|a_l^{TT}(v, t) - \tilde{a}_l^{TT}(v, t)| \rightarrow 0$. Finally, we conclude that $|a_l^{TT}(v, t)| \rightarrow 0$. \square

Lemma 3.10.

$$|a_l^{VT}(v, t)| \rightarrow 0.$$

Proof. Since \tilde{p}^l and \tilde{q}^l are independent exponential random variables,

$$\begin{aligned} \tilde{a}_l^{VT}(v, t) &= \frac{1}{l} \int_{E_l} (r+s)(r-s) \tilde{P}^{(v,t)}(dr, ds) \\ &\leq \frac{1}{l} E[(\tilde{p}^l)^2 + (\tilde{q}^l)^2] \\ &= \frac{1}{l} \left(\frac{2}{a^2(v, l)} + \frac{2}{b^2(v, l)} \right). \end{aligned}$$

Since $a(v, l)$ and $b(v, l)$ are monotone, $la(v, l)$ and $lb(v, l)$ converge to 1, and $a(v, l)$ and $b(v, l)$ converge to ∞ for all v , the last equation converges to 0 uniformly for $|v| \leq R$.

Next we compare a_l^{VT} and \tilde{a}_l^{VT} .

$$\begin{aligned} |a_l^{VT} - \tilde{a}_l^{VT}| &\leq \frac{1}{l} \int_{E_l} |r^2 - s^2| (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\ &= \frac{1}{l} \int_{E_l \cap \{r+s < \sqrt{l}\}} |r^2 - s^2| (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\ &\quad + \frac{1}{l} \int_{E_l \cap \{r+s > \sqrt{l}\}} |r^2 - s^2| (P^{(v,t)}(p^l \in dr, q^l \in ds) - P^{(v,t)}(\tilde{p}^l \in dr, \tilde{q}^l \in ds)) \\ &= I + II. \end{aligned}$$

Since $|r^2 - s^2| \leq 1/l$ on E_l , the second term II can be bounded as follows:

$$\begin{aligned} II &\leq \frac{1}{l^2} [P(p^l > \sqrt{l}/2) + P(q^l > \sqrt{l}/2, p^l < \sqrt{l}/2)] \\ &\quad + \frac{1}{l^2} [P(\tilde{p}^l > \sqrt{l}/2) + P(\tilde{q}^l > \sqrt{l}/2)] \end{aligned}$$

which we know converges to 0, since it is exactly the same as (15) on page 1526.

To bound the term I , we observe that on the set $E_l \cap \{r + s < \sqrt{l}\}$, \hat{V}^l must lie in the interval $(v - K\sqrt{l}, v + K\sqrt{l})$. Then

$$\begin{aligned} I &\leq C(v, l, \sqrt{l}) \int_{E_l \cap \{r+s < \sqrt{l}\}} (r+s)^2 \tilde{P}^{(v,t)}(dr, ds) \\ &\leq C(v, l, \sqrt{l}) \int_{r,s > 0} (r+s)^2 \tilde{P}^{(v,t)}(dr, ds) \\ &= C(v, l, \sqrt{l}) \left(\frac{2}{a^2(v, l)} + \frac{2}{a(v, l)b(v, l)} + \frac{2}{b^2(v, l)} \right). \end{aligned}$$

In this case, both terms converge to 0 as $l \rightarrow 0$, so $|a_l^{VT}(v, t) - \tilde{a}_l^{VT}(v, t)| \rightarrow 0$. Finally, we conclude by the triangle inequality that $|a_l^{VT}(v, t)| \rightarrow 0$. \square

Lemma 3.11.

$$\frac{1}{l^2} \Delta_l^\varepsilon(v, t) \rightarrow 0$$

uniformly for $|v| < R$.

Proof. In fact, we need only observe that for $l < 1/2$, the set $\{(u, v) : u + v < \sqrt{l}\} \subset E_l$. Then

$$\Delta_l^\varepsilon(v, t) \leq \frac{1}{l^2} [P(p^l > \sqrt{l}/2) + P(q^l > \sqrt{l}/2, p^l < \sqrt{l}/2)],$$

which we have shown to converge above (again, (15) on page 1526). \square

Proof of Lemma 3.5. We need to show that the martingale problem for

$$L = \frac{\partial^2}{\partial v^2} - 2v \frac{\partial}{\partial v} + 2 \frac{\partial}{\partial T}$$

has a unique solution. Since the coefficients are either bounded or linear, we can use Theorem 5.2.9 of [6]. Once uniqueness of the solution of the martingale problem for L is established, Lemma 3.5 follows directly from Theorem 11.2.3 of [11] and Lemmas 3.6–3.11. \square

An issue with Lemma 3.5 is that the measures $P_{v,t}^l$ are associated with $\hat{V}^l(t)$ rather than with the original $\tilde{V}^l(\tau)$. In fact, the convergence holds for $\tilde{V}^l(\tau)$ as well.

Theorem 3.12. *The process $\tilde{V}^l(\tau)$ converges weakly to the Ornstein-Uhlenbeck process.*

Proof. By the symmetry of a and b , the process interpolated on the other side (along σ_j 's) has the same limit as $\hat{V}^l(t)$. Since the processes interpolated along the top and the bottom of the sawtooth process converge to the same process, the whole sawtooth process must converge if we show that the distance between them is 0, or equivalently, that the distance between the two processes converges to 0 uniformly on finite time intervals. This follows from Lemmas 3.7 and 3.9.

The construction of $\hat{V}^l(t)$ involved a time change because we forced each switch of the process to have duration l^2 . The term $2 \frac{\partial}{\partial T}$ of the generator L indicates that in the limit, this is twice the duration of the actual time between switches. We can restore the original clock by dividing

the generator by two. When we do so, we find that the spatial component of the process has generator corresponding to solutions of the SDE

$$dX_t = dB_t - X_t dt.$$

This is an Ornstein-Uhlenbeck process. □

4 A Pair of Brownian Motions Separated by an Inert Particle

In this section, we consider an arrangement of two Brownian particles X_1 and X_2 separated by an inert particle Y in \mathbb{R} . More precisely, we construct processes $X_1(t) \leq Y(t) \leq X_2(t)$, where the interactions between $X_1(t)$ and $Y(t)$ and between $Y(t)$ and $X_2(t)$ are as described in Section 2.




Figure 3: Two Brownian particles separated by an inert particle

A method of construction different from that in Section 2 is needed if the two Brownian particles ever meet. Instead we introduce a random variable T_∞ to represent the first meeting of the two Brownian particles $X_1(t)$ and $X_2(t)$. In fact, we will show that with probability one, $T_\infty = \infty$.

Theorem 4.1. *Given independent Brownian motions B_1 and B_2 , with $B_j(0) = 0$, constants $x > 0$, $0 \leq y \leq x$, $v \in \mathbb{R}$, and $K > 0$, there exist unique processes $L_1(t)$ and $L_2(t)$, and a random time T_∞ , satisfying the following conditions:*

1. $X_1(t) \leq Y(t) \leq X_2(t)$, $0 \leq t \leq T_\infty$, where
 - (a) $X_1(t) \equiv B_1(t) - L_1(t)$,
 - (b) $X_2(t) \equiv x + B_2(t) + L_2(t)$,
 - (c) $V(t) \equiv v + K(L_1(t) - L_2(t))$, and
 - (d) $Y(t) \equiv y + \int_0^t V(s) ds$,
2. $L_1(t)$ and $L_2(t)$ are continuous, nondecreasing functions with $L_1(0) = L_2(0) = 0$,
3. $L_1(t)$ and $L_2(t)$ are flat off the sets $\{t : X_1(t) = Y(t)\}$ and $\{t : X_2(t) = Y(t)\}$, resp.

4. $T_\infty = \inf\{t : X_1(t) = X_2(t)\}$.

Proof. The construction method in the Section 2 and a sequence of stopping times can be used to construct this process up to the stopping time T_∞ , the limit of the stopping times used in the construction. After time T_∞ the process is not well-defined, but we show below that $P(T_\infty = \infty) = 1$. \square

We define $\mathcal{X}_x^K(t) \equiv (X_1(t), X_2(t), Y(t), V(t))$ for the processes constructed with initial state $y = 0$, $v = 0$, and constant K . The following lemma describes the scaling law of the process.

Lemma 4.2. $\varepsilon \mathcal{X}_x^K(t/\varepsilon^2) \stackrel{d}{=} \mathcal{X}_{\varepsilon x}^{K/\varepsilon}(t)$.

Proof. By Brownian scaling, the X_1 and X_2 components remain Brownian motions, and by uniqueness of local time, L_1 and L_2 have the same scaling. However, by the chain rule, the slope of the Y component has been multiplied by $1/\varepsilon$ for each t . \square

The rest of the section concerns the proof that $T_\infty = \infty$ a.s.

Theorem 4.3. Define a process $X^{\rho,T}(t)$ for $T > 0$ and $\rho \in (0, 1)$ as follows. Let

$$\Delta(t) = \begin{cases} 0 & t < T \\ -\rho L(T)(t - T) & t \geq T \end{cases}$$

By previous results, there are unique $L^{\rho,T}(t)$ and $L(t)$ such that

$$X^{\rho,T}(t) = [B(t) + \Delta(t)] + L^{\rho,T}(t) + \int_0^t L^{\rho,T}(s) ds \geq 0,$$

$$X(t) = B(t) + L(t) + \int_0^t L(s) ds \geq 0,$$

where $L^{\rho,T}(t)$ and $L(t)$ are the local times of $X^{\rho,T}(t)$ and $X(t)$, respectively, at zero. Define $L_\infty^{\rho,T} = \lim_{t \rightarrow \infty} L^{\rho,T}(t)$ and $L_\infty = \lim_{t \rightarrow \infty} L(t)$. Then

$$P(L_\infty^{\rho,T} > l) \leq P(L_\infty > l) = \exp(-l^2).$$

Proof. First note that $X(t) = X^{\rho,T}(t)$ and $L(t) = L^{\rho,T}(t)$ for $t \leq T$. Also note that the drift of term of $X^{\rho,T}(t)$ at time T is $(1 - \rho)L(T) > 0$. After time T , $X^{\rho,T}(t)$ may or may not return to the origin. If not, then $X(t)$ also would not have returned to the origin ($B(t) \geq B(t) + \Delta(t)$), so $L_\infty^{\rho,T} = (1 - \rho)L_\infty$.

Otherwise, $X^{\rho,T}$ returns to the origin at some time $\tau^{\rho,T}$. Define

$$S = \inf\{t \mid L(t) = (1 - \rho)L(T)\}$$

Notice that $X(S) = 0$ with probability 1. Construct a Brownian motion \tilde{B} by deleting the time interval $(S, \tau^{\rho,T})$ from $B(t)$:

$$\tilde{B}(t) = \begin{cases} B(t) & t \leq S \\ B(t - S + \tau^{\rho,T}) - B(\tau^{\rho,T}) + B(S) & t \geq S \end{cases}$$

and an associated local time:

$$\tilde{L}(t) = \begin{cases} L(t) & t \leq S \\ L^{\rho,T}(t - S + \tau^{\rho,T}) - \rho L(T) & t \geq S \end{cases}$$

and the associated reflected process with drift:

$$\tilde{X}(t) = \begin{cases} X(t) & t \leq S \\ X^{\rho,T}(t - S + \tau^{\rho,T}) & t \geq S \end{cases}.$$

Note that $\tilde{B}(t)$ is a Brownian motion because $\tau^{\rho,T}$ is a stopping time and S is depends only on $B[S, T]$ and so is independent of $B[0, S]$.

We will show that $\tilde{X}(t) = \tilde{B}(t) + \tilde{L}(t) + \int_0^t \tilde{L}(s) ds$. In fact, because of the pathwise uniqueness of solutions $L(t)$, we only need to check that \tilde{B} , \tilde{X} , and \tilde{L} are continuous at S .

$$\begin{aligned} \tilde{L}(S-) &= L(S) = (1 - \rho)L(S) \\ \tilde{L}(S+) &= L^{\rho,T}(\tau^{\rho,T}) - \rho L(T) \\ &= L^{\rho,T}(T) - \rho L(T) \\ &= L(T) - \rho L(T) \\ &= L(S-) \end{aligned}$$

The limit of $\tilde{L}(t)$ as $t \rightarrow \infty$ is $L_\infty^{\rho,T}$ (pathwise). But the limit of $\tilde{L}(t)$ will have the same distribution as L_∞ because $\tilde{B}(t)$ is a Brownian motion. Since we have either decreased L_∞ by a factor of ρ or replaced it with a new copy with identical distribution, the inequality holds. \square

Theorem 4.4. *For the process constructed in Theorem 4.1, $P(T = \infty) = 1$.*

Proof. By the previous lemma, we may assume that $K = 1$. We also assume that $v = 0$. We use slightly simplified versions of the process X_1 and X_2 below, which incorporate the drift term ($Y(t)$ in the definition), and which otherwise agree until time T with the definitions in Theorem 4.1.

$$\begin{aligned} X_1(t) &= B_1(t) + L_1(t) + \int_0^t V(s) ds \geq 0 \\ X_2(t) &= B_2(t) - L_2(t) + \int_0^t V(s) ds \leq 0, \end{aligned}$$

where $L_1(t)$, $L_2(t)$ are the local times of $X_1(t)$ and $X_2(t)$ at the origin, $B_1(0) > 0$ and $B_2(0) = 0$, and

$$V(t) = \begin{cases} L_1(t) - L_2(t) & t < T \\ 0 & t \geq T \end{cases},$$

with T a stopping time defined below.

Define $T_0 = 0$, $T_{j+1} = \inf\{t > T_j \mid V(t) = 0\}$, and define $T_\infty = \lim T_j$. On any of the intervals $[T_j, T_{j+1}]$ (say that $X_1(T_j) = 0$), the term $V(t)$ behaves exactly as in the case of one Brownian

particle and one inert particle, except that $V(t)$ may decrease when $X_2(t) = 0$. So $V(t)$ is dominated in distribution by L_∞ .

Using the previous theorem, we can check Novikov's condition and then apply the Girsanov theorem to (X_1, X_2) :

$$\begin{aligned}
E \exp\left(\frac{1}{2} \int_j^{j+1} (V(t))^2 dt\right) &\leq E \exp\left(\frac{1}{2} \int_j^{j+1} (L_\infty)^2 dt\right) \\
&\leq E \exp\left(\frac{1}{2} (L_\infty)^2\right) \\
&= \int_0^\infty \exp\left(\frac{1}{2} s^2\right) P(L_\infty \in ds) \\
&= \int_0^\infty \left[\int_0^s \exp\left(\frac{1}{2} r^2\right) r dr + 1 \right] P(L_\infty \in ds) \\
&= \int_0^\infty \exp\left(\frac{1}{2} r^2\right) r \int_r^\infty P(L_\infty \in ds) dr + 1 \\
&= \int_0^\infty \exp\left(\frac{1}{2} r^2\right) r \exp(-r^2) dr + 1 \\
&< \infty.
\end{aligned}$$

We can now apply Girsanov to see that under some measure, (X_1, X_2) is a standard reflected Brownian motion in the quadrant $\{(x, y) \mid x > 0, y > 0\}$. Observe that if $X_1(T_j) = 0$, then $X_2(T_{j+1}) = 0$ and $X_1(T_{j+2}) = 0$. Then $T_\infty < \infty$ implies that the reflected Brownian motion hits the origin, an event with probability zero. Therefore, $P(T_\infty = \infty) = 1$. \square

4.1 The limiting process is Bess(2)

In this section, we wish to determine the law of $X_2(t) - X_1(t)$ for the process described in Theorem 4.1, as the constant $K \rightarrow \infty$. Heuristically, one can see that if we apply regular Brownian rescaling to the process, the paths of the inert particle will get steeper, which is equivalent to increasing K . We can therefore get a picture of the long-term behavior of the process by letting $K \rightarrow \infty$.

As in the previous section, we approach the limit distribution through a Markov chain. We introduce the stopping times T_j defined by

$$T_0 = 0, \quad T_{j+1} = \inf\{t > T_j : V(t) = 0\},$$

and a two-dimensional Markov chain $\{\mathcal{Y}^K(j)\}_{j=0}^\infty$ defined by

$$\mathcal{Y}^K(j) = (X_2(T_j) - X_1(T_j), T_j).$$

We denote the transition probabilities of $\mathcal{Y}^K(j)$ by

$$\Pi_K(x, dy, dt) = P(\mathcal{Y}^K(j+1) \in (dy, dt) \mid \mathcal{Y}^K(j) = (x, 0)),$$

noting that $T_{j+1} - T_j$ is independent of the value of T_j .

Now that our processes are defined, we focus on the transition probabilities of $\{\mathcal{Y}_j^K\}_j$. The following definitions correspond to those in [11, section 11.2], with $h = 1/\sqrt{K}$.

$$\begin{aligned} b_K^X(x) &= \sqrt{K} \int_{|(y-x,t)| < 1} (y-x) \Pi_K(x, dy, dt), \\ b_K^T(x) &= \sqrt{K} \int_{|(y-x,t)| < 1} t \Pi_K(x, dy, dt), \\ a_K^{XX}(x) &= \sqrt{K} \int_{|(y-x,t)| < 1} (y-x)^2 \Pi_K(x, dy, dt), \\ a_K^{XT}(x) &= \sqrt{K} \int_{|(y-x,t)| < 1} (y-x)t \Pi_K(x, dy, dt), \\ a_K^{TT}(x) &= \sqrt{K} \int_{|(y-x,t)| < 1} t^2 \Pi_K(x, dy, dt), \\ \Delta_K^\varepsilon(x) &= \sqrt{K} \int_{|(y-x,t)| > \varepsilon} \Pi_K(x, dy, dt). \end{aligned}$$

In the calculations that follow, we focus on the first step in our Markov chain. We introduce two more random times between 0 and T_1 , defined by

$$S_1 = \sup\{t < T_1 : Y(t) = X_1(t)\},$$

$$S_2 = \inf\{t > 0 : Y(t) = X_2(t)\}.$$

The typical case will be that $0 < S_1 < S_2 < T_1$. Lemma 4.8 below makes this precise.

We also introduce the set $B(1) = \{\omega : |\mathcal{Y}^K(1) - \mathcal{Y}^K(0)| < 1\}$, which is the domain of the integrals above.

We define τ_∞^K as defined in the second section, to be the limit of $L_1(t)$ as $t \rightarrow \infty$ in the absence of the process $X_2(t)$. Applying Theorem 2.9, we compute

$$P(\tau_\infty^K > t) = \exp(-Kt^2).$$

Lemma 4.5.

$$\lim_{K \rightarrow \infty} \sqrt{K} P(L_1(S_1) > K^{-1/2+\delta}) = 0,$$

uniformly in x .

Proof. This follows from the inequality $L_1(S_1) \leq \tau_\infty^K$ and the explicit distribution for τ_∞^K (from Theorem 2.9):

$$\sqrt{K} P(\tau_\infty^K > K^{-1/2+\delta}) = \sqrt{K} \exp(-K^{2\delta}).$$

□

Next we need to show that S_1^K is sufficiently small. We do this first by examining the duration of excursions X_1 makes from the path of the inert particle. The measures are from Theorem 2.10.

Lemma 4.6. Define A_ε^K to be the number of excursions, of duration ε or larger, that X_1 makes from Y before time T_1 .

$$\lim_{K \rightarrow \infty} \sqrt{K} P(A_\varepsilon^K > 0; L_1(S_1) < K^{-1/2+\delta}) = 0.$$

Proof. $A_\varepsilon^K = \#\{(l, t) \in (\varepsilon, \infty) \times [0, L_1(S_1))\}$. If we condition the process X_1 not to make an infinite duration excursion from Y , then A_ε^K is a Poisson random variable with rate bounded above by

$$\int_{\tau=0}^{K^{-1/2+\delta}} \int_{l=\varepsilon}^{\infty} \frac{e^{-K^2\tau^2 l/2}}{\sqrt{2\pi l^3}} dl d\tau.$$

By a change of variables, we get

$$\begin{aligned} &= \int_{l=\varepsilon}^{\infty} \frac{1}{K l^2 \sqrt{\pi}} \int_{u=0}^{K^{1/2+\delta} \sqrt{l/2}} e^{-u^2} du dl \\ &< \frac{1}{K \sqrt{\pi}} \int_{l=\varepsilon}^{\infty} \frac{1}{l^2} \int_{u=0}^{\infty} e^{-u^2} du dl \\ &= \frac{2}{K\varepsilon}. \end{aligned}$$

Then

$$\sqrt{K} P(A_\varepsilon^K > 0; L_1(S_1) < K^{-1/2+\delta}) \leq \sqrt{K}(1 - e^{-2/K\varepsilon}),$$

which converges to 0 as $K \rightarrow \infty$. In fact, this will converge to 0 for $\varepsilon(K) = K^{-1/2+\delta}$, a fact we will use for the next lemma. \square

Lemma 4.7.

$$\sqrt{K} E(S_1^K \mathbf{1}_{B(1)}) \rightarrow 0$$

Proof. By the previous lemma, we need only consider excursions of length less than $K^{-1/2+\delta}$ on the set where $L_1(S_1) < K^{-1/2+\delta}$. Then

$$\begin{aligned} \sqrt{K} E(S_1^K \mathbf{1}_{B(1)}) &\leq \sqrt{K} \int_{l=0}^{K^{-1/2+\delta}} \int_{\sigma=0}^{K^{-1/2+\delta}} \frac{l e^{-K^2\sigma^2 l/2}}{\sqrt{2\pi l^3}} d\sigma dl \\ &\leq \sqrt{\frac{K}{2\pi}} \int_{l=0}^{K^{-1/2+\delta}} \int_{\sigma=0}^{K^{-1/2+\delta}} l^{-1/2} d\sigma dl \\ &= \frac{\sqrt{2}}{\sqrt{\pi}} K^{-\frac{1}{4} + \frac{3\delta}{2}} \rightarrow 0. \end{aligned}$$

\square

The next lemma allows us to work with the much nicer density of τ_∞^K instead of $L_1(S_1)$.

Lemma 4.8.

$$\lim_{K \rightarrow \infty} \sqrt{K} P(L_1(S_1) < \tau_\infty^K) = 0$$

Proof. For $L_1(S_1) < \tau_\infty^K$, the inert particle must cross the gap between X_t^1 and X_t^2 before S_1^K , the last meeting time of X_t^1 and the inert particle. Since the particle is in contact with X_t^1 at the instant S_1^K , it is sufficient to show that

$$\lim_{K \rightarrow \infty} \sqrt{K} P \left(\sup_{s, t < K^{-1/2+\delta}} (|X_1(s)| + |X_2(t) - x|) > x \right) = 0.$$

This is equivalent to showing that

$$\lim_{K \rightarrow \infty} \sqrt{K} P \left(\sup_{s, t < K^{-1/2+\delta}} (|B_1(s)| + |B_2(t)| + L_1(S_1)) > x \right) = 0.$$

We bound the LHS by

$$\lim_{K \rightarrow \infty} 3\sqrt{K} P \left(\sup_{t < K^{-1/2+\delta}} |B_1(t)| > x \right) \leq \lim_{K \rightarrow \infty} 12\sqrt{K} P \left(B_1(K^{-1/2+\delta}) > x \right),$$

which is 0 by a standard computation. □

We will also need a lower bound for $L_1(S_1)$, because the time it takes for the inert particle to cross the gap between the Brownian particles, $S_2 - S_1$, is approximately $x/KL_1(S_1)$.

Lemma 4.9.

$$\lim_{K \rightarrow \infty} \sqrt{K} P(L_1(S_1) < K^{-(3/4+\delta)}) = 0$$

Proof. By the previous lemma, we need only show this for τ_∞^K .

$$\begin{aligned} \lim_{K \rightarrow \infty} \sqrt{K} P(L_1(S_1) < K^{-(3/4+\delta)}) &= \lim_{K \rightarrow \infty} \sqrt{K} (1 - \exp(-K^{-1/2-2\delta})) \\ &= \lim_{K \rightarrow \infty} (1 + 4\delta) \exp(-K^{-1/2-2\delta}) \\ &= 0. \end{aligned}$$

□

Lemma 4.10.

$$\lim_{K \rightarrow \infty} \sqrt{K} E(L_1(S_1))^2 \rightarrow 0.$$

Proof. Because $L_1(S_1) < \tau_\infty^K$, it is enough to compute the expectation of $(\tau_\infty^K)^2$:

$$E(\tau_\infty^K)^2 = \int_0^\infty 2se^{-Ks^2} ds = \frac{1}{K}.$$

Multiplying by \sqrt{K} and taking the limit yields the result. □

Lemma 4.11.

$$\lim_{K \rightarrow \infty} \sqrt{K} E \left(L_1(S_1) - \frac{\sqrt{\pi}}{2} \right) = 0.$$

Proof. By Lemma 4.8, it is enough to compute the expectation of τ_∞^K :

$$E\tau_\infty^K = \int_0^\infty e^{-Ks^2} ds = \frac{\sqrt{\pi}}{2\sqrt{K}}.$$

Multiplying by \sqrt{K} , and taking the limit yields the result. \square

Lemma 4.12. *For all $R > 0$,*

$$\lim_{K \rightarrow \infty} \sup_{\frac{1}{R} < |x| < R} (b_K^T(x) - \sqrt{\pi}x) = 0.$$

Proof. Using Lemma 4.7 we disregard the contribution of S_1^K and $T_1^K - S_2^K$.

By [6, p. 196], we have that a Brownian motion with drift μ , starting at x , has hitting time density at zero

$$P^\mu(T \in dt) = \frac{x}{\sqrt{2\pi t^3}} \exp\left(\frac{-(x - \mu t)^2}{2t}\right) dt.$$

A table of integrals (e.g. [5, p. 353]) will reveal that

$$\int_0^\infty t P^\mu(T \in dt) = \frac{x}{\mu}.$$

In our case, assuming $X_1(S_1^K)$ is small, that is, the two Brownian particles remain close to distance x apart, we have

$$\begin{aligned} \sqrt{K} E S_2^K &= \sqrt{K} \int_{\tau=K^{-(3/4+\delta)}}^{K^{(-1/2+\delta)}} \int_{t=0}^\infty t P^{K\tau}(S_2^K \in dt) P(\tau_x^K \in d\tau) \\ &\rightarrow x\sqrt{\pi}. \end{aligned}$$

The assumption that $X_1(S_1^K)$ is small can be justified by noting that $X_1(S_1^K)$ will have mean $L_1(S_1)$ and variance S_1 , and then applying Lemmas 4.5 and 4.7. \square

Lemma 4.13. *For all $R > 0$,*

$$\lim_{K \rightarrow \infty} \sup_{\frac{1}{R} < |x| < R} a_K^{TT}(x) = 0.$$

Proof. Using the same densities as in the previous lemma, we compute

$$\begin{aligned} \sqrt{K} E (S_2^K)^2 &= \sqrt{K} \int_{\tau=K^{-(3/4+\delta)}}^{K^{(-1/2+\delta)}} \int_{t=0}^\infty t^2 P^{K\tau}(S_2^K \in dt) P(L_1(S_1) \in d\tau) \\ &\rightarrow 0. \end{aligned}$$

\square

Lemma 4.14. *For all $R > 0$,*

$$\lim_{K \rightarrow \infty} \sup_{\frac{1}{R} < |x| < R} (b_K^X(x) - \sqrt{\pi}) = 0.$$

Proof. The change in $X_2 - X_1$ can be expressed as

$$X_2 - X_1 - x = B_2(T_1) - B_1(T_1) + 2L_1(S_1).$$

Taking the expectation leaves

$$E(X_2 - X_1 - x) = 2EL_1(S_1),$$

and the result follows from Lemma 4.11. □

Lemma 4.15. *For all $R > 0$,*

$$\lim_{K \rightarrow \infty} \sup_{\frac{1}{R} < |x| < R} (a_K^{XX}(x) - 2x\sqrt{\pi}) = 0.$$

Proof. Following the proof of the previous lemma,

$$\begin{aligned} (X_2 - X_1 - x)^2 &= B_2(T_1)^2 + B_1(T_1)^2 + 4(L_1(S_1))^2 + 2B_2(T_1)B_1(T_1) \\ &\quad + 4L_1(S_1)(B_2(T_1) - B_1(T_1)). \end{aligned}$$

Taking the expectation leaves

$$E(X_2 - X_1 - x)^2 = 2ET_1 + 4E(L_1(S_1))^2 + 0 + 0,$$

and the result follows from Lemma 4.10 and Lemma 4.12. □

Lemma 4.16. *For all $R > 0$,*

$$\lim_{K \rightarrow \infty} \sup_{\frac{1}{R} < |x| < R} (a_h^{XT}(x)) = 0.$$

Proof. As in the preceding lemmas,

$$(X_2 - X_1 - x)T_1 = (B_2(T_1) - B_1(T_1) + 2L_1(S_1))T_1.$$

Taking the expectation leaves

$$E(X_2 - X_1 - x)T_1 = 2EL_1(S_1)T_1.$$

As in the previous lemmas, we discard the contribution from $T_1 - S_2$. Using the probability densities from Lemma 4.12, it is easy to see that

$$\int_0^\infty \int_0^\infty t\tau P^{K\tau}(S_2^K \in dt)P(\tau_x^K \in d\tau) = \frac{x}{K},$$

and the result follows. □

We define the process $\hat{\mathcal{Y}}^K(t)$ to be a piecewise linear process derived from the Markov chain $\mathcal{Y}^K(j)$ so that $\hat{\mathcal{Y}}^K(j/\sqrt{K}) = \mathcal{Y}^K(j)$.

We can take the domain of L to be $C^2((0, \infty) \times [0, \infty))$ in the following theorems.

Theorem 4.17. *The limiting process $\lim_{K \rightarrow \infty} \hat{\mathcal{Y}}^K(t)$ has infinitesimal generator*

$$L = X\sqrt{\pi} \frac{\partial^2}{\partial X^2} + \sqrt{\pi} \frac{\partial}{\partial X} + X\sqrt{\pi} \frac{\partial}{\partial T}.$$

Proof. Apply [11, Thm. 11.2.3] with the preceding lemmas. □

From the $X\sqrt{\pi}(\partial/\partial T)$ term above, we can see that the space-time process $\hat{\mathcal{Y}}^K(t)$ runs at a different rate than do the original Brownian motions which defined our process. We can perform a change of time to restore the original clock, by dividing the generator by $X\sqrt{\pi}$.

Theorem 4.18. *The limit of the process $(X_2(t/2) - X_1(t/2))$ as $K \rightarrow \infty$ is the 2-dimensional Bessel process.*

Proof. We actually change the clock by the factor $2X\sqrt{\pi}$ to get the correct Brownian motion term, because the original process is the difference of two Brownian motions. The generator of the space-time process, after the change of time, is

$$L = \frac{1}{2} \frac{\partial^2}{\partial X^2} + \frac{1}{2X} \frac{\partial}{\partial X} + \frac{1}{2} \frac{\partial}{\partial T}.$$

Since the process now has a linear clock rate, the first coordinate of the process will be the original $(X_2(t/2) - X_1(t/2))$, with the generator L with the T term omitted. This is exactly the generator of the two-dimensional Bessel process. □

5 A Process with Inert Drift in \mathbb{R}^d

The Brownian motion with inert drift constructed in Section 2, considered as a process on the domain $[0, \infty)$, can be generalized to higher dimensional domains. In this section, we will construct such a process in C^2 domains. Unfortunately, an error was found in the original proof of uniqueness for this process. A correct proof for bounded C^2 domains will appear in [2].

We would like to point out that S. Ramasubramanian has previously extended a more general version of the process to orthants; see [9].

We will rely on some results by P. Lions and A. Sznitman from [8]. Let D be an open set in \mathbb{R}^d , with a unit inward vector field n on ∂D satisfying (19) and (20) below. We make the following assumptions as in [8]:

$$\exists C_0, \forall x \in \partial D, \forall x' \in \bar{D}, \forall L \in n(x), (x - x', L) + C_0|x - x'|^2 \geq 0 \tag{19}$$

$$\begin{aligned} \forall x \in \partial D, \text{ if } \exists C \geq 0, \exists L \in \mathbb{R}^d : \forall x' \in \bar{D}, (x - x', L) + C|x - x'|^2 \geq 0, \\ \text{then } L = \theta n(x) \text{ for some } \theta \geq 0 \end{aligned} \tag{20}$$

A domain D is called admissible if there is a sequence $\{D_m\}$ of bounded smooth open sets in \mathbb{R}^d such that

1. D and D_m satisfy (19) and (20),
2. if $x_m \in \bar{D}_m$, $x_m \rightarrow x$, then $x \in \bar{D}$, and

3. if $K \subset D$ is compact, then $K \subset D_m$ for m large enough.

Note that C^2 domains are admissible by this definition.

We will call a pair (x_t, L_t) a solution of the Skorohod problem (w, D, n) if the following hold:

1. $x_t \in C([0, \infty), \bar{D})$,
2. $L_t \in C([0, \infty), \mathbb{R}^d)$, and L_t has bounded variation on every interval $(0, T)$ for all $T < \infty$,
3. $|L|_t = \int_0^t 1_{\{x_s \in \partial D\}} d|L|_s$,
4. $L_t = \int_0^t n(x_s) d|L|_s$,
5. $x_t = w_t + L_t$ for $t \geq 0$.

Notationally, x^j or $(x)^j$ will denote the j -th component of x when x is a vector or vector function.

We will call a function x_t a solution to the extended Skorohod problem if condition 5 above is replaced by

$$x_t = w_t + \int_0^t L_s ds + L_t. \quad (21)$$

5.1 Existence when D lies above the graph of a function

The results in this section will be very similar to the one-dimensional case. We assume that $D = \{x \in \mathbb{R}^d : x^d > f(x^1, \dots, x^{d-1})\}$ is admissible, with $f(0, \dots, 0) = 0$, and that there is an $0 < \alpha < 1$ so that $|f(x)| < 1 - \alpha$ and $n(x)^d > \alpha$ for all x . Again note that $f \in C^2$ will satisfy this in a neighborhood of any point with an appropriate choice of coordinates.

Lemma 5.1. *If $x_t = w_t + L_t$ is a solution to the Skorohod problem in D , then*

$$\alpha |L|_t < L_t^d < |L|_t < |L|_t / \alpha. \quad (22)$$

Proof. From [8] we have that

$$L_t = \int_0^t n(x_s) d|L|_s$$

so that

$$L_t^d = \int_0^t (n(x_s))^d d|L|_s > \alpha |L|_t.$$

Clearly, L_t^d is a nondecreasing function. Since $L_0 = 0$, we also have that $|L|_t < |L|_t$. Combining these, we get (22). \square

Details of uniqueness will appear in [2].

Lemma 5.2. *If $x_t = (w_t + I_t) + L_t$ solves the Skorohod problem in D for $(w_t + I_t)$ and I_t^d is an increasing function with $I_0^d = 0$, then*

$$L_t^d \leq \sup_{0 \leq s \leq t} (-w_s^d \wedge 0) + (1 - \alpha) \quad (23)$$

Proof. The proof is by contradiction. Let T be a time such that (23) does not hold, and let S be the largest time less than T such that (23) does hold. Then for $S < t \leq T$,

$$(w_t + I_t + L_t)^d > x_t^d + I_t^d + \sup_{0 \leq s \leq t} (-w_s^d \wedge 0) + (1 - \alpha) > 1 - \alpha \geq f(\tilde{x}_t).$$

This implies that L_t^d is constant on the interval $[S, T]$, a contradiction. \square

Theorem 5.3. *Given a continuous w_t with $w_0 \in D$, with D satisfying the conditions at the beginning of the section, there is an x_t satisfying (21).*

Proof. We will construct a solution as in the one-dimensional case.

We combine the results of Lions and Sznitman with Lemma 5.2 to construct, for any $\varepsilon > 0$, x_t^ε , L_t^ε , and I_t^ε satisfying

$$x_t^\varepsilon = w_t + I_t^\varepsilon + L_t^\varepsilon,$$

where L_t^ε is the local time of $w_t + I_t^\varepsilon$, and where

$$\begin{aligned} I_t^\varepsilon &= L_{T_n^\varepsilon}^\varepsilon(t - T_n^\varepsilon), & T_n^\varepsilon < t < T_{n+1}^\varepsilon, \\ T_n^\varepsilon &= \inf\{t > 0 : |L^\varepsilon|_t = n\varepsilon\}. \end{aligned}$$

By Lemma 5.2 and (22), the family $\{L_t^\varepsilon\}_{\varepsilon > 0}$ is bounded for each t , and by Lemma 5.4 below, is equicontinuous in t . We can therefore apply the Ascoli–Arzelà Theorem and find a subsequence converging uniformly on any $[0, T]$ to some L_t and x_t . The uniform convergence gives that x_t is a solution to (21). \square

5.2 Existence when D is a bounded domain

Lemma 5.4. *Let $x_t = w_t + L_t$ be a solution to the Skorohod problem in D , and let $\varepsilon > 0$. There is a $\delta > 0$ so that $|L|_T - |L|_S < \varepsilon$ whenever $\text{diam}(w_{[S, T]}) < \delta$.*

Proof. Let $x \in \partial D$, and choose coordinates where x is the origin and $n(x) = \langle 0, \dots, 0, 1 \rangle$. Because D is C^1 , we can find a $0 < \rho < \varepsilon$ so that for $y \in \partial D$ and $|y - x| < 3\rho$, we have $n(y)^d > 2/3$, and $|y^d| < \rho/3$. Let $\delta = \rho/3$. Suppose that $\rho < |L|_T - |L|_S < 2\rho$. Then $|x_T - x_S| < 2\rho + |w_T - w_S| < 3\rho$, and $x_T^d - x_S^d > w_T^d - w_S^d + \int_S^T n(x_s) d|L|_s > 2\rho/3 - \rho/3$, so that $x_T \notin \partial D$, a contradiction. \square

It is left to show that for a more general domain solutions to (21) exist. We can do this by piecing together graphs of functions.

Theorem 5.5. *Solutions to (21) exist for D an admissible, bounded, piecewise C^2 domain.*

Proof. The construction is standard. Divide D into neighborhoods N_1, \dots, N_m which are nice, in the sense that, under an appropriate rotation of the standard coordinate system, each $N_j \cap \partial D$ is a section of the graph of a function f_j satisfying the conditions at the beginning of the previous section. Assume that w_t first encounters N_1 . Construct the domain which lies above the graph of f_1 and construct $x_t^{(1)}$ satisfying (21) on this new domain. Let $T_1 = \inf\{t : x_t^{(1)} \notin N_1\}$. Repeat the process starting at T_1 for the function $w_t = w_t + L_{T_1} + L_{T_1}(t - T_1)$. Continue that construction, so that the limit x_t satisfies (21) on $[0, \lim_{n \rightarrow \infty} T_n]$.

We wish to show that $\lim_{n \rightarrow \infty} T_n = \infty$. If not, say $T_n \rightarrow T$, then by Lemma 5.4 we must have that $\lim_{t \rightarrow T} |L_t| = \infty$. Then there is some $1 \leq j \leq d$, $R, S < T$ so that $0 < L_R^j < L_t^j$ for $R < t < S$, and $L_S^j \geq L_R^j + \text{diam}(\partial D) + \text{diam}(w[0, T])$. But this contradicts that $x_R, x_S \in D$.

□

References

- [1] Martin Barlow, Krzysztof Burdzy, Haya Kaspri, and Avi Mandelbaum, *Variably skewed Brownian motion*, Electron. Comm. Probab. **5** (2000), 57–66 (electronic). MR1752008 (2001j:60146)
- [2] R. Bass, K. Burdzy, Z. Q. Chen, and M. Hairer, *Stationary distributions for diffusions with inert drift*, (forthcoming paper).
- [3] Krzysztof Burdzy and David White, *A Gaussian oscillator*, Electron. Comm. Probab. **9** (2004), 92–95 (electronic). MR2108855 (2005i:60173)
- [4] Stewart N. Ethier and Thomas G. Kurtz, *Markov processes*, Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics, John Wiley & Sons Inc., New York, 1986, Characterization and convergence. MR838085 (88a:60130)
- [5] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, sixth ed., Academic Press Inc., San Diego, CA, 2000, Translated from the Russian, Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger. MR1773820 (2001c:00002)
- [6] Ioannis Karatzas and Steven E. Shreve, *Brownian motion and stochastic calculus*, second ed., Graduate Texts in Mathematics, vol. 113, Springer-Verlag, New York, 1991. MR1121940 (92h:60127)
- [7] Frank B. Knight, *On the path of an inert object impinged on one side by a Brownian particle*, Probab. Theory Related Fields **121** (2001), no. 4, 577–598. MR1872429 (2002i:60148)
- [8] P.-L. Lions and A.-S. Sznitman, *Stochastic differential equations with reflecting boundary conditions*, Comm. Pure Appl. Math. **37** (1984), no. 4, 511–537. MR745330 (85m:60105)
- [9] S. Ramasubramanian, *A subsidy-surplus model and the Skorokhod problem in an orthant*, Math. Oper. Res. **25** (2000), no. 3, 509–538. MR1855180 (2002g:91107)
- [10] H. L. Royden, *Real analysis*, third ed., Macmillan Publishing Company, New York, 1988. MR1013117 (90g:00004)

- [11] Daniel W. Stroock and S. R. Srinivasa Varadhan, *Multidimensional diffusion processes*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 233, Springer-Verlag, Berlin, 1979. MR532498 (81f:60108)