

**SOLUTION OF LINEAR MATRIX EQUATIONS IN A  
\*CONGRUENCE CLASS<sup>§</sup>**

ROGER A. HORN\*, VLADIMIR V. SERGEICHUK<sup>†</sup>, AND NAOMI SHAKED-MONDERER<sup>‡</sup>

**Abstract.** The possible \*congruence classes of a square solution to the real or complex linear matrix equation  $AX = B$  are determined. The solution is elementary and self contained, and includes several known results as special cases, e.g.,  $X$  is Hermitian or positive semidefinite, and  $X$  is real with positive definite symmetric part.

**Key words.** Linear matrix equations, \*Congruence, Positive definite matrix, Positive semidefinite matrix, Hermitian part, Symmetric part.

**AMS subject classifications.** 15A04, 15A06, 15A21, 15A57, 15A63.

**1. Introduction.** Let  $\mathbb{F}$  be either  $\mathbb{R}$  or  $\mathbb{C}$ , let  $\mathbb{F}^{p \times q}$  denote the vector space (over  $\mathbb{F}$ ) of  $p$ -by- $q$  matrices with entries in  $\mathbb{F}$ , and let  $A, B \in \mathbb{F}^{k \times n}$  be given. We are interested in the linear matrix equation  $AX = B$ , which we assume to be *consistent*:  $\text{rank } A = \text{rank } [A \ B]$ .

For a given  $S \in \mathbb{F}^{n \times n}$  let  $S^* \equiv \bar{S}^T$  denote the conjugate transpose, so  $S^* = S^T$  if  $\mathbb{F} = \mathbb{R}$ . Matrices  $X, Y \in \mathbb{F}^{n \times n}$  are in the same *\*congruence class* if there is a nonsingular  $S \in \mathbb{F}^{n \times n}$  such that  $X = S^*YS$ . The *Hermitian part* of  $X \in \mathbb{F}^{n \times n}$  is  $H(X) \equiv (X + X^*)/2$ ; when  $\mathbb{F} = \mathbb{R}$ ,  $H(X)$  is also called the *symmetric part* of  $X$ . Let  $I_p$  (respectively,  $0_p$ ) denote the  $p$ -by- $p$  identity (respectively, zero) matrix.

When does  $AX = B$  have a solution  $X$  in a given \*congruence class? Special cases of this question involving positive semidefinite or Hermitian solutions were investigated in [1]; [2] asked an equivalent question: If  $\{\xi_1, \dots, \xi_k\}$  and  $\{\eta_1, \dots, \eta_k\}$  are given sets of real or complex vectors of the same size, when is there a Hermitian or positive definite matrix  $K$  such that  $K\xi_i = \eta_i$  for  $i = 1, \dots, k$ ?

**2. Solution of  $AX = B$  in a given \*congruence class.** Our main result is the following theorem.

**THEOREM 1.** *Let  $A, B \in \mathbb{F}^{k \times n}$  be given, and suppose the linear matrix equation  $AX = B$  is consistent. Let  $r = \text{rank } A$ , and let  $M = BA^*$ . Then there are matrices  $N \in \mathbb{F}^{r \times r}$  and  $E \in \mathbb{F}^{r \times (n-r)}$  such that:*

- (a)  *$M$  is \*congruent to  $N \oplus 0_{k-r}$ .*
- (b) *For each given  $F \in \mathbb{F}^{(n-r) \times r}$  and  $G \in \mathbb{F}^{(n-r) \times (n-r)}$  there is an  $X \in \mathbb{F}^{n \times n}$  such that  $AX = B$  and  $X$  is \*congruent to*

$$\begin{bmatrix} N & E \\ F & G \end{bmatrix}.$$

<sup>§</sup>Received by the editors 11 March 2005. Accepted for publication 31 May 2005. Handling Editor Ravindra B. Bapat.

\*Department of Mathematics, University of Utah, Salt Lake City, Utah 84103, USA (rhorn@math.utah.edu).

<sup>†</sup>Institute of Mathematics, Tereshchenkivska 3, Kiev, Ukraine (sergeich@imath.kiev.ua).

<sup>‡</sup>Emek Yezreel College, Emek Yezreel 19300, Israel (nomi@tx.technion.ac.il).

(c) If  $\text{rank } M = \text{rank } B$ , then for each given  $C \in \mathbb{F}^{(n-r) \times (n-r)}$  there is an  $X \in \mathbb{F}^{n \times n}$  such that  $AX = B$  and  $X$  is  $*$ -congruent to  $N \oplus C$  over  $\mathbb{F}$ .

*Proof.* Using the singular value decomposition, one can construct a unitary  $U \in \mathbb{F}^{n \times n}$  and a nonsingular  $R \in \mathbb{F}^{k \times k}$  such that

$$RAU = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}.$$

Consistency ensures that  $B = AC$  for some  $C \in \mathbb{F}^{n \times n}$ , so

$$RBU = (RAU)(U^*CU) = \begin{bmatrix} N & E \\ 0 & 0 \end{bmatrix},$$

in which  $N \in \mathbb{F}^{r \times r}$ . A matrix  $X = U\mathcal{X}U^*$  satisfies  $AX = B$  if and only if  $\mathcal{X} \in \mathbb{F}^{n \times n}$  has the property that  $(RAU)\mathcal{X} = RBU$  if and only if it has the form

$$(1) \quad \mathcal{X} = \begin{bmatrix} N & E \\ F & G \end{bmatrix}, \quad G \in \mathbb{F}^{(n-r) \times (n-r)};$$

the entries of  $F$  and  $G$  may be any elements of  $\mathbb{F}$ . Since  $RMR^* = RBU(RAU)^* = N \oplus 0_{k-r}$ ,  $M$  is  $*$ -congruent to  $N \oplus 0_{k-r}$ .

We have

$$\text{rank } M = \text{rank } N \leq \text{rank } [N \ E] = \text{rank } B,$$

so  $\text{rank } M = \text{rank } B$  if and only if  $\text{rank } B = \text{rank } N$  if and only if every column of  $E$  is in the range of  $N$ , that is, if and only if there is a matrix  $Z$  over  $\mathbb{F}$  such that  $E = NZ$ . If  $\text{rank } M = \text{rank } B$ , we may take  $X = U\mathcal{X}U^*$ , in which

$$\begin{aligned} \mathcal{X} &= \begin{bmatrix} N & NZ \\ Z^*N & Z^*NZ + C \end{bmatrix} \\ &= \begin{bmatrix} I_r & Z \\ 0 & I_{n-r} \end{bmatrix}^* \begin{bmatrix} N & 0 \\ 0 & C \end{bmatrix} \begin{bmatrix} I_r & Z \\ 0 & I_{n-r} \end{bmatrix}. \end{aligned}$$

Then  $AX = B$  and  $X$  is  $*$ -congruent to  $N \oplus C$  over  $\mathbb{F}$ .  $\square$

Several known results follow easily from our theorem. In each of the following corollaries, we use the notation of the theorem and assume that  $AX = B$  is consistent.

**COROLLARY 2** ([2, Theorem 2.1]). *Suppose  $\text{rank } A = k$ . There is a Hermitian positive definite matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  if and only if  $M$  is Hermitian positive definite.*

*Proof.* The rank condition implies that  $M$  is  $*$ -congruent to  $N$ , so  $N$  is Hermitian positive definite if  $M$  is. The theorem ensures that there is a matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  and  $X$  is  $*$ -congruent to  $N \oplus I_{n-k}$  over  $\mathbb{F}$ , so this  $X$  is Hermitian positive definite. Conversely, if  $X$  is Hermitian positive definite and  $AX = B$ , then  $B$  and  $AX^{1/2}$  have full row rank, so  $M = BA^* = AXA^* = (AX^{1/2})(AX^{1/2})^*$  is Hermitian positive definite.  $\square$

COROLLARY 3 ([1, Theorem 2.2]). *There is a Hermitian positive semidefinite matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  if and only if  $\text{rank } M = \text{rank } B$  and  $M$  is Hermitian positive semidefinite.*

*Proof.* If  $M$  is Hermitian positive semidefinite, then so is  $N$ . For any Hermitian positive semidefinite  $C \in \mathbb{F}^{(n-r) \times (n-r)}$ , the theorem ensures that there is a matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  and  $X$  is \*congruent to  $N \oplus C$  over  $\mathbb{F}$ ; such an  $X$  is Hermitian positive semidefinite. Conversely, if  $X$  is Hermitian positive semidefinite and  $AX = B$ , then  $M = BA^* = AXA^*$  is Hermitian positive semidefinite, and  $\text{rank } M = \text{rank } (AX^{1/2})(AX^{1/2})^* = \text{rank } (AX^{1/2}) = \text{rank } AX = \text{rank } B$ .  $\square$

The real case of part (b) in the following corollary was proved in [2, Theorem 2.1] with the restriction that  $A$  has full row rank.

COROLLARY 4. (a) *There is a square matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  and  $H(X)$  is positive semidefinite if and only if  $H(M)$  is positive semidefinite.*

(b) *There is a square matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  and  $H(X)$  is positive definite if and only if  $H(M)$  is positive semidefinite and  $\text{rank } H(M) = \text{rank } A$ .*

*Proof.* Necessity in both cases follows from observing that  $H(M) = AH(X)A^* = (AH(X)^{1/2})(AH(X)^{1/2})^*$ . Thus,  $\text{rank } H(M) = \text{rank } (AH(X)^{1/2}) = \text{rank } A$  if  $H(X)$  is nonsingular.

Conversely,  $H(M)$  is \*congruent to  $H(N) \oplus 0_{k-r}$  so  $H(N)$  is positive semidefinite and  $\text{rank } H(N) = \text{rank } H(M)$ . Take  $F = -E^*$  and  $G = I_{n-r}$  in (1), so that  $H(X)$  is \*congruent to  $H(\mathcal{X}) = H(N) \oplus I_{n-r}$ . For this  $X$ ,  $AX = B$ ,  $H(X)$  is positive semidefinite, and  $H(X)$  is positive definite if  $\text{rank } H(M) = r$ .  $\square$

Part (a) of the following corollary was proved in [1, Theorem 2.1].

COROLLARY 5. (a) *There is a square matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  and  $X$  is Hermitian if and only if  $M$  is Hermitian.*

(b) *There is a square matrix  $X$  over  $\mathbb{F}$  such that  $AX = B$  and  $X$  is skew-Hermitian if and only if  $M$  is skew-Hermitian.*

*Proof.* Necessity in both cases follows from observing that  $M = AXA^*$ . Conversely, choosing  $G = 0$  and  $F = \pm E^*$  in (1) proves sufficiency.  $\square$

The inertia of a Hermitian matrix  $H$  is  $\text{In } H = (\pi(H), \nu(H), \zeta(H))$ , in which  $\pi(H)$  is the number of positive eigenvalues of  $H$ ,  $\nu(H)$  is the number of negative eigenvalues, and  $\zeta(H)$  is the nullity. Since we know the general parametric form (1), the preceding corollaries can be made more specific in the Hermitian cases by describing the inertias that are possible for  $X$  given the inertia of  $M$ . Our final corollary is an example of such a result.

COROLLARY 6. *Suppose  $M$  is Hermitian and  $\text{rank } M = \text{rank } B$ . Then  $X$  may be chosen to be Hermitian with inertia  $(\alpha, \beta, \gamma)$  if and only if  $\alpha, \beta$ , and  $\gamma$  are nonnegative integers such that  $\alpha + \beta + \gamma = n$  and  $(\alpha, \beta, \gamma) \geq \text{In } M - (0, 0, k - r)$ .*

*Proof.* Since  $\text{rank } M = \text{rank } B$ , the theorem ensures for any  $C \in \mathbb{F}^{(n-r) \times (n-r)}$  the existence of an  $X$  that is \*congruent over  $\mathbb{F}$  to  $N \oplus C$ . Take  $C$  to be Hermitian, in which case  $\text{In } X = \text{In } N + \text{In } C \geq \text{In } M - (0, 0, k - r)$ , and all permitted inertias can be achieved by a suitable choice of  $C$ .  $\square$

If the rank condition in the preceding corollary is not satisfied, there may be

further restrictions on the possible set of inertias of  $A$ . Consider the example  $A = [1 \ 0]$ ,  $B = [0 \ 1]$ ,  $M = [0]$ . Any Hermitian solution to  $AX = B$  must have the form

$$X = \begin{bmatrix} 0 & 1 \\ 1 & t \end{bmatrix}$$

for some real  $t \in \mathbb{F}$ , and any such matrix has inertia  $(1, 1, 0) \not\asymp (0, 0, 1)$ .

#### REFERENCES

- [1] C.G. Khatri and S.K. Mitra. Hermitian and nonnegative definite solutions of linear matrix equations. *SIAM J. Appl. Math.*, 31:579–585, 1976.
- [2] A. Pinkus. Interpolation by matrices. *Electron. J. Linear Algebra*, 11:281–291, 2004.