

PERTURBATIONS OF FUNCTIONS OF DIAGONALIZABLE MATRICES*

MICHAEL I. GIL[†]

Abstract. Let A and \tilde{A} be $n \times n$ diagonalizable matrices and f be a function defined on their spectra. In the present paper, bounds for the norm of $f(A) - f(\tilde{A})$ are established. Applications to differential equations are also discussed.

Key words. Matrix valued functions, Perturbations, Similarity of matrices, Diagonalizable matrices.

AMS subject classifications. 15A54, 15A45, 15A60.

1. Introduction and statement of the main result. Let \mathbb{C}^n be a Euclidean space with the scalar product (\cdot, \cdot) , Euclidean norm $\|\cdot\| = \sqrt{(\cdot, \cdot)}$ and identity operator I . A and \tilde{A} are $n \times n$ matrices with eigenvalues λ_j and $\tilde{\lambda}_j$ ($j = 1, \dots, n$), respectively. $\sigma(A)$ denotes the spectrum of A , A^* is the adjoint to A , and $N_2(A)$ is the Hilbert-Schmidt (Frobenius) norm of A : $N_2^2(A) = \text{Trace}(A^*A)$.

In the sequel, it is assumed that each of the matrices A and \tilde{A} has n linearly independent eigenvectors, and therefore, these matrices are diagonalizable. In other words, the eigenvalues of these matrices are semi-simple.

Let f be a scalar function defined on $\sigma(A) \cup \sigma(\tilde{A})$. The aim of this paper is to establish inequalities for the norm of $f(A) - f(\tilde{A})$. The literature on perturbations of matrix valued functions is very rich but mainly, perturbations of matrix functions of a complex argument and matrix functions of Hermitian matrices were considered, cf. [1, 11, 13, 14, 16, 18]. The matrix valued functions of a non-Hermitian argument have been investigated essentially less, although they are very important for various applications; see the book [10].

The following quantity plays an essential role hereafter:

$$g(A) := \left[N_2^2(A) - \sum_{k=1}^n |\lambda_k|^2 \right]^{1/2}.$$

*Received by the editors December 8, 2009. Accepted for publication April 19, 2010. Handling Editor: Peter Lancaster.

[†]Department of Mathematics, Ben Gurion University of the Negev, P.O. Box 653, Beer-Sheva 84105, Israel (gilmi@cs.bgu.ac.il). Supported by the Kameah fund of the Israel.

$g(A)$ enjoys the following properties:

$$g^2(A) \leq 2N_2^2(A_I) \quad (A_I = (A - A^*)/2i) \quad \text{and} \quad g^2(A) \leq N_2^2(A) - |\text{Trace}(A^2)|, \quad (1.1)$$

cf. [8, Section 2.1]. If A is normal, then $g(A) = 0$. Denote by $\mu_j, j = 1, \dots, m \leq n$, the distinct eigenvalues of A , and by p_j the algebraic multiplicity of μ_j . In particular, one can write $\mu_1 = \lambda_1 = \dots = \lambda_{p_1}, \mu_2 = \lambda_{p_1+1} = \dots = \lambda_{p_1+p_2}$, etc. Similarly, $\tilde{\mu}_j, j = 1, \dots, \tilde{m} \leq n$, are the distinct eigenvalues of \tilde{A} , and \tilde{p}_j denotes the algebraic multiplicity of $\tilde{\mu}_j$.

Let δ_j be the half-distance from μ_j to the other eigenvalues of A , namely,

$$\delta_j := \min_{k=1, \dots, m; k \neq j} |\mu_j - \mu_k|/2 > 0.$$

Similarly,

$$\tilde{\delta}_j := \min_{k=1, \dots, \tilde{m}; k \neq j} |\tilde{\mu}_j - \tilde{\mu}_k|/2 > 0.$$

Put

$$\beta(A) := \sum_{j=1}^m p_j \sum_{k=0}^{n-1} \frac{g^k(A)}{\delta_j^k \sqrt{k!}} \quad \text{and} \quad \beta(\tilde{A}) := \sum_{j=1}^{\tilde{m}} \tilde{p}_j \sum_{k=0}^{n-1} \frac{g^k(\tilde{A})}{\tilde{\delta}_j^k \sqrt{k!}}.$$

Here $g^0(A) = \delta_j^0 = 1$ and $\tilde{g}^0(A) = \tilde{\delta}_j^0 = 1$. According to (1.1),

$$\beta(A) \leq \sum_{j=1}^m p_j \sum_{k=0}^{n-1} \frac{(\sqrt{2}N_2(A_I))^k}{\delta_j^k \sqrt{k!}}.$$

In what follows, we put

$$\frac{f(\lambda_k) - f(\tilde{\lambda}_j)}{\lambda_k - \tilde{\lambda}_j} = 0 \quad \text{if} \quad \lambda_k = \tilde{\lambda}_j.$$

Now we are in a position to formulate our main result.

THEOREM 1.1. *Let A and \tilde{A} be $n \times n$ diagonalizable matrices and f be a function defined on $\sigma(A) \cup \sigma(\tilde{A})$. Then the inequalities*

$$N_2(f(A) - f(\tilde{A})) \leq \beta(A)\beta(\tilde{A}) \max_{j,k} \left| \frac{f(\lambda_k) - f(\tilde{\lambda}_j)}{\lambda_k - \tilde{\lambda}_j} \right| N_2(A - \tilde{A}) \quad (1.2)$$

and

$$N_2(f(A) - f(\tilde{A})) \leq \beta(A)\beta(\tilde{A}) \max_{j,k} |f(\lambda_k) - f(\tilde{\lambda}_j)| \quad (1.3)$$

are valid.

The proof of this theorem is divided into lemmas which are presented in the next two sections. The importance of Theorem 1.1 lies in the fact that the right-hand sides of inequalities (1.2) and (1.3) only involve universal quantities calculated for A and \tilde{A} , and the values of the function f on the spectra $\sigma(A)$ and $\sigma(\tilde{A})$, but e.g. no matrices performing similarities of A and \tilde{A} to diagonal ones.

2. The basic lemma. Let T and \tilde{T} be the invertible matrices performing similarities of A and \tilde{A} to diagonal ones, that is,

$$T^{-1}AT = S, \tag{2.1}$$

and

$$\tilde{T}^{-1}\tilde{A}\tilde{T} = \tilde{S}, \tag{2.2}$$

where $S = \text{diag}(\lambda_j)$ and $\tilde{S} = \text{diag}(\tilde{\lambda}_j)$. Everywhere below, $\{d_k\}_{k=1}^n$ is the standard orthonormal basis and $\|B\| = \sup \{\|Bh\|/\|h\| : h \in \mathbb{C}^n\}$ for an $n \times n$ matrix B . Recall that $N_2(AB) \leq \|B\|N_2(A)$.

We begin with the following lemma.

LEMMA 2.1. *Let the hypotheses of Theorem 1.1 hold. Then*

$$N_2^2(f(A) - f(\tilde{A})) \leq \|\tilde{T}^{-1}\|^2 \|T\|^2 \sum_{j,k=1}^n |(f(\lambda_k) - f(\tilde{\lambda}_j))(T^{-1}\tilde{T}d_j, d_k)|^2. \tag{2.3}$$

Proof. We have

$$\begin{aligned} N_2^2(A - \tilde{A}) &= N_2^2(TST^{-1} - \tilde{T}\tilde{S}\tilde{T}^{-1}) = N_2^2(TST^{-1}\tilde{T}\tilde{T}^{-1} - TT^{-1}\tilde{T}\tilde{S}\tilde{T}^{-1}) \\ &= N_2^2(T(ST^{-1}\tilde{T} - T^{-1}\tilde{T}\tilde{S})\tilde{T}^{-1}) \leq \|\tilde{T}^{-1}\|^2 \|T\|^2 N_2^2(ST^{-1}\tilde{T} - T^{-1}\tilde{T}\tilde{S}). \end{aligned}$$

But

$$\begin{aligned} N_2^2(ST^{-1}\tilde{T} - T^{-1}\tilde{T}\tilde{S}) &= \sum_{j,k=1}^n |(ST^{-1}\tilde{T} - T^{-1}\tilde{T}\tilde{S})d_j, d_k|^2 \\ &= \sum_{j,k=1}^n |(T^{-1}\tilde{T}d_j, S^*d_k) - (T^{-1}\tilde{T}\tilde{S}d_j, d_k)|^2 = J, \end{aligned} \tag{2.4}$$

where

$$J := \sum_{j,k=1}^n |(\lambda_k - \tilde{\lambda}_j)(T^{-1}\tilde{T}d_j, d_k)|^2.$$

So,

$$N_2^2(A - \tilde{A}) \leq \|\tilde{T}^{-1}\|^2 \|T\|^2 J.$$

Similarly, taking into account that

$$T^{-1}f(A)T = f(S) = \text{diag}(f(\lambda_j)) \quad \text{and} \quad \tilde{T}^{-1}f(\tilde{A})\tilde{T} = f(\tilde{S}) = \text{diag}(f(\tilde{\lambda}_j)),$$

we get (2.3), as claimed. \square

From the previous lemma we obtain at once:

COROLLARY 2.2. *Under the hypotheses of Theorem 1.1, we have*

$$N_2(f(A) - f(\tilde{A})) \leq \kappa_T \tilde{\kappa}_T \left[\sum_{j,k=1}^n |f(\lambda_k) - f(\tilde{\lambda}_j)|^2 \right]^{1/2},$$

where

$$\kappa_T := \|T\| \|T^{-1}\| \quad \text{and} \quad \tilde{\kappa}_T := \|\tilde{T}\| \|\tilde{T}^{-1}\|.$$

Recall that

$$\frac{f(\lambda_k) - f(\tilde{\lambda}_j)}{\lambda_k - \tilde{\lambda}_j} = 0 \quad \text{if} \quad \lambda_k = \tilde{\lambda}_j.$$

Note that (2.3) implies

$$\begin{aligned} N_2^2(f(A) - f(\tilde{A})) &\leq \|\tilde{T}^{-1}\|^2 \|T\|^2 \sum_{j,k=1}^n \left| \frac{f(\lambda_k) - f(\tilde{\lambda}_j)}{\lambda_k - \tilde{\lambda}_j} (\lambda_k - \tilde{\lambda}_j) (T^{-1}\tilde{T}d_j, d_k) \right|^2 \\ &\leq \|\tilde{T}^{-1}\|^2 \|T\|^2 \max_{j,k} \left| \frac{f(\lambda_k) - f(\tilde{\lambda}_j)}{\lambda_k - \tilde{\lambda}_j} \right|^2 J. \end{aligned} \tag{2.5}$$

It follows from (2.4) that

$$J = N_2^2(T^{-1}A\tilde{T} - T^{-1}\tilde{A}\tilde{T}) \leq \|T^{-1}\|^2 \|\tilde{T}\|^2 N_2^2(A - \tilde{A}).$$

Thus, (2.5) yields:

COROLLARY 2.3. *Under the hypotheses of Theorem 1.1, the inequality*

$$N_2(f(A) - f(\tilde{A})) \leq \kappa_T \tilde{\kappa}_T N_2(A - \tilde{A}) \max_{j,k} \left| \frac{f(\lambda_k) - f(\tilde{\lambda}_j)}{\lambda_k - \tilde{\lambda}_j} \right| \tag{2.6}$$

is true.

Since

$$\left| \frac{f(\lambda_k) - f(\tilde{\lambda}_j)}{\lambda_k - \tilde{\lambda}_j} \right| \leq \sup_{s \in \text{co}(A, \tilde{A})} |f'(s)|,$$

where $co(A, \tilde{A})$ is the closed convex hull of the eigenvalues of the both matrices A and \tilde{A} , cf. [17], we have:

COROLLARY 2.4. *Let A and \tilde{A} be $n \times n$ diagonalizable matrices and f be a function defined and differentiable on $co(A, \tilde{A})$. Then*

$$N_2(f(A) - f(\tilde{A})) \leq \kappa_T \tilde{\kappa}_T \max_{s \in co(A, \tilde{A})} |f'(s)| N_2(A - \tilde{A}).$$

We need also the following corollary of Lemma 2.1.

COROLLARY 2.5. *Under the hypotheses of Theorem 1.1, it holds that*

$$N_2(f(A) - f(\tilde{A})) \leq \|\tilde{T}^{-1}\| \|T\| \max_{j,k} |f(\lambda_k) - f(\tilde{\lambda}_j)| \left[\sum_{j=1}^n \|\tilde{T}d_j\|^2 \sum_{k=1}^n \|(T^{-1})^*d_k\|^2 \right]^{1/2}. \tag{2.7}$$

3. Proof of Theorem 1.1. Let Q_j be the Riesz projection for μ_j :

$$Q_j = -\frac{1}{2\pi i} \int_{L_j} R_\lambda(A) d\lambda, \tag{3.1}$$

where $L_j := \{z \in \mathbb{C} : |z - \mu_j| = \delta_j\}$.

LEMMA 3.1. *The inequality*

$$\|Q_j\| \leq \sum_{k=0}^{n-1} \frac{g^k(A)}{\delta_j^k \sqrt{k!}}$$

is true.

Proof. By the Schur theorem, there is an orthonormal basis $\{e_k\}$, in which $A = D + V$, where $D = \text{diag}(\lambda_j)$ is a normal matrix (the diagonal part of A) and V is an upper triangular nilpotent matrix (the nilpotent part of A). Let P_j be the eigenprojection of D corresponding to μ_j . Then by (3.1),

$$Q_j - P_j = -\frac{1}{2\pi i} \int_{L_j} (R_\lambda(A) - R_\lambda(D)) d\lambda = \frac{1}{2\pi i} \int_{L_j} R_\lambda(A) V R_\lambda(D) d\lambda$$

since $V = A - D$. But

$$R_\lambda(A) = (D + V - I\lambda)^{-1} = (I + R_\lambda(D)V)^{-1} R_\lambda(D).$$

Consequently,

$$R_\lambda(A) = \sum_{k=0}^{n-1} (-1)^k (R_\lambda(D)V)^k R_\lambda(D).$$

So, we have

$$Q_j - P_j = \sum_{k=1}^{n-1} C_k, \tag{3.2}$$

where

$$C_k = (-1)^k \frac{1}{2\pi i} \int_{L_j} (R_\lambda(D)V)^k R_\lambda(D) d\lambda.$$

By Theorem 2.5.1 of [8], we get

$$\|(R_\lambda(D)V)^k\| \leq \frac{N_2^k(R_\lambda(D)V)}{\sqrt{k!}} \leq \frac{N_2^k(V)}{\rho^k(A, \lambda)\sqrt{k!}},$$

where $\rho(A, \lambda) = \min_k |\lambda - \lambda_k|$. In addition, directly from the definition of $g(A)$, when the Hilbert-Schmidt norm is calculated in the basis $\{e_k\}$, we have $N_2(V) = g(A)$. So,

$$\|(R_\lambda(D)V)^k\| \leq \frac{g^k(A)}{\delta_j^k \sqrt{k!}} \quad (\lambda \in L_j),$$

and therefore,

$$\|C_k\| \leq \frac{1}{2\pi} \int_{L_j} \|(R_\lambda(D)V)^k R_\lambda(D)\| |d\lambda| \leq \frac{g^k(A)}{\delta_j^{k+1} \sqrt{k!}} \frac{1}{2\pi} \int_{L_j} |d\lambda| = \frac{g^k(A)}{\delta_j^k \sqrt{k!}}.$$

Hence,

$$\|Q_j - P_j\| \leq \sum_{k=1}^{n-1} \|C_k\| \leq \sum_{k=1}^{n-1} \frac{g^k(A)}{\delta_j^k \sqrt{k!}}, \tag{3.3}$$

and therefore,

$$\|Q_j\| \leq \|P_j\| + \|Q_j - P_j\| \leq \sum_{k=0}^{n-1} \frac{g^k(A)}{\delta_j^k \sqrt{k!}},$$

as claimed. \square

Let $\{v_k\}_{k=1}^n$ be a sequence of the eigenvectors of A , and $\{u_k\}_{k=1}^n$ be the biorthogonal sequence: $(v_j, u_k) = 0$ ($j \neq k$), $(v_j, u_j) = 1$ ($j, k = 1, \dots, n$). So

$$A = \sum_{k=1}^m \mu_k Q_k = \sum_{k=1}^n \lambda_k(\cdot, u_k) v_k. \tag{3.4}$$

Rearranging these biorthogonal sequences, we can write

$$Q_j = \sum_{k=1}^{p_j} (\cdot, u_{jk}) v_{jk},$$

where $\{u_{jk}\}_{k=1}^{p_j}$ and $\{v_{jk}\}_{k=1}^{p_j}$ are biorthogonal: $(v_{jk}, u_{jm}) = 0$ ($m \neq k$), $(v_{jk}, u_{jk}) = 1$ ($m, k = 1, \dots, p_j$). Observe that we can always choose these systems so that $\|u_{jk}\| = \|v_{jk}\|$.

LEMMA 3.2. *Let $\|u_{jk}\| = \|v_{jk}\|$. Then we have*

$$\|u_{jk}\|^2 \leq \|Q_j\| \quad (k = 1, \dots, p_j).$$

Proof. Clearly, $Q_j u_{jk} = (Q_j u_{jk}, u_{jk}) v_{jk}$. So,

$$(Q_j u_{jk}, v_{jk}) = (u_{jk}, u_{jk})(v_{jk}, v_{jk}) = \|u_{jk}\|^4.$$

Hence, $\|u_{jk}\|^4 \leq \|Q_j\| \|u_{jk}\|^2$, as claimed. \square

Lemmas 3.1 and 3.2 imply:

COROLLARY 3.3. *The inequalities*

$$\|u_{js}\|^2 \leq \sum_{k=0}^{n-1} \frac{g^k(A)}{\delta_j^k \sqrt{k!}} \quad (s = 1, 2, \dots, p_j)$$

are valid.

Again, let $\{d_k\}$ be the standard orthonormal basis.

LEMMA 3.4. *Let A be an $n \times n$ diagonalizable matrix. Then the operator*

$$T = \sum_{k=1}^n (\cdot, d_k) v_k \tag{3.5}$$

has the inverse one defined by

$$T^{-1} = \sum_{k=1}^n (\cdot, u_k) d_k, \tag{3.6}$$

and (2.1) holds.

Proof. Indeed, we can write out

$$T^{-1}T = \sum_{j=1}^n d_j \sum_{k=1}^n (\cdot, d_k)(v_k, u_j) = \sum_{j=1}^n d_j (\cdot, d_j) = I$$

and

$$TT^{-1} = \sum_{k=1}^n (\cdot, u_k) \sum_{j=1}^n (d_k, d_j) v_j = \sum_{k=1}^n (\cdot, u_k) v_k = I.$$

Moreover,

$$AT = \sum_{j=1}^n v_j \lambda_j(\cdot, d_j) \quad \text{and} \quad S = \sum_{k=1}^n \lambda_k(\cdot, d_k) d_k.$$

Hence,

$$T^{-1}AT = \sum_{k=1}^n d_k \sum_{j=1}^n (v_j, u_k) \lambda_j(\cdot, d_j) = \sum_{j=1}^n d_j \lambda_j(\cdot, d_j) = S.$$

So, (2.1) really holds. \square

LEMMA 3.5. *Let T be defined by (3.5). Then*

$$\|T\|^2 \leq \sum_{j=1}^m p_j \|Q_j\| \quad \text{and} \quad \|T^{-1}\|^2 \leq \sum_{j=1}^m p_j \|Q_j\|,$$

and therefore,

$$\kappa_T \leq \sum_{j=1}^m p_j \|Q_j\|.$$

Proof. Due to the previous lemma

$$\|T^{-1}x\|^2 = \sum_{k=1}^n |(x, u_k)|^2 \leq \|x\|^2 \sum_{k=1}^n \|u_k\|^2 \quad (x \in \mathbb{C}^n). \quad (3.7)$$

By the Schwarz inequality,

$$\begin{aligned} (Tx, Tx) &= \sum_{k=1}^n \sum_{s=1}^n (x, d_k)(v_k, v_s)(x, d_s) \\ &\leq \sum_{k=1}^n |(x, d_k)|^2 \left[\sum_{s,k=1}^n |(v_k, v_s)|^2 \right]^{1/2} \leq \|x\|^2 \sum_{s=1}^n \|v_s\|^2. \end{aligned} \quad (3.8)$$

But by Lemma 3.2,

$$\sum_{s=1}^n \|u_s\|^2 = \sum_{s=1}^n \|v_s\|^2 = \sum_{j=1}^m \sum_{k=1}^{p_j} \|v_{jk}\|^2 \leq \sum_{j=1}^m p_j \|Q_j\|.$$

Now (3.7) and (3.8) yield the required result. \square

Lemmas 3.1 and 3.5 imply:

COROLLARY 3.6. *The inequality $\kappa_T \leq \beta(A)$ is true.*

LEMMA 3.7. *Under the hypotheses of Theorem 1.1, we have*

$$N_2(f(A) - f(\tilde{A})) \leq \|\tilde{T}^{-1}\| \|T\| \max_{j,k} |f(\lambda_k) - f(\tilde{\lambda}_j)| \left[\sum_{j=1}^{\tilde{m}} \tilde{p}_j \|\tilde{Q}_j\| \sum_{k=1}^m p_k \|Q_k\| \right]^{1/2},$$

where Q_k, \tilde{Q}_j are the eigenprojections of A and \tilde{A} , respectively.

Proof. By (3.6), we have

$$(T^{-1})^* = \sum_{k=1}^n (\cdot, d_k) u_k.$$

Moreover, due to Lemma 3.2,

$$\sum_{j=1}^n \|(T^{-1})^* d_j\|^2 = \sum_{k=1}^n \|u_k\|^2 \leq \sum_{k=1}^m p_k \|Q_k\|.$$

Similarly,

$$\sum_{k=1}^n \|\tilde{T} d_k\|^2 \leq \sum_{k=1}^m p_k \|\tilde{Q}_k\|.$$

Now (2.7) implies the required result. \square

Thanks to Lemmas 3.5 and 3.7 we get the inequality

$$N_2(f(A) - f(\tilde{A})) \leq \max_{j,k} |f(\lambda_k) - f(\tilde{\lambda}_j)| \sum_{j=1}^{\tilde{m}} \tilde{p}_j \|\tilde{Q}_j\| \sum_{k=1}^m p_k \|Q_k\|. \quad (3.9)$$

Proof of Theorem 1.1: Inequality (1.2) is due to (2.6) and Corollary 3.6. Inequality (1.3) is due to (3.9) and Lemma 3.1. \square

4. Applications. Let us consider the two differential equations

$$\frac{du}{dt} = Au \quad (t > 0) \quad \text{and} \quad \frac{d\tilde{u}}{dt} = \tilde{A}\tilde{u} \quad (t > 0),$$

whose solutions are $u(t)$ and $\tilde{u}(t)$, respectively. Let us take the initial conditions $u(0) = \tilde{u}(0) \in \mathbb{C}^n$. Then we have

$$u(t) - \tilde{u}(t) = (\exp [At] - \exp [\tilde{A}t])u(0).$$

By (1.2),

$$\|u(t) - \tilde{u}(t)\| \leq \|u(0)\| N_2(\exp [At] - \exp [\tilde{A}t])$$

$$\leq te^{t\alpha} N_2(A - \tilde{A})\beta(A)\beta(\tilde{A})\|u(0)\| \quad (t \geq 0),$$

where

$$\alpha = \max\left\{\max_k \operatorname{Re} \lambda_k, \max_j \operatorname{Re} \tilde{\lambda}_j\right\}.$$

Furthermore, let $0 \notin \sigma(A) \cup \sigma(\tilde{A})$. Then the function

$$A^{-1/2} \sin(A^{1/2}t) \quad (t > 0)$$

is the Green function to the Cauchy problem for the equation

$$\frac{d^2 w}{dt^2} + Aw = 0 \quad (t > 0).$$

Simultaneously, consider the equation

$$\frac{d^2 \tilde{w}}{dt^2} + \tilde{A}\tilde{w} = 0 \quad (t > 0),$$

and take the initial conditions

$$w(0) = \tilde{w}(0) \in \mathbb{C}^n \quad \text{and} \quad w'(0) = \tilde{w}'(0) = 0.$$

Then we have

$$w(t) - \tilde{w}(t) = [A^{-1/2} \sin(A^{1/2}t) - \tilde{A}^{-1/2} \sin(\tilde{A}^{1/2}t)]w(0).$$

By (1.2),

$$\|w(t) - \tilde{w}(t)\| \leq \beta(A)\beta(\tilde{A})N_2(A - \tilde{A}) \max_{j,k} \left| \frac{\frac{\sin(t\sqrt{\lambda_k})}{\sqrt{\lambda_k}} - \frac{\sin(t\sqrt{\tilde{\lambda}_j})}{\sqrt{\tilde{\lambda}_j}}}{\lambda_k - \tilde{\lambda}_j} \right| \quad (t > 0).$$

Here one can take an arbitrary branch of the roots.

Acknowledgment. I am very grateful to the referee of this paper for his really helpful remarks.

REFERENCES

- [1] R. Bhatia. *Matrix Analysis*. Springer, Berlin, 1997.
- [2] K.N. Boyadzhiev. Some inequalities for generalized commutators. *Publ. Res. Inst. Math. Sci.*, 26(3):521–527, 1990.

- [3] K. Du. Note on structured indefinite perturbations to Hermitian matrices. *J. Comput. Appl. Math.*, 202(2):258–265, 2007.
- [4] P. Forrester and E. Rains. Jacobians and rank 1 perturbations relating to unitary Hessenberg matrices. *Int. Math. Res. Not.* 2006, Article ID 48306, 2006.
- [5] B. Fritzsche, B. Kirstein, and A. Lasarow. On Schwarz-Pick-Potapov block matrices of matrix-valued functions which belong to the extended Potapov class. *Analysis (Munich)*, 22(3):243–263, 2002.
- [6] B. Fritzsche, B. Kirstein, and A. Lasarow. Orthogonal rational matrix-valued functions on the unit circle: Recurrence relations and a Favard-type theorem. *Math. Nachr.*, 279(5-6):513–542, 2006.
- [7] M.I. Gil'. Perturbations of simple eigenvectors of linear operators. *Manuscripta Mathematica*, 100:213–219, 1999.
- [8] M.I. Gil'. *Operator Functions and Localization of Spectra*. Lecture Notes In Mathematics, vol. 1830, Springer-Verlag, Berlin, 2003.
- [9] L. Han, M. Neumann, and M. Tsatsomeros. Spectral radii of fixed Frobenius norm perturbations of nonnegative matrices. *SIAM J. Matrix Anal. Appl.*, 21(1):79–92, 1999.
- [10] N.J. Higham. *Functions of Matrices: Theory and Computations*. Society for Industrial and Applied Mathematics, Philadelphia, 2008.
- [11] R. Hryniv and P. Lancaster. On the perturbation of analytic matrix functions. *Integral Equations Operator Theory*, 34(3):325–338, 1999.
- [12] T. Jiang, X. Cheng, and L. Chen. An algebraic relation between consimilarity and similarity of complex matrices and its applications. *J. Phys. A, Math. Gen.*, 39(29):9215–9222, 2006.
- [13] P. Lancaster, A.S. Markus, and F. Zhou. Perturbation theory for analytic matrix functions: The semisimple case. *SIAM J. Matrix Anal. Appl.*, 25(3):606–626, 2003.
- [14] H. Langer and B. Najman. Remarks on the perturbation of analytic matrix functions. III. *Integral Equations Operator Theory*, 15(5):796–806, 1992.
- [15] A. Lasarow. Dual Szego pairs of sequences of rational matrix-valued functions. *Int. J. Math. Math. Sci.* 2006, Article ID 23723, 2006.
- [16] A. Mikhailova, B. Pavlov, and L. Prokhorov. Intermediate Hamiltonian via Glazman's splitting and analytic perturbation for meromorphic matrix-functions. *Math. Nachr.*, 280(12):1376–1416, 2007.
- [17] A.M. Ostrowski. *Solution of Equations in Euclidean and Banach Spaces*. Academic Press, New York, 1973.
- [18] J.-G. Sun. Perturbation of the matrix sign function. *Linear Algebra Appl.*, 250:177–206, 1997.
- [19] L. Verde-Star. Functions of matrices. *Linear Algebra Appl.*, 406:285–300, 2005.