

## WEIGHTED-EP ELEMENTS IN $C^*$ -ALGEBRAS\*

DIJANA MOSIĆ<sup>†</sup> AND DRAGAN S. DJORDJEVIĆ<sup>†</sup>

**Abstract.** The weighted-EP elements in  $C^*$ -algebras are defined and characterized.

**Key words.** EP elements, Moore-Penrose inverse, Group inverse,  $C^*$ -algebra.

**AMS subject classifications.** 46L05, 47A05.

**1. Introduction.** There are many equivalent characterizations of EP elements in a ring or  $C^*$ -algebra (see, for example, [10, 19, 21, 23, 24, 27]), many more still for Banach or Hilbert space operators and matrices (see [1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 18, 22, 26]). In [30], Tian and Wang defined weighted-EP matrices and presented characterizations of weighted-EP matrices using various rank formulas for matrices. In this paper, weighted-EP elements of  $C^*$ -algebras are studied using different methods, extending the results from [30] to more general settings.

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra with the unit 1. An element  $a \in \mathcal{A}$  is *regular* if there exists some  $b \in \mathcal{A}$  satisfying  $aba = a$ . The set of all regular elements of  $\mathcal{A}$  is denoted by  $\mathcal{A}^-$ . An element  $a \in \mathcal{A}$  satisfying  $a^* = a$  is called *symmetric* (or *Hermitian*). An element  $x \in \mathcal{A}$  is positive if  $x = y^*y$  for some  $y \in \mathcal{A}$ . Alternatively,  $x \in \mathcal{A}$  is positive if  $x$  is Hermitian and  $\sigma(x) \subseteq [0, +\infty)$ , where the spectrum of element  $x$  is denoted by  $\sigma(x)$ .

An element  $a \in \mathcal{A}$  is *group invertible* if there exists  $a^\# \in \mathcal{A}$  such that

$$aa^\#a = a, \quad a^\#aa^\# = a^\#, \quad aa^\# = a^\#a.$$

Recall that  $a^\#$  is uniquely determined by these equations. The group inverse  $a^\#$  exists if and only if  $a\mathcal{A} = a^2\mathcal{A}$  and  $\mathcal{A}a = \mathcal{A}a^2$  if and only if  $a \in a^2\mathcal{A} \cap \mathcal{A}a^2$  (see [12, 28]). We use  $\mathcal{A}^\#$  to denote the set of all group invertible elements of  $\mathcal{A}$ . The group inverse  $a^\#$  double commutes with  $a$ , that is,  $ax = xa$  implies  $a^\#x = xa^\#$  [6, 11].

An element  $a^\dagger \in \mathcal{A}$  is the *Moore-Penrose inverse* (or *MP-inverse*) of  $a \in \mathcal{A}$ , if

---

\*Received by the editors on February 13, 2011. Accepted for publication on September 6, 2011.  
 Handling Editor: Michael Tsatsomeros.

<sup>†</sup>Faculty of Sciences and Mathematics, University of Niš, PO Box 224, 18000 Niš, Serbia (dijana@pmf.ni.ac.rs, dragan@pmf.ni.ac.rs). The authors are supported by the Ministry of Science, Republic of Serbia, grant no. 174007.

the following hold [29]:

$$aa^\dagger a = a, \quad a^\dagger aa^\dagger = a^\dagger, \quad (aa^\dagger)^* = aa^\dagger, \quad (a^\dagger a)^* = a^\dagger a.$$

There is at most one  $a^\dagger$  such that above conditions hold (see [14, 17]). The set of all Moore–Penrose invertible elements of  $\mathcal{A}$  will be denoted by  $\mathcal{A}^\dagger$ .

**THEOREM 1.1.** [14] *In a unital  $C^*$ -algebra  $\mathcal{A}$ ,  $a \in \mathcal{A}$  is MP-invertible if and only if  $a$  is regular.*

**DEFINITION 1.2.** Let  $\mathcal{A}$  be a unital  $C^*$ -algebra, and let  $e$  and  $f$  be invertible positive elements in  $\mathcal{A}$ . The element  $a \in \mathcal{A}$  has the *weighted MP-inverse* with weights  $e$  and  $f$  if there exists  $b \in \mathcal{A}$  such that

$$aba = a, \quad bab = b, \quad (eab)^* = eab, \quad (fba)^* = fba.$$

The unique weighted MP-inverse with weights  $e$  and  $f$  will be denoted by  $a_{e,f}^\dagger$  if it exists [6]. The set of all weighted MP-invertible elements of  $\mathcal{A}$  with weights  $e$  and  $f$  will be denoted by  $\mathcal{A}_{e,f}^\dagger$ .

**THEOREM 1.3.** [6] *Let  $\mathcal{A}$  be a unital  $C^*$ -algebra and let  $e$  and  $f$  be positive invertible elements of  $\mathcal{A}$ . If  $a \in \mathcal{A}$  is regular, then the unique weighted MP-inverse  $a_{e,f}^\dagger$  exists and*

$$a_{e,f}^\dagger = f^{-1/2}(e^{1/2}af^{-1/2})^\dagger e^{1/2}.$$

Define the mapping  $x \mapsto x^{*,e,f} = e^{-1}x^*f$ , for all  $x \in \mathcal{A}$ . Notice that  $(*,e,f) : \mathcal{A} \rightarrow \mathcal{A}$  is not an involution, because in general  $(xy)^{*,e,f} \neq y^{*,e,f}x^{*,e,f}$ . Now, we formulate the following result which can be proved directly by the definition of the weighted MP-inverse.

**THEOREM 1.4.** *Let  $\mathcal{A}$  be a unital  $C^*$ -algebra and let  $e$  and  $f$  be positive invertible elements of  $\mathcal{A}$ . For any  $a \in \mathcal{A}^-$ , the following hold:*

- (a)  $(a_{e,f}^\dagger)_{f,e}^\dagger = a$ ;
- (b)  $(a^{*,f,e})_{f,e}^\dagger = (a_{e,f}^\dagger)^{*,e,f}$ ;
- (c)  $a^{*,f,e} = a_{e,f}^\dagger aa^{*,f,e} = a^{*,f,e} aa_{e,f}^\dagger$ ;
- (d)  $a^{*,f,e}(a_{e,f}^\dagger)^{*,e,f} = a_{e,f}^\dagger a$ ;
- (e)  $(a_{e,f}^\dagger)^{*,e,f} a^{*,f,e} = aa_{e,f}^\dagger$ ;
- (f)  $(a^{*,f,e}a)_{f,f}^\dagger = a_{e,f}^\dagger (a_{e,f}^\dagger)^{*,e,f}$ ;
- (g)  $(aa^{*,f,e})_{e,e}^\dagger = (a_{e,f}^\dagger)^{*,e,f} a_{e,f}^\dagger$ ;
- (h)  $a_{e,f}^\dagger = (a^{*,f,e}a)_{f,f}^\dagger a^{*,f,e} = a^{*,f,e}(aa^{*,f,e})_{e,e}^\dagger$ ; and

$$(i) \ (a^{*e,f})_{f,e}^\dagger = a(a^{*f,e}a)_{f,f}^\dagger = (aa^{*f,e})_{e,e}^\dagger a.$$

For  $a \in \mathcal{A}$ , consider two annihilators

$$a^\circ = \{x \in \mathcal{A} : ax = 0\}, \quad {}^\circ a = \{x \in \mathcal{A} : xa = 0\}.$$

Note that

$$(a^*)^\circ = a^\circ \Leftrightarrow {}^\circ(a^*) = {}^\circ a, \quad a\mathcal{A} = a^*\mathcal{A} \Leftrightarrow \mathcal{A}a = \mathcal{A}a^*.$$

LEMMA 1.5. [10] For  $a \in \mathcal{A}$ ,  $a \in \mathcal{A}^- \Leftrightarrow \mathcal{A} = (a^*\mathcal{A}) \oplus a^\circ$ .

The following result is very useful in the rest of paper and can be verified by properties of the weighted MP-inverse.

LEMMA 1.6. Let  $a \in \mathcal{A}^-$ , and let  $e$  and  $f$  be invertible positive elements in  $\mathcal{A}$ . Then

- (a)  $a_{e,f}^\dagger \mathcal{A} = a_{e,f}^\dagger a \mathcal{A} = f^{-1} a^* \mathcal{A}$ ;
- (b)  $(a_{e,f}^\dagger)^* \mathcal{A} = (aa_{e,f}^\dagger)^* \mathcal{A} = ea \mathcal{A}$ ;
- (c)  $a^\circ = (ea)^\circ$ ;
- (d)  $(a^*)^\circ = (f^{-1} a^*)^\circ$ ;
- (e)  $(a_{e,f}^\dagger)^\circ = [(ea)^*]^\circ$ ; and
- (f)  $[(a_{e,f}^\dagger)^*]^\circ = (af^{-1})^\circ$ ;

Now, we state an important result related to the weighted Moore-Penrose inverse. In [19, Lemma 1.5], the following result is proved for the ordinary Moore-Penrose inverse. Observe that conditions (1.1) and (1.2) also appear in the proof of [15, Theorem 10] for the ordinary Moore-Penrose inverse.

LEMMA 1.7. Let  $a \in \mathcal{A}^-$ , and let  $e$  and  $f$  be invertible positive elements in  $\mathcal{A}$ . Then

$$(1.1) \quad a_{e,f}^\dagger = (a^{*f,e}a + 1 - a_{e,f}^\dagger a) a^{*f,e} = a^{*f,e} (aa^{*f,e} + 1 - aa_{e,f}^\dagger)^{-1},$$

$$(1.2) \quad a^{*f,e} \mathcal{A}^{-1} = a_{e,f}^\dagger \mathcal{A}^{-1} \text{ and } \mathcal{A}^{-1} a^{*f,e} = \mathcal{A}^{-1} a_{e,f}^\dagger,$$

$$(1.3) \quad (a^{*f,e})^\circ = (a_{e,f}^\dagger)^\circ \text{ and } {}^\circ(a^{*f,e}) = {}^\circ(a_{e,f}^\dagger).$$

*Proof.* By Theorem 1.4, we can verify

$$a^{*f,e} = (a^{*f,e}a + 1 - a_{e,f}^\dagger a) a_{e,f}^\dagger = a_{e,f}^\dagger (aa^{*f,e} + 1 - aa_{e,f}^\dagger),$$

$$(a^{*f,e}a + 1 - a_{e,f}^\dagger a)^{-1} = a_{e,f}^\dagger (a_{e,f}^\dagger)^{*e,f} + 1 - a_{e,f}^\dagger a$$

and

$$(aa^{*f,e} + 1 - aa_{e,f}^\dagger)^{-1} = (a_{e,f}^\dagger)^{*e,f} a_{e,f}^\dagger + 1 - aa_{e,f}^\dagger.$$

Thus, the part (1.1) holds and it implies the equalities (1.2) and (1.3).  $\square$

We recall the definition of EP elements.

DEFINITION 1.8. An element  $a \in \mathcal{A}^-$  is EP if  $aa^\dagger = a^\dagger a$ .

LEMMA 1.9. [19] An element  $a \in \mathcal{A}$  is EP, if  $a \in \mathcal{A}^-$  and  $a\mathcal{A} = a^*\mathcal{A}$  (or, equivalently, if  $a \in \mathcal{A}^-$  and  $a^\circ = (a^*)^\circ$ ).

The condition  $a\mathcal{A} = a^*\mathcal{A}$  gave the EP elements their name for *equal projections* onto the range of  $a$  and  $a^*$  in the case of matrices and closed range Hilbert space operators. These elements are important since they are characterized by commutativity with their Moore–Penrose inverse. Also notice that EP elements are those elements for which the group and the Moore–Penrose inverse exist and coincide.

In this paper, as an extension of EP elements, we are concerned with elements of a  $C^*$ -algebra which commute with their weighted Moore–Penrose inverse. These elements are called weighted-EP elements. In particular, we give several equivalent conditions for an element of  $C^*$ -algebra to be weighted-EP. The motivation for this paper is an interesting paper by Tian and Wang [30]. They studied such characterizations for weighted-EP complex square matrices.

We conclude this section with the following results on the reverse order law for the Moore–Penrose inverse of a product, which will be used later.

LEMMA 1.10. [25, Theorem 2.4] Let  $\mathcal{A}$  be a unital  $C^*$ -algebra and let  $a, b, ab \in \mathcal{A}^-$ . Then the following conditions are equivalent:

- (a)  $(ab)^\dagger = b^\dagger a^\dagger$ ;
- (b)  $a^*ab = bb^\dagger a^*ab$  and  $abb^* = abb^* a^\dagger a$ .

THEOREM 1.11. Let  $\mathcal{A}$  be a unital  $C^*$ -algebra and let  $a, b, ab \in \mathcal{A}^-$ . Then  $(ab)^\dagger = b^\dagger a^\dagger$  if and only if  $a^*ab\mathcal{A} \subseteq b\mathcal{A}$  and  $bb^*a^*\mathcal{A} \subseteq a^*\mathcal{A}$ .

*Proof.*  $\implies$ : If  $(ab)^\dagger = b^\dagger a^\dagger$ , then, by Lemma 1.10,  $a^*ab = bb^\dagger a^*ab$  and  $abb^* = abb^* a^\dagger a$  which imply

$$bb^*a^* = (abb^*) = (abb^* a^\dagger a)^* = a^*(a^\dagger)^* bb^* a^*.$$

Hence,  $a^*ab\mathcal{A} \subseteq b\mathcal{A}$  and  $bb^*a^*\mathcal{A} \subseteq a^*\mathcal{A}$ .

$\Leftarrow$ : Conversely, from  $a^*ab\mathcal{A} \subseteq b\mathcal{A}$  and  $bb^*a^*\mathcal{A} \subseteq a^*\mathcal{A}$ , we conclude that  $a^*ab = bx$ , for some  $x \in \mathcal{A}$ , and  $bb^*a^* = a^*y$ , for some  $y \in \mathcal{A}$ . Then the equalities

$$bb^\dagger a^*ab = bb^\dagger bx = bx = a^*ab$$

and

$$abb^*a^\dagger a = (a^\dagger abb^*a^*)^* = (a^\dagger aa^*y)^* = (a^*y)^* = (bb^*a^*)^* = abb^*$$

imply  $(ab)^\dagger = b^\dagger a^\dagger$ , by Lemma 1.10.  $\square$

**2. Weighted-EP elements in  $C^*$ -algebras.** First, we state the definition of weighted-EP elements in  $C^*$ -algebras.

**DEFINITION 2.1.** An element  $a \in \mathcal{A}$  is said to be *weighted-EP* with respect to two invertible positive elements  $e, f \in \mathcal{A}$  (or weighted-EP w.r.t.  $(e, f)$ ) if both  $ea$  and  $af^{-1}$  are EP, that is,  $a \in \mathcal{A}^-$ ,  $ea\mathcal{A} = (ea)^*\mathcal{A}$ , and  $af^{-1}\mathcal{A} = (af^{-1})^*\mathcal{A}$ .

In the following theorem, a number of necessary and sufficient conditions for an element to be weighted-EP are presented.

**THEOREM 2.2.** Let  $\mathcal{A}$  be a unital  $C^*$ -algebra, and let  $e$  and  $f$  be invertible positive elements in  $\mathcal{A}$ . For  $a \in \mathcal{A}^-$  the following statements are equivalent:

- (I)  $a$  is weighted-EP w.r.t.  $(e, f)$ ;
- (II)  $a$  is weighted-EP w.r.t.  $(f, e)$ ;
- (III)  $a$  is both weighted-EP w.r.t.  $(e, e)$  and w.r.t.  $(f, f)$ ;
- (IV)  $ea\mathcal{A} = fa\mathcal{A} = a^*\mathcal{A}$ ;
- (V)  $e^{-1}a^*\mathcal{A} = f^{-1}a^*\mathcal{A} = a\mathcal{A}$ ;
- (VI)  $a_{e,f}^\dagger \mathcal{A} = a\mathcal{A}$  and  $(a_{e,f}^\dagger)^*\mathcal{A} = a^*\mathcal{A}$ ;
- (VII)  $a^*$  is weighted-EP w.r.t.  $(e^{-1}, f^{-1})$ ;
- (VIII)  $aa_{e,f}^\dagger = a_{e,f}^\dagger a$ ;
- (IX)  $a \in \mathcal{A}^\#$  and  $a^k = a_{e,f}^\dagger a a^k = a^k a a_{e,f}^\dagger$ , for any/some integer  $k \geq 1$ ;
- (X)  $a_{e,f}^\dagger = a(a_{e,f}^\dagger)^2 = (a_{e,f}^\dagger)^2 a$ ;
- (XI)  $a \in \mathcal{A}^\#$  and  $a^\# = a_{e,f}^\dagger$ ;
- (XII)  $a \in \mathcal{A}^\#$  and both  $ea a^\#$  and  $f a a^\#$  are Hermitian;
- (XIII)  $a \in \mathcal{A}^\#$  and  $a^\# a_{e,f}^\dagger = a_{e,f}^\dagger a^\#$ ;
- (XIV)  $a \in \mathcal{A}^\#$  and  $aa^\# a_{e,f}^\dagger = a_{e,f}^\dagger a^\# a$ ;
- (XV)  $a \in a_{e,f}^\dagger \mathcal{A}^{-1} \cap \mathcal{A}^{-1} a_{e,f}^\dagger$ ;
- (XVI)  $a \in a_{e,f}^\dagger \mathcal{A} \cap \mathcal{A} a_{e,f}^\dagger$ ;
- (XVII)  $a\mathcal{A}^{-1} = f^{-1}a^*\mathcal{A}^{-1}$  and  $\mathcal{A}^{-1}a = \mathcal{A}^{-1}a^*e$ ;
- (xviii)  $\mathcal{A}^{-1}a^* = \mathcal{A}^{-1}af^{-1}$  and  $a^*\mathcal{A}^{-1} = ea\mathcal{A}^{-1}$ ;
- (XIX) there exists  $x \in \mathcal{A}$  such that  $a = e^{-1}a^*xa^*f$ ;

- (XX)  $a = (ae^{-1})^\dagger ae^{-1}afa(fa)^\dagger$ ;  
(XXI)  $a \in \mathcal{A}^\#$  and  $a^k$  is weighted-EP w.r.t.  $(e, f)$ , for any/some integer  $k \geq 1$ ;  
(XXII)  $aa^*a$  is weighted-EP w.r.t.  $(e, f)$ ;  
(XXIII)  $a^\circ = [(ea)^*]^\circ$  and  $(a^*)^\circ = (af^{-1})^\circ$ ;  
(XXIV)  $\mathcal{A} = e^{-1}a^*\mathcal{A} \oplus (a^*)^\circ = a^*\mathcal{A} \oplus (a^*f)^\circ$ ;  
(XXV)  $a^\dagger$  is weighted-EP w.r.t.  $(e^{-1}, f^{-1})$ ;  
(XXVI)  $a_{e,f}^\dagger$  is weighted-EP w.r.t.  $(e, f)$ ;  
(XXVII)  $a \in \mathcal{A}^\#$  and  $a^{2k-1} = a_{e,f}^\dagger a^{2k+1} a_{e,f}^\dagger$ , for any/some integer  $k \geq 1$ ;  
(XXVIII)  $a \in \mathcal{A}^\#$  and  $aa_{e,f}^\dagger a_{e,f}^\dagger a = a_{e,f}^\dagger aaa_{e,f}^\dagger$ ;  
(XXIX)  $a \in \mathcal{A}^\#$  and  $a^\#$  is weighted-EP w.r.t.  $(e, f)$ ;  
(XXX)  $a \in \mathcal{A}^\#$  and  $aa^\# = aa_{e,e}^\dagger = aa_{f,f}^\dagger$  (or  $aa^\# = a_{e,e}^\dagger a = a_{f,f}^\dagger a$ );  
(XXXI)  $a \in \mathcal{A}^\#$  and  $aa^\# = aa_{e,f}^\dagger = aa_{f,e}^\dagger$  (or  $aa^\# = a_{f,e}^\dagger a = a_{e,f}^\dagger a$ );  
(XXXII)  $a \in \mathcal{A}^\#$ ,  $aa_{e,e}^\dagger e^{-1}a^*a = e^{-1}a^*aaa_{e,e}^\dagger$  and  $aa_{f,f}^\dagger f^{-1}a^*a = f^{-1}a^*aaa_{f,f}^\dagger$ ;  
(XXXIII)  $a \in \mathcal{A}^\#$ ,  $aa_{e,f}^\dagger e^{-1}a^*a = e^{-1}a^*aaa_{e,f}^\dagger$  and  $aa_{f,e}^\dagger f^{-1}a^*a = f^{-1}a^*aaa_{f,e}^\dagger$ ;  
(XXXIV)  $a \in \mathcal{A}^\#$ ,  $a_{e,e}^\dagger aaa^*e = aa^*ea_{e,e}^\dagger a$  and  $a_{f,f}^\dagger aaa^*f = aa^*fa_{f,f}^\dagger a$ ;  
(XXXV)  $a \in \mathcal{A}^\#$ ,  $a_{f,e}^\dagger aaa^*e = aa^*ea_{f,e}^\dagger a$  and  $a_{e,f}^\dagger aaa^*f = aa^*fa_{e,f}^\dagger a$ ;  
(XXXVI)  $a \in \mathcal{A}^\#$  and  $a^k aa_{e,f}^\dagger + a_{e,f}^\dagger aa^k = 2a^k$ , for any/some integer  $k \geq 1$ ;  
(XXXVII)  $a \in \mathcal{A}^\#$  and  $a_{e,f}^\dagger a^\# a + aa^\# a_{e,f}^\dagger = 2a_{e,f}^\dagger$ ;  
(XXXVIII)  $a \in \mathcal{A}^\#$  and  $a^{*,f,e} = a^{*,f,e} aa^\# = a^\# aa^{*,f,e}$ ;  
(XXXIX)  $a \in \mathcal{A}^\#$  and  $a^{*,f,e} aa^\# + a^\# aa^{*,f,e} = 2a^{*,f,e}$ ;  
(XL)  $a \in \mathcal{A}^\#$  and  $a^k aa_{e,f}^\dagger + (a^k aa_{e,f}^\dagger)^* = a_{e,f}^\dagger aa^k + (a_{e,f}^\dagger aa^k)^* = a^k + (a^k)^*$ , for any/some integer  $k \geq 1$ ;  
(XLI)  $aa_{e,f}^\dagger(a + \lambda a_{e,f}^\dagger) = (a + \lambda a_{e,f}^\dagger)aa_{e,f}^\dagger$  and  $a_{e,f}^\dagger a(a + \lambda a_{e,f}^\dagger) = (a + \lambda a_{e,f}^\dagger)a_{e,f}^\dagger a$ , for any/some complex number  $\lambda \neq 0$ ;  
(XLII)  $ab = ba \Rightarrow a_{e,f}^\dagger b = ba_{e,f}^\dagger$ ;  
(XLIII)  $a_{e,f}^\dagger = f(a)$ , for some function  $f$  holomorphic in a neighbourhood of  $\sigma(a)$ ;  
(XLIV)  $(a + \lambda a_{e,e}^\dagger)\mathcal{A} = (a + \lambda a_{f,f}^\dagger)\mathcal{A} = (\lambda a + a^3)\mathcal{A}$  and  $\mathcal{A}(a + \lambda a_{e,e}^\dagger) = \mathcal{A}(a + \lambda a_{f,f}^\dagger) = \mathcal{A}(\lambda a + a^3)$ , for any/some complex number  $\lambda \neq 0$ ;  
(XLV)  $(a + \lambda a_{e,f}^\dagger)\mathcal{A} = (\lambda a + a^3)\mathcal{A}$  and  $\mathcal{A}(a + \lambda a_{e,f}^\dagger) = \mathcal{A}(\lambda a + a^3)$ , for any/some complex number  $\lambda \neq 0$ ;  
(XLVI)  $(a + \lambda a_{e,e}^\dagger)^\circ = (a + \lambda a_{f,f}^\dagger)^\circ = (\lambda a + a^3)^\circ$  and  $^\circ(a + \lambda a_{e,e}^\dagger) = ^\circ(a + \lambda a_{f,f}^\dagger) = ^\circ(\lambda a + a^3)$ , for any/some complex number  $\lambda \neq 0$ ;  
(XLVII)  $(a + \lambda a_{e,f}^\dagger)^\circ = (\lambda a + a^3)^\circ$  and  $^\circ(a + \lambda a_{f,e}^\dagger) = ^\circ(\lambda a + a^3)$ , for any/some complex number  $\lambda \neq 0$ ;  
(XLVIII)  $a \in \mathcal{A}^\#$  and  $(a_{e,f}^\dagger)^2 a^\# = a_{e,f}^\dagger a^\# a_{e,f}^\dagger = a^\# (a_{e,f}^\dagger)^2$ ;  
(XLIX)  $a \in \mathcal{A}^\#$  and  $a(a_{e,f}^\dagger)^2 = a^\# = (a_{e,f}^\dagger)^2 a$ ;  
(L)  $a \in \mathcal{A}^\#$ ,  $a^{*,f,e} a_{e,f}^\dagger = a^{*,f,e} a^\#$  and  $a_{e,f}^\dagger a^{*,f,e} = a^\# a^{*,f,e}$ ;  
(LI)  $a \in \mathcal{A}^\#$  and  $(a_{e,f}^\dagger)^2 = (a^\#)^2$ ;  
(LII)  $a^{*,e,f} = a^{*,e,f} a_{e,f}^\dagger a = aa_{e,f}^\dagger a^{*,e,f}$ ;

- (LIII)  $a \in \mathcal{A}^\#$  and  $(a^\#)^{*,f} = aa^\#(a^\#)^{*,f} = (a^\#)^{*,f}a^\#a$  (or  $(a^\#)^{*,f,e} = aa^\#(a^\#)^{*,f,e} = (a^\#)^{*,f,e}a^\#a$ );
- (LIV)  $a \in \mathcal{A}^\#$  and  $a_{e,f}^\dagger(a^\#)^2 = (a^\#)^2a_{e,f}^\dagger$ ;
- (LV)  $a \in \mathcal{A}^\#$  and  $a^ka_{e,f}^\dagger = a_{e,f}^\dagger a^k$ , for any/some integer  $k \geq 1$ ;
- (LVI)  $aa_{e,f}^\dagger(a + \lambda a^{*,f}) = (a + \lambda a^{*,f})aa_{e,f}^\dagger$  and  $a_{e,f}^\dagger a(a + \lambda a^{*,f}) = (a + \lambda a^{*,f})a_{e,f}^\dagger a$ , for any/some complex number  $\lambda \neq 0$ ;
- (LVII)  $a \in \mathcal{A}^\#$ ,  $aa_{e,e}^\dagger(aa^*e - e^{-1}a^*a) = (aa^*e - e^{-1}a^*a)aa_{e,e}^\dagger$  and  $aa_{f,f}^\dagger(aa^*f - f^{-1}a^*a) = (aa^*f - f^{-1}a^*a)aa_{f,f}^\dagger$ ;
- (LVIII)  $a \in \mathcal{A}^\#$ ,  $aa_{e,f}^\dagger(aa^*e - e^{-1}a^*a) = (aa^*e - e^{-1}a^*a)aa_{e,f}^\dagger$  and  $aa_{f,e}^\dagger(aa^*f - f^{-1}a^*a) = (aa^*f - f^{-1}a^*a)aa_{f,e}^\dagger$ ;
- (LIX)  $a \in \mathcal{A}^\#$ ,  $a_{e,e}^\dagger a(aa^*e - e^{-1}a^*a) = (aa^*e - e^{-1}a^*a)a_{e,e}^\dagger a$  and  $a_{f,f}^\dagger a(aa^*f - f^{-1}a^*a) = (aa^*f - f^{-1}a^*a)a_{f,f}^\dagger a$ ;
- (LX)  $a \in \mathcal{A}^\#$ ,  $a_{f,e}^\dagger a(aa^*e - e^{-1}a^*a) = (aa^*e - e^{-1}a^*a)a_{f,e}^\dagger a$  and  $a_{e,f}^\dagger a(aa^*f - f^{-1}a^*a) = (aa^*f - f^{-1}a^*a)a_{e,f}^\dagger a$ ;
- (LXI)  $a \in \mathcal{A}^\#$  and  $(a^{s+t})^\dagger = (a^s)_{e,1}^\dagger(a^t)_{1,e}^\dagger = (a^s)_{f,1}^\dagger(a^t)_{1,f}^\dagger$ , for any/some integers  $s, t \geq 1$ ;
- (LXII)  $a \in \mathcal{A}^\#$  and  $(a^{s+t})_{e,f}^\dagger = (a^s)_{f,f}^\dagger(a^t)_{e,f}^\dagger = (a^s)_{e,f}^\dagger(a^t)_{e,e}^\dagger$ , for any/some integers  $s, t \geq 1$ .

*Proof.* (I)  $\Rightarrow$  (II): Assume that  $a$  is weighted-EP w.r.t.  $(e, f)$ , i.e.,  $ea$  and  $af^{-1}$  are EP. From  $ea\mathcal{A} = (ea)^*\mathcal{A}$ , we obtain  $a\mathcal{A} = e^{-1}a^*e\mathcal{A} = (ae^{-1})^*e\mathcal{A}$  implying  $ae^{-1}e\mathcal{A} = (ae^{-1})^*e\mathcal{A}$ , that is,  $ae^{-1}\mathcal{A} = (ae^{-1})^*\mathcal{A}$ . In the same way  $af^{-1}\mathcal{A} = (af^{-1})^*\mathcal{A}$  implies  $fa\mathcal{A} = (fa)^*\mathcal{A}$ . Hence,  $fa$  and  $ae^{-1}$  are EP, i.e.,  $a$  is weighted-EP w.r.t.  $(f, e)$ .

(II)  $\Rightarrow$  (I): This implication can be proved in the same way as (I)  $\Rightarrow$  (II).

(III)  $\Leftrightarrow$  (I): Obviously, because (I)  $\Leftrightarrow$  (II).

(IV)  $\Leftrightarrow$  (I): Notice that  $(ea)^*\mathcal{A} = a^*\mathcal{A}$  and  $af^{-1}\mathcal{A} = a\mathcal{A}$ . Now,  $ea\mathcal{A} = fa\mathcal{A} = a^*\mathcal{A}$  is equivalent to  $ea\mathcal{A} = a^*e\mathcal{A}$  and  $a\mathcal{A} = f^{-1}a^*\mathcal{A}$ , that is,  $ea\mathcal{A} = (ea)^*\mathcal{A}$  and  $af^{-1}\mathcal{A} = (af^{-1})^*\mathcal{A}$ . These equalities mean that  $ea$  and  $af^{-1}$  are EP, i.e.,  $a$  is weighted-EP w.r.t.  $(e, f)$ .

(V)  $\Leftrightarrow$  (IV): This is easy to check.

(VI)  $\Leftrightarrow$  (I): By Lemma 1.6,  $a_{e,f}^\dagger\mathcal{A} = a\mathcal{A}$  and  $(a_{e,f}^\dagger)^*\mathcal{A} = a^*\mathcal{A}$  is equivalent to  $f^{-1}a^*\mathcal{A} = a\mathcal{A}$  and  $ea\mathcal{A} = a^*\mathcal{A}$  which is  $(af^{-1})^*\mathcal{A} = af^{-1}\mathcal{A}$  and  $ea\mathcal{A} = (ea)^*\mathcal{A}$ .

(VII)  $\Leftrightarrow$  (V): Using the equivalence (I)  $\Leftrightarrow$  (V) for  $a^*$ , we have that  $a^*$  is weighted-EP w.r.t.  $(e^{-1}, f^{-1})$  if and only if  $e^{-1}a^*\mathcal{A} = f^{-1}a^*\mathcal{A} = a\mathcal{A}$ .

(VIII)  $\Rightarrow$  (VI): The equality  $aa_{e,f}^\dagger = a_{e,f}^\dagger a$  gives

$$a\mathcal{A} = a_{e,f}^\dagger aa\mathcal{A} \subset a_{e,f}^\dagger a\mathcal{A} = aa_{e,f}^\dagger a_{e,f}^\dagger \mathcal{A} \subset a\mathcal{A},$$

and

$$a^*\mathcal{A} = (aaa_{e,f}^\dagger)^*\mathcal{A} \subset (a_{e,f}^\dagger)^*\mathcal{A} = (a_{e,f}^\dagger a_{e,f}^\dagger a)^*\mathcal{A} \subset a^*\mathcal{A},$$

i.e.,  $a_{e,f}^\dagger \mathcal{A} = a\mathcal{A}$  and  $(a_{e,f}^\dagger)^*\mathcal{A} = a^*\mathcal{A}$ .

(VI)  $\Rightarrow$  (IX): Since  $a_{e,f}^\dagger \mathcal{A} = a\mathcal{A}$ , then  $a = a_{e,f}^\dagger y$  for some  $y \in \mathcal{A}$ . Now,

$$a = a_{e,f}^\dagger y = a_{e,f}^\dagger a(a_{e,f}^\dagger y) = a_{e,f}^\dagger aa,$$

and  $a^k = a_{e,f}^\dagger aa^k$ , for any/some integer  $k \geq 1$ . Similarly, the assumption  $(a_{e,f}^\dagger)^*\mathcal{A} = a^*\mathcal{A}$  implies  $a^* = (a_{e,f}^\dagger)^*x$  for some  $x \in \mathcal{A}$  and

$$a^* = (a_{e,f}^\dagger)^*x = (a_{e,f}^\dagger aa_{e,f}^\dagger)^*x = (aa_{e,f}^\dagger)^*(a_{e,f}^\dagger)^*x = (aa_{e,f}^\dagger)^*a^* = (aaa_{e,f}^\dagger)^*.$$

Applying involution to this equality, we get  $a = aaa_{e,f}^\dagger$  and, for any/some integer  $k \geq 1$ ,  $a^k = a^k aa_{e,f}^\dagger$ . Notice that, from  $a \in a^2\mathcal{A} \cap \mathcal{A}a^2$ , it follows  $a^\#$  exists.

(IX)  $\Rightarrow$  (VIII): If  $a \in \mathcal{A}^\#$  and  $a^k = a_{e,f}^\dagger aa^k = a^k aa_{e,f}^\dagger$ , for any/some integer  $k \geq 1$ , then

$$aa_{e,f}^\dagger = (a^\#)^k (a^{k+1} a_{e,f}^\dagger) = (a^\#)^k a^k = a^k (a^\#)^k = a_{e,f}^\dagger a^{k+1} (a^\#)^k = a_{e,f}^\dagger a.$$

(X)  $\Rightarrow$  (VIII): Applying the equality  $a_{e,f}^\dagger = a(a_{e,f}^\dagger)^2 = (a_{e,f}^\dagger)^2 a$ , we obtain  $a_{e,f}^\dagger a = a((a_{e,f}^\dagger)^2 a) = aa_{e,f}^\dagger$ .

(VIII)  $\Rightarrow$  (X): Obviously.

(VIII)  $\Leftrightarrow$  (XI)  $\Leftrightarrow$  (XII): By the uniquely determined group and weighted-MP inverse.

(XIII)  $\Rightarrow$  (VIII): The hypothesis  $a^\# a_{e,f}^\dagger = a_{e,f}^\dagger a^\#$  implies

$$aa_{e,f}^\dagger = a^2(a^\# a_{e,f}^\dagger) = a^2 a_{e,f}^\dagger a^\# = a^2 a_{e,f}^\dagger a(a^\#)^2 = a^2 (a^\#)^2 = aa^\#$$

and

$$a_{e,f}^\dagger a = (a_{e,f}^\dagger a^\#)a^2 = a^\# a_{e,f}^\dagger a^2 = (a^\#)^2 aa_{e,f}^\dagger a^2 = a^\# a.$$

Therefore,  $aa_{e,f}^\dagger = a_{e,f}^\dagger a$ .



(XI)  $\Rightarrow$  (XIII): From the equality  $a^\# = a_{e,f}^\dagger$ , we have  $a^\# a_{e,f}^\dagger = (a_{e,f}^\dagger)^2 = a_{e,f}^\dagger a^\#$ . So, the condition (XIII) holds.

(XIV)  $\Rightarrow$  (VIII): Suppose that  $a \in \mathcal{A}^\#$  and  $aa^\# a_{e,f}^\dagger = a_{e,f}^\dagger a^\# a$ . Then we get the equality (VIII):

$$\begin{aligned} aa_{e,f}^\dagger &= a(aa^\# a_{e,f}^\dagger) = aa_{e,f}^\dagger a^\# a = aa_{e,f}^\dagger aa^\# = aa^\# \\ &= a^\# aa_{e,f}^\dagger a = (aa^\# a_{e,f}^\dagger) a = a_{e,f}^\dagger a^\# aa = a_{e,f}^\dagger a. \end{aligned}$$

(XI)  $\Rightarrow$  (XIV): By  $a^\# = a_{e,f}^\dagger$ , obviously, the condition (XIV) is satisfied.

(VIII)  $\Rightarrow$  (XV): Using  $aa_{e,f}^\dagger = a_{e,f}^\dagger a$ , we can verify that  $a = (a^2 + 1 - a_{e,f}^\dagger a) a_{e,f}^\dagger$  and  $(a^2 + 1 - a_{e,f}^\dagger a)^{-1} = (a_{e,f}^\dagger)^2 + 1 - a_{e,f}^\dagger a$ . Thus,  $a \in \mathcal{A}^{-1} a_{e,f}^\dagger$ . Since  $aa_{e,f}^\dagger = a_{e,f}^\dagger a$ , by previous equalities, we conclude that  $a = a_{e,f}^\dagger (a^2 + 1 - aa_{e,f}^\dagger)$  and  $(a^2 + 1 - aa_{e,f}^\dagger)^{-1} = (a_{e,f}^\dagger)^2 + 1 - aa_{e,f}^\dagger$  which yields  $a \in a_{e,f}^\dagger \mathcal{A}^{-1}$ . Hence, (XV) holds.

(XV)  $\Rightarrow$  (XVI): Obviously.

(XVI)  $\Rightarrow$  (IX): From the condition  $a \in a_{e,f}^\dagger \mathcal{A}$ , we have  $a = a_{e,f}^\dagger x$ , for some  $x \in \mathcal{A}$ , and  $a^k - a_{e,f}^\dagger a a^k = (a_{e,f}^\dagger - a_{e,f}^\dagger a a_{e,f}^\dagger) x a^{k-1} = 0$  for integer  $k \geq 1$ . In the similar way,  $a \in \mathcal{A} a_{e,f}^\dagger$  gives  $a^k = a^k a a_{e,f}^\dagger$  for integer  $k \geq 1$ . When  $k = 1$ , we observe that  $a \in a^2 \mathcal{A} \cap \mathcal{A} a^2$  and  $a^\#$  exists. So, the condition (IX) holds.

(XV)  $\Leftrightarrow$  (XVII): The assumption  $a \in a_{e,f}^\dagger \mathcal{A}^{-1} \cap \mathcal{A}^{-1} a_{e,f}^\dagger$  is equivalent to  $a \mathcal{A}^{-1} = a_{e,f}^\dagger \mathcal{A}^{-1}$  and  $\mathcal{A}^{-1} a = \mathcal{A}^{-1} a_{e,f}^\dagger$ . By Lemma 1.7, we observe that these equalities hold if and only if  $a \mathcal{A}^{-1} = a^{*f,e} \mathcal{A}^{-1} = f^{-1} a^* \mathcal{A}^{-1}$  and  $\mathcal{A}^{-1} a = \mathcal{A}^{-1} a^{*f,e} = \mathcal{A}^{-1} a^* e$ .

(XVII)  $\Leftrightarrow$  (XVIII): Applying the involution, we check this equivalence.

(XIX)  $\Rightarrow$  (II): Suppose that there exists  $x \in \mathcal{A}$  such that  $a = e^{-1} a^* x a^* f$ . Then, by Lemma 1.7,  $a \in e^{-1} a^* \mathcal{A} \cap \mathcal{A} a^* f = a^{*e,f} \mathcal{A} \cap \mathcal{A} a^{*e,f} = a_{f,e}^\dagger \mathcal{A} \cap \mathcal{A} a_{f,e}^\dagger$ . Now, by (xvi)  $\Leftrightarrow$  (i), we deduce that  $a$  is weighted-EP w.r.t.  $(f, e)$ .

(II)  $\Rightarrow$  (XIX): If  $a$  is weighted-EP w.r.t.  $(f, e)$ , by the equivalence (I)  $\Leftrightarrow$  (XVI),  $a \in a_{f,e}^\dagger \mathcal{A} \cap \mathcal{A} a_{f,e}^\dagger = e^{-1} a^* \mathcal{A} \cap \mathcal{A} a^* f$ . Therefore, for some  $y, z \in \mathcal{A}$ ,  $a = e^{-1} a^* y = z a^* f$  and  $a = aa_{f,e}^\dagger = e^{-1} a^* (y a_{f,e}^\dagger z) a^* f$ . For  $x = y a_{f,e}^\dagger z$ , the statement (XIX) is satisfied.

(XX)  $\Rightarrow$  (II): Since  $a = (ae^{-1})^\dagger ae^{-1} f a (fa)^\dagger$ , we conclude that  $a \in (ae^{-1})^\dagger \mathcal{A} \cap \mathcal{A} (fa)^\dagger = (ae^{-1})^* \mathcal{A} \cap \mathcal{A} (fa)^* = a^{*e,f} \mathcal{A} \cap \mathcal{A} a^{*e,f} = a_{f,e}^\dagger \mathcal{A} \cap \mathcal{A} a_{f,e}^\dagger$ . Using (XVI)  $\Leftrightarrow$  (I), we observe that  $a$  is weighted-EP w.r.t.  $(f, e)$ .

(II)  $\Rightarrow$  (XX): The condition (II) implies that  $ae^{-1}$  and  $fa$  are EP, and hence,

$$\begin{aligned} (ae^{-1})^\dagger ae^{-1} f a (fa)^\dagger &= ((ae^{-1})^\dagger ae^{-1} ae^{-1}) e f a (fa)^\dagger = ae^{-1} e f a (fa)^\dagger \\ &= f^{-1} (f a f a (fa)^\dagger) = f^{-1} f a = a. \end{aligned}$$

Thus, the condition (XX) holds.

(XXI)  $\Leftrightarrow$  (XII): Applying the equivalence (I)  $\Leftrightarrow$  (XII) for  $a^k$ ,  $k \geq 1$ , we see that  $a \in \mathcal{A}^\#$  and  $a^k$  is weighted-EP w.r.t.  $(e, f)$  if and only if  $a \in \mathcal{A}^\#$  and  $ea^k(a^k)^\#$ ,  $f a^k(a^k)^\#$  are Hermitian which is equivalent to  $a \in \mathcal{A}^\#$  and  $ea a^\#$ ,  $f a a^\#$  are Hermitian, by  $(a^k)^\# = (a^\#)^k$ .

(XXII)  $\Leftrightarrow$  (IV): Notice that, from  $a \in \mathcal{A}^-$ , it follows that  $aa^*a \in \mathcal{A}^-$  and  $(aa^*a)^\dagger = a^\dagger(a^*)^\dagger a^\dagger$ . Using the equivalence (I)  $\Leftrightarrow$  (IV) for  $aa^*a$ , we observe that  $aa^*a$  is weighted-EP w.r.t.  $(e, f)$  is equivalent to  $ea a^*a \mathcal{A} = f a a^*a \mathcal{A} = a^*aa^*\mathcal{A}$ . Since  $a \in \mathcal{A}^-$ ,  $a^\dagger$  exists and  $ea = eaa^\dagger a = eaa^*(a^\dagger)^* = eaa^*aa^\dagger(a^\dagger)^*$ . Consequently,  $ea a^*a \mathcal{A} = ea \mathcal{A}$  and in the same way  $f a a^*a \mathcal{A} = f a \mathcal{A}$ . By  $a^* = a^*aa^\dagger = a^*aa^\dagger aa^\dagger = a^*aa^*(a^\dagger)^*a^\dagger$ , we conclude  $a^*aa^*\mathcal{A} = a^*\mathcal{A}$ . Hence,  $ea a^*a \mathcal{A} = f a a^*a \mathcal{A} = a^*aa^*\mathcal{A}$  is equivalent to (IV).

(XXIII)  $\Leftrightarrow$  (I): Observe that  $a$  is weighted-EP w.r.t.  $(e, f)$  if and only if elements  $ea$  and  $af^{-1}$  are EP. By the definition of EP elements, this is equivalent to  $(ea)^\circ = [(ea)^*]^\circ$  and  $(af^{-1})^\circ = [(af^{-1})^*]^\circ = (f^{-1}a^*)^\circ$ , which can be written as  $a^\circ = [(ea)^*]^\circ$  and  $(af^{-1})^\circ = (a^*)^\circ$ , by Lemma 1.6.

(V)  $\Rightarrow$  (XXIV): The condition (V) gives  $e^{-1}a^*\mathcal{A} = a\mathcal{A}$  and its equivalent condition (IV) imply  $f a \mathcal{A} = a^*\mathcal{A}$ . For  $a^*, a^*f \in \mathcal{A}^-$ , Lemma 1.5 implies that  $\mathcal{A} = a\mathcal{A} \oplus (a^*)^\circ = f a \mathcal{A} \oplus (a^*f)^\circ$ . Thus,  $\mathcal{A} = e^{-1}a^*\mathcal{A} \oplus (a^*)^\circ = a^*\mathcal{A} \oplus (a^*f)^\circ$ .

(XXIV)  $\Rightarrow$  (VII): From  $\mathcal{A} = e^{-1}a^*\mathcal{A} \oplus (a^*)^\circ = a^*\mathcal{A} \oplus (a^*f)^\circ$ , we see that  $\mathcal{A} = e^{-1}a^*\mathcal{A} \oplus (e^{-1}a^*)^\circ = a^*f \mathcal{A} \oplus (a^*f)^\circ$ . Define the left regular representation  $L_a : \mathcal{A} \rightarrow \mathcal{A}$  by  $L_a(x) = ax$  for all  $x \in \mathcal{A}$ . Now,  $\mathcal{A} = R(L_{e^{-1}a^*}) \oplus N(L_{e^{-1}a^*}) = R(L_{a^*f}) \oplus N(L_{a^*f})$  which implies that  $L_{e^{-1}a^*}$  i  $L_{a^*f}$  are EP operators. According to [3, Remark 12], necessary and sufficient condition for  $a \in \mathcal{A}$  to be EP is that  $L_a \in \mathcal{L}(\mathcal{A})$  is EP. So, elements  $e^{-1}a^*$ ,  $a^*f$  are EP, and  $a^*$  is weighted-EP w.r.t.  $(e^{-1}, f^{-1})$ .

(XXV)  $\Leftrightarrow$  (V): By the equivalence (I)  $\Leftrightarrow$  (V) for  $a^\dagger$ , we get that  $a^\dagger$  is weighted-EP w.r.t.  $(e^{-1}, f^{-1})$  if and only if  $e(a^\dagger)^*\mathcal{A} = f(a^\dagger)^*\mathcal{A} = a^\dagger\mathcal{A}$ . Recall that  $a^\dagger\mathcal{A} = a^*\mathcal{A}$  and  $(a^\dagger)^*\mathcal{A} = a\mathcal{A}$ . Now,

$$e(a^\dagger)^*\mathcal{A} = a^\dagger\mathcal{A} \Leftrightarrow (a^\dagger)^*\mathcal{A} = e^{-1}a^*\mathcal{A} \Leftrightarrow a\mathcal{A} = e^{-1}a^*\mathcal{A}$$

and, similarly,  $f(a^\dagger)^*\mathcal{A} = a^\dagger\mathcal{A} \Leftrightarrow a\mathcal{A} = f^{-1}a^*\mathcal{A}$ .

(XXVI)  $\Leftrightarrow$  (VI): If we apply the equivalence (I)  $\Leftrightarrow$  (iv) for  $a_{e,f}^\dagger$ , then  $a_{e,f}^\dagger$  is weighted-EP w.r.t.  $(e, f) \Leftrightarrow ea_{e,f}^\dagger\mathcal{A} = fa_{e,f}^\dagger\mathcal{A} = (a_{e,f}^\dagger)^*\mathcal{A}$ . By Lemma 1.6, we obtain

$$ea_{e,f}^\dagger\mathcal{A} = (a_{e,f}^\dagger)^*\mathcal{A} \Leftrightarrow ea_{e,f}^\dagger\mathcal{A} = ea\mathcal{A} \Leftrightarrow a_{e,f}^\dagger\mathcal{A} = a\mathcal{A}$$

and

$$fa_{e,f}^\dagger\mathcal{A} = (a_{e,f}^\dagger)^*\mathcal{A} \Leftrightarrow a^*\mathcal{A} = (a_{e,f}^\dagger)^*\mathcal{A}.$$

(XXVII)  $\Rightarrow$  (VIII): Assume that  $a \in \mathcal{A}^\#$  and  $a^{2k-1} = a_{e,f}^\dagger a^{2k+1} a_{e,f}^\dagger$ , for any/some integer  $k \geq 1$ . Consequently, we have

$$aa_{e,f}^\dagger = (a^\#)^{2k} a^{2k+1} a_{e,f}^\dagger = (a^\#)^{2k} a(a_{e,f}^\dagger a^{2k+1} a_{e,f}^\dagger) = (a^\#)^{2k} aa^{2k-1} = a^\# a$$

and

$$a_{e,f}^\dagger a = (a_{e,f}^\dagger a^{2k+1} a_{e,f}^\dagger) a (a^\#)^{2k} = a^{2k-1} a (a^\#)^{2k} = aa^\#$$

implying  $aa_{e,f}^\dagger = a_{e,f}^\dagger a$ .

The implication (VIII)  $\Rightarrow$  (XXVII) is easy to check.

(XXVIII)  $\Rightarrow$  (IX): Suppose that  $a \in \mathcal{A}^\#$  and  $aa_{e,f}^\dagger a_{e,f}^\dagger a = a_{e,f}^\dagger a a a_{e,f}^\dagger$ . Now, observe that

$$\begin{aligned} a_{e,f}^\dagger a a a_{e,f}^\dagger &= aa_{e,f}^\dagger a_{e,f}^\dagger a = aa^\# (aa_{e,f}^\dagger a_{e,f}^\dagger a) a^\# a \\ (2.1) \quad &= a^\# aa_{e,f}^\dagger a a a_{e,f}^\dagger aa^\# = a^\# a. \end{aligned}$$

First, if  $k$  is a positive integer, then multiplying the equality (2.1) by  $a^k$  from the left side, we get  $a^k aa_{e,f}^\dagger = a^k$  and then multiplying the equality (2.1) by  $a^k$  from the right side, we obtain  $a_{e,f}^\dagger aa^k = a^k$ . So, the condition (IX) is satisfied.

The implication (VIII)  $\Rightarrow$  (XXVIII) is obvious.

(XXIX)  $\Leftrightarrow$  (IV): Using (I)  $\Leftrightarrow$  (IV) for  $a^\#$ , we deduce that  $a^\#$  is weighted-EP w.r.t.  $(e, f)$  if and only if  $ea^\# \mathcal{A} = fa^\# \mathcal{A} = (a^\#)^* \mathcal{A}$ . This is equivalent to (iv), because  $a^\# \mathcal{A} = a \mathcal{A}$ ,  $a^\# \mathcal{A} = a \mathcal{A}$  and  $(a^\#)^* \mathcal{A} = a^* \mathcal{A}$ .

(XXX)  $\Rightarrow$  (XII): From  $aa^\# = aa_{e,e}^\dagger = aa_{f,f}^\dagger$  we conclude that elements  $ea a^\# = eaa_{e,e}^\dagger$  and  $fa a^\# = faa_{f,f}^\dagger$  are Hermitian.

(III)  $\Rightarrow$  (XXX): Since  $a$  is weighted-EP w.r.t.  $(e, f)$ , it follows that  $a^\# = a_{e,f}^\dagger$ . Hence, the fact that  $a$  is both weighted-EP w.r.t.  $(e, e)$  and w.r.t.  $(f, f)$  gives  $a^\# = a_{e,e}^\dagger = a_{f,f}^\dagger$ . Thus,  $aa^\# = aa_{e,e}^\dagger = aa_{f,f}^\dagger$ .

(I)  $\Rightarrow$  (XXXI)  $\Rightarrow$  (XII): This part follows similarly as (III)  $\Rightarrow$  (XXX)  $\Rightarrow$  (XII), using the equivalence (I)  $\Leftrightarrow$  (II).

(XXXII)  $\Rightarrow$  (XXX): By the equality  $aa_{e,e}^\dagger e^{-1} a^* a = e^{-1} a^* a a a_{e,e}^\dagger$ , we have

$$\begin{aligned} (2.2) \quad a^* &= a^* aa^\dagger = a^* a a a^\# a^\dagger = e(e^{-1} a^* a a a_{e,e}^\dagger) a a^\# a^\dagger \\ &= eaa_{e,e}^\dagger e^{-1} a^* a a a^\# a^\dagger = eaa_{e,e}^\dagger e^{-1} a^*. \end{aligned}$$

Applying the involution to (2.2), we obtain  $a = ae^{-1} eaa_{e,e}^\dagger = a a a_{e,e}^\dagger$  which yields  $a^\# a = aa_{e,e}^\dagger$ . In the same way, the assumption  $aa_{f,f}^\dagger f^{-1} a^* a = f^{-1} a^* a a a_{f,f}^\dagger$  implies  $a^\# a = aa_{f,f}^\dagger$ . Therefore, the condition (XXX) holds.

(III)  $\Rightarrow$  (XXXII): The condition (III) gives that  $a^\# = a_{e,e}^\dagger = a_{f,f}^\dagger$ . Then we get

$$\begin{aligned} aa_{e,e}^\dagger e^{-1} a^* a &= e^{-1} eaa_{e,e}^\dagger e^{-1} a^* a = e^{-1} (ae^{-1} eaa_{e,e}^\dagger)^* a \\ &= e^{-1} (aaa^\#)^* a = e^{-1} a^* a = e^{-1} a^* a a a_{e,e}^\dagger \end{aligned}$$

and similarly  $aa_{f,f}^\dagger f^{-1} a^* a = f^{-1} a^* a a a_{f,f}^\dagger$ .

The implications (I)  $\Rightarrow$  (XXXIII)  $\Rightarrow$  (XXX), (III)  $\Rightarrow$  (XXXIV)  $\Rightarrow$  (XXX) and (I)  $\Rightarrow$  (XXXV)  $\Rightarrow$  (XXX) can be proved in the same way as (III)  $\Rightarrow$  (XXXII)  $\Rightarrow$  (XXX).

(XXXVI)  $\Rightarrow$  (VIII): Multiplying  $a^k aa_{e,f}^\dagger + a_{e,f}^\dagger aa^k = 2a^k$ ,  $k \geq 1$ , from the right side by  $(a^\#)^k$ , we obtain

$$a^k aa_{e,f}^\dagger (a^\#)^k + a_{e,f}^\dagger a a a^\# = 2a a^\#.$$

Further, the equality

$$a^k aa_{e,f}^\dagger a (a^\#)^{k+1} + a_{e,f}^\dagger a = 2a a^\#,$$

gives  $aa^\# + a_{e,f}^\dagger a = 2a a^\#$ , i.e.,  $a_{e,f}^\dagger a = aa^\#$ . Similarly, multiplying  $a^k aa_{e,f}^\dagger + a_{e,f}^\dagger aa^k = 2a^k$  from the left side by  $(a^\#)^k$ , we show that  $aa_{e,f}^\dagger = aa^\#$ . So,  $a_{e,f}^\dagger a = aa_{e,f}^\dagger$ .

(VIII)  $\Rightarrow$  (XXXVI)  $\wedge$  (XXXVII): We can easily check this implication.

(XXXVII)  $\Rightarrow$  (VIII): Multiplying the equality  $a_{e,f}^\dagger a^\# a + aa^\# a_{e,f}^\dagger = 2a_{e,f}^\dagger$  by  $a$  first from the right side, we get  $a^\# a = a_{e,f}^\dagger a$  and then from the left side, we obtain  $aa^\# = a_{e,f}^\dagger a$ . Hence, we deduce that  $a_{e,f}^\dagger a = aa_{e,f}^\dagger$ .

(XXXVIII)  $\Rightarrow$  (XII): The condition  $a^{*f,e} = a^{*f,e} aa^\# = a^\# aa^{*f,e}$  is equivalent to  $a^* = a^* eaa^\# e^{-1} = fa^\# af^{-1} a^*$ . Then, from

$$(eaa^\#)^* = (a^\#)^* a^* e = (a^\#)^* a^* eaa^\# e^{-1} e = (aa^\#)^* eaa^\#$$

and

$$(fa^\# a)^* = a^* (a^\#)^* f = fa^\# af^{-1} a^* (a^\#)^* f = fa^\# af^{-1} (a^\# a)^* f,$$

we conclude that elements  $eaa^\#$  and  $fa^\# a$  are Hermitian.

(XII)  $\Rightarrow$  (XXXVIII): If  $eaa^\#$  is Hermitian, then

$$a^{*f,e} aa^\# = f^{-1} a^* eaa^\# = f^{-1} (eaa^\# a)^* = f^{-1} (ea)^* = f^{-1} a^* e = a^{*f,e}.$$

In the same way, since  $faa^\#$  is Hermitian, it follows  $a^{*f,e} = a^\# aa^{*f,e}$ .

(XXXIX)  $\Rightarrow$  (XXXVIII): Multiplying the equality  $a^{*f,e} aa^\# + a^\# aa^{*f,e} = 2a^{*f,e}$  by  $aa_{e,f}^\dagger$  from the right side, we get

$$a^{*f,e} aa_{e,f}^\dagger + a^\# aa^{*f,e} aa_{e,f}^\dagger = 2a^{*f,e} aa_{e,f}^\dagger.$$

By Theorem 1.4, we have  $a^{*f,e} + a^\# aa^{*f,e} = 2a^{*f,e}$  which implies  $a^\# aa^{*f,e} = a^{*f,e}$ . Similarly, multiplying the equality  $a^{*f,e} aa^\# + a^\# aa^{*f,e} = 2a^{*f,e}$  from the left side by  $a$  and then by  $a_{e,f}^\dagger$ , we obtain  $a^{*f,e} aa^\# = a^{*f,e}$ . Hence, (XXIX) is satisfied.

The implication (XXXVIII)  $\Rightarrow$  (XXXIX) is obvious.

(XL)  $\Rightarrow$  (IX): Multiplying the condition  $a^k aa_{e,f}^\dagger + (a^k aa_{e,f}^\dagger)^* = a^k + (a^k)^*$ ,  $k \geq 1$ , by  $a$  from the right side, we see that

$$(2.3) \quad (a^k aa_{e,f}^\dagger)^* a = (a^k)^* a.$$

Applying the involution to (2.3), we obtain

$$a^* a^k aa_{e,f}^\dagger = a^* a^k,$$

which gives

$$a^k aa_{e,f}^\dagger = aa_{e,f}^\dagger a^k aa_{e,f}^\dagger = (a^\dagger)^* (a^* a^k aa_{e,f}^\dagger) = (a^\dagger)^* a^* a^k = a^k.$$

In the same way, multiplying the hypothesis  $a_{e,f}^\dagger aa^k + (a_{e,f}^\dagger aa^k)^* = a^k + (a^k)^*$ ,  $k \geq 1$ , by  $a$  from the left side, we show  $a_{e,f}^\dagger aa^k = a^k$ . Thus, (ix) holds.

The implication (VIII)  $\Rightarrow$  (XL)  $\wedge$  (XLI) is obvious.

(XLI)  $\Rightarrow$  (VIII): The equality  $aa_{e,f}^\dagger (a + \lambda a_{e,f}^\dagger) = (a + \lambda a_{e,f}^\dagger) aa_{e,f}^\dagger$  is equivalent to

$$(2.4) \quad a + \lambda aa_{e,f}^\dagger a_{e,f}^\dagger = a aa_{e,f}^\dagger + \lambda a_{e,f}^\dagger.$$

Multiplying (2.4) from the left side by  $a_{e,f}^\dagger$ , we get

$$a_{e,f}^\dagger a + \lambda a_{e,f}^\dagger a_{e,f}^\dagger a_{e,f}^\dagger = a_{e,f}^\dagger a aa_{e,f}^\dagger + \lambda a_{e,f}^\dagger a_{e,f}^\dagger,$$

which yields  $a_{e,f}^\dagger a = a_{e,f}^\dagger a aa_{e,f}^\dagger$ . Analogously,  $a_{e,f}^\dagger a (a + \lambda a_{e,f}^\dagger) = (a + \lambda a_{e,f}^\dagger) a_{e,f}^\dagger a$  implies  $aa_{e,f}^\dagger = a_{e,f}^\dagger a aa_{e,f}^\dagger$ . Therefore,  $a_{e,f}^\dagger a = aa_{e,f}^\dagger$ .

(XLII)  $\Rightarrow$  (VIII): Assume that  $ab = ba$  implies  $a_{e,f}^\dagger b = ba_{e,f}^\dagger$ . If  $b = a$ , then  $a_{e,f}^\dagger a = aa_{e,f}^\dagger$ .

(XI)  $\Rightarrow$  (XLII): By  $a^\# = a_{e,f}^\dagger$  and the double commutativity of  $a^\#$ , from  $ab = ba$  we obtain  $a^\# b = ba^\#$ , i.e.,  $a_{e,f}^\dagger b = ba_{e,f}^\dagger$ .

(XLIII)  $\Rightarrow$  (VIII): Let  $a_{e,f}^\dagger = f(a)$  for some function  $f$  holomorphic in a neighbourhood of  $\sigma(a)$ . By a property of the holomorphic calculus,  $a_{e,f}^\dagger$  commutes with  $a$ .

(XI)  $\Rightarrow$  (XLIII): By  $a_{e,f}^\dagger = a^\#$  along with [20, Theorem 4.4], we have  $a^\# = f(a)$ , where  $f$  is holomorphic in a neighbourhood of  $\sigma(a)$ , and  $f(\lambda) = 0$  in a neighbourhood of 0,  $f(\lambda) = \lambda^{-1}$  in a neighbourhood of  $\sigma(a) \setminus \{0\}$ , it follows (XLIII).

(XLIV)  $\Rightarrow$  (III): Since  $(a + \lambda a_{e,e}^\dagger)\mathcal{A} = (\lambda a + a^3)\mathcal{A}$  for  $\lambda \neq 0$ , we have  $a + \lambda a_{e,e}^\dagger = (\lambda a + a^3)x$  for some  $x \in \mathcal{A}$ . Now, from

$$\begin{aligned} a + \lambda a a_{e,e}^\dagger a_{e,e}^\dagger &= a a_{e,e}^\dagger (a + \lambda a_{e,e}^\dagger) = a a_{e,e}^\dagger (\lambda a + a^3)x \\ &= (\lambda a + a^3)x = a + \lambda a_{e,e}^\dagger, \end{aligned}$$

we conclude that  $a a_{e,e}^\dagger a_{e,e}^\dagger = a_{e,e}^\dagger$ . In the same way,  $\mathcal{A}(a + \lambda a_{e,e}^\dagger) = \mathcal{A}(\lambda a + a^3)$  gives  $a_{e,e}^\dagger a_{e,e}^\dagger a = a_{e,e}^\dagger$ . So,  $a_{e,e}^\dagger = a a_{e,e}^\dagger a_{e,e}^\dagger = a_{e,e}^\dagger a_{e,e}^\dagger a$ , which implies that  $a$  is weighted-EP w.r.t.  $(e, e)$ , by (I)  $\Leftrightarrow$  (X).

Similarly, from the equalities  $(a + \lambda a_{f,f}^\dagger)\mathcal{A} = (\lambda a + a^3)\mathcal{A}$  and  $\mathcal{A}(a + \lambda a_{f,f}^\dagger) = \mathcal{A}(\lambda a + a^3)$ , for  $\lambda \neq 0$ , we can show that  $a$  is weighted-EP w.r.t.  $(f, f)$ .

(III)  $\Rightarrow$  (XLIV): The condition (III) implies  $a^\# = a_{e,e}^\dagger = a_{f,f}^\dagger$ . Then, for  $\lambda \neq 0$ , by

$$(2.5) \quad a + \lambda a_{e,e}^\dagger = a + \lambda a^\# = (a^3 + \lambda a)(a^\#)^2 \in (a^3 + \lambda a)\mathcal{A}$$

and

$$(2.6) \quad a^3 + \lambda a = (a + \lambda a^\#)a^2 = (a + \lambda a_{e,e}^\dagger)a^2 \in (a + \lambda a_{e,e}^\dagger)\mathcal{A},$$

we deduce  $(a + \lambda a_{e,e}^\dagger)\mathcal{A} = (\lambda a + a^3)\mathcal{A}$ . In the same way, it follows the rest of condition (XLIV).

(XI)  $\Rightarrow$  (XLV)  $\Rightarrow$  (X): It follows in the same way as the part (III)  $\Leftrightarrow$  (XLIV).

(XLVI)  $\Rightarrow$  (XXIII): Assume that  $(a + \lambda a_{e,e}^\dagger)^\circ = (\lambda a + a^3)^\circ$  for  $\lambda \neq 0$ . If  $ax = 0$  for some  $x \in \mathcal{A}$ , then  $(\lambda a + a^3)x = 0$  implies  $(a + \lambda a_{e,e}^\dagger)x = 0$ . Now, we conclude  $a_{e,e}^\dagger x = 0$  and  $a^\circ \subset (a_{e,e}^\dagger)^\circ$ . Therefore, by Lemma 1.6,  $a^\circ \subset [(ea)^*]^\circ$ .

Let  ${}^\circ(a + \lambda a_{e,e}^\dagger) = {}^\circ(\lambda a + a^3)$  for  $\lambda \neq 0$ , and  $(ea)^*x = 0$  for some  $x \in \mathcal{A}$ . Applying the involution, we see that  $x^*ea = 0$  which gives  $x^*e(\lambda a + a^3) = 0$ . Then  $x^*e(a + \lambda a_{e,e}^\dagger) = 0$  and, consequently,  $x^*ea_{e,e}^\dagger = 0$ , i.e.,  $(a_{e,e}^\dagger)^*ex = 0$ . By this equality, we have

$$ax = ae^{-1}(ea_{e,e}^\dagger)^*x = ae^{-1}a^*(a_{e,e}^\dagger)^*ex = 0.$$

Hence,  $[(ea)^*]^\circ \subset a^\circ$  and  $a^\circ = [(ea)^*]^\circ$ .

The equalities  $(a + \lambda a_{f,f}^\dagger)^\circ = (\lambda a + a^3)^\circ$  and  ${}^\circ(a + \lambda a_{f,f}^\dagger) = {}^\circ(\lambda a + a^3)$  for  $\lambda \neq 0$ , imply  $(a^*)^\circ = (af^{-1})^\circ$  in the similar way.

(III)  $\Rightarrow$  (XLVI): The assumption (III) gives  $a^\# = a_{e,e}^\dagger = a_{f,f}^\dagger$ , so by (2.5) and (2.6), we deduce  ${}^\circ(a + \lambda a_{e,e}^\dagger) = {}^\circ(\lambda a + a^3)$ , for  $\lambda \neq 0$ . Similarly, we can prove the rest of (XLVI).

(III)  $\Rightarrow$  (XLVII)  $\Rightarrow$  (XXIII): Similarly as (III)  $\Rightarrow$  (XLVI)  $\Rightarrow$  (XXIII).

(XLVIII)  $\Rightarrow$  (VIII): Using the equality  $(a_{e,f}^\dagger)^2 a^\# = a_{e,f}^\dagger a^\# a_{e,f}^\dagger$ , first we get

$$\begin{aligned} (a_{e,f}^\dagger)^2 a^\# &= ((a_{e,f}^\dagger)^2 a^\#) a a^\# = a_{e,f}^\dagger a^\# a_{e,f}^\dagger a a^\# \\ &= a_{e,f}^\dagger (a^\#)^2 a a_{e,f}^\dagger a a^\# = a_{e,f}^\dagger (a^\#)^2 \end{aligned}$$

and then

$$\begin{aligned} a a_{e,f}^\dagger &= a^3 (a^\#)^2 a_{e,f}^\dagger = a^3 a_{e,f}^\dagger a (a^\#)^2 a_{e,f}^\dagger = a^3 (a_{e,f}^\dagger a^\# a_{e,f}^\dagger) \\ &= a^3 ((a_{e,f}^\dagger)^2 a^\#) = a^3 a_{e,f}^\dagger (a^\#)^2 = a^3 a_{e,f}^\dagger a (a^\#)^3 = a a^\#. \end{aligned}$$

We can show that  $a_{e,f}^\dagger a^\# a_{e,f}^\dagger = a^\# (a_{e,f}^\dagger)^2$  implies  $a_{e,f}^\dagger a = a a^\#$  in the same way. Thus,  $a_{e,f}^\dagger a = a a_{e,f}^\dagger$ .

The implication (XI)  $\Rightarrow$  (XLVIII)  $\wedge$  (XLIX) is obvious.

(XLIX)  $\Rightarrow$  (VIII): From the hypothesis  $a(a_{e,f}^\dagger)^2 = a^\# = (a_{e,f}^\dagger)^2 a$ , we have

$$a a_{e,f}^\dagger = a a^\# a_{e,f}^\dagger = a a (a_{e,f}^\dagger)^2 a a_{e,f}^\dagger = a (a (a_{e,f}^\dagger)^2) = a a^\#$$

and

$$a_{e,f}^\dagger a = a_{e,f}^\dagger a a^\# a = a_{e,f}^\dagger a (a_{e,f}^\dagger)^2 a a = ((a_{e,f}^\dagger)^2 a) a = a^\# a.$$

Therefore, we deduce that  $a_{e,f}^\dagger a = a a_{e,f}^\dagger$ .

(L)  $\Rightarrow$  (XLIX): The equalities  $a^{*f,e} a_{e,f}^\dagger = a^{*f,e} a^\#$  and  $a_{e,f}^\dagger a^{*f,e} = a^\# a^{*f,e}$  are equivalent to  $a^* e a_{e,f}^\dagger = a^* e a^\#$  and  $a_{e,f}^\dagger f^{-1} a^* = a^\# f^{-1} a^*$ . By  $a^* e a_{e,f}^\dagger = a^* e a^\#$ , we obtain

$$\begin{aligned} a(a_{e,f}^\dagger)^2 &= e^{-1} (e a a_{e,f}^\dagger)^* a_{e,f}^\dagger = e^{-1} (a_{e,f}^\dagger)^* (a^* e a_{e,f}^\dagger) = e^{-1} (a_{e,f}^\dagger)^* a^* e a^\# \\ &= e^{-1} e a a_{e,f}^\dagger a^\# = a a_{e,f}^\dagger a (a^\#)^2 = a^\#. \end{aligned}$$

Analogously, from  $a_{e,f}^\dagger f^{-1} a^* = a^\# f^{-1} a^*$ , we get  $a^\# = (a_{e,f}^\dagger)^2 a$ . So, the condition (XLIX) holds.

The implication (XI)  $\Rightarrow$  (L)  $\wedge$  (LI) is obvious.

(LI)  $\Rightarrow$  (XLIX): Multiplying  $(a_{e,f}^\dagger)^2 = (a^\#)^2$  by  $a$  from the left side and then from the right side, we observe that (XLIX) is satisfied.

(LII)  $\Leftrightarrow$  (VIII): The assumption  $a^{*e,f} = a^{*e,f} a_{e,f}^\dagger a = a a_{e,f}^\dagger a^{*e,f}$  is equivalent to  $a^* f = a^* f a_{e,f}^\dagger a$  and  $e^{-1} a^* = a a_{e,f}^\dagger e^{-1} a^*$ . Applying the involution to these equalities, we see that they are equivalent to  $f a = f a_{e,f}^\dagger a a$  and  $a e^{-1} = a e^{-1} e a a_{e,f}^\dagger e^{-1}$ , i.e.,  $a = a_{e,f}^\dagger a a = a a a_{e,f}^\dagger \Leftrightarrow a_{e,f}^\dagger a = a a_{e,f}^\dagger$ .

(LIII)  $\Rightarrow$  (XII): Since  $(a^\#)^{*e,f} = a a^\# (a^\#)^{*e,f} = (a^\#)^{*e,f} a^\# a$  can be written as  $(a^\#)^* = e a a^\# e^{-1} (a^\#)^* = (a^\#)^* f a^\# a f^{-1}$ , we get

$$(e a a^\#)^* = (a^\#)^* a^* e = e a a^\# e^{-1} (a^\#)^* a^* e = e a a^\# e^{-1} (e a a^\#)^*$$

and

$$(f a^\# a)^* = a^* (a^\#)^* f = a^* (a^\#)^* f a^\# a f^{-1} f = (a^\# a)^* f a^\# a.$$

So, we conclude that  $e a a^\#$  and  $f a^\# a$  are Hermitian.

(XI)  $\Rightarrow$  (LIII): It is easy to check this part, by Theorem 1.4.

(LIV)  $\Rightarrow$  (XII): The condition  $a_{e,f}^\dagger (a^\#)^2 = (a^\#)^2 a_{e,f}^\dagger$  gives

$$e a a^\# = e a^3 a_{e,f}^\dagger a (a^\#)^3 = e a^3 (a_{e,f}^\dagger (a^\#)^2) = e a^3 (a^\#)^2 a_{e,f}^\dagger = e a a_{e,f}^\dagger$$

and

$$f a^\# a = f (a^\#)^3 a a_{e,f}^\dagger a^3 = f ((a^\#)^2 a_{e,f}^\dagger) a^3 = f a_{e,f}^\dagger (a^\#)^2 a^3 = f a_{e,f}^\dagger a.$$

Therefore,  $e a a^\#$  and  $f a^\# a$  are Hermitian elements.

The implication (XI)  $\Rightarrow$  (LIV)  $\wedge$  (LV) is obvious.

(LV)  $\Rightarrow$  (VIII): Suppose that  $a \in \mathcal{A}^\#$  and  $a^k a_{e,f}^\dagger = a_{e,f}^\dagger a^k$ , for any/some integer  $k \geq 1$ . Then

$$a a_{e,f}^\dagger = (a^\#)^{k-1} (a^k a_{e,f}^\dagger) = (a^\#)^{k-1} a_{e,f}^\dagger a^k = (a^\#)^k a a_{e,f}^\dagger a^k = a^\# a$$

and

$$a_{e,f}^\dagger a = (a_{e,f}^\dagger a^k) (a^\#)^{k-1} = a^k a_{e,f}^\dagger (a^\#)^{k-1} = a^k a_{e,f}^\dagger a (a^\#)^k = a a^\#.$$

Hence,  $a a_{e,f}^\dagger = a_{e,f}^\dagger a$ .

(LVI)  $\Rightarrow$  (VIII): The equality  $a a_{e,f}^\dagger (a + \lambda a^{*e,f}) = (a + \lambda a^{*e,f}) a a_{e,f}^\dagger$ , for  $\lambda \neq 0$ , is equivalent to

$$(2.7) \quad a + \lambda a a_{e,f}^\dagger a^{*e,f} = a a a_{e,f}^\dagger + \lambda a^{*e,f} a a_{e,f}^\dagger.$$

Multiplying (2.7) from the right side by  $a$ , we observe that

$$(2.8) \quad a a_{e,f}^\dagger a^{*e,f} a = e^{-1} a^{*e,f}.$$



Multiplying (2.8) from the right side by  $a_{f,e}^\dagger$ , we get  $aa_{e,f}^\dagger a^{*e,f} = a^{*e,f}$ .

Similarly, from  $a_{e,f}^\dagger a(a + \lambda a^{*e,f}) = (a + \lambda a^{*e,f})a_{e,f}^\dagger a$ , for  $\lambda \neq 0$ , we obtain  $a^{*e,f} a_{e,f}^\dagger a = a^{*e,f}$ . Thus, the condition (LII) is satisfied.

(VIII)  $\Rightarrow$  (LVI): If  $a_{e,f}^\dagger a = aa_{e,f}^\dagger$ , then

$$\begin{aligned} (a + \lambda a^{*e,f})aa_{e,f}^\dagger &= a + \lambda e^{-1} a^* f a_{e,f}^\dagger a = a + e^{-1} (f a_{e,f}^\dagger a a)^* \\ &= a + \lambda e^{-1} a^* f = a + \lambda e^{-1} (a e^{-1} e a a_{e,f}^\dagger)^* f \\ &= a + \lambda a a_{e,f}^\dagger e^{-1} a^* f = aa_{e,f}^\dagger (a + \lambda a^{*e,f}). \end{aligned}$$

The second equality follows similarly.

(LVII)  $\Leftrightarrow$  (XXXII): Note that the assumption  $aa_{e,e}^\dagger (aa^*e - e^{-1}a^*a) = (aa^*e - e^{-1}a^*a)aa_{e,e}^\dagger$  is equivalent to  $aa^*e - aa_{e,e}^\dagger e^{-1}a^*a = aa^*e - e^{-1}a^*aaa_{e,e}^\dagger$ , i.e.,  $aa_{e,e}^\dagger e^{-1}a^*a = e^{-1}a^*aaa_{e,e}^\dagger$ .

In the same way, the equality  $aa_{f,f}^\dagger (aa^*f - f^{-1}a^*a) = (aa^*f - f^{-1}a^*a)aa_{f,f}^\dagger$  holds if and only if  $aa_{f,f}^\dagger f^{-1}a^*a = f^{-1}a^*aaa_{f,f}^\dagger$ .

The equivalences (LVIII)  $\Leftrightarrow$  (XXXIII), (LIX)  $\Leftrightarrow$  (XXXIV) and (L)  $\Leftrightarrow$  (XXXV) follow similarly as (LVII)  $\Leftrightarrow$  (XXXII).

(LXI)  $\Leftrightarrow$  (IV): For  $s, t \geq 1$  and  $a \in \mathcal{A}^\#$ , notice that  $a^s, a^t, a^{s+t} \in \mathcal{A}^\#$ , and hence,  $a^s, a^t, a^{s+t} \in \mathcal{A}^-$ . By Theorem 1.3,  $(a^{s+t})^\dagger = (a^s)_{e,1}^\dagger (a^t)_{1,e}^\dagger$  is equivalent to  $[(a^t e^{-1/2})(e^{1/2} a^s)]^\dagger = (e^{1/2} a^s)^\dagger (a^t e^{-1/2})^\dagger$  which holds, by Theorem 1.11, if and only if  $e^{-1/2}(a^t)^* a^{s+t} \mathcal{A} \subseteq e^{1/2} a^s \mathcal{A}$  and  $e^{1/2} a^s (a^{s+t})^* \mathcal{A} \subseteq e^{-1/2}(a^t)^* \mathcal{A}$ , i.e.,  $(a^t)^* a^{s+t} \mathcal{A} \subseteq ea^s \mathcal{A}$  and  $ea^s (a^{s+t})^* \mathcal{A} \subseteq (a^t)^* \mathcal{A}$ . By elementary computations, this is equivalent to  $a^* \mathcal{A} \subseteq ea \mathcal{A}$  and  $ea \mathcal{A} \subseteq a^* \mathcal{A}$ , that is,  $a^* \mathcal{A} = ea \mathcal{A}$ . Analogy,  $(a^{s+t})^\dagger = (a^s)_{f,1}^\dagger (a^t)_{1,f}^\dagger$ ,  $s, t \geq 1 \Leftrightarrow a^* \mathcal{A} = fa \mathcal{A}$ .

(LXII)  $\Leftrightarrow$  (IV): Observe that, for  $s, t \geq 1$ ,  $(a^{s+t})_{e,f}^\dagger = (a^s)_{f,f}^\dagger (a^t)_{e,f}^\dagger$  is equivalent to  $[(e^{1/2} a^t f^{-1/2})(f^{1/2} a^s f^{-1/2})]^\dagger = (f^{1/2} a^s f^{-1/2})^\dagger (e^{1/2} a^t f^{-1/2})^\dagger$ , by Theorem 1.3. Using Theorem 1.11, the previous equality is equivalent to  $f^{-1/2}(a^t)^* ea^{t+s} f^{-1/2} \mathcal{A} \subseteq f^{1/2} a^s f^{-1/2} \mathcal{A}$  and  $f^{1/2} a^s f^{-1/2} (a^{t+s})^* e^{1/2} \mathcal{A} \subseteq f^{-1/2}(a^t)^* e^{1/2} \mathcal{A}$ , that is,  $(a^t)^* ea^{t+s} \mathcal{A} \subseteq fa^s \mathcal{A}$  and  $fa^s f^{-1/2} (a^{t+s})^* \mathcal{A} \subseteq (a^t)^* \mathcal{A}$ . It follows by elementary computations that this is equivalent to  $a^* \mathcal{A} \subseteq fa \mathcal{A}$  and  $fa \mathcal{A} \subseteq a^* \mathcal{A}$ , i.e.,  $a^* \mathcal{A} = fa \mathcal{A}$ . Similarly,  $(a^{s+t})_{e,f}^\dagger = (a^s)_{e,f}^\dagger (a^t)_{e,e}^\dagger$ ,  $s, t \geq 1$  if and only if  $a^* \mathcal{A} = ea \mathcal{A}$ .  $\square$

From the previous theorem, we can get the following result.

**COROLLARY 2.3.** *Let  $\mathcal{A}$  be a unital  $C^*$ -algebra, and let  $e$  and  $f$  be invertible positive elements in  $\mathcal{A}$ . For  $a \in \mathcal{A}^-$  the following statements are equivalent:*

- (a)  *$a$  is weighted-EP w.r.t.  $(e, f)$ ;*

- (b)  $a^{*f,e}\mathcal{A} = a^{*e,f}\mathcal{A} = a\mathcal{A}$  (or  $a^{*f,e}\mathcal{A} = a\mathcal{A}$  and  $\mathcal{A}a^{*f,e} = \mathcal{A}a$ );
- (c)  $(a^{*f,e})^\circ = a^\circ$  and  ${}^\circ(a^{*f,e}) = {}^\circ a$ ;
- (d)  $a\mathcal{A}^{-1} = a^{*f,e}\mathcal{A}^{-1}$  and  $\mathcal{A}^{-1}a = \mathcal{A}^{-1}a^{*f,e}$ ; and
- (e)  $\mathcal{A}^{-1}a^* = \mathcal{A}^{-1}(a^{*f,e})^*$  and  $a^*\mathcal{A}^{-1} = (a^{*f,e})^*\mathcal{A}^{-1}$ ;

**Acknowledgment.** We are grateful to the referee for helpful comments concerning the paper.

## REFERENCES

- [1] O.M. Baksalary and G. Trenkler. Characterizations of EP, normal and Hermitian matrices. *Linear Multilinear Algebra*, 56:299–304, 2006.
- [2] A. Ben-Israel and T.N.E. Greville. *Generalized Inverses: Theory and Applications*, 2nd edition. Springer, New York, 2003.
- [3] E. Boasso. On the Moore–Penrose inverse, EP Banach space operators, and EP Banach algebra elements. *J. Math. Anal. Appl.*, 339:1003–1014, 2008.
- [4] S.L. Campbell and C.D. Meyer Jr. EP operators and generalized inverses. *Canad. Math. Bull.*, 18:327–333, 1975.
- [5] S. Cheng and Y. Tian. Two sets of new characterizations for normal and EP matrices. *Linear Algebra Appl.*, 375:181–195, 2003.
- [6] D. Cvetković, D.S. Djordjević, and J.J. Koliha. Moore–Penrose inverse in rings with involution. *Linear Algebra Appl.*, 426:371–381, 2007.
- [7] D.S. Djordjević. Products of EP operators on Hilbert spaces. *Proc. Amer. Math. Soc.*, 129(6):1727–1731, 2000.
- [8] D.S. Djordjević. Characterization of normal, hyponormal and EP operators. *J. Math. Anal. Appl.*, 329(2):1181–1190, 2007.
- [9] D.S. Djordjević and J.J. Koliha. Characterizing hermitian, normal and EP operators. *Filomat*, 21(1):39–54, 2007.
- [10] D.S. Djordjević, J.J. Koliha, and I. Straškraba. Factorization of EP elements in  $C^*$ -algebras. *Linear Multilinear Algebra*, 57(6):587–594, 2009.
- [11] D.S. Djordjević and V. Rakočević. *Lectures on generalized inverses*. Faculty of Sciences and Mathematics, University of Niš, 2008.
- [12] M.P. Drazin. Pseudoinverse in associative rings and semigroups. *Amer. Math. Monthly*, 65:506–514, 1958.
- [13] D. Drivaliaris, S. Karanasios, and D. Pappas. Factorizations of EP operators. *Linear Algebra Appl.*, 429(7):1555–1567, 2008.
- [14] R.E. Harte and M. Mbekhta. On generalized inverses in  $C^*$ -algebras. *Studia Math.*, 103:71–77, 1992.
- [15] R.E. Harte and M. Mbekhta. On generalized inverses in  $C^*$ -algebras II. *Studia Math.*, 106:129–138, 1993.
- [16] R.E. Hartwig and I.J. Katz. On products of EP matrices. *Linear Algebra Appl.*, 252:339–345, 1997.
- [17] J.J. Koliha. The Drazin and Moore–Penrose inverse in  $C^*$ -algebras. *Math. Proc. Royal Irish Acad.*, 99A:17–27, 1999.
- [18] J.J. Koliha. A simple proof of the product theorem for EP matrices. *Linear Algebra Appl.*, 294:213–215, 1999.
- [19] J.J. Koliha. Elements of  $C^*$ -algebras commuting with their Moore–Penrose inverse. *Studia Math.*, 139:81–90, 2000.

- [20] J.J. Koliha. A generalized Drazin inverse. *Glasgow Math. J.*, 38:367–381, 1996.
- [21] J.J. Koliha and P. Patrício. Elements of rings with equal spectral idempotents. *J. Australian Math. Soc.*, 72:137–152, 2002.
- [22] G. Lesnjak. Semigroups of EP linear transformations. *Linear Algebra Appl.*, 304(1-3):109–118, 2000.
- [23] D. Mosić, D.S. Djordjević, and J.J. Koliha. EP elements in rings. *Linear Algebra Appl.*, 431:527–535, 2009.
- [24] D. Mosić and D.S. Djordjević. Partial isometries and EP elements in rings with involution. *Electron. J. Linear Algebra*, 18:761–772, 2009.
- [25] D. Mosić and D.S. Djordjević. Reverse order law for the Moore–Penrose inverse in  $C^*$ -algebras. *Electron. J. Linear Algebra*, 22:92–111, 2011.
- [26] D. Mosić and D.S. Djordjević. EP elements in Banach algebras. *Banach J. Math. Anal.*, 5(2):25–32, 2011.
- [27] P. Patrício and R. Puystjens. Drazin–Moore–Penrose invertibility in rings. *Linear Algebra Appl.*, 389:159–173, 2004.
- [28] P. Patrício and A. Veloso da Costa. On the Drazin index of regular elements. *Cent. Eur. J. Math.*, 7(2):200–205, 2009.
- [29] R. Penrose. A generalized inverse for matrices. *Proc. Cambridge Philos. Soc.*, 51:406–413, 1955.
- [30] Y. Tian and H. Wang. Characterizations of EP matrices and weighted-EP matrices. *Linear Algebra Appl.*, 434(5):1295–1318, 2011.