

ROOTS OF THE PHASE OPERATORS

G. JORJADZE AND I. SARISHVILI

ABSTRACT. The k th root taken of the bosonic phase operator is considered. This leads to the extension of the Hilbert space. In the case $k = 2$ an ordinary fermionic extension arises. Particles whose statistics depends on k are introduced for other values of k .

1. PHASE OPERATORS

The problem of polar decomposition of the creation and annihilation operators (\mathbf{a}^* , \mathbf{a}) of a harmonic oscillator goes back to Dirac [1]. In this section some aspects of this problem will be reviewed [2].

Let h be the oscillator hamiltonian

$$h = (p^2 + q^2)/2 \tag{1.1}$$

on the phase plane Γ with coordinates p, q and Poisson brackets (PB)

$$\{q, p\} = 1. \tag{1.2}$$

If the polar angle φ is introduced by

$$p = \sqrt{2h} \sin \varphi, \quad q = \sqrt{2h} \cos \varphi, \tag{1.3}$$

then for the complex variables a and a^*

$$a = \frac{q + ip}{\sqrt{2}}, \quad a^* = \frac{q - ip}{\sqrt{2}}$$

(1.3) is equivalent to the representation

$$a = \sqrt{h} \exp(i\varphi), \quad a^* = \sqrt{h} \exp(-i\varphi). \tag{1.3'}$$

1991 *Mathematics Subject Classification.* 81Q10, 81Q60.

Key words and phrases. Creation and annihilation operators, fermion operators, commutation relations.

Although the variable φ is not global ($\varphi \in S^1$), it is clear that the functions $\exp(\pm i\varphi)$ are correctly defined on the whole phase space Γ (except the origin), and for the PB with hamiltonian (1.1) we have

$$\{h, \exp(\pm i\varphi)\} = \pm i \exp(\pm i\varphi) \quad (1.4)$$

which in local coordinates corresponds to

$$\{h, \varphi\} = 1. \quad (1.4')$$

In the quantum case for the operator \mathbf{h} (we use boldface notation) we choose the normal ordering

$$\mathbf{h} = \mathbf{a}^* \mathbf{a} \equiv \frac{\mathbf{p}^2 + \mathbf{q}^2}{2} - \frac{1}{2} \mathbf{I}. \quad (1.5)$$

Then this \mathbf{h} can be considered a particle (boson) number operator. It is well known that the eigenvalues of \mathbf{h} are nonnegative integers. The corresponding eigenvectors $|n\rangle$ ($n = 0, 1, 2, \dots$)

$$\mathbf{h}|n\rangle = n|n\rangle \quad (1.5')$$

can be constructed by applying the creation operator to the vacuum state $|0\rangle$

$$|n\rangle = \frac{1}{\sqrt{n!}} \mathbf{a}^{*n} |0\rangle$$

and we have

$$\mathbf{a}|n\rangle = \sqrt{n}|n-1\rangle, \quad \mathbf{a}^*|n\rangle = \sqrt{n+1}|n+1\rangle. \quad (1.5'')$$

The vectors $|n\rangle$ form the basis of the Hilbert space. For further convenience we denote this Hilbert space by H_B (bosonic space).

From the correspondence of PB with commutators, for the phase operators $\exp(\pm i\varphi)$ we get (see (1.4))

$$[\mathbf{h}, \exp(\pm i\varphi)] = \mp \exp(\pm i\varphi).$$

So the operator $\exp(i\varphi)$ ($\exp(-i\varphi)$) now decreases (increases) by 1 the eigenvalues of the operator \mathbf{h} , and thus we can write

$$\exp(i\varphi)|n\rangle = \begin{cases} |n-1\rangle, & n > 0, \\ 0, & n = 0, \end{cases} \quad \exp(-i\varphi)|n\rangle = |n+1\rangle \quad (1.6)$$

from which by virtue of (1.5') we have

$$\exp(i\varphi) = \frac{1}{\sqrt{\mathbf{h} + \mathbf{I}}} \mathbf{a}, \quad \exp(-i\varphi) = \mathbf{a}^* \frac{1}{\sqrt{\mathbf{h} + \mathbf{I}}}. \quad (1.6')$$

According to (1.3') these expressions actually correspond to $\exp(\pm i\varphi)$ in the case of some operator ordering while the operator function $\sqrt{\mathbf{h} + \mathbf{I}}$ acts on the basis vectors as a diagonal operator

$$\frac{1}{\sqrt{\mathbf{h} + \mathbf{I}}} |n\rangle = \frac{1}{\sqrt{n+1}} |n\rangle.$$

Thus (1.6) and (1.6') are equivalent and may be regarded as a definition of the phase operators $\mathbf{exp}(\pm i\varphi)$ in the Hilbert space H_B .

To conclude this section, note that although the operators $\mathbf{exp}(i\varphi)$ and $\mathbf{exp}(-i\varphi)$ are mutually conjugate, they satisfy only the one-side unitary relations

$$\mathbf{exp}(-i\varphi) \mathbf{exp}(i\varphi) = \mathbf{I} - |0\rangle\langle 0|, \quad \mathbf{exp}(i\varphi) \mathbf{exp}(-i\varphi) = \mathbf{I} \quad (1.7)$$

where $|0\rangle\langle 0|$ is the projection operator on the vacuum state.

2. SQUARE ROOT OF THE PHASE OPERATORS

In this section we consider a square root taken of the phase operators, i.e., expressions $\mathbf{exp}(\pm i\varphi/2)$.

First note that the corresponding classical functions $\exp(\pm i\varphi/2)$ are not determined on the phase plane as single-valued continuous functions and it is expected that there will be problems in their definition. Formally for the PB we get (see (1.4'))

$$\{h, \exp(\pm i\varphi/2)\} = \pm(i/2) \exp(\pm i\varphi/2)$$

which in the quantum case has to be written in the form

$$[\mathbf{h}, \mathbf{exp}(\pm i\varphi/2)] = \mp(1/2) \mathbf{exp}(\pm i\varphi/2). \quad (2.1)$$

From this follows that the operators $\mathbf{exp}(\pm i\varphi/2)$ must decrease (or increase) by $1/2$ the levels of the oscillator hamiltonian. However, since such states are absent in the spectrum, relations (2.1) cannot be fulfilled. Therefore a meaningful determination of a square root taken of the phase operators is impossible on the Hilbert space H_B .

As is wellknown, since Dirac's equation describes fermions and Dirac's operator $i\gamma^\mu \partial_\mu$ is connected with the square root taken of the d'Alembert's operator ∂^2 [3], we shall try to connect the square root of the phase operators with the fermionic operators. For this we artificially extend the space H_B by introducing the fermionic operators \mathbf{f} and \mathbf{f}^*

$$\mathbf{f}^2 = \mathbf{f}^{*2} = \mathbf{0}, \quad \mathbf{f} \mathbf{f}^* + \mathbf{f}^* \mathbf{f} = \mathbf{I}. \quad (2.2)$$

These relations can be represented in the two-dimensional space H_F with basis vectors $|n\rangle_F$ ($n = 0, 1$), where

$$\mathbf{f}|0\rangle_F = 0, \quad \mathbf{f}|1\rangle_F = |0\rangle_F, \quad \mathbf{f}^*|0\rangle_F = |1\rangle_F, \quad \mathbf{f}^*|1\rangle_F = 0 \quad (2.3)$$

and for the fermionic number operator $\mathbf{N}_F = \mathbf{f}^* \mathbf{f}$ we have

$$[\mathbf{N}_F, \mathbf{f}] = -\mathbf{f}, \quad [\mathbf{N}_F, \mathbf{f}^*] = \mathbf{f}^*, \quad \mathbf{N}_F |n\rangle_F = n|n\rangle_F.$$

Now we shall consider the exterior product of the spaces $\mathcal{H}_F = \mathbf{H}_B \otimes \mathbf{H}_F$ and modify the hamiltonian \mathbf{h} :

$$\mathbf{H} = \mathbf{h} \otimes \mathbf{I}_F + (1/2)\mathbf{I}_B \otimes \mathbf{N}_F \quad (2.4)$$

with \mathbf{I}_B and \mathbf{I}_F as identity operators on the spaces H_B and H_F , respectively.

Below, for simplicity, $\mathbf{h} \otimes \mathbf{I}_F$ and $\mathbf{I}_B \otimes \mathbf{N}_F$ will be replaced by \mathbf{h} and \mathbf{N}_F , respectively, i.e.,

$$\mathbf{H} = \mathbf{h} + (1/2)\mathbf{N}_F = \mathbf{a}^* \mathbf{a} + (1/2)\mathbf{f}^* \mathbf{f}. \quad (2.4')$$

The eigenstates of the new hamiltonian \mathbf{H} are characterized by two numbers $|n_B, n_F\rangle$,

$$\mathbf{H}|n_B, n_F\rangle = (n_B + \frac{1}{2}n_F)|n_B, n_F\rangle \quad (2.4'')$$

where $n_B = 0, 1, 2, \dots, n_F = 0, 1$.

Introducing the operators \mathbf{A}^\pm

$$\mathbf{A}^+ = \mathbf{f}^* + \exp(-i\varphi)\mathbf{f}, \quad \mathbf{A}^- = \mathbf{f} + \exp(+i\varphi)\mathbf{f}^*, \quad (2.5)$$

it is easy to verify that the operator \mathbf{A}^+ increases by 1/2 and the operator \mathbf{A}^- decreases by 1/2 every eigenvalue level of the operator \mathbf{H} (except the vacuum state which is canceled by the action of \mathbf{A}^-). So the following relations are satisfied:

$$[\mathbf{H}, \mathbf{A}^\pm] = \pm(1/2)\mathbf{A}^\pm. \quad (2.5')$$

Considering quadratic combinations of the operators \mathbf{A}^\pm , it is easy to verify (see (2.2) and (1.7)) that

$$\begin{aligned} (\mathbf{A}^+)^2 &= \exp(-i\varphi), & (\mathbf{A}^-)^2 &= \exp(i\varphi), \\ \mathbf{A}^+ \mathbf{A}^- &= \mathbf{I} - |0, 0\rangle\langle 0, 0|, & \mathbf{A}^- \mathbf{A}^+ &= \mathbf{I}. \end{aligned} \quad (2.6)$$

From this (also compare (2.1) and (2.5')) one concludes that the operators \mathbf{A}^\pm may be considered as a definition of the operators $\exp(\mp i\varphi/2)$:

$$\exp(\mp i\varphi/2) \equiv \mathbf{A}^\pm. \quad (2.7)$$

Thus on the extended space \mathcal{H}_F it is possible to define the square root of the phase operators which are connected with fermionic operators.

3. THE k TH ROOT OF PHASE OPERATORS

The arguments of Section 2 can be generalized to the case of arbitrary k ($k > 2$) if we consider the k th root taken of phase operators

$$\exp(\pm i\varphi/k).$$

For this purpose we introduce the k -dimensional unitary space H_k with the orthonormal basis

$$|0\rangle_k, |1\rangle_k, |2\rangle_k, \dots, |k-1\rangle_k \quad (3.1)$$

and define the operators $\mathbf{f}_k, \mathbf{f}_k^*$:

$$\mathbf{f}_k |n\rangle_k = \begin{cases} |n-1\rangle_k, & n \geq 1, \\ 0, & n = 0; \end{cases} \quad (3.2)$$

$$\mathbf{f}_k^* |n\rangle_k = \begin{cases} |n+1\rangle_k, & n \leq k-2, \\ 0, & n = k-1. \end{cases} \quad (3.2')$$

In the basis (3.1) the operators \mathbf{f}_k and \mathbf{f}_k^* have the representation

$$\mathbf{f}_k = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \cdot & \cdot & \cdot & \dots & \cdot & \cdot \\ 0 & 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{pmatrix}, \quad \mathbf{f}_k^* = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ \cdot & \cdot & \cdot & \dots & \cdot & \cdot \\ 0 & 0 & 0 & \dots & 1 & 0 \end{pmatrix}.$$

It is obvious that

$$(\mathbf{f}_k)^k = (\mathbf{f}_k^*)^k = \mathbf{0} \quad (3.3)$$

and it is also easy to verify that the k^2 number of “normally” ordered monomes $(\mathbf{f}_k^*)^j (\mathbf{f}_k)^m$, where $0 \leq j \leq k-1$, $0 \leq m \leq k-1$, give k^2 linearly independent matrices and form the basis in the space of linear operators on H_k . Therefore any monome can be written as a linear combination of normally ordered monomes. For example, $\mathbf{f}_k \mathbf{f}_k^*$ takes the form

$$\mathbf{f}_k \mathbf{f}_k^* = \mathbf{I} - (\mathbf{f}_k^*)^{k-1} (\mathbf{f}_k)^{k-1} \quad (3.4)$$

which is the generalization of the commutation relations (2.2) in the case of arbitrary k . It is possible to compute other operator relations; in particular we have

$$\sum_{j=0}^{k-1} (\mathbf{f})^j (\mathbf{f}^*)^{k-1} (\mathbf{f})^{k-j-1} = \mathbf{I}, \quad (3.5)$$

$$(\mathbf{f})^j (\mathbf{f}^*)^m = \mathbf{0}, \quad \text{if } j - m \geq k.$$

The particle number operator \mathbf{N}_k can be introduced on the space \mathbb{H}_k as

$$\mathbf{N}_k |n\rangle_k = n |n\rangle_k.$$

It is easy to verify that the operator \mathbf{N}_k can be written in the form

$$\mathbf{N}_k = \mathbf{f}_k^* \mathbf{f}_k + \mathbf{f}_k^{*2} \mathbf{f}_k^2 + \cdots + (\mathbf{f}_k^*)^{k-1} (\mathbf{f}_k)^{k-1} \quad (3.6)$$

and satisfies the relations

$$[\mathbf{N}_k, \mathbf{f}_k] = -\mathbf{f}_k, \quad [\mathbf{N}_k, \mathbf{f}_k^*] = \mathbf{f}_k^*. \quad (3.6')$$

After this let us consider the exterior product of the spaces \mathbb{H}_B and \mathbb{H}_k

$$\mathcal{H}_k = \mathbb{H}_B \otimes \mathbb{H}_k \quad (3.7)$$

and define the hamiltonian on \mathcal{H}_k

$$\mathbf{H} = \mathbf{h} + \frac{1}{k} \mathbf{N}_k \quad (3.8)$$

where we use the above-mentioned abbreviation (see (2.4)).

The eigenvectors of the new hamiltonian \mathbf{H} are characterized by two quantum numbers $|n_B, n_k\rangle$, where $n_B = 0, 1, 2, \dots$, $n_k = 0, 1, 2, \dots, k-1$, and we have

$$\mathbf{H} |n_B, n_k\rangle = (n_B + \frac{1}{k} n_k) |n_B, n_k\rangle. \quad (3.9)$$

The energy levels do not degenerate and are equidistant with an interval $1/k$.

Here, by analogy with (2.5), the operators \mathbf{A}_k^\pm can be introduced in the following way:

$$\begin{aligned} \mathbf{A}_k^+ &= \mathbf{f}_k^* + \exp(-i\varphi) (\mathbf{f}_k)^{k-1}, \\ \mathbf{A}_k^- &= \mathbf{f}_k + \exp(i\varphi) (\mathbf{f}_k^*)^{k-1}. \end{aligned} \quad (3.10)$$

They change any energy level by $1/k$ (\mathbf{A}_k^+ increases and \mathbf{A}_k^- decreases, except the vacuum state which is canceled by the action of \mathbf{A}_k^-) and satisfy the commutation relations

$$[\mathbf{H}, \mathbf{A}_k^\pm] = \pm \frac{1}{k} \mathbf{A}_k^\pm.$$

Now, using (3.4) and (3.5), we can verify that

$$(\mathbf{A}_k^+)^k = \exp(-i\varphi), \quad (\mathbf{A}_k^-)^k = \exp(i\varphi), \quad (3.11)$$

and also

$$\mathbf{A}_k^- \mathbf{A}_k^+ = \mathbf{I}, \quad \mathbf{A}_k^+ \mathbf{A}_k^- = \mathbf{I} - |0, 0\rangle \langle 0, 0|.$$

So after all this our result can be formulated as

Theorem. *The k th root of the phase operators is defined on the Hilbert space $l_2 \otimes H_k$ and has form (3.10).*

Finally, note that the correct definition of nonsingle-valued functions $\exp(\pm i\varphi/k)$ can be made on the k -sheet Riemann surface. In [4] the case $k = 2$ was considered. As a result, one can conclude that the quantization on the two-sheet Riemann surface is connected with the introduction of fermionic degrees of freedom. It is easy to verify that this situation is valid for arbitrary k . On the other hand, the case of arbitrary k can be related with anyon physics and fractional statistics [5].

REFERENCES

1. P. A. M. Dirac, The elimination of nodes in quantum mechanics. *Proc. R. Soc.* **A111**(1926), 281-305.
2. P. Carruthers and M. Nieto, Action-angle variables in quantum mechanics. *Rev. Mod. Phys.* **40** (1968), 411-441.
3. S. S. Schweber, An introduction to relativistic quantum field theory. *Row, Peterson and Co, New York*, 1961.
4. D. N. Gordeziani, G. P. Jorjadze, and I. T. Sarishvili, Approximation of relativistic strings. (Russian) *Proc. of XI Workshop "Problems of high energy Physics and Field Theory," Protvino - 1988*, 69-72, *Nauka, Moscow*, 1989.
5. D. P. Arovas, R. Schrieffer, F. Wilczek, and A. Zee, Statistical mechanics of anions. *Nucl. Phys. B* **251**(1985), 117-126.

(Received 15.09.1993)

Authors' address:
A. Razmadze Mathematical Institute
Georgian Academy of Sciences
1, Rukhadze St., Tbilisi 380093
Republic of Georgia