

CROSSED SEMIMODULES AND SCHREIER INTERNAL CATEGORIES IN THE CATEGORY OF MONOIDS

A. PATCHKORIA

ABSTRACT. We introduce the notion of a Schreier internal category in the category of monoids and prove that the category of Schreier internal categories in the category of monoids is equivalent to the category of crossed semimodules. This extends a well-known equivalence of categories between the category of internal categories in the category of groups and the category of crossed modules.

1. INTRODUCTION AND STATEMENT OF RESULTS

The description of internal categories in the category of groups as crossed modules is well known (see, e.g., [1]). There exist the same descriptions of internal categories in Rings, Lie algebras, etc. One can find different descriptions (not as crossed modules) of internal categories and groupoids in groups, Mal'tsev varieties of universal algebras, congruence modular varieties, and Mal'tsev categories in [2], [3], [4] and [5] respectively.

Let *Mon* denote the category of monoids. In this note we show that there exists an equivalence between the category of internal categories in *Mon* (note that monoids are not congruence modular) satisfying the so-called Schreier condition and the category of crossed semimodules. This answers a question posed by G. Janelidze (oral communication).

By a crossed semimodule we mean a pair

$$\Phi = (\mu : A \rightarrow X, \varphi : X \rightarrow \text{End}(A)),$$

where A, X are monoids and μ, φ are homomorphisms of monoids satisfying

$$\text{(CSM 1) } \mu(\varphi(x)(a)) + x = x + \mu(a), \quad x \in X, a \in A,$$

$$\text{(CSM 2) } \varphi(\mu(a))(a') + a = a + a', \quad a, a' \in A.$$

1991 *Mathematics Subject Classification.* 18B40, 20M50.

Key words and phrases. Monoid, crossed semimodule, internal category, internal groupoid, Schreier condition.

A crossed semimodule morphism is a commutative diagram

$$\begin{array}{ccc} \Phi : A & \xrightarrow{\mu} & X \\ \tau = (\beta, \lambda) \downarrow & & \downarrow \lambda \\ & \beta \downarrow & \\ \Phi' : A' & \xrightarrow{\mu'} & X' \end{array}$$

with β and λ monoid homomorphisms, which satisfies

$$\beta(\varphi(x)(a)) = \varphi'(\lambda(x))(\beta(a))$$

for all $x \in X$ and $a \in A$.

Recall that an internal category in the category of monoids is a diagram

$$M : \begin{array}{ccccc} & \xrightarrow{\pi_1} & & \xleftarrow{s_0} & \\ M_2 & \xrightarrow{m} & M_1 & \xrightarrow{d_0} & M_0 \\ & \xrightarrow{\pi_2} & & \xrightarrow{d_1} & \end{array} ,$$

where M_0 is the monoid of objects, M_1 the monoid of morphisms, and d_0, d_1, s_0 are homomorphisms of monoids called the domain, the codomain, and the identity respectively; M_2 together with π_1 and π_2 forms the pullback

$$\begin{array}{ccc} M_2 & \xrightarrow{\pi_2} & M_1 \\ \pi_1 \downarrow & & \downarrow d_1 \\ M_1 & \xrightarrow{d_0} & M_0 \end{array} ;$$

m is a homomorphism of monoids called the composition of M and usually written as $m(g, f) = g \circ f$; and the following conditions hold:

- (i) $d_0 s_0 = d_1 s_0 = 1_{M_0}$,
- (ii) $d_0 m = d_0 \pi_1, d_1 m = d_1 \pi_2$,
- (iii) $m(m(h, g), f) = m(h, m(g, f))$,
- (iv) $m(s_0 d_1(f), f) = f = m(f, s_0 d_0(f))$.

A morphism of internal categories in Mon is a commutative diagram

$$\begin{array}{ccccccc}
 & & \xrightarrow{\pi_1} & & \xleftarrow{s_0} & & \\
 M : & M_2 & \xrightarrow{m} & M_1 & \xrightarrow{d_0} & M_0 & \\
 & \downarrow & & \downarrow & & \downarrow & \\
 \gamma = (\gamma_1, \gamma_0) & \gamma_1 \times \gamma_1 & & \gamma_1 & & \gamma_0 & \\
 & & \xrightarrow{\pi_2} & & \xrightarrow{d_1} & & \\
 M' : & M'_2 & \xrightarrow{m'} & M'_1 & \xrightarrow{d'_0} & M'_0 & \\
 & & \xrightarrow{\pi'_1} & & \xleftarrow{s'_0} & & \\
 & & & & \xrightarrow{d'_1} & & \\
 & & \xrightarrow{\pi'_2} & & \xrightarrow{d'_1} & &
 \end{array}$$

where γ_0 and γ_1 are monoid homomorphisms.

An internal groupoid in Mon is an internal category M in Mon in which every morphism has an inverse, i.e., for every $f \in M_1$ there exists a unique $f^{-1} \in M_1$ such that $m(f, f^{-1}) = s_0 d_1(f)$ and $m(f^{-1}, f) = s_0 d_0(f)$.

Definition. A Schreier internal category in Mon is an internal category in Mon which satisfies the Schreier condition: for any $f \in M_1$ there exists a unique $g \in \text{Ker}(d_0)$ such that

$$f = g + s_0 d_0(f).$$

We prove

Theorem. *The category of Schreier internal categories in the category of monoids is equivalent to the category of crossed semimodules.*

Restricting this equivalence gives

Corollary 1. *The category of Schreier internal groupoids in the category of monoids is equivalent to the category of crossed semimodules $\Phi = (\mu : A \rightarrow X, \varphi : X \rightarrow \text{End}(A))$ such that A is a group.*

Note that Corollary 1 can be obtained as a special case of the theorem of [6].

Restricting this equivalence further, we obtain

Corollary 2 (R. Brown and C. B. Spencer [1]). *The category of internal categories (= groupoids) in the category of groups is equivalent to the category of crossed modules.*

2. THE PROOFS

Let

$$x \xrightarrow{f} y \xrightarrow{f'} z$$

be a composable pair in a Schreier internal category M in Mon . Then there exist uniquely defined $g, h \in \text{Ker}(d_0)$ such that

$$f = g + s_0 d_0(f) \quad \text{and} \quad f' = h + s_0(d_1(g) + d_0(f)).$$

Since m is a monoid homomorphism and (iv) holds, we obtain

$$\begin{aligned} m(f', f) &= m(h + s_0(d_1(g) + d_0(f)), g + s_0 d_0(f)) = \\ &= m(h, 0) + m(s_0(d_1(g) + d_0(f)), g + s_0 d_0(f)) = \\ &= m(h, s_0 d_0(h)) + m(s_0 d_1(g + s_0 d_0(f)), g + s_0 d_0(f)) = \\ &= h + g + s_0 d_0(f). \end{aligned}$$

So the composition m is uniquely defined,

$$m(h + s_0(d_1(g) + d_0(f)), g + s_0 d_0(f)) = h + g + s_0 d_0(f). \quad (1)$$

Proof of the theorem. Let

$$M : \quad \begin{array}{ccccc} & & \xrightarrow{\pi_1} & & \\ & & \xrightarrow{m} & & \\ & & \xrightarrow{\pi_2} & & \\ M_2 & & & M_1 & & M_0 \\ & & & \xleftarrow{s_0} & \\ & & & \xrightarrow{d_0} & \\ & & & \xrightarrow{d_1} & \end{array}$$

be a Schreier internal category in Mon . By the Schreier condition, for any $x \in M_0$ and any $g \in \text{Ker}(d_0)$ there exists a unique $\alpha(x, g) \in \text{Ker}(d_0)$ such that

$$s_0(x) + g = \alpha(x, g) + s_0(x).$$

We have

$$\begin{aligned} \alpha(x, g_1 + g_2) + s_0(x) &= s_0(x) + g_1 + g_2 = \\ &= \alpha(x, g_1) + s_0(x) + g_2 = \alpha(x, g_1) + \alpha(x, g_2) + s_0(x), \\ \alpha(x, 0) + s_0(x) &= s_0(x), \\ \alpha(x + y, g) + s_0(x + y) &= s_0(x + y) + g = s_0(x) + s_0(y) + g = \\ &= s_0(x) + \alpha(y, g) + s_0(y) = \alpha(x, \alpha(y, g)) + s_0(x) + s_0(y) = \\ &= \alpha(x, \alpha(y, g)) + s_0(x + y), \\ \alpha(0, g) + s(0) &= s(0) + g. \end{aligned}$$

Hence, in view of the Schreier condition, we obtain

$$\alpha(x, g_1 + g_2) = \alpha(x, g_1) + \alpha(x, g_2), \quad \alpha(x, 0) = 0,$$

$$\alpha(x + y, g) = \alpha(x, \alpha(y, g)), \quad \alpha(0, g) = g.$$

This means that we have a homomorphism of monoids

$$\psi : M_0 \rightarrow \text{End}(\text{Ker}(d_0)), \quad \psi(x)(g) = \alpha(x, g), \quad x \in M_0, \quad g \in \text{Ker}(d_0).$$

Consider the pair of monoid homomorphisms

$$\Psi = \left(d_1 : \text{Ker}(d_0) \rightarrow M_0, \psi : M_0 \rightarrow \text{End}(\text{Ker}(d_0)) \right).$$

It is clear that

$$d_1(\psi(x)(g)) + x = x + d_1(g), \quad x \in M_0, \quad g \in \text{Ker}(d_0)$$

since $s_0(x) + g = \psi(x)(g) + s_0(x)$ and $d_1 s_0 = 1_{M_0}$.

On the other hand,

$$s_0 d_1(g) + g' = \psi(d_1(g))(g') + s_0 d_1(g), \quad g, g' \in \text{Ker}(d_0).$$

This and (1) give

$$\begin{aligned} \psi(d_1(g))(g') + g &= m(\psi(d_1(g))(g') + s_0 d_1(g), g) = \\ &= m(s_0 d_1(g) + g', g) = m(s_0 d_1(g), g) + m(g', 0) = g + g', \end{aligned}$$

i.e.,

$$\psi(d_1(g))(g') + g = g + g', \quad g, g' \in \text{Ker}(d_0).$$

So Ψ is a crossed semimodule.

Let $\gamma = (\gamma_1, \gamma_0) : M \rightarrow M'$ be a morphism of Schreier internal categories in *Mon*. The Schreier condition allows us to write

$$\begin{aligned} \gamma_1(\psi(x)(g)) + s'_0 \gamma_0(x) &= \gamma_1(\psi(x)(g)) + \gamma_1 s_0(x) = \\ &= \gamma_1(\psi(x)(g) + s_0(x)) = \gamma_1(s_0(x) + g) = s'_0 \gamma_0(x) + \gamma_1(g) = \\ &= \psi'(\gamma_0(x))(\gamma_1(g)) + s'_0 \gamma_0(x), \end{aligned}$$

i.e.,

$$\gamma_1(\psi(x)(g)) + s'_0 \gamma_0(x) = \psi'(\gamma_0(x))(\gamma_1(g)) + s'_0 \gamma_0(x).$$

From this, by the same condition, we obtain

$$\gamma_1(\psi(x)(g)) = \psi'(\gamma_0(x))(\gamma_1(g)), \quad x \in M_0, \quad g \in \text{Ker}(d_0).$$

This means that the commutative diagram

$$\begin{array}{ccc} \text{Ker}(d_0) & \xrightarrow{d_1} & M_0 \\ \gamma_1 \downarrow & & \gamma_0 \downarrow \\ \text{Ker}(d'_0) & \xrightarrow{d'_1} & M'_0 \end{array}$$

is a crossed semimodules morphism.

So we have the functor

$$S : \text{Schreier internal categories in Mon} \rightarrow \text{Crossed semimodules},$$

defined by $S(M) = \Psi$ and $S(\gamma_1, \gamma_0) = (\gamma_1|_{\text{Ker}(d_0)}, \gamma_0) : \Psi \rightarrow \Psi'$.

There is a standard way to show that S is an equivalence of categories. Let

$$\Phi = \left(\mu : A \rightarrow X, \varphi : X \rightarrow \text{End}(A) \right)$$

be a crossed semimodule. Using the semidirect product $A \times_{\varphi} X$, we obtain the diagram

$$C : (A \times_{\varphi} X)_{d_0} \times_{d_1} (A \times_{\varphi} X) \begin{array}{ccc} \xrightarrow{\pi_1} & & \xleftarrow{s_0} \\ \xrightarrow{m} & A \times_{\varphi} X & \xrightarrow{d_0} X \\ \xrightarrow{\pi_2} & & \xrightarrow{d_1} \end{array}$$

where $d_0(a, x) = x$, $d_1(a, x) = \mu(a) + x$, $s_0(x) = (0, x)$, $m((a', \mu(a) + x), (a, x)) = (a' + a, x)$ and π_1, π_2 are the projections. It is plain to see that C is a Schreier internal category in *Mon*. On the other hand, any crossed semimodule morphism $\tau = (\beta, \lambda) : \Phi \rightarrow \Phi'$ defines the morphism of Schreier internal categories $\gamma = (\gamma_1, \lambda) : C \rightarrow C'$, where $\gamma_1(a, x) = (\beta(a), \lambda(x))$.

Thus we have the functor

$$T : \text{Crossed semimodules} \rightarrow \text{Schreier internal categories in Mon},$$

defined by $T(\Phi) = C$ and $T(\tau) = \gamma : C \rightarrow C'$.

It is straightforward to see that $1 \cong ST$. Due to the Schreier condition, $1 \cong TS$ holds, too. \square

Proof of Corollary 1. Suppose M is a Schreier internal groupoid in *Mon*. Consider the crossed semimodule

$$S(M) = \left(d_1 : \text{Ker}(d_0) \rightarrow M_0, \psi : M_0 \rightarrow \text{End}(\text{Ker}(d_0)) \right).$$

For any $g \in \text{Ker}(d_0)$ there exists a unique $h \in \text{Ker}(d_0)$ such that

$$m(h + s_0 d_1(g), g) = 0 \quad \text{and} \quad m(g, h + s_0 d_1(g)) = s_0 d_1(g).$$

Using (1), we obtain

$$h + g = 0 \quad \text{and} \quad g + h + s_0 d_1(g) = s_0 d_1(g).$$

Then the Schreier condition gives

$$h + g = 0 \quad \text{and} \quad g + h = 0,$$

i.e., $\text{Ker}(d_0)$ is a group.

Conversely, suppose $\Phi = (\mu : A \rightarrow X, \varphi : X \rightarrow \text{End}(A))$ is a semimodule such that A is a group. Then it is clear that for any morphism (a, x) of $T(\Phi)$ we have $(a, x)^{-1} = (-a, \mu(a) + x)$, i.e., $T(\Phi)$ is a Schreier internal groupoid in *Mon*. \square

ACKNOWLEDGMENT

This research was supported by INTAS Grant 93-436 "Algebraic K -theory, groups and categories."

REFERENCES

1. R. Brown and C. B. Spencer, G -groupoids, crossed modules and the fundamental groupoid of a topological group. *Proc. Konink. Nederl. Akad. Wetensch.* **79**(1976) 296–302.
2. J.-L. Loday, Spaces with finitely many non-trivial homotopy groups. *J. Pure Appl. Algebra* **24**(1982) 179–202.
3. G. Janelidze, Internal categories in Mal'cev varieties. *Preprint, The University of North York, Toronto*, 1990.
4. G. Janelidze and M. C. Pedicchio, Internal categories and groupoids in congruence modular varieties. *Preprint, McGill University, Montreal*, 1995.
5. A. Carboni, M. C. Pedicchio, and N. Pirovano, Internal graphs and internal groupoids in Mal'cev categories. *Category Theory 1991 (Montreal, PQ, 1991)*, 97–109, *CMS Conf. Proc.* 13, Amer. Math. Soc., Providence, RI, 1992.
6. T. Porter, Crossed modules in Cat and a Brown–Spencer theorem for 2-categories. *Cahiers Topologie Géom. Différentielle Catégoriques* **26** (1985), 381–388.

(Received 29.07.1996)

Author's address:

A. Razmadze Mathematical Institute
Georgian Academy of Sciences
1, M. Aleksidze St., Tbilisi 380093
Georgia