

Research Article

Nonlocal Cauchy Problem for Nonautonomous Fractional Evolution Equations

Fei Xiao

Department of Mathematics, University of Science and Technology of China, Hefei, Anhui 230026, China

Correspondence should be addressed to Fei Xiao, sheaf@mail.ustc.edu.cn

Received 28 November 2010; Accepted 29 January 2011

Academic Editor: Toka Diagana

Copyright © 2011 Fei Xiao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the mild solutions of a nonlocal Cauchy problem for nonautonomous fractional evolution equations $d^q u(t)/dt^q = -A(t)u(t) + f(t, (K_1 u)(t), (K_2 u)(t), \dots, (K_n u)(t))$, $t \in I = [0, T]$, $u(0) = A^{-1}(0)g(u) + u_0$, in Banach spaces, where $T > 0$, $0 < q < 1$. New results are obtained by using Sadovskii's fixed point theorem and the Banach contraction mapping principle. An example is also given.

1. Introduction

During the past decades, the fractional differential equations have been proved to be valuable tools in the investigation of many phenomena in engineering and physics; they attracted many researchers (cf., e.g., [1–9]). On the other hand, the autonomous and nonautonomous evolution equations and related topics were studied in, for example, [6, 7, 10–20], and the nonlocal Cauchy problem was considered in, for example, [2, 5, 18, 21–26].

In this paper, we consider the following nonlocal Cauchy problem for nonautonomous fractional evolution equations

$$\begin{aligned} \frac{d^q u(t)}{dt^q} &= -A(t)u(t) + f(t, (K_1 u)(t), (K_2 u)(t), \dots, (K_n u)(t)), \quad t \in I = [0, T], \\ u(0) &= A^{-1}(0)g(u) + u_0, \end{aligned} \tag{1.1}$$

in Banach spaces, where $0 < q < 1$, $g : C(I; X) \rightarrow X$. The terms $(K_i u)(t)$, $i = 1, \dots, n$ are

defined by

$$(K_i u)(t) = \int_0^t k_i(t, s)u(s)ds, \quad (1.2)$$

the positive functions $k_i(t, s)$ are continuous on $D = \{(t, s) \in \mathbb{R}^2 : 0 \leq s \leq t \leq T\}$ and

$$K_i^* = \sup_{t \in [0, T]} \int_0^t k_i(t, s)ds < \infty. \quad (1.3)$$

Let us assume that $u \in L([0, T]; X)$ and $A(t)$ is a family linear closed operator defined in a Banach space X . The fractional order integral of the function u is understood here in the Riemann-Liouville sense, that is,

$$I^q u(t) = \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} u(s)ds. \quad (1.4)$$

In this paper, we denote that C is a positive constant and assume that a family of closed linear $\{A(t) : t \in [0, T]\}$ satisfying

- (A1) the domain $D(A)$ of $\{A(t) : t \in [0, T]\}$ is dense in the Banach space X and independent of t ,
- (A2) the operator $[A(t) + \lambda]^{-1}$ exists in $L(X)$ for any λ with $\text{Re } \lambda \leq 0$ and

$$\| [A(t) + \lambda]^{-1} \| \leq \frac{C}{|\lambda + 1|}, \quad t \in [0, T]. \quad (1.5)$$

- (A3) There exists constant $\gamma \in (0, 1]$ and C such that

$$\| [A(t_1) - A(t_2)]A^{-1}(s) \| \leq C|t_1 - t_2|^\gamma, \quad t_1, t_2, s \in [0, T]. \quad (1.6)$$

Under condition (A2), each operator $-A(s)$, $s \in [0, T]$ generates an analytic semigroup $\exp(-tA(s))$, $t > 0$, and there exists a constant C such that

$$\| A^n(s) \exp(-tA(s)) \| \leq \frac{C}{t^n}, \quad (1.7)$$

where $n = 0, 1, t > 0, s \in [0, T]$ ([11]).

We study the existence of mild solution of (1.1) and obtain the existence theorem based on the measures of noncompactness. An example is given to show an application of the abstract results.

2. Preliminaries

Throughout this work, we set $I = [0, T]$. We denote by X a Banach space, $L(X)$ the space of all linear and bounded operators on X , and $C(I, X)$ the space of all X -valued continuous functions on I .

Lemma 2.1 (see [9]). (1) $I^q : L^1[0, T] \rightarrow L^1[0, T]$.

(2) For $g \in L^1[0, T]$, we have

$$\int_0^t \int_0^\eta (t - \eta)^{q-1} (\eta - s)^{\gamma-1} g(s) ds d\eta = B(q, \gamma) \int_0^t (t - s)^{q+\gamma-1} g(s) ds, \quad (2.1)$$

where $B(q, \gamma)$ is a Beta function.

Definition 2.2. Let B be a bounded set of seminormed linear space Y . The Kuratowski's measure of noncompactness (for brevity, α -measure) of B is defined as

$$\alpha(B) = \inf\{d > 0 : B \text{ has a finite cover by sets of diameter } \leq d\}. \quad (2.2)$$

From the definition, we can get some properties of α -measure immediately, see ([27]).

Lemma 2.3 (see [27]). *Let A and B be bounded sets of X . Then*

- (1) $\alpha(A) \leq \alpha(B)$, if $A \subseteq B$.
- (2) $\alpha(A) = \alpha(A^{\text{cl}})$, where A^{cl} denotes the closure of A .
- (3) $\alpha(A) = 0$ if and only if A is precompact.
- (4) $\alpha(\lambda A) = |\lambda| \alpha(A)$, $\lambda \in \mathbb{R}$.
- (5) $\alpha(A \cup B) = \max\{\alpha(A), \alpha(B)\}$.
- (6) $\alpha(A + B) \leq \alpha(A) + \alpha(B)$, where $A + B = \{x + y : x \in A, y \in B\}$.
- (7) $\alpha(A + x_0) = \alpha(A)$, for any $x_0 \in X$.

For $H \subset C(I, X)$ we define

$$\int_0^t H(s) ds = \left\{ \int_0^t u(s) ds : u \in H \right\}, \quad (2.3)$$

for $t \in I$, where $H(s) = \{u(s) \in X : u \in H\}$.

The following lemma will be needed.

Lemma 2.4 (see [27]). *If $H \subset C(I, X)$ is a bounded, equicontinuous set, then*

- (1) $\alpha(H) = \sup_{t \in I} \alpha(H(t))$.
- (2) $\alpha(\int_0^t H(s) ds) \leq \int_0^t \alpha(H(s)) ds$, for $t \in I$.

Lemma 2.5 (see [28]). *If $\{u_n\}_{n=1}^\infty \subset L^1(I, X)$ and there exists a $m(\cdot) \in L^1(I, \mathbb{R}^+)$ such that*

$$\|u_n(t)\| \leq m(t), \quad \text{a.e } t \in I, \quad (2.4)$$

then $\alpha(\{u_n(t)\}_{n=1}^\infty)$ is integrable and

$$\alpha\left(\left\{\int_0^t u_n(s) ds\right\}_{n=1}^\infty\right) \leq 2 \int_0^t \alpha(\{u_n(s)\}_{n=1}^\infty) ds. \quad (2.5)$$

We need to use the following Sadovskii's fixed point theorem.

Definition 2.6 (see [29]). *Let P be an operator in Banach space X . If P is continuous and takes bounded sets into bounded sets, and $\alpha(P(H)) < \alpha(H)$ for every bounded set H of X with $\alpha(H) > 0$, then P is said to be a condensing operator on X .*

Lemma 2.7 (Sadovskii's fixed point theorem [29]). *Let P be a condensing operator on Banach space X . If $P(B) \subseteq B$ for a convex, closed, and bounded set B of X , then P has a fixed point in B .*

According to [4], a mild solution of (1.1) can be defined as follows.

Definition 2.8. A function $u \in C(I, X)$ satisfying the equation

$$\begin{aligned} u(t) = & A^{-1}(0)g(u) + u_0 + \int_0^t \varphi(t-\eta, \eta)U(\eta)A(0)\left[A^{-1}(0)g(u) + u_0\right]d\eta \\ & + \int_0^t \varphi(t-\eta, \eta)f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta))d\eta \\ & + \int_0^t \int_0^\eta \varphi(t-\eta, \eta)\varphi(\eta, s)f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s))ds d\eta, \end{aligned} \quad (2.6)$$

is called a mild solution of (1.1), where

$$\varphi(t, s) = q \int_0^\infty \theta t^{q-1} \xi_q(\theta) \exp(-t^q \theta A(s)) d\theta, \quad (2.7)$$

and ξ_q is a probability density function defined on $[0, \infty)$ such that its Laplace transform is given by

$$\int_0^\infty e^{-\sigma x} \xi_q(\sigma) d\sigma = \sum_{j=0}^\infty \frac{(-x)^j}{\Gamma(1+qj)}, \quad q \in (0, 1], x > 0, \quad (2.8)$$

$$\varphi(t, \tau) = \sum_{k=1}^\infty \varphi_k(t, \tau),$$

where

$$\begin{aligned} \varphi_1(t, \tau) &= [A(t) - A(\tau)]\varphi(t - \tau, \tau), \\ \varphi_{k+1}(t, \tau) &= \int_{\tau}^t \varphi_k(t, s)\varphi_1(s, \tau)ds, \quad k = 1, 2, \dots, \\ U(t) &= -A(t)A^{-1}(0) - \int_0^t \varphi(t, s)A(s)A^{-1}(0)ds. \end{aligned} \tag{2.9}$$

To our purpose, the following conclusions will be needed. For the proofs refer to [4].

Lemma 2.9 (see [4]). *The operator-valued functions $\varphi(t - \eta, \eta)$ and $A(t)\varphi(t - \eta, \eta)$ are continuous in uniform topology in the variables t, η , where $0 \leq \eta \leq t - \varepsilon, 0 \leq t \leq T$, for any $\varepsilon > 0$. Clearly,*

$$\|\varphi(t - \eta, \eta)\| \leq C(t - \eta)^{q-1}. \tag{2.10}$$

Moreover, we have

$$\|\varphi(t, \eta)\| \leq C(t - \eta)^{\gamma-1}. \tag{2.11}$$

Remark 2.10. From the proof of Theorem 2.5 in [4], we can see

- (1) $\|U(t)\| \leq C + Ct^\gamma$.
- (2) For $t \in I, \int_0^t \varphi(t - \eta, \eta)U(\eta)d\eta$ is uniformly continuous in the norm of $L(X)$ and

$$\left\| \int_0^t \varphi(t - \eta, \eta)U(\eta)d\eta \right\| \leq C^2 t^q \left(\frac{1}{q} + t^\gamma B(q, \gamma + 1) \right) := \widetilde{M}(t). \tag{2.12}$$

3. Existence of Solution

Assume that

- (B1) $f : I \times X \times X \times \dots \times X \rightarrow X$ satisfies $f(\cdot, v_1, v_2, \dots, v_n) : I \rightarrow X$ is measurable for all $v_i \in X, i = 1, 2, \dots, n$ and $f(t, \cdot, \cdot, \dots, \cdot) : X \times X \times \dots \times X \rightarrow X$ is continuous for a.e $t \in I$, and there exist a positive function $\mu(\cdot) \in L^p(I, R^+)$ ($p > 1/q > 1$) and a continuous nondecreasing function $\omega : [0, \infty) \rightarrow [0, \infty)$ such that

$$\|f(t, v_1, v_2, \dots, v_n)\| \leq \mu(t)\omega\left(\sum_{i=1}^n \|v_i\|\right), \quad (t, v_1, v_2, \dots, v_n) \in I \times X \times X \times \dots \times X, \tag{3.1}$$

and set $T_{p,q} = \max\{T^{q-1/p}, T^q\}$.

(B2) For any bounded sets $D, D_1, D_2, \dots, D_n \subset X$, and $0 \leq \tau \leq s \leq t \leq T$,

$$\begin{aligned} \alpha(g(D)) &\leq \beta(t)\alpha(D), \\ \alpha(\psi(t-s, s)f(s, D_1, D_2, \dots, D_n)) \\ &\leq \beta_1(t, s)\alpha(D_1) + \beta_2(t, s)\alpha(D_2) + \dots + \beta_n(t, s)\alpha(D_n), \\ \alpha(\varphi(t-s, s)\varphi(s, \tau)f(\tau, D_1, D_2, \dots, D_n)) \\ &\leq \zeta_1(t, s, \tau)\alpha(D_1) + \zeta_2(t, s, \tau)\alpha(D_2) + \dots + \zeta_n(t, s, \tau)\alpha(D_n), \end{aligned} \quad (3.2)$$

where $\beta(t)$ is a nonnegative function, and $\sup_{t \in I} \beta(t) := \beta < \infty$,

$$\begin{aligned} \sup_{t \in I} \int_0^t \beta_i(t, s) ds &:= \beta_i < \infty, \quad i = 1, 2, \dots, n, \\ \sup_{t \in I} \int_0^t \int_0^s \zeta_j(t, s, \tau) d\tau ds &:= \zeta_j < \infty, \quad j = 1, 2, \dots, n. \end{aligned} \quad (3.3)$$

(B3) $g : C(I; X) \rightarrow X$ is continuous and there exists

$$0 < \alpha_1 < (C + \widetilde{M}(T))^{-1}, \quad \alpha_2 \geq 0 \quad (3.4)$$

such that

$$\|g(u)\| \leq \alpha_1 \|u\| + \alpha_2. \quad (3.5)$$

(B4) The functions μ and ω satisfy the following condition:

$$C(1 + CB(q, \gamma)) T_{p,q}^\gamma \Omega_{p,q} \left(\sum_{i=1}^n K_i^* \right) \|\mu\|_{L^p} \liminf_{\tau \rightarrow \infty} \frac{\omega(\tau)}{\tau} < 1 - \alpha_1 (C + \widetilde{M}(T)), \quad (3.6)$$

where $\Omega_{p,q} = ((p-1)/(pq-1))^{(p-1)/p}$, and $T_{p,q}^\gamma = \max\{T_{p,q}, T_{p,q+\gamma}\}$.

Theorem 3.1. Suppose that (B1)–(B4) are satisfied, and if $(C + \widetilde{M}(T))\beta + 4(\sum_{i=1}^n (\beta_i + 2\zeta_i)K_i^*) < 1$, then (1.1) has a mild solution on $[0, T]$.

Proof. Define the operator $F : C(I; X) \rightarrow C(I; X)$ by

$$\begin{aligned}
 F(u)(t) &= A^{-1}(0)g(u) + u_0 + \int_0^t \varphi(t - \eta, \eta)U(\eta)A(0) \left[A^{-1}(0)g(u) + u_0 \right] d\eta \\
 &\quad + \int_0^t \varphi(t - \eta, \eta) f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta)) d\eta \\
 &\quad + \int_0^t \int_0^\eta \varphi(t - \eta, \eta) \varphi(\eta, s) f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s)) ds d\eta, \quad t \in I.
 \end{aligned}
 \tag{3.7}$$

Then we proceed in five steps.

Step 1. We show that F is continuous.

Let u_i be a sequence that $u_i \rightarrow u$ as $i \rightarrow \infty$. Since f satisfies (B1), we have

$$f(t, (K_1u_i)(t), (K_2u_i)(t), \dots, (K_nu_i)(t)) \rightarrow f(t, (K_1u)(t), (K_2u)(t), \dots, (K_nu)(t)), \quad \text{as } i \rightarrow \infty.
 \tag{3.8}$$

Then

$$\begin{aligned}
 &\|F(u_i)(t) - F(u)(t)\| \\
 &\leq \|A^{-1}(0)\| \|g(u_i) - g(u)\| + \int_0^t \|\varphi(t - \eta, \eta)U(\eta)\| \|g(u_i) - g(u)\| d\eta \\
 &\quad + \int_0^t \|\varphi(t - \eta, \eta) [f(\eta, (K_1u_i)(\eta), (K_2u_i)(\eta), \dots, (K_nu_i)(\eta)) \\
 &\quad \quad \quad - f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta))]\| d\eta \\
 &\quad + \int_0^t \int_0^\eta \|\varphi(t - \eta, \eta) \varphi(\eta, s) [f(s, (K_1u_i)(s), (K_2u_i)(s), \dots, (K_nu_i)(s)) \\
 &\quad \quad \quad - f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s))]\| ds d\eta.
 \end{aligned}
 \tag{3.9}$$

According to the condition (A2), (2.12), and the continuity of g , we have

$$\begin{aligned}
 &\|A^{-1}(0)\| \|g(u_i) - g(u)\| \rightarrow 0, \quad \text{as } i \rightarrow \infty; \\
 &\int_0^t \|\varphi(t - \eta, \eta)U(\eta)\| \|g(u_i) - g(u)\| d\eta \rightarrow 0, \quad \text{as } i \rightarrow \infty.
 \end{aligned}
 \tag{3.10}$$

Noting that $u_i \rightarrow u$ in $C(I, X)$, there exists $\varepsilon > 0$ such that $\|u_i - u\| \leq \varepsilon$ for i sufficiently large. Therefore, we have

$$\begin{aligned} & \| [f(t, (K_1 u_i)(t), (K_2 u_i)(t), \dots, (K_n u_i)(t)) - f(t, (K_1 u)(t), (K_2 u)(t), \dots, (K_n u)(t))] \| \\ & \leq \mu(t) \left[\omega \left(\sum_{j=1}^n \| (K_j u_i)(t) \| \right) + \omega \sum_{j=1}^n \| (K_j u)(t) \| \right] \\ & \leq \mu(t) \left[\omega \left(\sum_{j=1}^n K_j^* (\|u\| + \varepsilon) \right) + \omega \left(\sum_{j=1}^n K_j^* \|u\| \right) \right]. \end{aligned} \quad (3.11)$$

Using (2.10) and by means of the Lebesgue dominated convergence theorem, we obtain

$$\begin{aligned} & \int_0^t \|\varphi(t - \eta, \eta) [f(\eta, (K_1 u_i)(\eta), (K_2 u_i)(\eta), \dots, (K_n u_i)(\eta)) \\ & \quad - f(\eta, (K_1 u)(\eta), (K_2 u)(\eta), \dots, (K_n u)(\eta))] \| d\eta \\ & \leq C \int_0^t (t - \eta)^{q-1} \| [f(\eta, (K_1 u_i)(\eta), (K_2 u_i)(\eta), \dots, (K_n u_i)(\eta)) \\ & \quad - f(\eta, (K_1 u)(\eta), (K_2 u)(\eta), \dots, (K_n u)(\eta))] \| d\eta, \\ & \rightarrow 0, \quad \text{as } i \rightarrow \infty. \end{aligned} \quad (3.12)$$

Similarly, by (2.10) and (2.11), we have

$$\begin{aligned} & \int_0^t \int_0^\eta \|\varphi(t - \eta, \eta) \varphi(\eta, s) \\ & \quad \times [f(s, (K_1 u_i)(t), (K_2 u_i)(t), \dots, (K_n u_i)(t)) \\ & \quad - f(s, (K_1 u)(s), (K_2 u)(s), \dots, (K_n u)(s))] \| ds d\eta \\ & \leq C^2 \int_0^t \int_0^\eta (t - \eta)^{q-1} (\eta - s)^{r-1} \\ & \quad \times \| [f(s, (K_1 u_i)(t), (K_2 u_i)(t), \dots, (K_n u_i)(t)) \\ & \quad - f(s, (K_1 u)(s), (K_2 u)(s), \dots, (K_n u)(s))] \| ds d\eta \\ & \rightarrow 0, \quad \text{as } i \rightarrow \infty. \end{aligned} \quad (3.13)$$

Therefore, we deduce that

$$\lim_{i \rightarrow \infty} \|F(u_i) - F(u)\| = 0. \quad (3.14)$$

Step 2. We show that F maps bounded sets of $C(I, X)$ into bounded sets in $C(I, X)$.

For any $r > 0$, we set $B_r = \{u \in C(I, X) : \|u\| \leq r\}$. Now, for $u \in B_r$, by (B1), we can see

$$\|f(t, (K_1u)(t), (K_2u)(t), \dots, (K_nu)(t))\| \leq \mu(t)\omega\left(\sum_{j=1}^n K_j^*r\right). \tag{3.15}$$

Based on (2.12), we denote that $S(t) := \int_0^t \varphi(t - \eta, \eta)U(\eta)d\eta$, we have

$$\|S(t)A(0)u_0\| \leq C^2t^q\left(\frac{1}{q} + t^\gamma B(q, \gamma + 1)\right)\|A(0)u_0\| = \widetilde{M}(t)\|A(0)u_0\|. \tag{3.16}$$

Then for any $u \in B_r$, by (A2), (2.10), (2.11), and Lemma 2.1, we have

$$\begin{aligned} \|(Fu)(t)\| &\leq \|A^{-1}(0)g(u)\| + \|u_0\| + \|S(t)g(u)\| + \|S(t)A(0)u_0\| \\ &\quad + \int_0^t \|\varphi(t - \eta, \eta)f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta))\|d\eta \\ &\quad + \int_0^t \int_0^\eta \|\varphi(t - \eta, \eta)\varphi(\eta, s)f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s))\|ds d\eta \\ &\leq (C + \widetilde{M}(t))\|g(u)\| + \|u_0\| + \widetilde{M}(t)\|A(0)u_0\| \\ &\quad + C \int_0^t (t - \eta)^{q-1}\mu(\eta)\omega\left(\sum_{j=1}^n K_j^*r\right)d\eta \\ &\quad + C^2 \int_0^t \int_0^\eta (t - \eta)^{q-1}(\eta - s)^{\gamma-1}\mu(s)\omega\left(\sum_{j=1}^n K_j^*r\right)ds d\eta \\ &\leq \alpha_1(C + \widetilde{M}(t))\|u\| + \alpha_2(C + \widetilde{M}(t)) + \|u_0\| + \widetilde{M}(t)\|A(0)u_0\| \\ &\quad + M_1\left[C \int_0^t (t - \eta)^{q-1}\mu(\eta)d\eta + C^2B(q, \gamma) \int_0^t (t - \eta)^{q+\gamma-1}\mu(\eta)d\eta\right], \end{aligned} \tag{3.17}$$

where $M_1 = \omega(\sum_{j=1}^n K_j^*r)$.

By means of the Hölder inequality, we have

$$\begin{aligned} \int_0^t (t - \eta)^{q-1}\mu(\eta)d\eta &= t^{(pq-1)/p}M_{p,q}\|\mu\|_{L^p} \leq T_{p,q}\Omega_{p,q}\|\mu\|_{L^p}, \\ \int_0^t (t - \eta)^{\gamma+q-1}\mu(\eta)d\eta &\leq T_{p,q+\gamma}\Omega_{p,q+\gamma}\|\mu\|_{L^p}. \end{aligned} \tag{3.18}$$

Thus

$$\begin{aligned} \|(Fu)(t)\| &\leq \alpha_1(C + \widetilde{M}(T))r + \alpha_2(C + \widetilde{M}(T)) + \|u_0\| + \widetilde{M}(T)\|A(0)u_0\| \\ &\quad + M_1\Omega_{p,q}T_{p,q}^\gamma [C + C^2B(q,\gamma)] \|\mu\|_{L^p} := \tilde{r}. \end{aligned} \quad (3.19)$$

This means $F(B_r) \subset B_{\tilde{r}}$.

Step 3. We show that there exists $m \in N$ such that $F(B_m) \subset B_m$.

Suppose the contrary, that for every $m \in N$, there exists $u_m \in B_m$ and $t_m \in I$, such that $\|(Fu_m)(t_m)\| > m$. However, on the other hand

$$\|f(t, (K_1u_m)(t), (K_2u_m)(t), \dots, (K_nu_m)(t))\| \leq \mu(t)\omega\left(\sum_{j=1}^n K_j^*m\right), \quad (3.20)$$

we have

$$\begin{aligned} m &< \|(Fu_m)(t_m)\| \leq \alpha_1(C + \widetilde{M}(T))\|u_m\| + \alpha_2(C + \widetilde{M}(T)) + \|u_0\| \\ &\quad + \widetilde{M}(T)\|A(0)u_0\| + M_1\left[C \int_0^{t_m} (t_m - \eta)^{q-1}\mu(\eta)d\eta + C^2B(q,\gamma) \int_0^{t_m} (t_m - \eta)^{q+\gamma-1}\mu(\eta)d\eta\right] \\ &\leq \alpha_1(C + \widetilde{M}(T))\|u_m\| + \alpha_2(C + \widetilde{M}(T)) + \|u_0\| \\ &\quad + \widetilde{M}(T)\|A(0)u_0\| + M_1\Omega_{p,q}T_{p,q}^\gamma [C + C^2B(q,\gamma)] \|\mu\|_{L^p} \\ &\leq \alpha_1(C + \widetilde{M}(T))m + \alpha_2(C + \widetilde{M}(T)) + \|u_0\| \\ &\quad + \widetilde{M}(T)\|A(0)u_0\| + M_1\Omega_{p,q}T_{p,q}^\gamma [C + C^2B(q,\gamma)] \|\mu\|_{L^p}. \end{aligned} \quad (3.21)$$

Dividing both sides by m and taking the lower limit as $m \rightarrow \infty$, we obtain

$$C(1 + CB(q,\gamma))T_{p,q}^\gamma \Omega_{p,q} \sum_{j=1}^n K_j^* \|\mu\|_{L^p} \liminf_{m \rightarrow \infty} \frac{\omega(m)}{m} \geq 1 - \alpha_1(C + \widetilde{M}(T)) \quad (3.22)$$

which contradicts (B4).

Step 4. Denote

$$F(u)(t) = A^{-1}(0)g(u) + u_0 + \int_0^t \varphi(t - \eta, \eta)U(\eta)A(0)[A^{-1}(0)g(u) + u_0]d\eta + G(u)(t), \quad (3.23)$$

where

$$\begin{aligned}
 G(u)(t) &= \int_0^t \varphi(t-\eta, \eta) f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta)) d\eta \\
 &\quad + \int_0^t \int_0^\eta \varphi(t-\eta, \eta) \varphi(\eta, s) f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s)) ds d\eta.
 \end{aligned}
 \tag{3.24}$$

We show that $G(u)(\cdot)$ is equicontinuous.

Let $0 < t_2 < t_1 < T$ and $u \in B_m$. Then

$$\|(Gu)(t_1) - (Gu)(t_2)\| \leq I_1 + I_2 + I_3 + I_4,
 \tag{3.25}$$

where

$$\begin{aligned}
 I_1 &= \int_0^{t_2} \|\varphi(t_1 - \eta, \eta) - \varphi(t_2 - \eta, \eta)\| f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta)) \|d\eta, \\
 I_2 &= \int_{t_2}^{t_1} \|\varphi(t_1 - \eta, \eta) f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta))\| d\eta, \\
 I_3 &= \int_0^{t_2} \int_0^\eta \|\varphi(t_1 - \eta, \eta) - \varphi(t_2 - \eta, \eta)\| \varphi(\eta, s) f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s)) \|ds d\eta, \\
 I_4 &= \int_{t_2}^{t_1} \int_0^\eta \|\varphi(t_1 - \eta, \eta) \varphi(\eta, s) f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s))\| ds d\eta.
 \end{aligned}
 \tag{3.26}$$

It follows from Lemma 2.9, (B1), and (3.20) that $I_1, I_3 \rightarrow 0$, as $t_2 \rightarrow t_1$.

For I_2 , from (2.10), (3.20), and (B1), we have

$$\begin{aligned}
 I_2 &= \int_{t_2}^{t_1} \|\varphi(t_1 - \eta, \eta) f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta))\| d\eta \\
 &\leq CM_1 \int_{t_2}^{t_1} (t_1 - \eta)^{q-1} \mu(\eta) d\eta \rightarrow 0, \quad \text{as } t_2 \rightarrow t_1.
 \end{aligned}
 \tag{3.27}$$

Similarly, by (2.10), (2.11), (B1), and Lemma 2.1, we have

$$\begin{aligned}
 I_4 &= \int_{t_2}^{t_1} \int_0^\eta \|\varphi(t_1 - \eta, \eta) \varphi(\eta, s) f(s, (K_1u)(s), (K_2u)(s), \dots, (K_nu)(s))\| ds d\eta \\
 &\leq C^2 M_1 \int_{t_2}^{t_1} (t_1 - \eta)^{q-1} \int_0^\eta (\eta - s)^{r-1} \mu(s) ds d\eta \rightarrow 0, \quad \text{as } t_2 \rightarrow t_1.
 \end{aligned}
 \tag{3.28}$$

Step 5. We show that $\alpha(F(H)) < \alpha(H)$ for every bounded set $H \subset B_m$. For any $\varepsilon > 0$, we can take a sequence $\{h_v\}_{v=1}^\infty \subset H$ such that

$$\alpha(H) \leq 2\alpha(\{h_v\}) + \varepsilon, \quad (3.29)$$

(cf. [30]). So it follows from Lemmas 2.3–2.5, 2.9, (2) in Remark 2.10, and (B2) that

$$\begin{aligned} \alpha(F(H)) &\leq C\alpha(g(H)) + \widetilde{M(T)}\alpha(g(H)) + 2\alpha(G\{h_v\}) + \varepsilon \\ &\leq C\alpha(g(H)) + \widetilde{M(T)}\alpha(g(H)) \\ &\quad + 2 \sup_{t \in I} \alpha \left(\left\{ \int_0^t \varphi(t-\eta, \eta) f(\eta, (K_1 h_v)(\eta), (K_2 h_v)(\eta), \dots, (K_n h_v)(\eta)) d\eta \right\} \right. \\ &\quad \left. + \left\{ \int_0^t \int_0^\eta \varphi(t-\eta, \eta) \varphi(\eta, s) \right. \right. \\ &\quad \left. \left. \times f(s, (K_1 h_v)(s), (K_2 h_v)(s), \dots, (K_n h_v)(s)) ds d\eta \right\} \right) + \varepsilon \\ &\leq C\beta\alpha(H) + \widetilde{M(T)}\beta\alpha(H) \\ &\quad + 4 \sup_{t \in I} \left(\int_0^t \alpha(\{\varphi(t-\eta, \eta) f(\eta, (K_1 h_v)(\eta), (K_2 h_v)(\eta), \dots, (K_n h_v)(\eta))\}) d\eta \right) \\ &\quad + 8 \sup_{t \in I} \left(\int_0^t \int_0^\eta \alpha(\{\varphi(t-\eta, \eta) \varphi(\eta, s) f(s, (K_1 h_v)(s), (K_2 h_v)(s), \dots, (K_n h_v)(s))\}) \right) \\ &\quad + \varepsilon \leq C\beta\alpha(H) + \widetilde{M(T)}\beta\alpha(H) + 4 \sup_{t \in I} \left(\int_0^t \left(\sum_{i=1}^n \beta_i(t, \eta) K_i^* \right) \alpha(\{h_v\}) d\eta \right) \\ &\quad + 8 \sup_{t \in I} \left(\int_0^t \int_0^\eta \left(\sum_{i=1}^n \zeta_i(t, \eta, s) K_i^* \right) \alpha(\{h_v\}) ds d\eta \right) + \varepsilon \\ &\leq C\beta\alpha(H) + \widetilde{M(T)}\beta\alpha(H) + \left(4 \sum_{i=1}^n \beta_i K_i^* + 8 \sum_{i=1}^n \zeta_i K_i^* \right) \alpha(\{h_v\}) + \varepsilon \\ &= \left[(C + \widetilde{M(T)})\beta + 4 \left(\sum_{i=1}^n (\beta_i + 2\zeta_i) K_i^* \right) \right] \alpha(H) + \varepsilon. \end{aligned} \quad (3.30)$$

Since ε is arbitrary, we can obtain

$$\alpha(F(H)) \leq \left[(C + \widetilde{M(T)})\beta + 4 \left(\sum_{i=1}^n (\beta_i + 2\zeta_i) K_i^* \right) \right] \alpha(H) < \alpha(H). \quad (3.31)$$

In summary, we have proven that F has a fixed point $\tilde{u} \in B_m$. Consequently, (1.1) has at least one mild solution. \square

Our next result is based on the Banach’s fixed point theorem.

(G1) There exists a positive function $l(\cdot) \in L^1(I, R^+)$ and a constant $\mu > 0$ such that

$$\begin{aligned} & \|g(u) - g(u^*)\| \leq \mu \|u - u^*\|, \\ & \|f(t, v_1, v_2, \dots, v_n) - f(t, w_1, w_2, \dots, w_n)\| \\ & \leq l(t) \left(\sum_{i=1}^n \|v_i - w_i\| \right), \quad (v_i, w_i) \in X^2, i = 1, 2, \dots, n. \end{aligned} \tag{3.32}$$

(G2) There exists a constant $0 < \delta < 1$ such that the function $\Lambda : I \rightarrow R^+$ defined by

$$\Lambda(t) = \mu(C + \widetilde{M}(T)) + C \left(\sum_{i=1}^n K_i^* \right) \Gamma(q) I^{q_l}(t) + C^2 \left(\sum_{i=1}^n K_i^* \right) \Gamma(q) \Gamma(\gamma) I^{q+\gamma_l}(t) \leq \delta, \quad t \in I. \tag{3.33}$$

Theorem 3.2. Assume that (G1), (G2) are satisfied, then (1.1) has a unique mild solution.

Proof. Let F be defined as in Theorem 3.1. For any $u, u^* \in C(I, X)$, we have

$$\begin{aligned} & \|f(t, (K_1u)(t), (K_2u)(t), \dots, (K_nu)(t)) - f(t, (K_1u^*)(t), (K_2u^*)(t), \dots, (K_nu^*)(t))\| \\ & \leq l(t) \left(\sum_{i=1}^n \|(K_iu)(t) - (K_iu^*)(t)\| \right) \\ & \leq l(t) \sum_{i=1}^n K_i^* \|u - u^*\|. \end{aligned} \tag{3.34}$$

Thus, from (A2), (2.10), (2.11), Lemma 2.1, we have

$$\begin{aligned} & \|(Fu)(t) - (Fu^*)(t)\| \\ & \leq \mu C \|u - u^*\| + \mu \int_0^t \|\varphi(t - \eta, \eta)U(\eta)\| \|u - u^*\| d\eta \\ & + \int_0^t \|\varphi(t - \eta, \eta)\| \|f(\eta, (K_1u)(\eta), (K_2u)(\eta), \dots, (K_nu)(\eta)) \\ & \quad - f(\eta, (K_1u^*)(\eta), (K_2u^*)(\eta), \dots, (K_nu^*)(\eta))\| d\eta \\ & + \int_0^t \int_0^\eta \|\varphi(t - \eta, \eta)\varphi(\eta, s)\| \|f(s, (Ku)(s), (Hu)(s)) - f(s, (Ku^*)(s), (Hu^*)(s))\| ds d\eta \end{aligned}$$

$$\begin{aligned}
&\leq \|u - u^*\| \left[\mu(C + \widetilde{M}(T)) + C \left(\sum_{i=1}^n K_i^* \right) \int_0^t (t - \eta)^{q-1} l(\eta) d\eta \right. \\
&\quad \left. + C^2 \left(\sum_{i=1}^n K_i^* \right) \int_0^t \int_0^\eta (t - \eta)^{q-1} (\eta - s)^{\gamma-1} l(s) ds d\eta \right] \\
&= \left[\mu(C + \widetilde{M}(T)) + C \left(\sum_{i=1}^n K_i^* \right) \Gamma(q) I^q l(t) + C^2 \left(\sum_{i=1}^n K_i^* \right) \Gamma(q) \Gamma(\gamma) I^{q+\gamma} l(t) \right] \|u - u^*\| \\
&= \Lambda(t) \|u - u^*\|.
\end{aligned} \tag{3.35}$$

We get

$$\|F(u) - F(u^*)\| \leq \delta \|u - u^*\|. \tag{3.36}$$

By the Banach contraction mapping principle, F has a unique fixed point, which is a mild solution of (1.1). \square

4. An Example

To illustrate the usefulness of our main result, we consider the following fractional differential equation:

$$\begin{aligned}
\frac{\partial^q}{\partial t^q} u(t, \xi) &= b(t, \xi) \frac{\partial^2}{\partial \xi^2} u(t, \xi) + \frac{t^n}{n} \int_0^t (t - s) u(s, \xi) ds + \frac{t^n}{n} \int_0^t e^{-(t+s)} u(s, \xi) ds, \quad \xi \in [0, 1], \\
u(t, 0) &= u(t, 1) = 0,
\end{aligned} \tag{4.1}$$

$$u(0, \xi) = - \int_0^\xi \int_0^y b^{-1}(0, x) \sin \left| \frac{u}{\lambda} \right| dx dy,$$

where $0 < q < 1$, $0 \leq t \leq 1$, $\lambda > C + \widetilde{M}(1)$, $n \in N$, $b(t, \xi)$ is continuous function and is uniformly Hölder continuous in t , that is, there exist $C > 0$ and $\gamma \in (0, 1)$ such that

$$\|b(t_1, \xi) - b(t_2, \xi)\| \leq C |t_1 - t_2|^\gamma, \quad 0 \leq t_1 \leq t_2 \leq 1. \tag{4.2}$$

Let $X = L^2([0, 1], R)$ and define $A(t)$ by

$$\begin{aligned}
D(A(t)) &= H^2(0, 1) \cap H_0^1(0, 1) = \{H^2(0, 1) : z(0) = z(1) = 0\}, \\
-A(t)(z) &= b(t, \xi) z''.
\end{aligned} \tag{4.3}$$

Then $-A(s)$ generates an analytic semigroup $\exp(-tA(s))$.

For $t \in [0, 1], \xi \in [0, 1]$, we set

$$\begin{aligned}
 u(t)(\xi) &= u(t, \xi), \\
 g(u) &= \sin\left|\frac{u}{\lambda}\right|, \\
 A^{-1}(0)g(u) &= -\int_0^\xi \int_0^y b^{-1}(0, x) \sin\left|\frac{u}{\lambda}\right| dx dy,
 \end{aligned} \tag{4.4}$$

$$f(t, (K_1u)(t), (K_2u)(t))(\xi) = \frac{t^n}{n} \int_0^t (t-s)u(s, \xi) ds + \frac{t^n}{n} \int_0^t e^{-(t+s)} u(s, \xi) ds,$$

where

$$\begin{aligned}
 (K_1u(t))(\xi) &= \int_0^t (t-s)u(s, \xi) ds, \\
 (K_2u(t))(\xi) &= \int_0^t e^{-(t+s)} u(s, \xi) ds, \\
 K_1^* &= \sup_{t \in I} \int_0^t (t-s) ds < \frac{1}{2} < \infty, \\
 K_2^* &= \sup_{t \in I} \int_0^t e^{-(t+s)} ds = \frac{1}{4} < \infty.
 \end{aligned} \tag{4.5}$$

Moreover, we can get

$$\begin{aligned}
 \|g(u)\| &\leq \frac{1}{\lambda} \|u\|, \\
 \alpha(g(D)) &\leq \frac{1}{\lambda} \alpha(D)
 \end{aligned} \tag{4.6}$$

for any $D \subset X$. Then the above equation (4.1) can be written in the abstract form as (1.1). On the other hand,

$$\begin{aligned}
 \|f(t, (Ku)(t), (Hu)(t))(\xi)\| &\leq \frac{t^n}{n} \|(K_1u)(t, \xi)\| + \|(K_2u)(t, \xi)\| \\
 &\leq \frac{t^n}{n} (K_1^* \|u\| + K_2^* \|u\|) \\
 &= \mu(t) \omega(K_1^* \|u\| + K_2^* \|u\|),
 \end{aligned} \tag{4.7}$$

where $\mu(t) = t^n$, $\omega(z) = z/n$ satisfying (B1). For any $u_1, u_2 \in X$,

$$\begin{aligned} & \|\varphi(t-s, s)f(s, (K_1u_1)(s), (K_2u_1)(s))(\xi) - \varphi(t-s, s)f(s, (K_1u_2)(s), (K_2u_2)(s))(\xi)\| \\ & \leq \frac{Cs^n}{n}(t-s)^{q-1}(\|(K_1u_1)(s)(\xi) - (K_1u_2)(s)(\xi)\| + \|(K_2u_1)(s)(\xi) - (K_2u_2)(s)(\xi)\|). \end{aligned} \quad (4.8)$$

Therefore, for any bounded sets $D_1, D_2 \subset X$, we have

$$\alpha(\varphi(t-s, s)f(s, D_1, D_2)) \leq \frac{Cs^n}{n}(t-s)^{q-1}(\alpha(D_1) + \alpha(D_2)). \quad (4.9)$$

Moreover,

$$\frac{C}{n} \sup_{t \in [0,1]} \int_0^t (t-s)^{q-1} s^n ds = \frac{C}{n} \sup_{t \in [0,1]} t^{n+q} B(q, n+1) = \frac{C}{n} B(q, n+1) := \beta_1 = \beta_2. \quad (4.10)$$

Similarly, we obtain

$$\begin{aligned} \alpha(\varphi(t-s, s)\varphi(s, \tau)f(\tau, D_1, D_2)) & \leq \frac{C^2}{n}(t-s)^{q-1}(s-\tau)^{\gamma-1}\tau^n(\alpha(D_1) + \alpha(D_2)), \\ \frac{C^2}{n} \sup_{t \in [0,1]} \int_0^t \int_0^s (t-s)^{q-1}(s-\tau)^{\gamma-1}\tau^n d\tau ds & \leq \frac{C^2}{n} B(q, \gamma) B(q+\gamma, n+1) := \zeta_1 = \zeta_2. \end{aligned} \quad (4.11)$$

Suppose further that

- (1) $(3/4n)C(1 + CB(q, \gamma))((p-1)/(pq-1))^{(p-1)/p} \|\mu\|_{L^p} < 1 - (C + \widetilde{M}(1))/\lambda$,
- (2) $(1/\lambda)(C + \widetilde{M}(1)) + 3(\beta_1 + 2\zeta_1) < 1$.

Then (4.1) has a mild solution by Theorem 3.1.

References

- [1] R. P. Agarwal, M. Belmekki, and M. Benchohra, "A survey on semilinear differential equations and inclusions involving Riemann-Liouville fractional derivative," *Advances in Difference Equations*, vol. 2009, Article ID 981728, 47 pages, 2009.
- [2] A. Anguraj, P. Karthikeyan, and G. M. N'Guérékata, "Nonlocal Cauchy problem for some fractional abstract integro-differential equations in Banach spaces," *Communications in Mathematical Analysis*, vol. 6, no. 1, pp. 31–35, 2009.
- [3] M. M. El-Borai, "Some probability densities and fundamental solutions of fractional evolution equations," *Chaos, Solitons and Fractals*, vol. 14, no. 3, pp. 433–440, 2002.
- [4] M. M. El-Borai, "The fundamental solutions for fractional evolution equations of parabolic type," *Journal of Applied Mathematics and Stochastic Analysis*, no. 3, pp. 197–211, 2004.
- [5] F. Li, "Mild solutions for fractional differential equations with nonlocal conditions," *Advances in Difference Equations*, vol. 2010, Article ID 287861, 9 pages, 2010.
- [6] F. Li, "Solvability of nonautonomous fractional integrodifferential equations with infinite delay," *Advances in Difference Equations*, vol. 2011, Article ID 806729, 18 pages, 2011.

- [7] J. Liang and T.-J. Xiao, "Solutions to nonautonomous abstract functional equations with infinite delay," *Taiwanese Journal of Mathematics*, vol. 10, no. 1, pp. 163–172, 2006.
- [8] G. M. Mophou, "Existence and uniqueness of mild solutions to impulsive fractional differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 3-4, pp. 1604–1615, 2010.
- [9] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach Science, New York, NY, USA, 1993.
- [10] T. Diagana, "Pseudo-almost automorphic solutions to some classes of nonautonomous partial evolution equations," *Differential Equations & Applications*, vol. 1, no. 4, pp. 561–582, 2009.
- [11] E. Hille and R. S. Phillips, *Functional Analysis and Semi-Groups*, vol. 31 of *American Mathematical Society Colloquium Publications*, American Mathematical Society, Providence, RI, USA, 1957.
- [12] K. Josić and R. Rosenbaum, "Unstable solutions of nonautonomous linear differential equations," *SIAM Review*, vol. 50, no. 3, pp. 570–584, 2008.
- [13] M. Kunze, L. Lorenzi, and A. Lunardi, "Nonautonomous Kolmogorov parabolic equations with unbounded coefficients," *Transactions of the American Mathematical Society*, vol. 362, no. 1, pp. 169–198, 2010.
- [14] J. Liang, R. Nagel, and T.-J. Xiao, "Approximation theorems for the propagators of higher order abstract Cauchy problems," *Transactions of the American Mathematical Society*, vol. 360, no. 4, pp. 1723–1739, 2008.
- [15] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, vol. 44 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1983.
- [16] T.-J. Xiao and J. Liang, *The Cauchy Problem for Higher-Order Abstract Differential Equations*, vol. 1701 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1998.
- [17] T.-J. Xiao and J. Liang, "Approximations of Laplace transforms and integrated semigroups," *Journal of Functional Analysis*, vol. 172, no. 1, pp. 202–220, 2000.
- [18] T.-J. Xiao and J. Liang, "Existence of classical solutions to nonautonomous nonlocal parabolic problems," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 63, no. 5–7, pp. e225–e232, 2005.
- [19] T.-J. Xiao and J. Liang, "Second order differential operators with Feller-Wentzell type boundary conditions," *Journal of Functional Analysis*, vol. 254, no. 6, pp. 1467–1486, 2008.
- [20] T.-J. Xiao, J. Liang, and J. van Casteren, "Time dependent Desch-Schappacher type perturbations of Volterra integral equations," *Integral Equations and Operator Theory*, vol. 44, no. 4, pp. 494–506, 2002.
- [21] L. Byszewski and V. Lakshmikantham, "Theorem about the existence and uniqueness of a solution of a nonlocal abstract Cauchy problem in a Banach space," *Applicable Analysis*, vol. 40, no. 1, pp. 11–19, 1991.
- [22] Z. Fan, "Impulsive problems for semilinear differential equations with nonlocal conditions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 2, pp. 1104–1109, 2010.
- [23] J. Liang and T.-J. Xiao, "Semilinear integrodifferential equations with nonlocal initial conditions," *Computers & Mathematics with Applications*, vol. 47, no. 6-7, pp. 863–875, 2004.
- [24] J. Liang, J. H. Liu, and T.-J. Xiao, "Nonlocal problems for integrodifferential equations," *Dynamics of Continuous, Discrete & Impulsive Systems. Series A*, vol. 15, no. 6, pp. 815–824, 2008.
- [25] J. Liang, J. van Casteren, and T.-J. Xiao, "Nonlocal Cauchy problems for semilinear evolution equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 50, no. 2, pp. 173–189, 2002.
- [26] H. Liu and J.-C. Chang, "Existence for a class of partial differential equations with nonlocal conditions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 9, pp. 3076–3083, 2009.
- [27] J. Banaś and K. Goebel, *Measures of Noncompactness in Banach Spaces*, vol. 60 of *Lecture Notes in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 1980.
- [28] H.-P. Heinz, "On the behaviour of measures of noncompactness with respect to differentiation and integration of vector-valued functions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 7, no. 12, pp. 1351–1371, 1983.
- [29] B. Sadovskii, "On a fixed point principle," *Functional Analysis and Its Applications*, vol. 2, pp. 151–153, 1967.
- [30] D. Bothe, "Multivalued perturbations of m -accretive differential inclusions," *Israel Journal of Mathematics*, vol. 108, pp. 109–138, 1998.