

Research Article

Monotone Positive Solution of Nonlinear Third-Order BVP with Integral Boundary Conditions

Jian-Ping Sun and Hai-Bao Li

Department of Applied Mathematics, Lanzhou University of Technology, Lanzhou, Gansu 730050, China

Correspondence should be addressed to Jian-Ping Sun, jpsun@lut.cn

Received 7 September 2010; Accepted 31 October 2010

Academic Editor: Michel C. Chipot

Copyright © 2010 J.-P. Sun and H.-B. Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the following third-order boundary value problem with integral boundary conditions $u'''(t) + f(t, u(t), u'(t)) = 0, t \in [0, 1]; u(0) = u'(0) = 0, u'(1) = \int_0^1 g(t)u'(t)dt$, where $f \in C([0, 1] \times [0, +\infty) \times [0, +\infty), [0, +\infty))$ and $g \in C([0, 1], [0, +\infty))$. By using the Guo-Krasnoselskii fixed-point theorem, some sufficient conditions are obtained for the existence and nonexistence of monotone positive solution to the above problem.

1. Introduction

Third-order differential equations arise in a variety of different areas of applied mathematics and physics, for example, in the deflection of a curved beam having a constant or varying cross section, a three-layer beam, electromagnetic waves or gravity driven flows and so on [1].

Recently, third-order two-point or multipoint boundary value problems (BVPs for short) have attracted a lot of attention [2–17]. It is known that BVPs with integral boundary conditions cover multipoint BVPs as special cases. Although there are many excellent works on third-order two-point or multipoint BVPs, a little work has been done for third-order BVPs with integral boundary conditions. It is worth mentioning that, in 2007, Anderson and Tisdell [18] developed an interval of λ values whereby a positive solution exists for the following third-order BVP with integral boundary conditions

$$\begin{aligned}(pu'')'(t) &= \lambda f(t, u(t)), \quad t \in [t_1, t_3], \\ au(t_1) - \beta u'(t_1) &= \int_{\xi_1}^{\xi_2} g(t)u(t)dt,\end{aligned}$$

$$\begin{aligned}
 u'(t_2) &= 0, \\
 (pu'')(t_3) &= \int_{\eta_1}^{\eta_2} h(t)(pu'')(t)dt
 \end{aligned}
 \tag{1.1}$$

by using the Guo-Krasnoselskii fixed-point theorem. In 2008, Graef and Yang [19] studied the third-order BVP with integral boundary conditions

$$\begin{aligned}
 u'''(t) &= g(t)f(u(t)), \quad t \in [0, 1], \\
 u(0) = u'(p) &= \int_q^1 w(t)u''(t)dt = 0.
 \end{aligned}
 \tag{1.2}$$

For second-order or fourth-order BVPs with integral boundary conditions, one can refer to [20–24].

In this paper, we are concerned with the following third-order BVP with integral boundary conditions

$$\begin{aligned}
 u'''(t) + f(t, u(t), u'(t)) &= 0, \quad t \in [0, 1], \\
 u(0) = u'(0) = 0, \quad u'(1) &= \int_0^1 g(t)u'(t)dt.
 \end{aligned}
 \tag{1.3}$$

Throughout this paper, we always assume that $f \in C([0, 1] \times [0, +\infty) \times [0, +\infty), [0, +\infty))$ and $g \in C([0, 1], [0, +\infty))$. Some sufficient conditions are established for the existence and nonexistence of monotone positive solution to the BVP (1.3). Here, a solution u of the BVP (1.3) is said to be monotone and positive if $u'(t) \geq 0$, $u(t) \geq 0$ and $u(t) \neq 0$ for $t \in [0, 1]$. Our main tool is the following Guo-Krasnoselskii fixed-point theorem [25].

Theorem 1.1. *Let E be a Banach space and let K be a cone in E . Assume that Ω_1 and Ω_2 are bounded open subsets of E such that $\theta \in \Omega_1$, $\overline{\Omega}_1 \subset \Omega_2$, and let $T : K \cap (\overline{\Omega}_2 \setminus \Omega_1) \rightarrow K$ be a completely continuous operator such that either*

- (1) $\|Tu\| \leq \|u\|$ for $u \in K \cap \partial\Omega_1$ and $\|Tu\| \geq \|u\|$ for $u \in K \cap \partial\Omega_2$, or
- (2) $\|Tu\| \geq \|u\|$ for $u \in K \cap \partial\Omega_1$ and $\|Tu\| \leq \|u\|$ for $u \in K \cap \partial\Omega_2$.

Then T has a fixed point in $K \cap (\overline{\Omega}_2 \setminus \Omega_1)$.

2. Preliminaries

For convenience, we denote $\mu = \int_0^1 tg(t)dt$.

Lemma 2.1. Let $\mu \neq 1$. Then for any $h \in C[0, 1]$, the BVP

$$\begin{aligned} -u'''(t) &= h(t), \quad t \in [0, 1], \\ u(0) = u'(0) &= 0, \quad u'(1) = \int_0^1 g(t)u'(t)dt \end{aligned} \quad (2.1)$$

has a unique solution

$$u(t) = \int_0^1 \left[G_1(t, s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau, s)g(\tau)d\tau \right] h(s)ds, \quad t \in [0, 1], \quad (2.2)$$

where

$$\begin{aligned} G_1(t, s) &= \frac{1}{2} \begin{cases} (2t - t^2 - s)s, & 0 \leq s \leq t \leq 1, \\ (1-s)t^2, & 0 \leq t \leq s \leq 1, \end{cases} \\ G_2(t, s) &= \begin{cases} (1-t)s, & 0 \leq s \leq t \leq 1, \\ (1-s)t, & 0 \leq t \leq s \leq 1. \end{cases} \end{aligned} \quad (2.3)$$

Proof. Let u be a solution of the BVP (2.1). Then, we may suppose that

$$u(t) = \int_0^1 G_1(t, s)h(s)ds + At^2 + Bt + C, \quad t \in [0, 1]. \quad (2.4)$$

By the boundary conditions in (2.1), we have

$$A = \frac{1}{2(1-\mu)} \int_0^1 h(s) \int_0^1 G_2(\tau, s)g(\tau)d\tau ds \quad \text{and} \quad B = C = 0. \quad (2.5)$$

Therefore, the BVP (2.1) has a unique solution

$$u(t) = \int_0^1 \left[G_1(t, s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau, s)g(\tau)d\tau \right] h(s)ds, \quad t \in [0, 1]. \quad (2.6)$$

□

Lemma 2.2 (see [12]). For any $(t, s) \in [0, 1] \times [0, 1]$,

$$\frac{t^2}{2}(1-s)s \leq G_1(t, s) \leq \frac{1}{2}(1-s)s. \quad (2.7)$$

Lemma 2.3 (see [26]). For any $(t, s) \in [0, 1] \times [0, 1]$,

$$0 \leq G_2(t, s) \leq (1 - s)s. \quad (2.8)$$

In the remainder of this paper, we always assume that $\mu < 1$, $\alpha \in (0, 1)$ and $\beta = \alpha^2/2$.

Lemma 2.4. If $h \in C[0, 1]$ and $h(t) \geq 0$ for $t \in [0, 1]$, then the unique solution u of the BVP (2.1) satisfies

$$(1) \ u(t) \geq 0, \ t \in [0, 1],$$

$$(2) \ u'(t) \geq 0, \ t \in [0, 1] \text{ and } \min_{t \in [\alpha, 1]} u(t) \geq \beta \|u\|, \text{ where } \|u\| = \max\{\|u\|_\infty, \|u'\|_\infty\}.$$

Proof. Since (1) is obvious, we only need to prove (2). By (2.2), we get

$$u'(t) = \int_0^1 \left[G_2(t, s) + \frac{t}{1 - \mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] h(s) ds, \quad t \in [0, 1], \quad (2.9)$$

which indicates that $u'(t) \geq 0$ for $t \in [0, 1]$.

On the one hand, by (2.9) and Lemma 2.3, we have

$$\|u'\|_\infty \leq \int_0^1 \left[(1 - s)s + \frac{1}{1 - \mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] h(s) ds. \quad (2.10)$$

On the other hand, in view of (2.2) and Lemma 2.2, we have

$$\|u\|_\infty \leq \int_0^1 \left[(1 - s)s + \frac{1}{1 - \mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] h(s) ds. \quad (2.11)$$

It follows from (2.10) and (2.11) that

$$\|u\| \leq \int_0^1 \left[(1 - s)s + \frac{1}{1 - \mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] h(s) ds, \quad (2.12)$$

which together with Lemma 2.2 implies that

$$\begin{aligned}
 \min_{t \in [\alpha, 1]} u(t) &= \min_{t \in [\alpha, 1]} \int_0^1 \left[G_1(t, s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] h(s) ds \\
 &\geq \min_{t \in [\alpha, 1]} \frac{t^2}{2} \int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] h(s) ds \\
 &= \frac{\alpha^2}{2} \int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] h(s) ds \\
 &\geq \beta \|u\|.
 \end{aligned} \tag{2.13}$$

□

Let $E = C^1[0, 1]$ be equipped with the norm $\|u\| = \max\{\|u\|_\infty, \|u'\|_\infty\}$. Then E is a Banach space. If we denote

$$K = \left\{ u \in E : u(t) \geq 0, u'(t) \geq 0, t \in [0, 1], \min_{t \in [\alpha, 1]} u(t) \geq \beta \|u\| \right\}, \tag{2.14}$$

then it is easy to see that K is a cone in E . Now, we define an operator T on K by

$$(Tu)(t) = \int_0^1 \left[G_1(t, s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u(s), u'(s)) ds, \quad t \in [0, 1]. \tag{2.15}$$

Obviously, if u is a fixed point of T , then u is a monotone nonnegative solution of the BVP (1.3).

Lemma 2.5. $T : K \rightarrow K$ is completely continuous.

Proof. First, by Lemma 2.4, we know that $T(K) \subset K$.

Next, we assume that $D \subset K$ is a bounded set. Then there exists a constant $M_1 > 0$ such that $\|u\| \leq M_1$ for any $u \in D$. Now, we will prove that $T(D)$ is relatively compact in K . Suppose that $\{y_k\}_{k=1}^\infty \subset T(D)$. Then there exist $\{x_k\}_{k=1}^\infty \subset D$ such that $Tx_k = y_k$. Let

$$\begin{aligned}
 M_2 &= \sup \{ f(t, x, y) : (t, x, y) \in [0, 1] \times [0, M_1] \times [0, M_1] \}, \\
 M_3 &= \frac{1}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau ds.
 \end{aligned} \tag{2.16}$$

Then for any k , by Lemma 2.2, we have

$$\begin{aligned}
 |y_k(t)| &= |(Tx_k)(t)| \\
 &= \left| \int_0^1 \left[G_1(t,s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau,s)g(\tau)d\tau \right] f(s, x_k(s), x'_k(s)) ds \right| \\
 &\leq \frac{M_2}{2} \int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau,s)g(\tau)d\tau \right] ds \\
 &= \frac{M_2}{2} \left(\frac{1}{6} + M_3 \right), \quad t \in [0,1],
 \end{aligned} \tag{2.17}$$

which implies that $\{y_k\}_{k=1}^\infty$ is uniformly bounded. At the same time, for any k , in view of Lemma 2.3, we have

$$\begin{aligned}
 |y'_k(t)| &= |(Tx_k)'(t)| \\
 &= \left| \int_0^1 \left[G_2(t,s) + \frac{t}{1-\mu} \int_0^1 G_2(\tau,s)g(\tau)d\tau \right] f(s, x_k(s), x'_k(s)) ds \right| \\
 &\leq M_2 \left(\int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau,s)g(\tau)d\tau \right] ds \right) \\
 &= M_2 \left(\frac{1}{6} + M_3 \right), \quad t \in [0,1],
 \end{aligned} \tag{2.18}$$

which shows that $\{y'_k\}_{k=1}^\infty$ is also uniformly bounded. This indicates that $\{y_k\}_{k=1}^\infty$ is equicontinuous. It follows from Arzela-Ascoli theorem that $\{y_k\}_{k=1}^\infty$ has a convergent subsequence in $C[0,1]$. Without loss of generality, we may assume that $\{y_k\}_{k=1}^\infty$ converges in $C[0,1]$. On the other hand, by the uniform continuity of $G_2(t,s)$, we know that for any $\varepsilon > 0$, there exists $\delta_1 > 0$ such that for any $t_1, t_2 \in [0,1]$ with $|t_1 - t_2| < \delta_1$, we have

$$|G_2(t_1, s) - G_2(t_2, s)| < \frac{\varepsilon}{2(M_2 + 1)}, \quad s \in [0,1]. \tag{2.19}$$

Let $\delta = \min\{\delta_1, \varepsilon/2(M_2M_3 + 1)\}$. Then for any k , $t_1, t_2 \in [0,1]$ with $|t_1 - t_2| < \delta$, we have

$$\begin{aligned}
 |y'_k(t_1) - y'_k(t_2)| &= |(Tx_k)'(t_1) - (Tx_k)'(t_2)| \\
 &\leq \int_0^1 \left[|G_2(t_1, s) - G_2(t_2, s)| + \frac{|t_1 - t_2|}{1-\mu} \int_0^1 G_2(\tau,s)g(\tau)d\tau \right] f(s, x_k(s), x'_k(s)) ds \\
 &\leq M_2 \int_0^1 |G_2(t_1, s) - G_2(t_2, s)| ds + M_2M_3|t_1 - t_2| \\
 &\leq \frac{M_2\varepsilon}{2(M_2 + 1)} + M_2M_3|t_1 - t_2| \\
 &< \varepsilon,
 \end{aligned} \tag{2.20}$$

which implies that $\{y'_k\}_{k=1}^\infty$ is equicontinuous. Again, by Arzela-Ascoli theorem, we know that $\{y'_k\}_{k=1}^\infty$ has a convergent subsequence in $C[0, 1]$. Therefore, $\{y_k\}_{k=1}^\infty$ has a convergent subsequence in $C^1[0, 1]$. Thus, we have shown that T is a compact operator.

Finally, we prove that T is continuous. Suppose that $u_m, u \in K$ and $\|u_m - u\| \rightarrow 0$ ($m \rightarrow \infty$). Then there exists $M_4 > 0$ such that for any m , $\|u_m\| \leq M_4$. Let

$$M_5 = \sup\{f(t, x, y) : (t, x, y) \in [0, 1] \times [0, M_4] \times [0, M_4]\}. \quad (2.21)$$

Then for any m and $t \in [0, 1]$, in view of Lemmas 2.2 and 2.3, we have

$$\begin{aligned} & \left[G_1(t, s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u_m(s), u'_m(s)) \\ & \leq \frac{M_5}{2} \left[1 + \frac{1}{1-\mu} \int_0^1 g(\tau) d\tau \right] (1-s)s, \quad s \in [0, 1], \\ & \left[G_2(t, s) + \frac{t}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u_m(s), u'_m(s)) \\ & \leq M_5 \left[1 + \frac{1}{1-\mu} \int_0^1 g(\tau) d\tau \right] (1-s)s, \quad s \in [0, 1]. \end{aligned} \quad (2.22)$$

By applying Lebesgue Dominated Convergence theorem, we get

$$\begin{aligned} \lim_{m \rightarrow \infty} (Tu_m)(t) &= \lim_{m \rightarrow \infty} \int_0^1 \left[G_1(t, s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u_m(s), u'_m(s)) ds \\ &= \int_0^1 \left[G_1(t, s) + \frac{t^2}{2(1-\mu)} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u(s), u'(s)) ds \\ &= (Tu)(t), \quad t \in [0, 1], \end{aligned} \quad (2.23)$$

$$\begin{aligned} \lim_{m \rightarrow \infty} (Tu_m)'(t) &= \lim_{m \rightarrow \infty} \int_0^1 \left[G_2(t, s) + \frac{t}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u_m(s), u'_m(s)) ds \\ &= \int_0^1 \left[G_2(t, s) + \frac{t}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u(s), u'(s)) ds \\ &= (Tu)'(t), \quad t \in [0, 1], \end{aligned}$$

which indicates that T is continuous. Therefore, $T : K \rightarrow K$ is completely continuous. \square

3. Main Results

For convenience, we define

$$\begin{aligned}
 f^0 &= \limsup_{x+y \rightarrow 0^+} \max_{t \in [0,1]} \frac{f(t, x, y)}{x+y}, & f_0 &= \liminf_{x+y \rightarrow 0^+} \min_{t \in [\alpha, 1]} \frac{f(t, x, y)}{x+y}, \\
 f^\infty &= \limsup_{x+y \rightarrow +\infty} \max_{t \in [0,1]} \frac{f(t, x, y)}{x+y}, & f_\infty &= \liminf_{x+y \rightarrow +\infty} \min_{t \in [\alpha, 1]} \frac{f(t, x, y)}{x+y}, \\
 H_1 &= 2 \int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s)g(\tau) d\tau \right] ds, \\
 H_2 &= \frac{\beta}{2} \int_\alpha^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s)g(\tau) d\tau \right] ds.
 \end{aligned} \tag{3.1}$$

Theorem 3.1. *If $H_1 f^0 < 1 < H_2 f_\infty$, then the BVP (1.3) has at least one monotone positive solution.*

Proof. In view of $H_1 f^0 < 1$, there exists $\varepsilon_1 > 0$ such that

$$H_1(f^0 + \varepsilon_1) \leq 1. \tag{3.2}$$

By the definition of f^0 , we may choose $\rho_1 > 0$ so that

$$f(t, x, y) \leq (f^0 + \varepsilon_1)(x+y), \text{ for } t \in [0, 1], (x+y) \in [0, \rho_1]. \tag{3.3}$$

Let $\Omega_1 = \{u \in E : \|u\| < \rho_1/2\}$. Then for any $u \in K \cap \partial\Omega_1$, in view of (3.2) and (3.3), we have

$$\begin{aligned}
 (Tu)'(t) &= \int_0^1 \left[G_2(t, s) + \frac{t}{1-\mu} \int_0^1 G_2(\tau, s)g(\tau) d\tau \right] f(s, u(s), u'(s)) ds \\
 &\leq \int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s)g(\tau) d\tau \right] (f^0 + \varepsilon_1)(u(s) + u'(s)) ds \\
 &\leq H_1(f^0 + \varepsilon_1) \|u\| \\
 &\leq \|u\|, \quad t \in [0, 1].
 \end{aligned} \tag{3.4}$$

By integrating the above inequality on $[0, t]$, we get

$$(Tu)(t) \leq \|u\|, \quad t \in [0, 1], \tag{3.5}$$

which together with (3.4) implies that

$$\|Tu\| \leq \|u\|, \quad u \in K \cap \partial\Omega_1. \quad (3.6)$$

On the other hand, since $1 < H_2 f_\infty$, there exists $\varepsilon_2 > 0$ such that

$$H_2(f_\infty - \varepsilon_2) \geq 1. \quad (3.7)$$

By the definition of f_∞ , we may choose $\rho_2 > \rho_1$, so that

$$f(t, x, y) \geq (f_\infty - \varepsilon_2)(x + y), \quad \text{for } t \in [\alpha, 1], \quad (x + y) \in [\rho_2, +\infty). \quad (3.8)$$

Let $\Omega_2 = \{u \in E : \|u\| < \rho_2/\beta\}$. Then for any $u \in K \cap \partial\Omega_2$, in view of (3.7) and (3.8), we have

$$\begin{aligned} (Tu)(1) &= \int_0^1 \left[G_1(1, s) + \frac{1}{2(1-\mu)} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u(s), u'(s)) ds \\ &\geq \frac{1}{2} \int_\alpha^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] (f_\infty - \varepsilon_2)(u(s) + u'(s)) ds \\ &\geq H_2(f_\infty - \varepsilon_2) \|u\| \\ &\geq \|u\|, \end{aligned} \quad (3.9)$$

which implies that

$$\|Tu\| \geq \|u\|, \quad u \in K \cap \partial\Omega_2. \quad (3.10)$$

Therefore, it follows from (3.6), (3.10), and Theorem 1.1 that the operator T has one fixed point $u \in K \cap (\overline{\Omega_2} \setminus \Omega_1)$, which is a monotone positive solution of the BVP (1.3). \square

Theorem 3.2. *If $H_1 f_\infty < 1 < H_2 f_0$, then the BVP (1.3) has at least one monotone positive solution.*

Proof. The proof is similar to that of Theorem 3.1 and is therefore omitted. \square

Theorem 3.3. *If $H_1 f(t, x, y) < (x + y)$ for $t \in [0, 1]$ and $(x + y) \in [0, +\infty)$, then the BVP (1.3) has no monotone positive solution.*

Proof. Suppose on the contrary that u is a monotone positive solution of the BVP (1.3). Then $u(t) \geq 0$ and $u'(t) \geq 0$ for $t \in [0, 1]$, and

$$\begin{aligned} u'(t) &= \int_0^1 \left[G_2(t, s) + \frac{t}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u(s), u'(s)) ds \\ &\leq \int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] f(s, u(s), u'(s)) ds \\ &< \frac{1}{H_1} \int_0^1 \left[(1-s)s + \frac{1}{1-\mu} \int_0^1 G_2(\tau, s) g(\tau) d\tau \right] (u(s) + u'(s)) ds \\ &\leq \|u\|, \quad t \in [0, 1]. \end{aligned} \tag{3.11}$$

By integrating the above inequality on $[0, t]$, we get

$$u(t) < \|u\|, \quad t \in [0, 1], \tag{3.12}$$

which together with (3.11) implies that

$$\|u\| < \|u\|. \tag{3.13}$$

This is a contradiction. Therefore, the BVP (1.3) has no monotone positive solution. \square

Similarly, we can prove the following theorem.

Theorem 3.4. *If $H_2 f(t, x, y) > (x + y)$ for $t \in [\alpha, 1]$ and $(x + y) \in [0, +\infty)$, then the BVP (1.3) has no monotone positive solution.*

Example 3.5. Consider the following BVP:

$$\begin{aligned} u'''(t) + \frac{1}{1+t} \left[\frac{u(t) + u'(t)}{e^{u(t)+u'(t)}} + \frac{1000(u(t) + u'(t))^2}{1 + u(t) + u'(t)} \right] &= 0, \quad t \in [0, 1], \\ u(0) = u'(0) = 0, \quad u'(1) &= \int_0^1 t u'(t) dt. \end{aligned} \tag{3.14}$$

Since $f(t, x, y) = 1/(1+t)[((x+y)/e^{x+y}) + (1000(x+y)^2/(1+x+y))]$ and $g(t) = t$, if we choose $\alpha = 1/2$, then it is easy to compute that

$$f^0 = 1, \quad f_\infty = 500, \quad H_1 = \frac{11}{24}, \quad H_2 = \frac{91}{12288}, \tag{3.15}$$

which shows that

$$H_1 f^0 < 1 < H_2 f_\infty. \tag{3.16}$$

So, it follows from Theorem 3.1 that the BVP (3.14) has at least one monotone positive solution.

Acknowledgment

This work was supported by the National Natural Science Foundation of China (10801068).

References

- [1] M. Gregus, *Third Order Linear Differential Equations*, Mathematics and Its Applications, Reidel, Dordrecht, the Netherlands, 1987.
- [2] D. R. Anderson, "Green's function for a third-order generalized right focal problem," *Journal of Mathematical Analysis and Applications*, vol. 288, no. 1, pp. 1–14, 2003.
- [3] Z. Du, W. Ge, and X. Lin, "Existence of solutions for a class of third-order nonlinear boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 294, no. 1, pp. 104–112, 2004.
- [4] Y. Feng, "Solution and positive solution of a semilinear third-order equation," *Journal of Applied Mathematics and Computing*, vol. 29, no. 1-2, pp. 153–161, 2009.
- [5] Y. Feng and S. Liu, "Solvability of a third-order two-point boundary value problem," *Applied Mathematics Letters*, vol. 18, no. 9, pp. 1034–1040, 2005.
- [6] L.-J. Guo, J.-P. Sun, and Y.-H. Zhao, "Existence of positive solutions for nonlinear third-order three-point boundary value problems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 10, pp. 3151–3158, 2008.
- [7] J. Henderson and C. C. Tisdale, "Five-point boundary value problems for third-order differential equations by solution matching," *Mathematical and Computer Modelling*, vol. 42, no. 1-2, pp. 133–137, 2005.
- [8] B. Hopkins and N. Kosmatov, "Third-order boundary value problems with sign-changing solutions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 1, pp. 126–137, 2007.
- [9] Z. Liu, L. Debnath, and S. M. Kang, "Existence of monotone positive solutions to a third order two-point generalized right focal boundary value problem," *Computers & Mathematics with Applications*, vol. 55, no. 3, pp. 356–367, 2008.
- [10] Z. Liu, J. S. Ume, and S. M. Kang, "Positive solutions of a singular nonlinear third order two-point boundary value problem," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 1, pp. 589–601, 2007.
- [11] R. Ma, "Multiplicity results for a third order boundary value problem at resonance," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 32, no. 4, pp. 493–499, 1998.
- [12] Y. Sun, "Positive solutions for third-order three-point nonhomogeneous boundary value problems," *Applied Mathematics Letters*, vol. 22, no. 1, pp. 45–51, 2009.
- [13] Y. Sun, "Positive solutions of singular third-order three-point boundary value problem," *Journal of Mathematical Analysis and Applications*, vol. 306, no. 2, pp. 589–603, 2005.
- [14] B. Yang, "Positive solutions of a third-order three-point boundary-value problem," *Electronic Journal of Differential Equations*, vol. 2008, no. 99, pp. 1–10, 2008.
- [15] Q. Yao, "Positive solutions of singular third-order three-point boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 354, no. 1, pp. 207–212, 2009.
- [16] Q. Yao, "Successive iteration of positive solution for a discontinuous third-order boundary value problem," *Computers & Mathematics with Applications*, vol. 53, no. 5, pp. 741–749, 2007.
- [17] Q. Yao and Y. Feng, "The existence of solution for a third-order two-point boundary value problem," *Applied Mathematics Letters*, vol. 15, no. 2, pp. 227–232, 2002.
- [18] D. R. Anderson and C. C. Tisdell, "Third-order nonlocal problems with sign-changing nonlinearity on time scales," *Electronic Journal of Differential Equations*, vol. 2007, no. 19, pp. 1–12, 2007.
- [19] J. R. Graef and B. Yang, "Positive solutions of a third order nonlocal boundary value problem," *Discrete and Continuous Dynamical Systems. Series S*, vol. 1, no. 1, pp. 89–97, 2008.
- [20] A. Boucherif, "Second-order boundary value problems with integral boundary conditions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 1, pp. 364–371, 2009.
- [21] M. Feng, D. Ji, and W. Ge, "Positive solutions for a class of boundary-value problem with integral boundary conditions in Banach spaces," *Journal of Computational and Applied Mathematics*, vol. 222, no. 2, pp. 351–363, 2008.
- [22] L. Kong, "Second order singular boundary value problems with integral boundary conditions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 5, pp. 2628–2638, 2010.

- [23] X. Zhang, M. Feng, and W. Ge, "Existence result of second-order differential equations with integral boundary conditions at resonance," *Journal of Mathematical Analysis and Applications*, vol. 353, no. 1, pp. 311–319, 2009.
- [24] X. Zhang and W. Ge, "Positive solutions for a class of boundary-value problems with integral boundary conditions," *Computers & Mathematics with Applications*, vol. 58, no. 2, pp. 203–215, 2009.
- [25] D. J. Guo and V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, vol. 5 of *Notes and Reports in Mathematics in Science and Engineering*, Academic Press, Boston, Mass, USA, 1988.
- [26] L. H. Erbe and H. Wang, "On the existence of positive solutions of ordinary differential equations," *Proceedings of the American Mathematical Society*, vol. 120, no. 3, pp. 743–748, 1994.