

*Research Article*

# Common Fixed Point Theorem for Four Non-Self Mappings in Cone Metric Spaces

Xianjiu Huang,<sup>1</sup> Chuanxi Zhu,<sup>1</sup> and Xi Wen<sup>2</sup>

<sup>1</sup> Department of Mathematics, Nanchang University, Nanchang, Jiangxi 330031, China

<sup>2</sup> Department of Computer Science, Nanchang University, Nanchang, Jiangxi 330031, China

Correspondence should be addressed to Xianjiu Huang, xjhuangxwen@163.com

Received 13 June 2009; Revised 1 March 2010; Accepted 18 April 2010

Academic Editor: Lai Jiu Lin

Copyright © 2010 Xianjiu Huang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We extend a common fixed point theorem of Radenović and Rhoades for four non-self-mappings in cone metric spaces.

## 1. Introduction and Preliminaries

Recently, Huang and Zhang [1] generalized the concept of a metric space, replacing the set of real numbers by ordered Banach space and obtained some fixed point theorems for mappings satisfying different contractive conditions. Subsequently, the study of fixed point theorems in such spaces is followed by some other mathematicians; see [2–8]. The aim of this paper is to prove a common fixed point theorem for four non-self-mappings on cone metric spaces in which the cone need not be normal. This result generalizes the result of Radenović and Rhoades [5].

Consistent with Huang and Zhang [1], the following definitions and results will be needed in the sequel.

Let  $E$  be a real Banach space. A subset  $P$  of  $E$  is called a cone if and only if

- (a)  $P$  is closed, nonempty and  $P \neq \{\theta\}$ ;
- (b)  $a, b \in \mathbb{R}$ ,  $a, b \geq 0$ ,  $x, y \in P$  implies  $ax + by \in P$ ;
- (c)  $P \cap (-P) = \{\theta\}$ .

Given a cone  $P \subset E$ , we define a partial ordering  $\leq$  with respect to  $P$  by  $x \leq y$  if and only if  $y - x \in P$ . A cone  $P$  is called normal if there is a number  $K > 0$  such that for all  $x, y \in E$ ,

$$\theta \leq x \leq y \quad \text{implies} \quad \|x\| \leq K\|y\|. \quad (1.1)$$

The least positive number satisfying the above inequality is called the normal constant of  $P$ , while  $x \ll y$  stands for  $y - x \in \text{int } P$  (interior of  $P$ ).

*Definition 1.1* (see [1]). Let  $X$  be a nonempty set. Suppose that the mapping  $d : X \times X \rightarrow E$  satisfies

- (d1)  $\theta \leq d(x, y)$  for all  $x, y \in X$  and  $d(x, y) = \theta$  if and only if  $x = y$ ;
- (d2)  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ;
- (d3)  $d(x, y) \leq d(x, z) + d(z, y)$  for all  $x, y, z \in X$ .

Then  $d$  is called a cone metric on  $X$ , and  $(X, d)$  is called a cone metric space.

The concept of a cone metric space is more general than that of a metric space.

*Definition 1.2* (see [1]). Let  $(X, d)$  be a cone metric space. One says that  $\{x_n\}$  is

- (e) a Cauchy sequence if for every  $c \in E$  with  $\theta \ll c$ , there is an  $N$  such that for all  $n, m > N$ ,  $d(x_n, x_m) \ll c$ ;
- (f) a Convergent sequence if for every  $c \in E$  with  $\theta \ll c$ , there is an  $N$  such that for all  $n > N$ ,  $d(x_n, x) \ll c$  for some fixed  $x \in X$ .

A cone metric space  $X$  is said to be complete if every Cauchy sequence in  $X$  is convergent in  $X$ . It is known that  $\{x_n\}$  converges to  $x \in X$  if and only if  $d(x_n, x) \rightarrow \theta$  as  $n \rightarrow \infty$ . It is a Cauchy sequence if and only if  $d(x_n, x_m) \rightarrow \theta$  ( $n, m \rightarrow \infty$ ).

*Remark 1.3* (see [9]). Let  $E$  be an ordered Banach (normed) space. Then  $c$  is an interior point of  $P$  if and only if  $[-c, c]$  is a neighborhood of  $\theta$ .

**Corollary 1.4** (see [10]). (1) If  $a \leq b$  and  $b \ll c$ , then  $a \ll c$ .

Indeed,  $c - a = (c - b) + (b - a) \geq c - b$  implies  $[-(c - a), c - a] \supseteq [-(c - b), c - b]$ .

(2) If  $a \ll b$  and  $b \ll c$ , then  $a \ll c$ .

Indeed,  $c - a = (c - b) + (b - a) \geq c - b$  implies  $[-(c - a), c - a] \supseteq [-(c - b), c - b]$ .

(3) If  $\theta \leq u \ll c$  for each  $c \in \text{int } P$ , then  $u = \theta$ .

*Remark 1.5* (see [5, 11]). If  $c \in \text{int } P$ ,  $\theta \leq a_n$ , and  $a_n \rightarrow \theta$ , then there exists an  $n_0$  such that for all  $n > n_0$  we have  $a_n \ll c$ .

*Remark 1.6* (see [6, 10]). If  $E$  is a real Banach space with cone  $P$  and if  $a \leq ka$  where  $a \in P$  and  $0 < k < 1$ , then  $a = \theta$ .

We find it convenient to introduce the following definition.

*Definition 1.7* (see [5]). Let  $(X, d)$  be a complete cone metric space and  $C$  a nonempty closed subset of  $X$ , and  $f, g : C \rightarrow X$  satisfying

$$d(fx, fy) \leq \lambda u, \quad (1.2)$$

where

$$u \in \left\{ \frac{d(gx, gy)}{2}, d(fx, gx), d(fy, gy), \frac{d(fx, gy) + d(fy, gx)}{q} \right\}, \quad (1.3)$$

for all  $x, y \in C$ ,  $0 < \lambda < 1/2$ ,  $q \geq 2 - \lambda$ , then  $f$  is called a generalized  $g$ -contractive mapping of  $C$  into  $X$ .

*Definition 1.8* (see [2]). Let  $f$  and  $g$  be self-maps on a set  $X$  (i.e.,  $f, g : X \rightarrow X$ ). If  $w = fx = gx$  for some  $x$  in  $X$ , then  $x$  is called a coincidence point of  $f$  and  $g$ , and  $w$  is called a point of coincidence of  $f$  and  $g$ . Self-maps  $f$  and  $g$  are said to be weakly compatible if they commute at their coincidence point; that is, if  $fx = gx$  for some  $x \in X$ , then  $fgx = gfx$ .

## 2. Main Result

The following theorem is Radenović and Rhoades [5] generalization of Imdad and Kumar's [12] result in cone metric spaces.

**Theorem 2.1.** *Let  $(X, d)$  be a complete cone metric space and  $C$  a nonempty closed subset of  $X$  such that for each  $x \in C$  and  $y \notin C$  there exists a point  $z \in \partial C$  (the boundary of  $C$ ) such that*

$$d(x, z) + d(z, y) = d(x, y). \quad (2.1)$$

*Suppose that  $f, g : C \rightarrow X$  are such that  $f$  is a generalized  $g$ -contractive mapping of  $C$  into  $X$ , and*

- (i)  $\partial C \subseteq gC$ ,  $fC \cap C \subseteq gC$ ,
- (ii)  $gx \subseteq \partial C \Rightarrow fx \in C$ ,
- (iii)  $gC$  is closed in  $X$ .

*Then the pair  $(f, g)$  has a coincidence point. Moreover, if pair  $(f, g)$  is weakly compatible, then  $f$  and  $g$  have a unique common fixed point.*

The purpose of this paper is to extend the above theorem for four non-self-mappings in cone metric spaces. We begin with the following definition.

*Definition 2.2.* Let  $(X, d)$  be a complete cone metric space and  $C$  a nonempty closed subset of  $X$ , and  $F, G, S, T : C \rightarrow X$  satisfying

$$d(Fx, Gy) \leq \lambda u, \quad (2.2)$$

where

$$u \in \left\{ \frac{d(Tx, Sy)}{2}, d(Tx, Fx), d(Sy, Gy), \frac{d(Tx, Gy) + d(Sy, Fx)}{q} \right\}, \quad (2.3)$$

for all  $x, y \in C$ ,  $0 < \lambda < 1/2$ ,  $q \geq 2 - \lambda$ , then  $(F, G)$  is called a generalized  $(T, S)$ -contractive mappings pair of  $C$  into  $X$ .

Notice that by setting  $G = F = f$  and  $T = S = g$  in (2.2), one deduces the slightly generalized form of (1.3).

We state and prove our main result as follows.

**Theorem 2.3.** *Let  $(X, d)$  be a complete cone metric space and  $C$  a nonempty closed subset of  $X$  such that for each  $x \in C$  and  $y \notin C$  there exists a point  $z \in \partial C$  (the boundary of  $C$ ) such that*

$$d(x, z) + d(z, y) = d(x, y). \quad (2.4)$$

Suppose that  $F, G, S, T : C \rightarrow X$  are such that  $(F, G)$  is a generalized  $(T, S)$ -contractive mappings pair of  $C$  into  $X$ , and

- (I)  $\partial C \subseteq SC \cap TC, FC \cap C \subseteq SC, GC \cap C \subseteq TC,$
- (II)  $Tx \subseteq \partial C \Rightarrow Fx \in C, Sx \subseteq \partial C \Rightarrow Gx \in C,$
- (III)  $SC$  and  $TC$  (or  $FC$  and  $GC$ ) are closed in  $X$ .

Then

- (IV)  $(F, T)$  has a point of coincidence,
- (V)  $(G, S)$  has a point of coincidence.

Moreover, if  $(F, T)$  and  $(G, S)$  are weakly compatible pairs, then  $F, G, S,$  and  $T$  have a unique common fixed point.

*Proof.* Firstly, we proceed to construct two sequences  $\{x_n\}$  and  $\{y_n\}$  in the following way.

Let  $x \in \partial C$  be arbitrary. Then (due to  $\partial C \subseteq TC$ ) there exists a point  $x_0 \in C$  such that  $x = Tx_0$ . Since  $Tx \subseteq \partial C \Rightarrow Fx \in C$ , one concludes that  $Fx_0 \in FC \cap C \subseteq SC$ . Thus, there exists  $x_1 \in C$  such that  $y_1 = Sx_1 = Fx_0 \in C$ . Since  $y_1 = Fx_0$  there exists a point  $y_2 = Gx_1$  such that

$$d(y_1, y_2) = d(Fx_0, Gx_1). \quad (2.5)$$

Suppose that  $y_2 \in C$ . Then  $y_2 \in GC \cap C \subseteq TC$  which implies that there exists a point  $x_2 \in C$  such that  $y_2 = Tx_2$ . Otherwise, if  $y_2 \notin C$ , then there exists a point  $p \in \partial C$  such that

$$d(Sx_1, p) + d(p, y_2) = d(Sx_1, y_2). \quad (2.6)$$

Since  $p \in \partial C \subseteq TC$  there exists a point  $x_2 \in C$  with  $p = Tx_2$ , so that

$$d(Sx_1, Tx_2) + d(Tx_2, y_2) = d(Sx_1, y_2). \quad (2.7)$$

Let  $y_3 = Fx_2$  be such that  $d(y_2, y_3) = d(Gx_1, Fx_2)$ . Thus, repeating the foregoing arguments, one obtains two sequences  $\{x_n\}$  and  $\{y_n\}$  such that

- (a)  $y_{2n} = Gx_{2n-1}, y_{2n+1} = Fx_{2n},$
- (b)  $y_{2n} \in C \Rightarrow y_{2n} = Tx_{2n}$  or  $y_{2n} \notin C \Rightarrow Tx_{2n} \in \partial C,$

$$d(Sx_{2n-1}, Tx_{2n}) + d(Tx_{2n}, y_{2n}) = d(Sx_{2n-1}, y_{2n}). \quad (2.8)$$

(c)  $y_{2n+1} \in C \Rightarrow y_{2n+1} = Sx_{2n+1}$  or  $y_{2n+1} \notin C \Rightarrow Sx_{2n+1} \in \partial C$ ,

$$d(Tx_{2n}, Sx_{2n+1}) + d(Sx_{2n+1}, y_{2n+1}) = d(Tx_{2n}, y_{2n+1}). \quad (2.9)$$

We denote that

$$\begin{aligned} P_0 &= \{Tx_{2i} \in \{Tx_{2n}\} : Tx_{2i} = y_{2i}\}, \\ P_1 &= \{Tx_{2i} \in \{Tx_{2n}\} : Tx_{2i} \neq y_{2i}\}, \\ Q_0 &= \{Sx_{2i+1} \in \{Sx_{2n+1}\} : Sx_{2i+1} = y_{2i+1}\}, \\ Q_1 &= \{Sx_{2i+1} \in \{Sx_{2n+1}\} : Sx_{2i+1} \neq y_{2i+1}\}. \end{aligned} \quad (2.10)$$

Note that  $(Tx_{2n}, Sx_{2n+1}) \notin P_1 \times Q_1$ , as if  $Tx_{2n} \in P_1$ , then  $y_{2n} \neq Tx_{2n}$ , and one infers that  $Tx_{2n} \in \partial C$  which implies that  $y_{2n+1} = Fx_{2n} \in C$ . Hence  $y_{2n+1} = Sx_{2n+1} \in Q_0$ . Similarly, one can argue that  $(Sx_{2n-1}, Tx_{2n}) \notin Q_1 \times P_1$ .

Now, we distinguish the following three cases.  $\square$

*Case 1.* If  $(Tx_{2n}, Sx_{2n+1}) \in P_0 \times Q_0$ , then from (2.2)

$$d(Tx_{2n}, Sx_{2n+1}) = d(Fx_{2n}, Gx_{2n-1}) \leq \lambda u_{2n-1}, \quad (2.11)$$

where

$$\begin{aligned} u_{2n-1} &\in \left\{ \frac{d(Sx_{2n-1}, Tx_{2n})}{2}, d(Sx_{2n-1}, Gx_{2n-1}), d(Tx_{2n}, Fx_{2n}), \frac{d(Tx_{2n}, Gx_{2n-1}) + d(Sx_{2n-1}, Fx_{2n})}{q} \right\} \\ &= \left\{ \frac{d(y_{2n-1}, y_{2n})}{2}, d(y_{2n-1}, y_{2n}), d(y_{2n}, y_{2n+1}), \frac{d(y_{2n-1}, y_{2n+1})}{q} \right\}. \end{aligned} \quad (2.12)$$

Clearly, there are infinite many  $n$  such that at least one of the following four cases holds:

(1)

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda \frac{d(y_{2n-1}, y_{2n})}{2} \leq \lambda d(Sx_{2n-1}, Tx_{2n}), \quad (2.13)$$

(2)

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda d(y_{2n-1}, y_{2n}) = \lambda d(Sx_{2n-1}, Tx_{2n}), \quad (2.14)$$

(3)

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda d(y_{2n}, y_{2n+1}) \implies d(Tx_{2n}, Sx_{2n+1}) = \theta \leq \lambda d(Sx_{2n-1}, Tx_{2n}), \quad (2.15)$$

(4)

$$\begin{aligned} d(Tx_{2n}, Sx_{2n+1}) &\leq \lambda \frac{d(y_{2n-1}, y_{2n+1})}{q} \\ &\leq \lambda \frac{d(y_{2n-1}, y_{2n}) + d(y_{2n}, y_{2n+1})}{q} \\ &= \lambda \frac{d(Sx_{2n-1}, Tx_{2n}) + d(Tx_{2n}, Sx_{2n+1})}{q}, \end{aligned} \quad (2.16)$$

which implies  $(1 - \lambda/q)d(Tx_{2n}, Sx_{2n+1}) \leq (\lambda/q)d(Sx_{2n-1}, Tx_{2n})$ , that is,

$$d(Tx_{2n}, Sx_{2n+1}) \leq \frac{\lambda}{q - \lambda} d(Sx_{2n-1}, Tx_{2n}) \leq \lambda d(Sx_{2n-1}, Tx_{2n}). \quad (2.17)$$

From (1), (2), (3), and (4) it follows that

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda d(Sx_{2n-1}, Tx_{2n}). \quad (2.18)$$

Similarly, if  $(Sx_{2n+1}, Tx_{2n+2}) \in Q_0 \times P_0$ , we have

$$d(Sx_{2n+1}, Tx_{2n+2}) = d(Fx_{2n}, Gx_{2n+1}) \leq \lambda d(Tx_{2n}, Sx_{2n+1}). \quad (2.19)$$

If  $(Sx_{2n-1}, Tx_{2n}) \in Q_0 \times P_0$ , we have

$$d(Sx_{2n-1}, Tx_{2n}) = d(Fx_{2n-2}, Gx_{2n-1}) \leq \lambda d(Tx_{2n-2}, Sx_{2n-1}). \quad (2.20)$$

*Case 2.* If  $(Tx_{2n}, Sx_{2n+1}) \in P_0 \times Q_1$ , then  $Sx_{2n+1} \in Q_1$  and

$$d(Tx_{2n}, Sx_{2n+1}) + d(Sx_{2n+1}, y_{2n+1}) = d(Tx_{2n}, y_{2n+1}) \quad (2.21)$$

which in turn yields

$$d(Tx_{2n}, Sx_{2n+1}) \leq d(Tx_{2n}, y_{2n+1}) = d(y_{2n}, y_{2n+1}) \quad (2.22)$$

and hence

$$d(Tx_{2n}, Sx_{2n+1}) \leq d(y_{2n}, y_{2n+1}) = d(Fx_{2n}, Gx_{2n-1}). \quad (2.23)$$

Now, proceeding as in Case 1, we have that (2.18) holds.

If  $(Sx_{2n+1}, Tx_{2n+2}) \in Q_1 \times P_0$ , then  $Tx_{2n} \in P_0$ . We show that

$$d(Sx_{2n+1}, Tx_{2n+2}) \leq \lambda d(Tx_{2n}, Sx_{2n-1}). \quad (2.24)$$

Using (2.21), we get

$$\begin{aligned} d(Sx_{2n+1}, Tx_{2n+2}) &\leq d(Sx_{2n+1}, y_{2n+1}) + d(y_{2n+1}, Tx_{2n+2}) \\ &= d(Tx_{2n}, y_{2n+1}) - d(Tx_{2n}, Sx_{2n+1}) + d(y_{2n+1}, Tx_{2n+2}). \end{aligned} \quad (2.25)$$

By noting that  $Tx_{2n+2}, Tx_{2n} \in P_0$ , one can conclude that

$$\begin{aligned} d(y_{2n+1}, Tx_{2n+2}) &= d(y_{2n+1}, y_{2n+2}) = d(Fx_{2n}, Gx_{2n+1}) \leq \lambda d(Tx_{2n}, Sx_{2n+1}), \\ d(Tx_{2n}, y_{2n+1}) &= d(y_{2n}, y_{2n+1}) = d(Fx_{2n}, Gx_{2n-1}) \leq \lambda d(Sx_{2n-1}, Tx_{2n}), \end{aligned} \quad (2.26)$$

in view of Case 1.

Thus,

$$d(Sx_{2n+1}, Tx_{2n+2}) \leq \lambda d(Sx_{2n-1}, Tx_{2n}) - (1 - \lambda)d(Tx_{2n}, Sx_{2n+1}) \leq \lambda d(Sx_{2n-1}, Tx_{2n}), \quad (2.27)$$

and we proved (2.24).

*Case 3.* If  $(Tx_{2n}, Sx_{2n+1}) \in P_1 \times Q_0$ , then  $Sx_{2n-1} \in Q_0$ . We show that

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda d(Sx_{2n-1}, Tx_{2n-2}). \quad (2.28)$$

Since  $Tx_{2n} \in P_1$ , then

$$d(Sx_{2n-1}, Tx_{2n}) + d(Tx_{2n}, y_{2n}) = d(Sx_{2n-1}, y_{2n}). \quad (2.29)$$

From this, we get

$$\begin{aligned} d(Tx_{2n}, Sx_{2n+1}) &\leq d(Tx_{2n}, y_{2n}) + d(y_{2n}, Sx_{2n+1}) \\ &= d(Sx_{2n-1}, y_{2n}) - d(Sx_{2n-1}, Tx_{2n}) + d(y_{2n}, Sx_{2n+1}). \end{aligned} \quad (2.30)$$

By noting that  $Sx_{2n+1}, Sx_{2n-1} \in Q_0$ , one can conclude that

$$\begin{aligned} d(y_{2n}, Sx_{2n+1}) &= d(y_{2n}, y_{2n+1}) = d(Fx_{2n}, Gx_{2n-1}) \leq \lambda d(Sx_{2n-1}, Tx_{2n}), \\ d(Sx_{2n-1}, y_{2n}) &= d(y_{2n-1}, y_{2n}) = d(Fx_{2n-2}, Gx_{2n-1}) \leq \lambda d(Sx_{2n-1}, Tx_{2n-2}), \end{aligned} \quad (2.31)$$

in view of Case 1.

Thus,

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda d(Sx_{2n-1}, Tx_{2n-2}) - (1 - \lambda)d(Sx_{2n-1}, Tx_{2n}) \leq \lambda d(Sx_{2n-1}, Tx_{2n-2}), \quad (2.32)$$

and we proved (2.28).

Similarly, if  $(Sx_{2n+1}, Tx_{2n+2}) \in Q_0 \times P_1$ , then  $Tx_{2n+2} \in P_1$ , and

$$d(Sx_{2n+1}, Tx_{2n+2}) + d(Tx_{2n+2}, y_{2n+2}) = d(Sx_{2n+1}, y_{2n+2}). \quad (2.33)$$

From this, we have

$$\begin{aligned} d(Sx_{2n+1}, Tx_{2n+2}) &\leq d(Sx_{2n+1}, y_{2n+2}) + d(y_{2n+2}, Tx_{2n+2}) \\ &\leq d(Sx_{2n+1}, y_{2n+2}) + d(Sx_{2n+1}, y_{2n+2}) - d(Sx_{2n+1}, Tx_{2n+2}) \\ &= 2d(Sx_{2n+1}, y_{2n+2}) - d(Sx_{2n+1}, Tx_{2n+2}) \implies d(Sx_{2n+1}, Tx_{2n+2}) \\ &\leq d(Sx_{2n+1}, y_{2n+2}). \end{aligned} \quad (2.34)$$

By noting that  $Sx_{2n+1} \in Q_0$ , one can conclude that

$$d(Sx_{2n+1}, Tx_{2n+2}) \leq d(Sx_{2n+1}, y_{2n+2}) = d(Fx_{2n}, Gx_{2n+1}) \leq \lambda d(Tx_{2n}, Sx_{2n+1}), \quad (2.35)$$

in view of Case 1.

Thus, in all Cases 1–3, there exists  $w_{2n} \in \{d(Sx_{2n-1}, Tx_{2n}), d(Tx_{2n-2}, Sx_{2n-1})\}$  such that

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda w_{2n}, \quad (2.36)$$

and there exists  $w_{2n+1} \in \{d(Sx_{2n-1}, Tx_{2n}), d(Tx_{2n}, Sx_{2n+1})\}$  such that

$$d(Sx_{2n+1}, Tx_{2n+2}) \leq \lambda w_{2n+1}. \quad (2.37)$$

Following the procedure of Assad and Kirk [13], it can easily be shown by induction that, for  $n \geq 1$ , there exists  $w_2 \in \{d(Tx_0, Sx_1), d(Sx_1, Tx_2)\}$  such that

$$d(Tx_{2n}, Sx_{2n+1}) \leq \lambda^{n-1/2} w_2, \quad d(Sx_{2n+1}, Tx_{2n+2}) \leq \lambda^n w_2. \quad (2.38)$$

From (2.38) and by the triangle inequality, for  $n > m$ , we have

$$\begin{aligned} d(Tx_{2n}, Sx_{2m+1}) &\leq d(Tx_{2n}, Sx_{2n-1}) + d(Sx_{2n-1}, Tx_{2n-2}) + \cdots + d(Tx_{2m+2}, Sx_{2m+1}) \\ &\leq (\lambda^m + \lambda^{m+1/2} + \cdots + \lambda^{n-1}) w_2 \leq \frac{\lambda^m}{1 - \sqrt{\lambda}} w_2 \longrightarrow \theta, \quad \text{as } m \longrightarrow \infty. \end{aligned} \quad (2.39)$$

From Remark 1.5 and Corollary 1.4(1),  $d(Tx_{2n}, Sx_{2m+1}) \ll c$ .



Thus, the sequence  $\{Tx_0, Sx_1, Tx_2, Sx_3, \dots, Sx_{2n-1}, Tx_{2n}, Sx_{2n-1}, \dots\}$  is a Cauchy sequence. Then, as noted in [14], there exists at least one subsequence  $\{Tx_{2n_k}\}$  or  $\{Sx_{2n_k+1}\}$  which is contained in  $P_0$  or  $Q_0$ , respectively, and finds its limit  $z \in C$ . Furthermore, subsequences  $\{Tx_{2n_k}\}$  and  $\{Sx_{2n_k+1}\}$  both converge to  $z \in C$  as  $C$  is a closed subset of complete cone metric space  $(X, d)$ . We assume that there exists a subsequence  $\{Tx_{2n_k}\} \subseteq P_0$  for each  $k \in \mathbb{N}$ , then  $Tx_{2n_k} = y_{2n_k} = Gx_{2n_k-1} \in C \cap GC \subseteq TC$ . Since  $TC$  as well as  $SC$  are closed in  $X$ , and  $\{Tx_{2n_k}\}$  is Cauchy in  $TC$ , it converges to a point  $z \in TC$ . Let  $w \in T^{-1}z$ , then  $Tw = z$ . Similarly,  $\{Sx_{2n_k+1}\}$  a subsequence of Cauchy sequence  $\{Tx_0, Sx_1, Tx_2, Sx_3, \dots, Sx_{2n-1}, Tx_{2n}, Sx_{2n-1}, \dots\}$  also converges to  $z$  as  $SC$  is closed. Using (2.2), one can write

$$d(Fw, z) \leq d(Fw, Gx_{2n_k-1}) + d(Gx_{2n_k-1}, z) \leq \lambda u_{2n_k-1} + d(Gx_{2n_k-1}, z), \quad (2.40)$$

where

$$\begin{aligned} u_{2n_k-1} &\in \left\{ \frac{d(Tw, Sx_{2n_k-1})}{2}, d(Tw, Fw), d(Sx_{2n_k-1}, Gx_{2n_k-1}), \frac{d(Tw, Gx_{2n_k-1}) + d(Fw, Sx_{2n_k-1})}{q} \right\} \\ &= \left\{ \frac{d(z, Sx_{2n_k-1})}{2}, d(z, Fw), d(Sx_{2n_k-1}, Gx_{2n_k-1}), \frac{d(z, Gx_{2n_k-1}) + d(Fw, Sx_{2n_k-1})}{q} \right\}. \end{aligned} \quad (2.41)$$

Let  $\theta \ll c$ . Clearly at least one of the following four cases holds for infinitely many  $n$ :

(1)

$$d(Fw, z) \leq \lambda \frac{d(z, Sx_{2n_k-1})}{2} + d(Gx_{2n_k-1}, z) \ll \lambda \frac{c}{2\lambda} + \frac{c}{2} = c; \quad (2.42)$$

(2)

$$\begin{aligned} d(Fw, z) &\leq \lambda d(z, Fw) + d(Gx_{2n_k-1}, z) \implies d(Fw, z) \\ &\leq \frac{1}{1-\lambda} d(Gx_{2n_k-1}, z) \ll \frac{1}{1-\lambda} (1-\lambda)c = c; \end{aligned} \quad (2.43)$$

(3)

$$\begin{aligned} d(Fw, z) &\leq \lambda d(Sx_{2n_k-1}, Gx_{2n_k-1}) + d(Gx_{2n_k-1}, z) \\ &\leq \lambda (d(Sx_{2n_k-1}, z) + d(z, Gx_{2n_k-1})) + d(Gx_{2n_k-1}, z) \\ &\leq (\lambda + 1) d(Gx_{2n_k-1}, z) + \lambda d(Sx_{2n_k-1}, z) \\ &\ll (\lambda + 1) \frac{c}{2(\lambda + 1)} + \lambda \frac{c}{2\lambda} = c; \end{aligned} \quad (2.44)$$

(4)

$$\begin{aligned}
d(Fw, z) &\leq \lambda \frac{d(z, Gx_{2n_k-1}) + d(Fw, Sx_{2n_k-1})}{q} + d(Gx_{2n_k-1}, z) \\
&\leq \lambda \frac{d(z, Gx_{2n_k-1}) + d(Fw, z) + d(z, Sx_{2n_k-1})}{q} + d(Gx_{2n_k-1}, z) \implies d(Fw, z) \\
&\leq \frac{q+\lambda}{q-\lambda} d(Gx_{2n_k-1}, z) + \frac{\lambda}{q-\lambda} d(z, Sx_{2n_k-1}) \\
&\ll \frac{q+\lambda}{q-\lambda} \frac{c}{2((q+\lambda)/(q-\lambda))} + \frac{\lambda}{q-\lambda} \frac{c}{2(\lambda/(q-\lambda))} = c.
\end{aligned} \tag{2.45}$$

In all cases we obtain  $d(Fw, z) \ll c$  for each  $c \in \text{int } P$ . Using Corollary 1.4(3) it follows that  $d(Fw, z) = \theta$  or  $Fw = z$ . Thus,  $Fw = z = Tw$ , that is,  $z$  is a coincidence point of  $F, T$ .

Further, since Cauchy sequence  $\{Tx_0, Sx_1, Tx_2, Sx_3, \dots, Sx_{2n-1}, Tx_{2n}, Sx_{2n-1}, \dots\}$  converges to  $z \in C$  and  $z = Fw, z \in FC \cap C \subseteq SC$ , there exists  $v \in C$  such that  $Sv = z$ . Again using (2.2), we get

$$d(Sv, Gv) = d(z, Gv) = d(Fw, Gv) \leq \lambda u, \tag{2.46}$$

where

$$\begin{aligned}
u &\in \left\{ \frac{d(Tw, Sv)}{2}, d(Tw, Fw), d(Sv, Gv), \frac{d(Tw, Gv) + d(Fw, Sv)}{q} \right\} \\
&= \left\{ \theta, \theta, d(Sv, Gv), \frac{d(z, Gv) + \theta}{q} \right\} \\
&= \left\{ \theta, d(Sv, Gv), \frac{d(Sv, Gv)}{q} \right\}.
\end{aligned} \tag{2.47}$$

Hence, we get the following cases:

$$d(Sv, Gv) \leq \lambda\theta = \theta, \quad d(Sv, Gv) \leq \lambda d(Sv, Gv), \quad d(Sv, Gv) \leq \frac{\lambda}{q} d(Sv, Gv). \tag{2.48}$$

Since  $\lambda/q \leq \lambda/(2-\lambda) = \lambda/(1+(1-\lambda)) < \lambda$ , using Remark 1.6 and Corollary 1.4(3), it follows that  $Sv = Gv$ ; therefore,  $Sv = z = Gv$ , that is,  $z$  is a coincidence point of  $(G, S)$ .

In case  $FC$  and  $GC$  are closed in  $X$ ,  $z \in FC \cap C \subseteq SC$  or  $z \in GC \cap C \subseteq TC$ . The analogous arguments establish (IV) and (V). If we assume that there exists a subsequence  $\{Sx_{2n_k+1}\} \subseteq Q_0$  with  $TC$  as well  $SC$  being closed in  $X$ , then noting that  $\{Sx_{2n_k+1}\}$  is a Cauchy sequence in  $SC$ , foregoing arguments establish (IV) and (V).

Suppose now that  $(F, T)$  and  $(G, S)$  are weakly compatible pairs, then

$$z = Fw = Tw \implies Fz = FTw = TFw = Tz, \quad z = Gv = Sv \implies Gz = GSv = SGv = Sz. \tag{2.49}$$

Then, from (2.2),

$$d(Fz, z) = d(Fz, Gv) \leq \lambda u, \quad (2.50)$$

where

$$\begin{aligned} u &\in \left\{ \frac{d(Sv, Tz)}{2}, d(Tz, Fz), d(Sv, Gv), \frac{d(Tz, Gv) + d(Sv, Fz)}{q} \right\} \\ &= \left\{ \frac{d(z, Fz)}{2}, d(Fz, Fz), d(z, z), \frac{d(Fz, z) + d(z, Fz)}{q} \right\} \\ &= \left\{ \frac{d(z, Fz)}{2}, \theta, \frac{2d(z, Fz)}{q} \right\}. \end{aligned} \quad (2.51)$$

Hence, we get the following cases:

$$d(Fz, z) \leq \lambda \frac{d(z, Fz)}{2}, \quad d(Fz, z) \leq \lambda \theta = \theta \quad \text{and} \quad d(Fz, z) \leq \frac{2\lambda d(z, Fz)}{q}, \quad (2.52)$$

Since  $2\lambda/q \leq 2\lambda/(2 - \lambda) = 2\lambda/(1 + (1 - \lambda)) < 2\lambda < 1$ , using Remark 1.6 and Corollary 1.4(3), it follows that  $Fz = z$ . Thus,  $Fz = z = Tz$ .

Similarly, we can prove that  $Gz = z = Sz$ . Therefore  $z = Fz = Gz = Sz = Tz$ , that is,  $z$  is a common fixed point of  $F, G, S$ , and  $T$ .

Uniqueness of the common fixed point follows easily from (2.2).

The following example shows that in general  $F, G, S$ , and  $T$  satisfying the hypotheses of Theorem 2.3 need not have a common coincidence justifying two separate conclusions (IV) and (V).

*Example 2.4.* Let  $E = C^1([0, 1], \mathbb{R})$ ,  $P = \{\varphi \in E : \varphi(t) \geq 0, t \in [0, 1]\}$ ,  $X = [0, +\infty)$ ,  $C = [0, 2]$ , and  $d : X \times X \rightarrow E$  defined by  $d(x, y) = |x - y|\varphi$ , where  $\varphi \in P$  is a fixed function, for example,  $\varphi(t) = e^t$ . Then  $(X, d)$  is a complete cone metric space with a nonnormal cone having the nonempty interior. Define  $F, G, S$ , and  $T : C \rightarrow X$  as

$$Fx = x + \frac{4}{5}, \quad Gx = x^2 + \frac{4}{5}, \quad Tx = 5x, \quad Sx = 5x^2, \quad x \in C. \quad (2.53)$$

Since  $\partial C = \{0, 2\}$ . Clearly, for each  $x \in C$  and  $y \notin C$  there exists a point  $z = 2 \in \partial C$  such that  $d(x, z) + d(z, y) = d(x, y)$ . Further,  $SC \cap TC = [0, 20] \cap [0, 10] = [0, 10] \supset \{0, 2\} = \partial C$ ,  $FC \cap C = [4/5, 14/5] \cap [0, 2] = [4/5, 2] \subset SC$ ,  $GC \cap C = [4/5, 24/5] \cap [0, 2] = [4/5, 2] \subset TC$ , and  $SC, TC, FC$ , and  $GC$  are closed in  $X$ .

Also,

$$\begin{aligned} T0 = 0 \in \partial C \implies F0 = \frac{4}{5} \in C, \quad S0 = 0 \in \partial C \implies G0 = \frac{4}{5} \in C, \\ T\left(\frac{2}{5}\right) = 2 \in \partial C \implies F\left(\frac{2}{5}\right) = \frac{6}{5} \in C, \quad S\left(\sqrt{\frac{2}{5}}\right) = 2 \in \partial C \implies G\left(\sqrt{\frac{2}{5}}\right) = \frac{6}{5} \in C. \end{aligned} \quad (2.54)$$

Moreover, for each  $x, y \in C$ ,

$$d(Fx, Gy) = |x - y^2| \varphi = \frac{2}{5} \left( \frac{1}{2} d(Tx, Sy) \right), \quad (2.55)$$

that is, (2.2) is satisfied with  $\lambda = 2/5$ .

Evidently,  $1 = T(1/5) = F(1/5) \neq 1/5$  and  $1 = S(1/\sqrt{5}) = G(1/\sqrt{5}) \neq 1/\sqrt{5}$ . Notice that two separate coincidence points are not common fixed points as  $FT(1/5) \neq TF(1/5)$  and  $SG(1/\sqrt{5}) \neq GS(1/\sqrt{5})$ , which shows necessity of weakly compatible property in Theorem 2.3.

Next, we furnish an illustrate example in support of our result. In doing so, we are essentially inspired by Imdad and Kumar [12].

*Example 2.5.* Let  $E = C^1([0, 1], R)$ ,  $P = \{\varphi \in E : \varphi(t) \geq 0, t \in [0, 1]\}$ ,  $X = [1, +\infty)$ ,  $C = [1, 3]$ , and  $d : X \times X \rightarrow E$  defined by  $d(x, y) = |x - y| \varphi$ , where  $\varphi \in P$  is a fixed function, for example,  $\varphi(t) = e^t$ . Then  $(X, d)$  is a complete cone metric space with a nonnormal cone having the nonempty interior. Define  $F, G, S$ , and  $T : C \rightarrow X$  as

$$\begin{aligned} Fx = \begin{cases} x^2 & \text{if } 1 \leq x \leq 2, \\ 2 & \text{if } 2 < x \leq 3, \end{cases} \quad Tx = \begin{cases} 4x^4 - 3 & \text{if } 1 \leq x \leq 2, \\ 13 & \text{if } 2 < x \leq 3, \end{cases} \\ Gx = \begin{cases} x^3 & \text{if } 1 \leq x \leq 2, \\ 2 & \text{if } 2 < x \leq 3, \end{cases} \quad Sx = \begin{cases} 4x^6 - 3 & \text{if } 1 \leq x \leq 2, \\ 13 & \text{if } 2 < x \leq 3. \end{cases} \end{aligned} \quad (2.56)$$

Since  $\partial C = \{1, 3\}$ . Clearly, for each  $x \in C$  and  $y \notin C$  there exists a point  $z = 3 \in \partial C$  such that  $d(x, z) + d(z, y) = d(x, y)$ . Further,  $SC \cap TC = [1, 253] \cap [1, 61] = [1, 61] \supset \{1, 3\} = \partial C$ ,  $FC \cap C = [1, 4] \cap [1, 3] = [1, 3] \subset SC$ , and  $GC \cap C = [1, 8] \cap [1, 3] = [1, 3] \subset TC$ .

Also,

$$\begin{aligned} T1 = 1 \in \partial C \implies F1 = 1 \in C, \quad S1 = 1 \in \partial C \implies G1 = 1 \in C, \\ T\left(\sqrt[4]{\frac{3}{2}}\right) = 3 \in \partial C \implies F\left(\sqrt[4]{\frac{3}{2}}\right) = \sqrt{\frac{3}{2}} \in C, \quad S\left(\sqrt[6]{\frac{3}{2}}\right) = 3 \in \partial C \implies G\left(\sqrt[6]{\frac{3}{2}}\right) = \sqrt{\frac{3}{2}} \in C. \end{aligned} \quad (2.57)$$

Moreover, if  $x \in [1, 2]$  and  $y \in [2, 3]$ , then

$$d(Fx, Gy) = |x^2 - 2|\varphi = \frac{|x^4 - 4|}{|x^2 + 2|}\varphi = \frac{4|x^4 - 4|/2}{2|x^2 + 2|}\varphi = \frac{1}{2(x^2 + 2)} \frac{d(Tx, Sy)}{2}. \quad (2.58)$$

Next, if  $x, y \in (2, 3]$ , then

$$d(Fx, Gy) = 0 = \lambda \cdot \frac{d(Tx, Sy)}{2}. \quad (2.59)$$

Finally, if  $x, y \in [1, 2]$ , then

$$d(Fx, Gy) = |x^2 - y^3|\varphi = \frac{|x^4 - y^6|}{|x^2 + y^3|}\varphi = \frac{4|x^4 - y^6|/2}{2|x^2 + y^3|}\varphi = \frac{1}{2(x^2 + y^3)} \frac{d(Tx, Sy)}{2}. \quad (2.60)$$

Therefore, condition (2.2) is satisfied if we choose  $\lambda = \max\{1/2(x^2 + 2), 1/2(x^2 + y^3)\} \in (0, 1/2)$ . Moreover 1 is a point of coincidence as  $T1 = F1$  as well as  $S1 = G1$  whereas both the pairs  $(F, T)$  and  $(G, S)$  are weakly compatible as  $TF1 = 1 = FT1$  and  $SG1 = 1 = GS1$ . Also,  $SC, TC, FC$ , and  $GC$  are closed in  $X$ . Thus, all the conditions of Theorem 2.3 are satisfied and 1 is the unique common fixed point of  $F, G, S$ , and  $T$ . One may note that 1 is also a point of coincidence for both the pairs  $(F, T)$  and  $(G, S)$ .

*Remark 2.6.* (1) Setting  $G = F = f$  and  $T = S = g$  in Theorem 2.3, one deduces Theorem 2.1 due to [5].

(2) Setting  $G = F = f$  and  $T = S = I_X$  in Theorem 2.3, we obtain the following result.

**Corollary 2.7.** *Let  $(X, d)$  be a complete cone metric space and  $C$  a nonempty closed subset of  $X$  such that for each  $x \in C$  and  $y \notin C$  there exists a point  $z \in \partial C$  (the boundary of  $C$ ) such that*

$$d(x, z) + d(z, y) = d(x, y). \quad (2.61)$$

Suppose that  $f : C \rightarrow X$  satisfies the condition

$$d(fx, fy) \leq \lambda u(x, y), \quad (2.62)$$

where

$$u(x, y) \in \left\{ \frac{d(x, y)}{2}, d(x, fx), d(y, fy), \frac{d(x, fy) + d(y, fx)}{q} \right\} \quad (2.63)$$

for all  $x, y \in C$ ,  $0 < \lambda < 1/2$ ,  $q \geq 2 - \lambda$ , and  $f$  has the additional property that for each  $x \in \partial C$ ,  $fx \in C$ ,  $f$  has a unique fixed point.

## Acknowledgments

The authors would like to express their sincere appreciation to the referees for their very helpful suggestions and many kind comments. This project was supported by the National Natural Science Foundation of China (10461007 and 10761007) and supported partly by the Provincial Natural Science Foundation of Jiangxi, China (2008GZS0076 and 2009GZS0019).

## References

- [1] L.-G. Huang and X. Zhang, "Cone metric spaces and fixed point theorems of contractive mappings," *Journal of Mathematical Analysis and Applications*, vol. 332, no. 2, pp. 1468–1476, 2007.
- [2] M. Abbas and G. Jungck, "Common fixed point results for noncommuting mappings without continuity in cone metric spaces," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 416–420, 2008.
- [3] M. Abbas and B. E. Rhoades, "Fixed and periodic point results in cone metric spaces," *Applied Mathematics Letters*, vol. 22, no. 4, pp. 511–515, 2009.
- [4] D. Ilić and V. Rakočević, "Common fixed points for maps on cone metric space," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 2, pp. 876–882, 2008.
- [5] S. Radenović and B. E. Rhoades, "Fixed point theorem for two non-self mappings in cone metric spaces," *Computers & Mathematics with Applications*, vol. 57, no. 10, pp. 1701–1707, 2009.
- [6] Sh. Rezapour and R. Hamlbarani, "Some notes on the paper: "Cone metric spaces and fixed point theorems of contractive mappings"," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 2, pp. 719–724, 2008.
- [7] P. Vetro, "Common fixed points in cone metric spaces," *Rendiconti del Circolo Matematico di Palermo*, vol. 56, no. 3, pp. 464–468, 2007.
- [8] D. Wardowski, "Endpoints and fixed points of set-valued contractions in cone metric spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 1-2, pp. 512–516, 2009.
- [9] Y. C. Wong and K. F. Ng, *Partially Ordered Topological Vector Spaces*, Oxford Mathematical Monographs, Clarendon Press, Oxford, UK, 1973.
- [10] Sh. Rezapour, "A review on topological properties of cone metric spaces," in *Analysis, Topology and Applications (ATA '08)*, Vrnjacka Banja, Serbia, May-June 2008.
- [11] G. Jungck, S. Radenović, S. Radojević, and V. Rakočević, "Common fixed point theorems for weakly compatible pairs on cone metric spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 643840, 13 pages, 2009.
- [12] M. Imdad and S. Kumar, "Rhoades-type fixed-point theorems for a pair of nonself mappings," *Computers & Mathematics with Applications*, vol. 46, no. 5-6, pp. 919–927, 2003.
- [13] N. A. Assad and W. A. Kirk, "Fixed point theorems for set-valued mappings of contractive type," *Pacific Journal of Mathematics*, vol. 43, pp. 553–562, 1972.
- [14] O. Hadžić and L. Gajić, "Coincidence points for set-valued mappings in convex metric spaces," *Univerzitet u Novom Sadu. Zbornik Radova Prirodno-Matematičkog Fakulteta. Serija za Matemati*, vol. 16, no. 1, pp. 13–25, 1986.