

Research Article

Bi-Lipschitz Mappings and Quasilinearly Subharmonic Functions

Oleksiy Dovgoshey¹ and Juhani Riihenta²

¹ Institute of Applied Mathematics and Mechanics, NASU, R. Luxemburg Street 74,
Donetsk 83114, Ukraine

² Department of Physics and Mathematics, University of Joensuu, P.O. Box 111,
80101 Joensuu, Finland

Correspondence should be addressed to Juhani Riihenta, juhani.riihenta@gmail.com

Received 30 November 2009; Accepted 25 December 2009

Academic Editor: Stanisława R. Kanas

Copyright © 2010 O. Dovgoshey and J. Riihenta. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

After considering a variant of the generalized mean value inequality of quasilinearly subharmonic functions, we consider certain invariance properties of quasilinearly subharmonic functions. Kojić has shown that in the plane case both the class of quasilinearly subharmonic functions and the class of regularly oscillating functions are invariant under conformal mappings. We give partial generalizations to her results by showing that in \mathbb{R}^n , $n \geq 2$, these both classes are invariant under bi-Lipschitz mappings.

1. Introduction

Notation. Our notation is rather standard; see, for example, [1–3] and the references therein. We recall here only the following. The Lebesgue measure in \mathbb{R}^n , $n \geq 2$, is denoted by m_n . We write $B^n(x, r)$ for the ball in \mathbb{R}^n , with center x and radius r . Recall that $m_n(B^n(x, r)) = v_n r^n$, where $v_n := m_n(B^n(0, 1))$. If D is an open set in \mathbb{R}^n , and $x \in D$, then we write $\delta_D(x)$ for the distance between the point x and the boundary ∂D of D . Our constants C are nonnegative, mostly ≥ 1 , and may vary from line to line.

1.1. Subharmonic Functions and Generalizations

Let Ω be an open set in \mathbb{R}^n , $n \geq 2$. Let $u : \Omega \rightarrow [-\infty, +\infty)$ be a Lebesgue measurable function. We adopt the following definitions.

(i) u is *subharmonic* if u is upper semicontinuous and if

$$u(x) \leq \frac{1}{v_n r^n} \int_{B^n(x,r)} u(y) dm_n(y) \quad (1.1)$$

for all balls $\overline{B^n(x,r)} \subset \Omega$. A subharmonic function may be $\equiv -\infty$ on any component of Ω ; see [3, page 9] and [4, page 60].

(ii) u is *nearly subharmonic* if $u^+ \in \mathcal{L}_{\text{loc}}^1(\Omega)$ and

$$u(x) \leq \frac{1}{v_n r^n} \int_{B^n(x,r)} u(y) dm_n(y) \quad (1.2)$$

for all balls $\overline{B^n(x,r)} \subset \Omega$. Observe that this definition, see [5, page 51], is slightly more general than the standard one [3, page 14].

(iii) Let $K \geq 1$. Then u is *K-quasilinearly subharmonic* if $u^+ \in \mathcal{L}_{\text{loc}}^1(\Omega)$ and

$$u_L(x) \leq \frac{K}{v_n r^n} \int_{B^n(x,r)} u_L(y) dm_n(y) \quad (1.3)$$

for all $L \geq 0$ and for all balls $\overline{B^n(x,r)} \subset \Omega$. Here $u_L := \max\{u, -L\} + L$.

The function u is *quasilinearly subharmonic* if u is K -quasilinearly subharmonic for some $K \geq 1$. For the definition and properties of quasilinearly subharmonic functions, see, for example, [1, 4–7] and the references therein.

Proposition 1.1 (cf. [5, Proposition 2.1, pages 54–55]). *The following holds.*

- (i) *A subharmonic function is nearly subharmonic but not conversely.*
- (ii) *A function is nearly subharmonic if and only if it is 1-quasilinearly subharmonic.*
- (iii) *A nearly subharmonic function is quasilinearly subharmonic but not conversely.*
- (iv) *If $u : \Omega \rightarrow [0, +\infty)$ is Lebesgue measurable, then u is K -quasilinearly subharmonic if and only if $u \in \mathcal{L}_{\text{loc}}^1(\Omega)$ and*

$$u(x) \leq \frac{K}{v_n r^n} \int_{B^n(x,r)} u(y) dm_n(y) \quad (1.4)$$

for all balls $\overline{B^n(x,r)} \subset \Omega$.

1.2. Bi-Lipschitz Mappings

Let D be an open set in \mathbb{R}^n , $n \geq 2$. Let $M \geq 1$ be arbitrary. A function $f : D \rightarrow \mathbb{R}^n$ is *M-bi-Lipschitz* if

$$\frac{|y-x|}{M} \leq |f(y) - f(x)| \leq M|y-x| \quad (1.5)$$

for all $x, y \in D$. A function is *bi-Lipschitz* if it is M -bi-Lipschitz for some $M \geq 1$. It is easy to see that if $f : D \rightarrow \mathbb{R}^n$ is M -bi-Lipschitz, then also $f^{-1} : D' \rightarrow \mathbb{R}^n$ is M -bi-Lipschitz, where $D' := f(D)$.

Let Ω be an open subset of \mathbb{R}^n . Let $p_D \in D$ and $x_\Omega \in \Omega$. We write

$$M\text{-BiLip}(p_D, x_\Omega, D, \Omega) := \{h : D \rightarrow \mathbb{R}^n : h \text{ is } M\text{-bi-Lipschitz, } h(p_D) = x_\Omega, h(D) \subset \Omega\}. \quad (1.6)$$

2. On the Generalized Mean Value Inequality

Lemma 2.1. *Let D be a bounded open set in \mathbb{R}^n , $n \geq 2$. Fix a point $p_D \in D$. Let Ω be a domain in \mathbb{R}^n . Let $u : \Omega \rightarrow [0, +\infty)$ be a K -quasilinearly subharmonic function. Then there is $C = C(K, n, D, M, p_D) \geq 1$ such that*

$$u(x_\Omega) \leq \frac{C}{m_n(h(D))} \int_{h(D)} u(y) dm_n(y) \quad (2.1)$$

for every point $x_\Omega \in \Omega$ and all $h \in M\text{-BiLip}(p_D, x_\Omega, D, \Omega)$, $M \geq 1$.

Proof. Take $x_\Omega \in \Omega$ and $h \in M\text{-BiLip}(p_D, x_\Omega, D, \Omega)$, $M \geq 1$, arbitrarily. (Observe that the set of bi-Lipschitz mappings is (in general) nonempty.) Write

$$R_D := \sup_{y \in D} |p_D - y|, \quad r_D := \delta_D(p_D). \quad (2.2)$$

Using the fact that $h|_{B^n(p_D, r_D)} : B^n(p_D, r_D) \rightarrow h(B^n(p_D, r_D))$ is a homeomorphism, one sees easily that $B^n(x_\Omega, r_D/M) \subset h(D)$. Since h is M -bi-Lipschitz, it follows from a result of Radó-Reichelderfer, see, for example, [8, Theorem 2.2, page 99], that $m_n(h(D)) \leq n!M^n m_n(D)$. (Observe that bi-Lipschitz mappings satisfy the property N and are differentiable almost everywhere, see, for example, [9, Theorem 33.2, page 112, Theorem 32.1, page 109].) Therefore,

$$\begin{aligned} u(x_\Omega) &\leq \frac{K}{v_n(r_D/M)^n} \int_{B^n(x_\Omega, r_D/M)} u(y) dm_n(y) \\ &\leq \frac{KM^n(R_D/r_D)^n}{m_n(B^n(p_D, R_D))} \int_{h(D)} u(y) dm_n(y) \\ &\leq \frac{KM^n(R_D/r_D)^n}{m_n(D)} \int_{h(D)} u(y) dm_n(y) \\ &\leq \frac{KM^n(R_D/r_D)^n}{m_n(h(D))/n!M^n} \int_{h(D)} u(y) dm_n(y) \\ &\leq \frac{n!KM^{2n}(R_D/r_D)^n}{m_n(h(D))} \int_{h(D)} u(y) dm_n(y). \end{aligned} \quad (2.3)$$

Thus (2.1) holds with $C = C(K, n, M, D, p_D)$. \square

Theorem 2.2. *Let D be an open set in \mathbb{R}^n , $n \geq 2$, with $m_n(D) < +\infty$. Fix a point $p_D \in D$. Let Ω be an open set in \mathbb{R}^n . Let $u : \Omega \rightarrow [0, +\infty)$ be a K -quasilinearly subharmonic function. Then there is a constant $C = C(K, n, D, M, p_D) \geq 1$ such that (2.1) holds for every point $x_\Omega \in \Omega$ and all $h \in M\text{-BiLip}(p_D, x_\Omega, D, \Omega)$, $M \geq 1$.*

Proof. Let $t > 1$ be arbitrary. It is easy to see that $tm_n(D \cap B^n(p_D, r_t)) \geq m_n(D)$ for some $r_t > 0$. Write $D_t := D \cap B^n(p_D, r_t)$ and $p_{D_t} = p_D$. One sees easily that D_t satisfies the assumptions of Lemma 2.1; that is, D_t is a bounded domain, $h(D_t) \subset h(D) \subset \Omega$ and $h(p_{D_t}) = h(p_D) = x_\Omega$. Hence there is a constant $C_1 = C_1(K, n, D, M, p_D) \geq 1$ such that

$$u(x_\Omega) \leq \frac{C_1}{m_n(h(D_t))} \int_{h(D_t)} u(y) dm_n(y) \quad (2.4)$$

for every point $x_\Omega \in \Omega$ and all $h \in M\text{-BiLip}(p_{D_t}, x_\Omega, D_t, \Omega)$. Since h and h^{-1} are M -bi-Lipschitz, it follows that $m_n(h(D)) \leq n!M^n m_n(D)$ and $m_n(D_t) \leq n!M^n m_n(h(D_t))$; see again [8, Theorem 2.2, page 99]. Thus for $C_2 = C_2(n, M) = (n!)^2 M^{2n}$,

$$\frac{m_n(D_t)}{m_n(D)} \leq C_2 \cdot \frac{m_n(h(D_t))}{m_n(h(D))}. \quad (2.5)$$

Proceed then as follows:

$$\begin{aligned} \frac{1}{m_n(h(D_t))} \int_{h(D_t)} u(y) dm_n(y) &\leq C_2 \cdot \frac{m_n(D)}{m_n(D_t)} \cdot \frac{1}{m_n(h(D))} \int_{h(D_t)} u(y) dm_n(y) \\ &\leq C_2 \cdot \frac{t}{m_n(h(D))} \int_{h(D_t)} u(y) dm_n(y) \\ &\leq C_2 \cdot \frac{t}{m_n(h(D))} \int_{h(D)} u(y) dm_n(y). \end{aligned} \quad (2.6)$$

Therefore

$$u(x_\Omega) \leq \frac{C_1 C_2 t}{m_n(h(D))} \int_{h(D)} u(y) dm_n(y), \quad (2.7)$$

concluding the proof. \square

3. An Invariance of the Class of Quasilinearly Subharmonic Functions

Suppose that G and U are open sets in the complex plane \mathbb{C} . If $f : U \rightarrow G$ is analytic and $u : G \rightarrow [-\infty, +\infty)$ is subharmonic, then $u \circ f$ is subharmonic; see, for example, [3, page 37] and [4, Corollary 3.3.4, page 70]. Using Koebe's one-quarter and distortion theorems, Kojic proved the following partial generalization.

Theorem 3.1 (see [6, Theorem 1, page 245]). *Let Ω and G be open sets in \mathbb{C} . Let $u : \Omega \rightarrow [0, +\infty)$ be a K -quasilinearly subharmonic function. If $\varphi : G \rightarrow \Omega$ is conformal, then the composition mapping $u \circ \varphi : G \rightarrow [0, +\infty)$ is C -quasilinearly subharmonic for some $C = C(K)$.*

For the definition and properties of conformal mappings, see, for example, [9, pages 13–15] and [8, pages 171–172].

Below we give a partial generalization to Kojić's result. Our result gives also a partial generalization to the standard result according to which in \mathbb{R}^n , $n \geq 2$, the class of subharmonic functions is invariant under orthogonal transformations; see [10, page 55].

Theorem 3.2. *Let Ω and U be open sets in \mathbb{R}^n , $n \geq 2$. Let $u : \Omega \rightarrow [0, +\infty)$ be a K -quasilinearly subharmonic function. If $f : U \rightarrow \Omega$ is M -bi-Lipschitz, then the composition mapping $u \circ f : U \rightarrow [0, +\infty)$ is C -quasilinearly subharmonic for some $C = C(K, n, M)$.*

Proof. It is sufficient to show that there exists a constant $C = C(K, n, f) > 0$ such that

$$(u \circ f)(x_0) \leq \frac{C}{m_n(B^n(x_0, r_0))} \int_{B^n(x_0, r_0)} (u \circ f)(x) dm_n(x) \quad (3.1)$$

for all $\overline{B^n(x_0, r_0)} \subset U$. To see this, observe first that

$$B^n\left(x'_0, \frac{r_0}{M}\right) \subset f(B^n(x_0, r_0)) \subset B^n(x'_0, Mr_0), \quad (3.2)$$

where $x'_0 = f(x_0)$.

Then

$$\begin{aligned} (u \circ f)(x_0) &= u(x'_0) \leq \frac{K}{m_n(B^n(x'_0, r_0/M))} \int_{B^n(x'_0, r_0/M)} u(y) dm_n(y) \\ &\leq \frac{K}{v_n(r_0/M)^n} \int_{B^n(x'_0, r_0/M)} (u \circ f)(f^{-1}(y)) dm_n(y) \\ &\leq \frac{KM^{2n}}{v_n(Mr_0)^n} \int_{B^n(x'_0, r_0/M)} (u \circ f)(f^{-1}(y)) dm_n(y) \\ &\leq \frac{KM^{2n}}{m_n(f(B^n(x_0, r_0)))} \int_{f(B^n(x_0, r_0))} (u \circ f)(f^{-1}(y)) dm_n(y) \\ &\leq \frac{KM^{2n}}{m_n(f(B^n(x_0, r_0)))} \int_{f(B^n(x_0, r_0))} (u \circ f)(f^{-1}(y)) |J_{f^{-1}}(y)| \cdot \frac{1}{|J_{f^{-1}}(y)|} dm_n(y) \\ &\leq \frac{KM^{2n}}{m_n(f(B^n(x_0, r_0)))} \int_{f(B^n(x_0, r_0))} (u \circ f)(f^{-1}(y)) |J_{f^{-1}}(y)| \cdot |J_f(f^{-1}(y))| dm_n(y) \\ &\leq \frac{KM^{2n}}{m_n(f(B^n(x_0, r_0)))} \int_{f(B^n(x_0, r_0))} (u \circ f)(f^{-1}(y)) |J_{f^{-1}}(y)| \cdot n! M^n dm_n(y) \\ &\leq \frac{n! KM^{3n}}{m_n(f(B^n(x_0, r_0)))} \int_{f(B^n(x_0, r_0))} (u \circ f)(f^{-1}(y)) |J_{f^{-1}}(y)| dm_n(y) \end{aligned}$$

$$\begin{aligned}
&\leq \frac{n!KM^{3n}}{m_n(f(B^n(x_0, r_0)))} \int_{B^n(x_0, r_0)} (u \circ f)(x) dm_n(x) \\
&\leq \frac{n!KM^{3n}}{m_n(B^n(x'_0, r_0/M))} \int_{B^n(x_0, r_0)} (u \circ f)(x) dm_n(x) \\
&\leq \frac{n!KM^{4n}}{m_n(B^n(x_0, r_0))} \int_{B^n(x_0, r_0)} (u \circ f)(x) dm_n(x).
\end{aligned} \tag{3.3}$$

Above we have used the routineous fact that for M -bi-Lipschitz mappings,

$$|J_f(f^{-1}(y))| \leq n!M^n, \tag{3.4}$$

and the already cited change of variable result of Radó-Reichelderfer; see [8, Theorem 2.2, page 99]. (Recall again that bi-Lipschitz mappings satisfy the property N and are differentiable almost everywhere.) \square

4. An Invariance of Regularly Oscillating Functions

Let Ω be an open set in \mathbb{R}^n , $n \geq 2$. Let $f : \Omega \rightarrow \mathbb{R}^m$ be continuous. Write

$$L(x, f) := \limsup_{y \rightarrow x} \frac{|f(y) - f(x)|}{|y - x|}. \tag{4.1}$$

The function $x \mapsto L(x, f)$ is a Borel function in Ω . If f is differentiable at x , then $L(x, f) = |f'(x)|$; see [9, page 11], [11, page 19], and [12, page 93].

A function $f : \Omega \rightarrow \mathbb{R}$ is *regularly oscillating*, if there is $K \geq 1$ such that

$$L(x, f) \leq Kr^{-1} \sup_{y \in B^n(x, r)} |f(y) - f(x)|, \quad \overline{B^n(x, r)} \subset \Omega. \tag{4.2}$$

The class of such functions is denoted by $\text{OC}_K^1(\Omega)$. The class of all regularly oscillating functions is denoted by $\text{RO}(\Omega)$; see [11, page 19], [13, page 17], [14], [6, page 245], and [12, page 96].

Using again Koebe's results, Kojić proved also the following result.

Theorem 4.1 (see [6, Theorem 2, page 245]). *Let Ω and G be open sets in \mathbb{C} . Let $u \in \text{OC}_K^1(\Omega)$. If $f : G \rightarrow \Omega$ is conformal, then $u \circ f \in \text{OC}_C^1(G)$, where C depends only on K .*

Below we give a partial generalization to Kojić's above result.

Theorem 4.2. *Let Ω and U be open sets in \mathbb{R}^n , $n \geq 2$. Let $u \in \text{OC}_K^1(\Omega)$. If $\varphi : U \rightarrow \Omega$ is M -bi-Lipschitz, $M \geq 1$, then $u \circ \varphi \in \text{OC}_{KM^2}^1(U)$.*

Proof. Let $\varphi : U \rightarrow \Omega$ be M -bi-Lipschitz. Take $x_0 \in U$ and $r_0 > 0$ arbitrarily such that $\overline{B^n(x_0, r_0)} \subset U$. Write $x'_0 = \varphi(x_0)$ and $x' = \varphi(x)$ for $x \in U$. Then

$$\begin{aligned}
 L(x_0, u \circ \varphi) &= \limsup_{x \rightarrow x_0} \frac{|u(\varphi(x)) - u(\varphi(x_0))|}{|x - x_0|} \\
 &= \limsup_{x \rightarrow x_0} \frac{|u(\varphi(x)) - u(\varphi(x_0))|}{|\varphi(x) - \varphi(x_0)|} \cdot \frac{|\varphi(x) - \varphi(x_0)|}{|x - x_0|} \\
 &\leq \limsup_{x' \rightarrow x'_0} \frac{|u(x') - u(x'_0)|}{|x' - x'_0|} \cdot \limsup_{x \rightarrow x_0} \frac{|\varphi(x) - \varphi(x_0)|}{|x - x_0|} \\
 &= L(x'_0, u) \cdot \limsup_{x \rightarrow x_0} \frac{|\varphi(x) - \varphi(x_0)|}{|x - x_0|}.
 \end{aligned} \tag{4.3}$$

Using (3.2) (for $f = \varphi$), we get

$$\begin{aligned}
 L(x'_0, u) &\leq \frac{K}{r_0/M} \sup_{x' \in B^n(x'_0, r_0/M)} |u(x') - u(x'_0)| \\
 &\leq \frac{KM}{r_0} \sup_{x' \in B^n(x'_0, r_0/M)} |u(x') - u(x'_0)| \\
 &\leq \frac{KM}{r_0} \sup_{x' \in \varphi(B^n(x_0, r_0))} |u(x') - u(x'_0)| \\
 &\leq \frac{KM}{r_0} \sup_{x \in B^n(x_0, r_0)} |u(\varphi(x)) - u(\varphi(x_0))| \\
 &\leq \frac{KM}{r_0} \sup_{x \in B^n(x_0, r_0)} |(u \circ \varphi)(x) - (u \circ \varphi)(x_0)|.
 \end{aligned} \tag{4.4}$$

On the other hand, since φ is M -bi-Lipschitz,

$$\limsup_{x \rightarrow x_0} \frac{|\varphi(x) - \varphi(x_0)|}{|x - x_0|} \leq \limsup_{x \rightarrow x_0} \frac{M|x - x_0|}{|x - x_0|} = M < +\infty. \tag{4.5}$$

Therefore,

$$\begin{aligned}
 L(x_0, u \circ \varphi) &\leq \frac{KM}{r_0} \sup_{x \in B^n(x_0, r_0)} |(u \circ \varphi)(x) - (u \circ \varphi)(x_0)| \cdot M \\
 &\leq \frac{KM^2}{r_0} \sup_{x \in B^n(x_0, r_0)} |(u \circ \varphi)(x) - (u \circ \varphi)(x_0)|.
 \end{aligned} \tag{4.6}$$

Thus $u \circ \varphi \in \text{OC}_{KM^2}^1(U)$. □

In addition of regularly oscillating functions, one sometimes considers so-called HC^1 functions, too; see [11, page 19], [13, page 16], and [12, page 93]. Their definition reads as follows. Let Ω be an open set in \mathbb{R}^n , $n \geq 2$. Let $K \geq 1$. A function $f : \Omega \rightarrow \mathbb{R}$ is in $HC_K^1(\Omega)$ if

$$L(x, f) \leq Kr^{-1} \sup_{y \in B^n(x, r)} |f(y)|, \quad \overline{B^n(x, r)} \subset \Omega. \quad (4.7)$$

The class $HC^1(\Omega)$ is the union of all $HC_K^1(\Omega)$, $K \geq 1$. Clearly, $HC_{2K}^1(\Omega) \subset OC_K^1(\Omega)$.

Proceeding as above in the proof of Theorem 4.2 one gets the following result.

Theorem 4.3. *Let Ω and U be open sets in \mathbb{R}^n , $n \geq 2$. Let $u \in HC_K^1(\Omega)$. If $\varphi : U \rightarrow \mathbb{R}^n$ is M -bi-Lipschitz, $M \geq 1$, then $u \circ \varphi \in HC_{KM^2}^1(U)$.*

Acknowledgment

The first author was partially supported by the Academy of Finland.

References

- [1] O. Dovgoshey and J. Riihenta, "A remark concerning generalized mean value inequalities for subharmonic functions," submitted.
- [2] O. Dovgoshey and J. Riihenta, "Mean value type inequalities for quasi-nearly subharmonic functions," manuscript.
- [3] M. Hervé, *Analytic and Plurisubharmonic Functions in Finite and Infinite Dimensional Spaces*, vol. 198 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1971.
- [4] D. H. Armitage and S. J. Gardiner, *Classical Potential Theory*, Springer, London, UK, 2001.
- [5] J. Riihenta, "Subharmonic functions, generalizations and separately subharmonic functions," in *Proceedings of the 14th Conference on Analytic Functions*, Chelm, Poland, July, 2007.
- [6] V. Kojić, "Quasi-nearly subharmonic functions and conformal mappings," *Filomat*, vol. 21, no. 2, pp. 243–249, 2007.
- [7] O. Djordjević and M. Pavlović, " L^p -integrability of the maximal function of a polyharmonic function," *Journal of Mathematical Analysis and Applications*, vol. 336, no. 1, pp. 411–417, 2007.
- [8] Yu. G. Reshetnyak, *Space Mappings with Bounded Distortion*, vol. 73 of *Translations of Mathematical Monographs*, American Mathematical Society, Providence, RI, USA, 1989.
- [9] J. Väisälä, *Lectures on n -Dimensional Quasiconformal Mappings*, vol. 229 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1971.
- [10] W. K. Hayman and P. B. Kennedy, *Subharmonic Functions. Vol. I*, London Mathematical Society Monographs, Academic Press, London, UK, 1976.
- [11] M. Pavlović, "On subharmonic behaviour and oscillation of functions on balls in \mathbb{R}^n ," *Institut Mathématique. Publications. Nouvelle Série*, vol. 55, no. 69, pp. 18–22, 1994.
- [12] M. Pavlović and J. Riihenta, "Classes of quasi-nearly subharmonic functions," *Potential Analysis*, vol. 29, no. 1, pp. 89–104, 2008.
- [13] M. Pavlović, "Subharmonic behaviour of smooth functions," *Matematički Vesnik*, vol. 48, no. 1-2, pp. 15–21, 1996.
- [14] M. Pavlović, *Introduction to Function Spaces on the Disk*, vol. 20 of *Posebna Izdanja*, Matematički Institut SANU, Belgrade, Serbia, 2004.