

ON HILBERT POLYNOMIAL OF CERTAIN DETERMINANTAL IDEALS

SHRINIVAS G. UDPIKAR

Department of Mathematics
S.P. College
Pune - 411030, India

(Received February 2, 1989 and in revised form July 10, 1989)

ABSTRACT. Let $X = (X_{ij})$ be an $m(1)$ by $m(2)$ matrix whose entries X_{ij} , $1 < i < m(1)$, $1 < j < m(2)$; are indeterminates over a field K . Let $K[X]$ be the polynomial ring in these $m(1)m(2)$ variables over K . A part of the second fundamental theorem of Invariant Theory says that the ideal $I[p+1]$ in $K[X]$, generated by $(p+1)$ by $(p+1)$ minors of X is prime. More generally in [1], Abhyankar defines an ideal $I[p,a]$ in $K[X]$, generated by different size minors of X and not only proves its primeness but also calculates the Hilbert function as well as the Hilbert polynomial of this ideal. The said Hilbert polynomial is completely determined by certain integer valued functions $F_D(m,p,a)$. In this paper we prove some important properties of these integer valued functions.

KEY WORDS AND PHRASES. Standard bi-tableaux, monomials in minors, Hilbert polynomial, arithmetic genus, irreducible variety, recursive formula.

1980 AMS SUBJECT CLASSIFICATION CODE. 05A15, 13B25.

1. INTRODUCTION.

We assume all notations and conventions of Abhyankar [1]. Let Z, N, N^* and Q denote the set of all integers, nonnegative integers, positive integers and rational numbers respectively.

For any two integers A and B , we let $[A,B] = \{D \in Z: A < D < B\}$.

For any $p \in N$, we denote by $Z(p)$, $N(p)$, and $N^*(p)$ the set of all maps from $[1,p]$ to Z , N and N^* respectively.

For any $p \in N^*$ and $D \in Z$, we let $Z(p,D) = \{d \in Z(p): d(1) + d(2) + \dots + d(p) = D\}$ and $N(p,D) = \{d \in N(p): d(1) + d(2) + \dots + d(p) = D\}$.

For any $p \in N$, we let $\text{rec}(2,p) = \{(k,i): k \in [1,2] \text{ and } i \in [1,p]\}$.

For any $p \in N$, by a bi-vector of length p , we mean a map $a: \text{rec}(2,p) \rightarrow N^*$, which for every $k \in [1,2]$ and $i \in [1,p]$, associates $a(k,i) \in N^*$ such that $a(k,1) < a(k,2) < \dots < a(k,p)$. In this case we put $\text{len}(a) = p$.

By a bi-vector, we mean a bi-vector of length p for some $p \in N$.

For any $p \in N$, $m \in N^*(2)$ and a bi-vector a of length p , we say that a is bounded by m and denote this by $a < m$, to mean that

$a(k,i) < m(k)$ for $k \in [1,2]$ and $i \in [1,p]$.

Given any two bi-vectors a and a' of lengths p and p' respectively, we define $a < a'$ to mean that $p > p'$ and an $a(k,i) < a'(k,i)$ for $k \in [1,2]$ and $i \in [1,p']$. We note that this defines a partial order on the set $\text{vec}(2)$ of all bi-vectors.

Given any $d \in \mathbb{N}$, by a bi-tableau of depth d , we mean a mapping

$T: [1,d] \rightarrow \text{vec}(2)$, which to every $e \in [1,d]$, associates $T[e] \in \text{vec}(2)$.

A bi-tableau T of depth d is said to be standard if $\text{len}(T[e])$ is positive for each $e \in [1,d]$ and

$$T[1] < T[2] < \dots < T[d].$$

Given any $m \in \mathbb{N}^*(2)$, $a \in \text{vec}(2)$ and a bi-tableau T of depth d , we say that

- (i) T is bounded by m if $T[e] < m$ for $e = 1, 2, \dots, d$
- (ii) T is predominated by a if $a < T[e]$ for $e = 1, 2, \dots, d$.

The area of a bi-tableau T of depth d is denoted by $\text{are}(T)$ and is defined as

$$\begin{aligned} \text{are}(T) &= \sum_{e=1}^d \text{len}(T[e]) \quad \text{if } d \neq 0 \text{ and} \\ &= 0 \quad \text{if } d = 0. \end{aligned}$$

Finally for any element V in an overring of \mathbb{Q} and any integer r , we set

$$\binom{V}{r} = \begin{cases} \frac{V(V-1) \dots (V-r+1)}{r!} & \text{if } r > 0 \\ 0 & \text{if } r < 0 \end{cases}$$

and
$$\begin{bmatrix} V \\ r \end{bmatrix} = \begin{cases} \binom{V+r}{r} & \text{if } r > 0 \\ 0 & \text{if } r < 0. \end{cases}$$

Let $X = (X_{ij})$ be any $m(1)$ by $m(2)$ matrix whose entries X_{ij} , $1 \leq i \leq m(1)$, $1 \leq j \leq m(2)$, are indeterminates over a field K . Let $K[X]$ be the polynomial ring in these $m(1)m(2)$ variables. Clearly to every bi-vector a of length p and bounded by m , there corresponds a uniquely determined $p \times p$ minor of X , say $\text{mor}(X,a)$, formed by the rows with row numbers $a(1,1), a(1,2), \dots, a(1,p)$ and by the columns with column numbers $a(2,1), a(2,2), \dots, a(2,p)$. By convention we define $\text{mor}(X,a) = 1$ if $\text{len}(a) = 0$.

Further, to a bi-tableau T of depth d , there corresponds a product of minors of X , which we call a monomial in minors of X . A monomial in minors of X is said to be standard if the corresponding bi-tableau is standard.

Using straightening formula, DeConcini, Eisenbud, and Procesi [2], proved that the standard monomials in minors of X form a base for the vector space $K[x]$ over K . Abhyankar proves this by enumerating the number of standard bi-tableaux with certain conditions. In fact, Abhyankar [3] proves that the number of standard bi-tableaux of area V , bounded by m and predominated by a given fixed bi-vector a of length p is equal to

$$H(V) = \sum_{D \in \mathbb{Z}} (-1)^D F_D(m,p,a) \begin{bmatrix} V \\ C-D \end{bmatrix}$$

where $C = C[m, p, a] = \sum_{i=1}^p (m(1) - a(1, i)) + \sum_{i=1}^p (m(2) - a(2, i)) + p - 1$

and $F_D(m, p, a)$, $D \in \mathbb{Z}$, are integer valued functions defined in section 2.

He further proves that if a is a given bi-vector of length p , bounded by m and if $I[p, a]$ is the ideal in $K[X]$ generated by all minors of X corresponding to the bi-vectors b not predominated by a , then $I[p, a]$ is a homogeneous prime ideal in $K[X]$ and $H(V)$ is the Hilbert function as well as the Hilbert polynomial of $I[p, a]$ in $K[X]$. In particular, if $a(k, i) = i$ for $k \in [1, 2]$ and $i \in [1, p]$, then $I[p, a]$ is the ideal in $K[X]$ generated by $(p+1)$ by $(p+1)$ minors of X and hence it follows by the above theorem that it is a prime ideal in $K[X]$. This forms a part of the second fundamental theorem of invariant theory and was originally proved by Pascal in 1888 and then reproved by Mount [4], Eagon and Hochster [5] and others.

Now $H(V)$ is a polynomial in V with rational coefficients and of degree C . Further $C!$ times the coefficient of V^C in $H(V)$ is $F_0(m, p, a)$, which equals the order of the irreducible variety defined by $I[p, a]$ in the $(m(1) m(2)-1)$ dimensional projective space over K (Refer to remark (20.18) of [1]). From this it follows that $F_0(m, p, a)$ is a positive integer (see theorem (2.1) of section 2). Further arithmetic genus of $I[p, a]$ is $(-1)^C [(F_0 - F_1 + F_2 - \dots + (-1)^C F_C) - 1] = 0$. Thus $F_D(m, p, a)$, $D \in \mathbb{Z}$ determine important geometric characters of the variety defined by $I[p, a]$ from Zariski and Samuel [6]. Hence it is interesting to study the properties of $F_D(m, p, a)$, $D \in \mathbb{Z}$.

2. INTEGER VALUED FUNCTIONS $F_D(m, p, a)$.

For any $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$, we put $\text{vec}(2) =$ the set of all bi-vectors, $\text{vec}[2, p] =$ the set of all bi-vectors of length p and $\text{vec}(2, m, p) = \{a \in \text{vec}[2, p] : a \leq m\}$.

Let $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$, $a \in \text{vec}(2, m, p)$ and $k \in [1, 2]$. Let $k' = 2$ if $k = 1$ and $k' = 1$ if $k = 2$.

Let $C[m, p, a, k] = \sum_{i=1}^p (m(k) - a(k, i))$,
 $C[m, p, a, k'] = \sum_{i=1}^p (m(k') - a(k', i))$, and
 $C = C[m, p, a] = C[m, p, a, k] + C[m, p, a, k'] + p - 1$.

We now define integer valued functions $F_D(m, p, a)$, $D \in \mathbb{Z}$.

For every $i, j \in [1, p]$, $e \in \mathbb{Z}(p)$ we denote by $G(m, p, a, e)$, the $p \times p$ matrix

whose $(i, j)^{\text{th}}$ entry is $G_{ij}(m, p, a, e) = \begin{pmatrix} m(k) - a(k, i) + i - j & & \\ & \dots & \\ m(k) - a(k, i) - e_i & & \end{pmatrix} \begin{pmatrix} m(k') - a(k', j) + j - i \\ & \dots & \\ & & e_i \end{pmatrix}$

where $e_i = e(i)$ for $i = 1, 2, \dots, p$. Further let for every $E \in \mathbb{Z}$, $H_E(m, p, a) = \sum_{e \in \mathbb{Z}(p, E)} \det G(m, p, a, e)$. Then for every $D \in \mathbb{Z}$, we define

$F_D(m, p, a) = \sum_{E \in \mathbb{Z}} \binom{E}{D} H_E(m, p, a)$.

In view of the usual properties of binomial coefficients and determinants, we may assume that $0 < e_i < (m(k) - a(k, i))$ for $i = 1, 2, \dots, p$. Hence the above sum is

essentially finite.

In [1], Abhyankar proves the following:

THEOREM 2.1. Let there be given any $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$ and $a \in \text{vec}(2, m, p)$. Then we have

- (i) $F_0(m, p, a)$ is a positive integer,
- (ii) for all $D \in \mathbb{Z}$, $F_D(m, p, a)$ is a nonnegative integer and
- (iii) $\{D \in \mathbb{Z} : F_D(m, p, a) \neq 0\} \subseteq [0, C(m, p, a)]$ where
 $C(m, p, a) = \min\{C[m, p, a, k], C[m, p, a, k']\}$.

PROOF. Refer to Lemma 9.15 and Theorems 9.13, 9.17 of [1].

He further proposes the following problems. Given any $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$ and $a \in \text{vec}(2, m, p)$, does there exist a nonnegative integer $C^*(m, p, a)$ such that
 Problem (I) $\{D \in \mathbb{Z} : F_D(m, p, a) \neq 0\} = [0, C^*(m, p, a)]$?
 Problem (II) If so, then is it possible that

$$C^*(m, p, a) = C(m, p, a)?$$

(Refer to remark 9.18 of [1]).

This paper settles both these problems in an affirmative manner.

3. THE MAIN RESULT.

We follow the usual convention that the sum over an empty family is zero. For any $p \in \mathbb{N}$ and $A \in \mathbb{Z}$, we let $J(p) =$ all subsets of $[1, p]$ and $J(p, A) =$ all subsets of $[1, p]$ with cardinality A .

Let there be given any $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$, $b \in \text{vec}[2, p+1]$, $u \in J(p)$, $U \in [0, p]$ and $k \in [1, 2]$. Further let

$$k' = \begin{cases} 2 & \text{if } k = 1 \\ 1 & \text{if } k = 2. \end{cases}$$

We define the sets $M[p, b, k, u]$ and $M(p, b, k, U)$ as follows:

$$M[p, b, k, u] = \left\{ \begin{array}{l} a \in \text{vec}[2, p] : a(k, i^*) = b(k, i^*) \text{ for all } i^* \in [1, p], \\ a(k', i) \in [b(k', i)+1, b(k', i+1)-1], \text{ for all } i \in u \\ \text{and } a(k', j) = b(k', j), \text{ for all } j \in [1, p] \setminus u \end{array} \right\}$$

and $M(p, b, k, U) = \cup M[p, b, k, u]$, $u \in J(p, U)$

Given any $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$, $b \in \text{vec}[2, p+1]$ and $k \in [1, 2]$, let $m^* \in \mathbb{N}^*(2)$ be such that

- (1) $m^*(k') = m(k')$ and $m^*(k) = m(k) + 1$ holds. Further let $p^* \in \mathbb{N}^*$ and $a^* \in \text{vec}(2, m^*, p^*)$ be such that either
- (2a) $p^* = p$ and $m^*(k) - a^*(k, p^*) \neq 0$ and $a^*(\hat{k}, i) = b(\hat{k}, i)$ for all $\hat{k} \in [1, 2]$ and $i \in [1, p]$ and $b(\hat{k}, p+1) = m(\hat{k}) + 1$ for all $\hat{k} \in [1, 2]$ holds or
- (2b) $p^* = p+1$, $m^*(k) - a^*(k, p^*) = 0$ and $a^* = b$ holds.

Finally for every $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$, $a \in \text{vec}[2, p]$, $U \in [0, p]$ and $D \in \mathbb{Z}$, we

set

$$S_U(m, p, a, D) = \sum_{I \in [0, U]} \binom{U}{I} F_{D-I}(m, p, a).$$

We now state a useful recursive formula in the following.

THEOREM 3.1. Let there be given any $m \in N^*(2)$, $p \in N^*$, $k \in [1, 2]$, $b \in \text{vec}[2, p+1]$ and $m^* \in N^*(2)$, $p^* \in N^*$, $a^* \in \text{vec}(2, m^*, p^*)$ such that (1) holds and either (2a) or (2b) holds. Then for every $D \in Z$, we have

$$F_D(m^*, p^*, a^*) = \sum_{I \in [0, p]} \sum_{U \in [0, p]} \sum_{a \in M(p, b, k, U)} \binom{U}{I} F_{D-I}(m, p, a).$$

PROOF. Refer to Theorem (9.8*), section 9 of [1].

In the above recursive formula, by interchanging the summations over I and U and by noting the fact that $\binom{U}{I} = 0$ if $I > U > 0$, we see that

$$F_D(m^*, p^*, a^*) = \sum_{U \in [0, p]} \sum_{a \in M(p, b, k, U)} S_U(m, p, a; D)$$

We shall use this form of the recursive formula later in this section.

In what follows, we shall prove some results which enable one to answer the first problem posed by Abhyankar, in an affirmative manner [Refer to problem (I) at the end of section 2].

THEOREM 3.2. Let $k \in [1, 2]$, $p \in N^*$ and $b \in \text{vec}[2, p+1]$ be given. Then we have the following.

(I) $M(p, b, k, 0) = M[p, b, k, \phi]$ is a nonempty set consisting of the unique element $\tilde{b} \in \text{vec}[2, p]$ obtained from b by putting

$$\tilde{b}(\hat{k}, i) = b(\hat{k}, i) \text{ for all } \hat{k} \in [1, 2] \text{ and } i \in [1, p].$$

(II) If $M(p, b, k, 1) = \phi$ then $\{b(k', j)\}$, $j = 1, 2, \dots, p$, is a sequence of consecutive integers i.e. $b(k', j+1) = b(k', j) + 1$, for all $j \in [1, p-1]$ and consequently $M(p, b, k, U) = \phi$ for every $U > 1$.

PROOF. (I) Since ϕ is the only subset of $[1, p]$ of cardinality zero, we have $M(p, b, k, 0) = M[p, b, k, \phi]$. The rest follows from the definition of $M[p, b, k, \phi]$. (II) Let $u_1 = \{i\}$ where i is an arbitrarily chosen element of $[1, p-1]$. Then by definition of $\mu[p, b, k_1, u_1]$ we have,

$$\begin{aligned} M[p, b, k, u_1] = \phi &\iff [b(k', i) + 1, b(k', i + 1) - 1] = \phi \\ &\iff b(k', i+1) = b(k', i) + 1, \text{ so that } \{b(k', j)\}, \end{aligned}$$

$j=1, 2, \dots, p$, is an increasing sequence of consecutive integers.

Finally, if $U > 1$ and $u \in J(p)$ is any set of cardinality U , then there is an $i \in u$ such that $i \in [1, p-1]$. Now if $a \in M(p, b, k, U)$, then we must have $a(k', i) \in [b(k', i) + 1, b(k', i + 1) - 1]$, which is a contradiction since $\{b(k', j)\}$, $j = 1, 2, \dots, p$, is a sequence of consecutive integers. Hence $M(p, b, k, U) = \phi$ for every $U > 1$ if $M(p, b, k, 1) = \phi$. Hence the theorem.

LEMMA 3.1. Let $m \in N^*(2)$, $p \in N^*$, $b \in \text{vec}[2, p+1]$ and $k \in [1, 2]$ be given. Let $m^* \in N^*(2)$, $p^* \in N^*$ and $a^* \in \text{vec}(2, m^*, p^*)$ be such that (1) is satisfied and either (2a) or (2b) is satisfied. Let \tilde{b} be the unique element of $\text{vec}[2, p]$ obtained from b by putting $\tilde{b}(\hat{k}, i) = b(\hat{k}, i)$ for all $\hat{k} \in [1, 2]$ and $i \in [1, p]$.

Then we have the following:

$$F_1(m^*, p^*, a^*) = 0 \Rightarrow F_1(m, p, \tilde{b}) = 0 \text{ and}$$

$$F_{1+j}(m^*, p^*, a^*) = F_{1+j}(m, p, \tilde{b}) \text{ for every } j \in \mathbb{N}.$$

PROOF. In view of theorem (3.1) and (I) of theorem (3.2), we have

$$F_1(m^*, p^*, a^*) = F_1(m, p, \tilde{b}) + \sum_{a \in M(p, b, k, l)} [F_1(m, p, a) + F_0(m, p, a)] \\ + \sum_{U \in [2, p]} \sum_{a \in M(p, b, k, U)} S_U(m, p, a; l).$$

But by theorem (2.1) of section 2, we have $F_0(m, p, a) > 0$ and $F_D(m, p, a) > 0$ for every $D \in \mathbb{Z}$. Hence $F_1(m^*, p^*, a^*) = 0 \Rightarrow F_1(m, p, \tilde{b}) = 0$ and $M(p, b, k, l) = \phi$. But then by (II) of theorem (3.2), we have $M(p, b, k, U) = \phi$ for every $U > 2$. Hence given any $j \in \mathbb{N}$, replacing l by $l + j$ in the above formula we get

$$F_{1+j}(m^*, p^*, a^*) = F_{1+j}(m, p, \tilde{b}). \text{ Hence the lemma.}$$

LEMMA 3.2. Let $D \in \mathbb{N}^*$. Assume that for all $m \in N^*(2)$, $p \in \mathbb{N}^*$, $b \in \text{vec}[2, p+1]$, $k \in [1, 2]$ and for all $m^* \in N^*(2)$, $p^* \in \mathbb{N}^*$ and $a^* \in \text{vec}(2, m^*, p^*)$ satisfying (1) and satisfying (2a) or (2b), we have

$$F_D(m^*, p^*, a^*) = 0 \Rightarrow (i) F_D(m, p, \tilde{b}) = 0 \text{ and}$$

$$(ii) F_{D+j}(m^*, p^*, a^*) = F_{D+j}(m, p, \tilde{b}) \text{ for all } j \in \mathbb{N}.$$

Then for every $a \in \text{vec}(2, m, p)$, we have

$$F_D(m, p, a) = 0 \Rightarrow F_{D+j}(m, p, a) = 0 \text{ for every } j \in \mathbb{N}.$$

PROOF. We prove this by double induction on p and $r(p) = m(k) - a(k, p)$. Suppose that for $a^* \in \text{vec}(2, m^*, p^*)$, $F_D(m^*, p^*, a^*) = 0$. Hence by assumption, (i) and (ii) hold.

If $r^*(p^*) = m^*(k) - a^*(k, p^*) \neq 0$, then $p = p^*$ so that (2a) holds and $r(p) < r^*(p^*)$. Hence by (i) and inductive hypothesis, we have $F_{D+j}(m, p, \tilde{b}) = 0$, for all $j \in \mathbb{N}$.

Hence by (ii) $F_{D+j}(m^*, p^*, a^*) = 0$ for all $j \in \mathbb{N}$. On the other hand if $r^*(p^*) = 0$ and $p^* = p+1$, then (2b) holds and as above, we get

$$F_{D+j}(m^*, p^*, a^*) = 0, \text{ for all } j \in \mathbb{N}.$$

Thus to complete the proof, we only need to prove the result when $p = 1$ and $r(p) = m(k) - a(k, p) = 0$. But in this case $C(m, p, a) = 0$ and hence by theorem (2.1), we see that $F_{D+j}(m, p, a) = 0$, for all $D \in \mathbb{N}^*$ and $j \in \mathbb{N}$. Hence the lemma.

Now we are in a position to prove the main result of this section.

THEOREM 3.3. For every $D, j \in \mathbb{N}^*$, we have the following:

$$F_D(m, p, a) = 0 \Rightarrow F_{D+j}(m, p, a) = 0, \text{ for every triplet } (m, p, a) \text{ such that } m \in N^*(2), \\ p \in \mathbb{N}^* \text{ and } a \in \text{vec}(2, m, p).$$

PROOF. By induction on D . In view of lemma 3.1 and lemma 3.2, the theorem is clearly true if $D = 1$. Now assume that D is any integer greater than 1 and the theorem holds for every $D' \in \mathbb{N}^*$ such that $D' < D$. To prove the theorem for this fixed D , we shall use lemma 3.2.

So let there be given $m \in N^*(2)$, $p \in \mathbb{N}^*$, $b \in \text{vec}[2, p+1]$, $k \in [1, 2]$ and $m^* \in N^*(2)$, $p^* \in \mathbb{N}^*$, $a^* \in \text{vec}(2, m^*, p^*)$ such that (1) is satisfied and (2a) or (2b) is

satisfied.

Assume that $F_D(m^*, p^*, a^*) = 0$. By theorem 3.1, we have

$$F_D(m^*, p^*, a^*) = F_D(m, p, \tilde{b}) + \sum_{U \in [1, p]} \sum_{a \in M(p, b, k, U)} S_U(m, p, a; D) = 0. \tag{3.1}$$

Note that $F_D(m, p, a) > 0$ for every $D \in Z$. Hence, if for some $U \in [1, p]$, $M(p, b, k, U)$ is nonempty, then by (3.1), we must necessarily have $S_U(m, p, a; D) = 0$.

Hence by definition of $S_U(m, p, a; D)$ and the fact that $\binom{U}{1} > 0$ if $1 \in [0, U]$, we have

$F_{D-1}(m, p, a) = 0$ for every $1 \in [0, U]$. In particular, $F_{D-U}(m, p, a) = 0$. But $U > 1$. Hence by inductive hypothesis, we get $F_{D+j-U}(m, p, a) = 0$ for every $j \in N$. Hence, given any $j \in N$, by using the above identity as j is successively replaced by $j + U - 1 \in N$, as 1 varies from 0 to U , we get $S_U(m, p, a; D + j) = 0$. Thus for $U > 1$, in view of (3.1), we have either $M(p, b, k, U) = \emptyset$ or $S_U(m, p, a; D + j) = 0$, for all $j \in N$.

Hence by recursive formula, we get $F_{D+j}(m^*, p^*, a^*) = F_{D+j}(m, p, \tilde{b})$ for all $j \in N$. But by (3.1), we certainly have $F_D(m, p, \tilde{b}) = 0$.

Hence by Lemma 3.2, we get the required result.

From this theorem, it is clear that, given any $m \in N^*(2)$, $p \in N^*$ and $a \in \text{vec}(2, m, p)$, the set $\{D \in Z : F_D(m, p, a) \neq 0\}$ is an interval of the form $[0, C^*(m, p, a)]$ for some nonnegative integer $C^*(m, p, a)$.

Thus, Abhyankar's conjecture is true.

4. SOLUTION TO SECOND PROBLEM.

From section 2, we know that for every $D \in Z$.

$$F_D(m, p, a) = \sum_{E \in Z} \binom{E}{D} \sum_{e \in Z(p, E)} \det \left\{ \begin{pmatrix} m(k) - a(k, i) + i - j \\ \vdots \\ m(k) - a(k, i) - e_i \end{pmatrix} \begin{pmatrix} m(k') - a(k', j) + j - i \\ \vdots \\ e_i \end{pmatrix} \right\}$$

where $m \in N^*(2)$, $p \in N^*$, $a \in \text{vec}(2, m, p)$, $k \in [1, 2]$ and

$$k' = \begin{cases} 2 & \text{if } k = 1 \\ 1 & \text{if } k = 2. \end{cases}$$

Noting that $0 < e_i < (m(k) - a(k, i))$ for $i = 1, 2, \dots, p$; we see that

$$F_D(m, p, a) = \det \left\{ \begin{pmatrix} m(k') - a(k', j) + j - i \\ \vdots \\ m(k) - a(k, i) \end{pmatrix} \right\}$$

if $D = C[m, p, a, k] = \sum_{i=1}^p (m(k) - a(k, i))$.

Hence in particular, $F_D(m, p, a) = 1$, if no two of $(m(k') - a(k', j))$, $j = 1, 2, \dots, p$ are consecutive and $(m(k') - a(k', j)) = (m(k) - a(k, j))$ for $j = 1, 2, \dots, p$; where $D = C[m, p, a, k]$.

Hence by the main theorem of section 3, we see that $F_D(m, p, a) > 0$ for every $D \in [0, C(m, p, a)]$ where $C(m, p, a) = \min\{C[m, p, a, k], C[m, p, a, k']\}$.

Hence there exist $m \in \mathbb{N}^*(2)$, $p \in \mathbb{N}^*$ and $a \in \text{vec}(2, m, p)$ such that $F_D(m, p, a) > 0$ for every $D \in [0, C(m, p, a)]$.

Affirmative answers to problems (I) and (II) enable us to write the Hilbert Polynomial of $I[p, a]$ in $K[X]$ in much tidier form as follows:

$$H(V) = \sum_{D=0}^{C^*(m, p, a)} (-1)^D F_D(m, p, a) \begin{bmatrix} V \\ C-D \end{bmatrix}$$

where $C^*(m, p, a)$ is some integer such that $0 < C^*(m, p, a) < C(m, p, a)$ and $F_D(m, p, a) > 0$ for every $D \in [0, C^*(m, p, a)]$.

ACKNOWLEDGEMENTS. I wish to thank Professor Shreeram S. Abhyankar for his valuable guidance while working with him for my Ph.D Degree. I wish also to thank Panjab University, Chandigarh for the award of the U.G.C. fellowship and Bhaskaracharya Pratishthana, Pune, for providing necessary facilities.

REFERENCES

1. ABHYANKAR, S., Enumerative Combinatorics of Young Tableaux, Marcel Dekker, Inc. 1988.
2. DECONCINI, EISENBUD, PROCESI, Yong Diagrams and Determinantal Varieties; Inventiones Mathematicae 56, (1980), 129-165
3. ABHYANKAR, S., Combinatorie des tableaux de Young, varietes determinantielles et calcul de fonctions de Hilbert; Nice lecture notes by A. Galligo; Universita E. Politecnico di Torino: Rendiconti del Seminario Matematico 1985.
4. MOUNT, K.R., A Remark on Determinantal Loci; Journal London Math. Soc. 42, (1967), 595-598.
5. EAGON, J. and HOCHSTER, M., Cohen Macaulay Rings, Invariant Theory and the Generic Perfection of Determinantal Loci; American J. Math. 93, (1971), 1020-1058.
6. ZARISKI, O and SAMUEL, P., Commutative Algebra, Volume II, Van Nostrand Princeton 1969.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

| | |
|------------------------|-----------------|
| Manuscript Due | July 1, 2009 |
| First Round of Reviews | October 1, 2009 |
| Publication Date | January 1, 2010 |

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br