

ON THE ARENS PRODUCTS AND REFLEXIVE BANACH ALGEBRAS

PAK-KEN WONG

Department of Mathematics
Seton Hall University
South Orange, New Jersey 07079

(Received March 29, 1990)

ABSTRACT. We give a characterization of reflexive Banach algebras involving the Arens product.

KEY WORDS AND PHRASES. Arens products, Arens regularity, conjugate space, weakly completely continuous (w.c.c.) algebra.

1980 AMS SUBJECT CLASSIFICATION CODES. Primary 46H10; Secondary 46H99.

1. INTRODUCTION.

Let A be a semisimple Banach algebra and A^{**} the second conjugate space of A with the Arens product \circ . If (A^{**}, \circ) is semisimple and it has a dense socle, then we show that the following statements are equivalent: (1) A is reflexive. (2) A^{**} is w.c.c. (3) A is w.c.c. (4) A and A^{**} have the same socle. This is a generalization of a result by Duncan and Hosseini [1, p.319, Theorem 6(ii)]. We also show that if (A^{**}, \circ) is semisimple and A is l.w.c.c., then A is Arens regular.

2. NOTATION AND PRELIMINARIES. Definitions not explicitly given are taken from Rickart's book [2].

Let A be a Banach algebra. Then A^* and A^{**} will denote the first and second conjugate spaces of A , and π the canonical map of A into A^{**} . The two Arens products on A^{**} are defined in stages according to the following rules (see [3] and [4]). Let $x, y \in A, f \in A^*$, and $F, G \in A^{**}$.

Define fox by $(\text{fox})(y) = f(xy)$. Then $\text{fox} \in A^{**}$.

Define Gof by $(\text{Gof})(x) = G(\text{fox})$. Then $\text{Gof} \in A^*$.

Define FoG by $(\text{FoG})(f) = F(\text{Gof})$. Then $\text{FoG} \in A^{**}$.

Define xo'f by $(\text{xo'f})(y) = f(yx)$. Then $\text{xo'f} \in A^*$.

Define fo'F by $(\text{fo'F})(x) = F(\text{xo'f})$. Then $\text{fo'F} \in A^*$.

Define Fo'G by $(\text{Fo'G})(f) = G(\text{fo'F})$. Then $\text{Fo'G} \in A^{**}$.

A^{**} is a Banach algebra under the products FoG and Fo'G and π is an algebra isomorphism of A into (A^{**}, \circ) and (A^{**}, \circ') . In general, \circ and \circ' are distinct on A^{**} . If they agree on A^{**} , then A is called Arens regular.

LEMMA 2.1. Let A be a Banach algebra. Then, for all $x \in A, f \in A^*$, and $F, G \in A^{**}$, we have

- (1) $\pi(x) \circ F = \pi(x) \circ' F$ and $F \circ \pi(x) = F \circ' \pi(x)$.
- (2) If $\{F_i\} \subset A^{**}$ and $F_i \rightarrow F$ weakly in A^{**} , then $F_i \circ G \rightarrow F \circ G$ and $G \circ' F_i \rightarrow G \circ' F$ weakly.

PROOF. See [3, p.842 and p. 843].

Let A be a Banach algebra. An element $a \in A$ is called left weakly completely continuous (1.w.c.c.) if the mapping L_a defined by $L_a(x) = ax (x \in A)$ is weakly completely continuous. We say that A is 1.w.c.c. if each $a \in A$ is 1.w.c.c. If A is both 1.w.c.c. and r.w.c.c., then A is called w.c.c.

In this paper, all algebras and linear spaces under consideration are over the field C of complex numbers.

3. THE MAIN RESULT.

LEMMA 3.1. Let A be a Banach algebra. Then A is 1.w.c.c. (resp. r.w.c.c.) if and only $\pi(A)$ is a right (resp. left) ideal of (A^{**}, \circ) .

PROOF. This result is well known (see [1, p.318, Lemma 3] or [2, p.443, Lemma]).

In the rest of this section, we shall assume that A and (A^{**}, \circ) are semisimple Banach algebras.

THEOREM 3.3. Suppose that (A^{**}, \circ) has a dense socle. Then the following statements are equivalent:

- (1) A is reflexive.
- (2) A^{**} is w.c.c.
- (3) A is w.c.c.
- (4) $\pi(A)$ and A^{**} have the same socle.

PROOF.

(1) \Rightarrow (2). Assume that A is reflexive. Then $A^{(4)} = A^{**} = A$; in particular, $\pi(A)^{**}$ is a two-sided ideal of $A^{(4)}$. Hence by Lemma 3.1, A^{**} is w.c.c.

(2) \Rightarrow (3). Assume that A^{**} is w.c.c. Then $\pi(A^{**})$ is a two-sided ideal of $A^{(4)}$. As observed in [1, p.319, Theorem 6(ii)], $\pi(A)$ is a two-sided ideal of A^{**} . Hence A is w.c.c.

(3) \Rightarrow (4). Assume that A is w.c.c. Then $\pi(A)$ is a two-sided ideal of A^{**} . Let E be a minimal idempotent of A^{**} . Since $E \circ A^{**} \circ E = E \circ \pi(A) \circ E = CE$, it follows that $E \in \pi(A)$. Consequently, E is a minimal idempotent of $\pi(A)$. If e is a minimal idempotent of A , then $\pi(e) \circ A^{**} \subset \pi(A)$ and so $\pi(e) \circ A^{**} = \pi(eA)$. Hence, $\pi(e) \circ A^{**} \circ \pi(e) = \pi(eAe) = C\pi(e)$ and so $\pi(e)$ is a minimal idempotent of A^{**} . Therefore, $\pi(A)$ and A^{**} have the same socle.

(4) \Rightarrow (1). Assume that $\pi(A)$ and A^{**} have the same socle. Since $\pi(S)$ is dense in A^{**} , it follows that $\pi(A)$ is dense in A^{**} and so $\pi^*(A) = A^{**}$. Therefore A is reflexive. This completes the proof of the theorem.

REMARK. It is well known that a semisimple annihilator Banach algebra A is w.c.c. (see [5]). Also, A has a dense socle. Therefore, Theorem 3.2 generalizes [1, p.319, Theorem 6(ii)].

THEOREM 3.3. If A is 1.w.c.c., then A is Arens regular.

PROOF. Since A is 1.w.c.c., by Lemma 3.1, $\pi(A)$ is a right ideal of A^{**} . Let F and $G \in A^{**}$ and $x \in A$. Then

$$\begin{aligned} \pi(x) \circ (F \circ G - F \circ' G) &= \pi(x) \circ F \circ G - \pi(x) \circ (F \circ' G) \\ &= \pi(x) \circ F \circ G - \pi(x) \circ' (f \circ' G) \quad \text{By Lemma 2.1(1)} \end{aligned}$$

$$\begin{aligned}
&= \pi(x) \circ F \circ G - (\pi(x) \circ F) \circ G \\
&= \pi(x) \circ F \circ G - (\pi(x) \circ F) \circ G \quad (\text{because } \pi(x) \circ F \in \pi(A)) \\
&= 0
\end{aligned}$$

Hence $\pi(A) \circ (F \circ G - F \circ G) = (0)$. Therefore, by Lemma 2.1 (2), we have $A^{**} \circ (F \circ G - F \circ G) = (0)$. Since (A^{**}, \circ) is semisimple, it follows that $F \circ G - F \circ G = 0$ and so $F \circ G = F \circ G$. Therefore, A is Arens regular. This completes the proof.

REFERENCES

1. DUNCAN, J. and HOSSEINIUM, S.A.R. The Second Dual of a Banach Algebra, Proc. Royal Soc. of Edinburgh, **84A** (1979), 309-325.
2. WONG, P.K. Arens Product and the Algebra of Double Multipliers, Proc. Amer. Math. Soc. **94** (1985), 441-444.
3. ARENS, R.F. The Adjoint of a Bilinear Operation, Proc. Amer. Math. Soc. **2** (1951), 839-848.
4. BONSALL, F.F. and DUNCAN, J. Complete Normed Algebras, Springer, Berlin, 1973.
5. WONG, P.K. On the Arens Product and Certain Banach Algebras, Trans. Amer. Math. Soc. **180** (1973), 437-438.
6. RICKART, C.E. General Theory of Banach Algebras, University Series in Higher Math., Van Nostrand, Princeton, N.J., 1960.
7. WONG, P.K. Arens Product and the Algebra of Double Multipliers II, Proc. Amer. Soc. **100** (1987), 447-453.

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.otero@usc.es