

A REMARK ON THE WEIGHTED AVERAGES FOR SUPERADDITIVE PROCESSES

DOGAN CÖMEZ

Department of Mathematics
300 Minard Hall, P.O. BOX 5075
North Dakota State University
Fargo, ND 58105

(Received November 24, 1987 and in revised form August 22, 1988)

ABSTRACT. A decomposition of a superadditive process into a difference of an additive and a positive purely superadditive process is obtained. This result is used to prove an ergodic theorem for weighted averages of superadditive processes.

KEY WORDS AND PHRASES. Ergodic theorem, bounded sequences, superadditive process, additive process.

1980 AMS SUBJECT CLASSIFICATION CODES. Primary 47A35, Secondary 28D99, 60G10.

1. **INTRODUCTION.** Let (X, μ) be a probability space and let $L_p(X, \mu)$, $1 < p < \infty$ be the classical Banach space of real valued functions f with $\int |f|^p d\mu = \|f\|_p^p < \infty$.

Let $T: L_p \rightarrow L_p$ be a linear operator. A family of L_p functions $F = \{F_n\}_{n \geq 1}$ is called a

T -superadditive process if

$$F_{n+m} \geq F_n + T^n F_m, \text{ for all } n, m \geq 1, \quad (1.1)$$

and F is called a T -additive process if equality holds in (1.1). Notice that if we

let $f_i = F_{i+1} - F_i$, for $i \geq 0$ we have $F_n = \sum_{i=0}^{n-1} f_i$, where $F_0 \equiv 0$, for all $n \geq 1$. Consider

a sequence $A = \{a_n\}_{n \geq 0}$ of complex numbers and a T -(super) additive process F . We

define a family of L_p -functions $(F, A) = \{a_n f_n\}_{n \geq 0}$, and set $S_n(F, A) = \sum_{i=0}^{n-1} a_i f_i$. If A is

the constant sequence $l=(1,1,\dots)$, then $S_n(F,A) = F_n$.

In the following, we observe that the weighted and subsequential ergodic theorems for T-superadditive processes are direct consequences of their T-additive counterparts.

2. THE DECOMPOSITION OF F.

In this section a decomposition of a T-superadditive process F into a difference of a T-additive process G and a positive, purely T-superadditive process H (that is, H is a positive T-superadditive process that does not dominate any non-zero positive T-additive process) is obtained.

CASE p=1. Let T be a positive Dunford-Schwartz operator i.e., T is an L_1 -contraction with $\|T\|_\infty < 1$. We will also assume that T is Markovian, that

$\int Tf \, d\mu = \int f \, d\mu$. In this case, if $\sup_{n \geq 1} \frac{1}{n} \|F_n\|_1 < \infty$, then the decomposition result is

obtained by M.A. Akcoglu and L. Sucheston [1]. Namely, they obtained that for all $n \geq 1$,

$$F_n = G_n - H_n \tag{2.1}$$

where $G_n = \sum_{i=0}^{n-1} T^i \delta$ for some $\delta \in L_1$, and $H_n = \sum_{i=0}^{n-1} h_i$ with $h_i = f_i - T^i \delta$. Using this

result they showed that $\lim_{n \rightarrow \infty} \frac{1}{n} F_n$ exists a.e., and moreover it is a consequence of

the same result that $\lim_{n \rightarrow \infty} \frac{1}{n} H_n = 0$ a.e.

CASE $1 < p < \infty$. In this case we let T be a positive L_p -contraction and F a T-

superadditive process with $\lim_{n \rightarrow \infty} \inf \left\| \frac{1}{n} \sum_{i=0}^n (F_i - TF_{i-1}) \right\|_p < \infty$. Under these

conditions B. Hachem [2] showed that $\lim_{n \rightarrow \infty} \frac{1}{n} F_n$ exists a.e. by reducing the problem to

a problem in an appropriate L_1 -space and employing Akcoglu-Sucheston's result in case p=1 above. Here we observe that the same technique can be applied to yield to a decomposition result similar to (2.1).

Using a result of M.A. Akcoglu and L. Sucheston [3] one can decompose X uniquely into disjoint union of sets E and E^c where:

(i) E is the support of a T-invariant function $h \in L_p^+$, and $\text{supp } g \subset E$ for all T-invariant $g \in L_p^+$.

(ii) $L_p(E)$ and $L_p(E^c)$ are invariant subspaces for T.

Then the following results are obtained [2,3]:

$$1_{E^c} \left(\frac{F}{n} \right) \rightarrow 0 \text{ a.e.}, \text{ so one can assume that } X=E. \tag{2.2}$$

The operator $P:L_p(m) \rightarrow L_p(m)$ defined as $Pf = \frac{T(fh)}{h}$, $f \in L_p(m)$, is a positive $L_p(m)$ -contraction and $Pl=1$, where $m=h^p \cdot \mu$. (2.3)

In particular, $\int Pfdm = \int f dm$ for all $f \in L_p^+(m)$. So P can be extended to a Markovian

operator on $L_1(m)$. Consequently $F'=\{h^{-1}F_n\}$ is a bounded P -superadditive process [2] in $L_1(m)$. Now by applying the Akcoglu-Sucheston's result [1] we can decompose F' into a difference of a P -additive process and a positive, purely P -superadditive process as $h^{-1}F_n = G_n - H_n$, $n > 1$. Also we see that:

$$\lim_{n \rightarrow \infty} \frac{1}{n} (h^{-1}F_n) \text{ exists } m\text{-a.e. and } \lim_{n \rightarrow \infty} \frac{1}{n} H_n = 0 \text{ } m\text{-a.e.}, \tag{2.4}$$

so $F_n = hG_n - hH_n$, and that $G=\{hG_n\}$ and $H=\{hH_n\}$ are T -additive and T -superadditive processes respectively by (2.3). Consequently (2.4) gives that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} F_n \text{ exists } \mu\text{-a.e. } X \text{ and} \\ \lim_{n \rightarrow \infty} \frac{1}{n} H_n = 0 \text{ } \mu\text{-a.e. } X \end{aligned} \tag{2.5}$$

by (2.2) and (2.3)

3. WEIGHTED AVERAGES.

Given a linear operator T on L_p , $1 < p < \infty$, and a sequence $A=\{a_n\}_{n>0}$ of complex numbers if

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} a_k T^k f \text{ exists a.e.}$$

for all $f \in L_p$, then we say that A is a good weight for T [4], or (A,T) is Birkhoff [5].

R. Sato [6] showed that the uniform sequences are good for $1 < p < \infty$. C. Ryll-Nardzewski [7] proved that the bounded Besicovitch sequences are good for T induced by a measure preserving transformation $\phi: X \rightarrow X$ by $Tf(x)=f(\phi(x))$ for any $f \in L_1$. This result combined with the remarkable theorem of J. Baxter and J. Olsen [5, Theorem 2.19] imply that bounded Besicovitch sequences are good for Dunford-Schwartz operators.

Now we observe the following: Let T be an operator on L_p and F be a T -superadditive process. If $F_n = G_n - H_n$, then for any sequence A

$$S_n(F,A) = S_n(G,A) - S_n(H,A) \tag{3.1}$$

Also

Also

$$0 < \lim_{n \rightarrow \infty} \sup_{n > 1} \frac{1}{n} |S_n(A, H)| < M \cdot \lim_{n \rightarrow \infty} \sup \frac{1}{n} H_n$$

where $M = \sup_{n > 1} |a_n|$. Therefore if $\lim_{n \rightarrow \infty} \frac{1}{n} H_n = 0$ a.e., $\lim_{n \rightarrow \infty} \frac{1}{n} S_n(A, H) = 0$ a.e. for

any bounded sequence A. We summarize this discussion as

THEOREM 3.2. Let T be a positive Dunford-Schwartz operator on L_p , $1 < p < \infty$, and F be a T-superadditive process. Assume also that

$$(i) \quad T \text{ is Markovian and } \sup_{n > 1} \left\| \frac{1}{n} F_n \right\|_1 < \infty \text{ when } p=1,$$

$$(ii) \quad \lim_{n \rightarrow \infty} \inf \left\| \frac{1}{n} \sum_{i=1}^n (F_i - T F_{i-1}) \right\|_p < \infty \quad (F_0 = 0) \text{ when } 1 < p < \infty.$$

If A is a bounded sequence such that (A, T) is Birkhoff, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} S_n(F, A) \text{ exists a.e.} \quad (3.3)$$

REMARK 3.4. The limit in (3.3) exists a.e. when A is a uniform sequence or a bounded Besicovitch sequence of $A \in \overline{W}_4$ [5]. In particular the subsequence theorem [5, 4] is valid for superadditive processes.

ACKNOWLEDGEMENTS. The author would like to thank Professor J.H. Olsen of North Dakota State University for his valuable comments on the subject.

REFERENCES

1. AKCOGLU, M.A. and SUCHESTON, L., A Ratio Ergodic Theorem for Superadditive Processes, Z. Wahr., 44, (1978), 269-278.
2. HACHEM, B., Un Theoreme Ergodique Sur-additif dans L_p , $1 < p < \infty$, C.R. Acad. Sci. Paris, Serie. I, 292, 531-534, (1981).
3. AKCOGLU, M.A. and SUCHESTON, On Convergence of Iterates of Positive Contractions in L_p -spaces, J. Approx. Theory, 13, (1975), 348-362.
4. BELLOW, A. and LOSERT, V., The Weighted Ergodic Theorem and the Individual Ergodic Theorem Along Subsequences, Trans. A.M.S., 288, (1985), 307-3435.
5. BAXTER, J. and OLSEN, J., Weighted and Subsequential Ergodic Theorems, Can. J. Math., 35, (1983), 145-166.
6. SATO, R., Operator Averages for Subsequences, Math. J. Okayama Univ., 22, (1980), 161-168.
7. RYLL-NARDZEWSKI, C., Topics in Ergodic Theory, Proc. of the Winter School in Prob., Karpacz, Poland, 131-156, Lecture Notes in Math., #472, Springer-Verlag, Berlin, (1975).

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru