

**FRACTIONAL DERIVATIVES OF HOLOMORPHIC FUNCTIONS ON  
BOUNDED SYMMETRIC DOMAINS OF  $C^n$**

**ZENGJIAN LOU**

Department of Mathematics  
Qufu Normal University  
Qufu Shandong, 273165  
P R CHINA

(Received March 8, 1994 and in revised form January 13, 1995)

**ABSTRACT.** Let  $f \in H(B_n)$   $f^{[\beta]}$  denotes the  $\beta$ th fractional derivative of  $f$  If  $f^{[\beta]} \in A^{p,q,\alpha}(B_n)$ , we show that

- (I) If  $\beta < \frac{\alpha+1}{p} + \frac{n}{q} = \delta$ , then  $f \in A^{s,t,\alpha}(B_n)$ , and  $\|f\|_{s,t,\alpha} \leq C \|f^{[\beta]}\|_{p,q,\alpha}$ ,  $s = \frac{\delta p}{\delta - \beta}$ ,  $t = \frac{\delta q}{\delta - \beta}$
- (II) If  $\beta = \frac{\alpha+1}{p} + \frac{n}{q}$ , then  $f \in B(B_n)$  and  $\|f\|_B \leq C \|f^{[\beta]}\|_{p,q,\alpha}$
- (III) If  $\beta > \frac{\alpha+1}{p} + \frac{n}{q}$ , then  $f \in \Lambda_{\beta - \frac{\alpha+1}{p} - \frac{n}{q}}(B_n)$  especially If  $\beta = 1$  then  $\|f\|_{\Lambda_{1 - \frac{\alpha+1}{p} - \frac{n}{q}}} \leq C \|f^{[1]}\|_{p,q,\alpha}$  where  $B_n$  is the unit ball of  $C^n$

**KEY WORDS AND PHRASES.** Fractional derivative, Bergman space, Bloch space, Lipschitz space  
**1991 AMS SUBJECT CLASSIFICATION CODE.** 32A10

Let  $\Omega$  be a bounded symmetric domain in the complex vector space  $C^n$ ,  $o \in \Omega$ , with Bergman-Silov boundary  $b$ ,  $\Gamma$  the group of holomorphic automorphisms of  $\Omega$  and  $\Gamma_0$  its isotropy group. It is known that  $\Omega$  is circular and star-shaped with respect to  $o$  and  $b$  is circular. The group  $\Gamma_0$  is transitive on  $b$  and  $b$  has a unique normalized  $\Gamma_0$ -invariant measure  $\sigma$  with  $\sigma(b) = 1$ . Hua [2] constructed by group representation theory a system  $\{\phi_{kv}\}$  of homogeneous polynomials,  $k = 0, 1, \dots, v = 1, \dots, m_k, m_k = \binom{n+k-1}{k}$ , complete and orthogonal on  $\Omega$  and orthonormal on  $b$ .

By  $H(\Omega)$  we denote the class of all holomorphic functions on  $\Omega$ . Every  $f \in H(\Omega)$  has a series expansion

$$f(z) = \sum_{k,v} a_{kv} \phi_{kv}(z), \quad a_{kv} = \lim_{r \rightarrow 1} \int_b f(r\xi) \overline{\phi_{kv}(\xi)} d\sigma(\xi) \quad (0)$$

where  $\sum_{k,v} = \sum_{k=0}^{\infty} \sum_{v=1}^{m_k}$  and the convergence is uniform on a compact subset of  $\Omega$ .

Let  $f \in H(\Omega)$  with the expansion (0) and  $\beta > 0$ . The  $\beta$ th fractional derivatives of  $f$  are defined, respectively, by

$$f^{[\beta]}(z) = \sum_{k,v} \frac{\Gamma(k+1+\beta)}{\Gamma(k+1)} a_{kv} \phi_{kv}(z)$$

$$f_{[\beta]}(z) = \sum_{k,v} \frac{\Gamma(k+1)}{\Gamma(k+1+\beta)} a_{kv} \phi_{kv}(z)$$

It is known that  $f^{[\beta]}, f_{[\beta]} \in H(\Omega)$  and

$$f(\tau\xi) = \frac{1}{\Gamma(\beta)} \int_0^1 (1-\rho)^{\beta-1} f^{[\beta]}(r\rho\xi) d\rho. \quad (1)$$

Let  $f \in H(\Omega)$ . It will be said that  $f$  belongs to the Bergman spaces  $A^{p,q,\alpha}(\Omega)$ ,  $0 < p, q \leq \infty, \alpha > -1$  if

$$\|f\|_{p,q,\alpha} = \begin{cases} \left( \int_0^1 (1-r)^\alpha M_q(r, f)^p dr \right)^{\frac{1}{p}}, & p < \infty \\ \sup_{0 < r < 1} (1-r)^\alpha M_q(r, f), & p = \infty \end{cases}$$

is finite, where

$$M_q(r, f) = \left( \int_b |f(r\xi)|^q d\sigma(\xi) \right)^{1/q}, \quad 0 < q < \infty$$

and

$$M_\infty(r, f) = \sup_{\xi \in b} |f(r\xi)|$$

see [1,3,5,6,7] for more on  $A^{p,q,\alpha}(\Omega)$  For  $0 < p \leq \infty$ , let  $A^p(\Omega)$  denote  $A^{p,p,0}(\Omega)$  (see [10,12]),  $H^p(\Omega)$  denote  $A^{\infty,p,0}(\Omega)$  (see [9])

Let  $B_n$  denote the unit ball in  $C^n$  A function  $f \in H(B_n)$  is called a Bloch function, that is  $f \in B(B_n)$ , if ([8,11])

$$\|f\|_B = \sup_{z \in B_n} (1 - |z|)|f^{[1]}(z)| < \infty$$

For  $0 < \alpha < \infty$ , the definition of Lipschitz space  $\Lambda_\alpha(B_n)$  can be found in [4, §8.8]

In [10] and [12], Watanabe and Stojan considered the problem If  $f' \in A^p(D)$  ( $D$  is the unit disc of  $C^1$ ), then  $q = ?$  such that  $f \in A^q(D)$  In this paper we consider and solve the same problem in  $A^{p,q,\alpha}(\Omega)$

The main results of this paper are the following

**THEOREM 1.** Let  $0 < p, q \leq \infty, \alpha > -1, 0 < \beta < \delta \leq \frac{\alpha+1}{p} + \frac{n}{q}$ , if  $f^{[\beta]} \in A^{p,q,\alpha}(\Omega)$  and  $f^{[\beta]}(r\xi) = O\left(\|f^{[\beta]}\|_{p,q,\alpha}(1-r)^{-\delta}\right)$ , then  $f \in A^{s,t,\alpha}(\Omega)$  and  $\|f\|_{s,t,\alpha} \leq C\|f^{[\beta]}\|_{p,q,\alpha}$ , where  $s = \frac{\delta p}{\delta - \beta}, t = \frac{\delta q}{\delta - \beta}$

**THEOREM 2.** Let  $0 < p, q \leq \infty, \alpha > -1, 0 < \beta < \infty, f^{[\beta]} \in A^{p,q,\alpha}(B_n)$ .

- (I) If  $\beta < \frac{\alpha+1}{p} + \frac{n}{q} = \delta$ , then  $f \in A^{s,t,\alpha}(B_n)$ , and  $\|f\|_{s,t,\alpha} \leq C\|f^{[\beta]}\|_{p,q,\alpha}$ , where  $s, t$  are the same as above
- (II) If  $\beta = \frac{\alpha+1}{p} + \frac{n}{q}$ , then  $f \in B(B_n)$  and  $\|f\|_B \leq C\|f^{[\beta]}\|_{p,q,\alpha}$
- (III) If  $\beta > \frac{\alpha+1}{p} + \frac{n}{q}$ , then  $f \in \Lambda_{\beta - \frac{\alpha+1}{p} - \frac{n}{q}}(B_n)$ , especially If  $\beta = 1$ , then  $\|f\|_{\Lambda_{1 - \frac{\alpha+1}{p} - \frac{n}{q}}} \leq C\|f^{[1]}\|_{p,q,\alpha}$

**REMARK.** (i) Theorem 2(I) ( $p = q, \alpha = 0, \beta = n = 1$ ) extends the results of Watanabe's and Stojan's (ii) Theorem 1 ( $p = \infty$ ) extends the results of Shi's ([9]) and Lou's ([6,7])

**REFERENCES**

- [1] AHERN, P AND JEVTIC, M , Duality and multipliers for mixed norm spaces, *Mich. Math. J.*, **30** (1983), 53-63
- [2] HUA, L K , Harmonic analysis of functions of several variables in the classical domains, *Trans. Amer. Math. Soc.*, **6** (1963).
- [3] JEVTIC, M , Projection theorems, fractional derivatives and inclusion theorems for mixed-norm spaces on the ball, *Analysis*, **9** (1989), 83-105
- [4] KRANTZ, S.G , *Function Theory of Several Complex Variables*, John Wiley & Sons, New York, 1982
- [5] LOU, Z J , A note on a conjecture of S Axler, *J. Math. Res. & Exp.*, **11** (4) (1991), 629-630 (Chinese)
- [6] LOU, Z J , Hardy-Littlewood type theorem on Bergman spaces, *J. Math. Res. & Exp.*, **11** (3) (1991) (Chinese)
- [7] LOU, Z J , Hardy-Littlewood type theorem on Bergman spaces of several complex variables, *J. Qufu Normal Univ.*, **1** (1992), 111-112 (Chinese)
- [8] LOU, Z J., Characterizations of Bloch functions on the unit ball of  $C^n$ , *Kodai Math. J.*, **16** (1993), 74-78
- [9] SHI, J H , Hardy-Littlewood theorems on bounded symmetric domains, *Science in China*, **4** (1988), 366-375
- [10] STOJAN, D , Some new properties of the spaces  $A^p$ , *Mat. Vesnik*, **5** 18,33 (1981), 151-157
- [11] TIMONEY, R M , Bloch functions in several complex variables, *Bull. London. Math. Soc.*, **12** (1980), 241-267