

LOCAL STABILITY OF THE ADDITIVE FUNCTIONAL EQUATION AND ITS APPLICATIONS

SOON-MO JUNG and BYUNGBAE KIM

Received 5 October 2001

The main purpose of this paper is to prove the Hyers-Ulam stability of the additive functional equation for a large class of unbounded domains. Furthermore, by using the theorem, we prove the stability of Jensen's functional equation for a large class of restricted domains.

2000 Mathematics Subject Classification: 39B82, 39B62.

1. Introduction. The starting point of studying the stability of functional equations seems to be the famous talk of Ulam [14] in 1940, in which he discussed a number of important unsolved problems. Among those was the question concerning the stability of group homomorphisms: let G_1 be a group and let G_2 be a metric group with a metric $d(\cdot, \cdot)$. Given $\varepsilon > 0$, does there exist a $\delta > 0$ such that if a mapping $h : G_1 \rightarrow G_2$ satisfies the inequality $d(h(xy), h(x)h(y)) < \delta$ for all $x, y \in G_1$, then there exists a homomorphism $H : G_1 \rightarrow G_2$ with $d(h(x), H(x)) < \varepsilon$ for all $x \in G_1$?

The case of approximately additive mappings was solved by Hyers [3] under the assumption that G_1 and G_2 are Banach spaces. Later, the result of Hyers was significantly generalized by Rassias [11]. It should be remarked that we can find in [4] a lot of references concerning the stability of functional equations (see also [2, 5, 6]).

In [12, 13], Skof investigated the Hyers-Ulam stability of the additive functional equation for many cases of restricted domains in \mathbb{R} . Later, Losonczi [9] proved the local stability of the additive equation for more general cases and applied the result to the proof of stability of the Hosszú's functional equation.

In Section 2, the Hyers-Ulam stability of the additive equation will be investigated for a large class of unbounded domains. Moreover, in Section 3, we will apply the previous result to the proof of the local stability of the Jensen's functional equation on unbounded domains.

Throughout this paper, let E_1 and E_2 be a real (or complex) normed space and a Banach space, respectively.

2. Stability of additive equation on restricted domains. Assume that $\varphi : (0, \infty) \rightarrow [0, \infty)$ is a decreasing mapping for which there exists a $d > 0$ such

that

$$\varphi(s) \leq s, \quad (2.1)$$

for any $s \geq d$.

We now define

$$\begin{aligned} B_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| < \varphi(\|x\|)\} \cup \{(0, y) \in E_1^2 : y \in E_1\}, \\ B_2 &= \{(x, y) \in E_1^2 : \|x + y\| < d\}. \end{aligned} \quad (2.2)$$

In the following theorem, we generalize the theorems of Skof [12, 13] and of Losonczi [9] concerning the stability of the additive equation on restricted domains.

THEOREM 2.1. *If a mapping $f : E_1 \rightarrow E_2$ with $\|f(0)\| \leq \varepsilon$ satisfies the inequality*

$$\|f(x + y) - f(x) - f(y)\| \leq \varepsilon, \quad (2.3)$$

for some $\varepsilon \geq 0$ and all $(x, y) \in E_1^2 \setminus (B_1 \cup B_2)$, then there exists a unique additive mapping $A : E_1 \rightarrow E_2$ such that

$$\|f(x) - A(x)\| \leq 39\varepsilon, \quad \forall x \in E_1. \quad (2.4)$$

PROOF. First, we assume that $(x, y) \in B_2$ satisfies $x \neq 0$, $y \neq 0$, and $x + y \neq 0$. For this case, we can choose a $z_1 \in E_1$ with

$$\begin{aligned} \|z_1\| \geq \varphi(\|x + y\|), \quad \|z_1\| \geq \varphi(\|x\|), \quad \|x + z_1\| \geq \varphi(\|y\|), \\ \|x + y + z_1\| \geq d, \quad \|x + z_1\| \geq d. \end{aligned} \quad (2.5)$$

Thus, the pairs $(x + y, z_1)$, (x, z_1) , and $(y, x + z_1)$ do not belong to $B_1 \cup B_2$. Hence, it follows from (2.3) that

$$\begin{aligned} \|f(x + y) - f(x) - f(y)\| &\leq \| -f(x + y + z_1) + f(x + y) + f(z_1) \| \\ &\quad + \|f(x + z_1) - f(x) - f(z_1)\| \\ &\quad + \|f(x + y + z_1) - f(y) - f(x + z_1)\| \\ &\leq 3\varepsilon, \end{aligned} \quad (2.6)$$

for any $(x, y) \in B_2$ with $x \neq 0$, $y \neq 0$, and $x + y \neq 0$.

When $x = 0$ or $y = 0$, we have

$$\|f(x + y) - f(x) - f(y)\| = \|f(0)\| \leq \varepsilon. \quad (2.7)$$

Taking this fact into account, we see that inequality (2.6) is valid for all $(x, y) \in B_2$ with $x + y \neq 0$.

We now assume that $(x, y) \in B_2$ satisfies $x + y = 0$ and $\|x\| \geq d$. (In this case, $\|y\| = \|-x\| \geq d$.) In view of (2.1), both the pairs $(-x, -x)$ and $(x, -2x)$ do not belong to $B_1 \cup B_2$. Hence, it follows from (2.3) that

$$\|f(-2x) - 2f(-x)\| \leq \varepsilon, \quad \|f(-x) - f(x) - f(-2x)\| \leq \varepsilon. \quad (2.8)$$

From the last two inequalities we get

$$\begin{aligned} \|f(x+y) - f(x) - f(y)\| &= \|f(0) - f(x) - f(-x)\| \\ &\leq \|f(0)\| + \|f(-2x) - 2f(-x)\| \\ &\quad + \|f(-x) - f(x) - f(-2x)\| \\ &\leq 3\varepsilon. \end{aligned} \quad (2.9)$$

Considering all the previous inequalities including (2.3), we conclude that f satisfies the inequality

$$\|f(x+y) - f(x) - f(y)\| \leq 3\varepsilon, \quad (2.10)$$

for all $(x, y) \in E_1^2 \setminus (B_1 \cup B_2) \cup \{(u, v) \in B_2 : \|u\| \geq d\}$.

Now, let $(x, y) \in E_1^2$ be arbitrarily given with $\|x\| \geq d$ and $\|y\| \geq d$. Since φ is decreasing, we see by (2.1) that

$$\varphi(\|x\|) \leq \varphi(d) \leq d \leq \|y\|, \quad (2.11)$$

and this implies that $(x, y) \notin B_1$. If, moreover, the given pair (x, y) belongs to B_2 , then $(x, y) \in \{(u, v) \in B_2 : \|u\| \geq d\}$. Otherwise, $(x, y) \in E_1^2 \setminus (B_1 \cup B_2)$. Hence, it follows from (2.10) that

$$\|f(x+y) - f(x) - f(y)\| \leq 3\varepsilon, \quad (2.12)$$

for all $(x, y) \in E_1^2$ with $\|x\| \geq d$ and $\|y\| \geq d$.

Assume that $(x, y) \in E_1^2$ with $\|x\| < d$ and $\|y\| \geq 4d$. In this case, we may choose a $z_2 \in E_1$ with $2d \leq \|z_2\| < 3d$. Then, it holds that

$$\begin{aligned} \|x - z_2\| \geq d, \quad \|y + z_2\| \geq d, \quad \|x - z_2\| \geq d, \quad \|z_2\| \geq 2d, \\ \|-z_2\| \geq 2d, \quad \|y + z_2\| \geq d, \quad \|z_2\| \geq 2d, \quad \|-z_2\| \geq 2d. \end{aligned} \quad (2.13)$$

It then follows from (2.12) and (2.13) that

$$\begin{aligned}
\|f(x+y) - f(x) - f(y)\| &\leq \|f(x+y) - f(x-z_2) - f(y+z_2)\| \\
&\quad + \|-f(x) + f(x-z_2) + f(z_2)\| \\
&\quad + \|-f(y) + f(-z_2) + f(y+z_2)\| \\
&\quad + \|f(0) - f(z_2) - f(-z_2)\| + \|-f(0)\| \\
&\leq 13\varepsilon,
\end{aligned} \tag{2.14}$$

for $(x, y) \in E_1^2$ with $\|x\| < d$ and $\|y\| \geq 4d$.

Combining (2.12) and (2.14), we have

$$\|f(x+y) - f(x) - f(y)\| \leq 13\varepsilon, \tag{2.15}$$

for all $(x, y) \in E_1^2$ with $\|y\| \geq 4d$. Since the Cauchy difference $f(x+y) - f(x) - f(y)$ is symmetric with respect to x and y , we conclude that inequality (2.15) is true for all $(x, y) \in E_1^2$ with $\|x\| \geq 4d$ or $\|y\| \geq 4d$.

If $(x, y) \in E_1^2$ satisfies $\|x\| < 4d$ and $\|y\| < 4d$, then we can choose a $z_3 \in E_1$ with $\|z_3\| \geq 8d$. Then, we have $\|x+z_3\| \geq 4d$. Since inequality (2.15) holds true for all $(x, y) \in E_1^2$ with $\|x\| \geq 4d$ or $\|y\| \geq 4d$, we get

$$\begin{aligned}
\|f(x+y) - f(x) - f(y)\| &\leq \|-f(x+y+z_3) + f(x+y) + f(z_3)\| \\
&\quad + \|f(x+z_3) - f(x) - f(z_3)\| \\
&\quad + \|f(x+y+z_3) - f(y) - f(x+z_3)\| \\
&\leq 39\varepsilon,
\end{aligned} \tag{2.16}$$

for any $(x, y) \in E_1^2$ with $\|x\| < 4d$ and $\|y\| < 4d$.

Inequality (2.16) together with (2.15) yields

$$\|f(x+y) - f(x) - f(y)\| \leq 39\varepsilon, \quad \forall x, y \in E_1. \tag{2.17}$$

According to [1], there exists a unique additive mapping $A : E_1 \rightarrow E_2$ that satisfies inequality (2.4) for each x in E_1 . \square

COROLLARY 2.2. *Let $d > 0$ and $\varepsilon \geq 0$ be given. If a mapping $f : E_1 \rightarrow E_2$ with $\|f(0)\| \leq \varepsilon$ satisfies inequality (2.3) for all $x, y \in E_1$ with $\max\{\|x\|, \|y\|\} \geq d$ and $\|x+y\| \geq d$, then there exists a unique additive mapping $A : E_1 \rightarrow E_2$ that satisfies inequality (2.4) for each $x \in E_1$.*

PROOF. Because of the symmetry property of the Cauchy difference with respect to x and y , we can, without loss of generality, assume that f satisfies inequality (2.3) for all $x, y \in E_1$ with $\|y\| \geq d$ and $\|x+y\| \geq d$.

For a constant mapping $\varphi(s) = d$ ($s > 0$), define

$$\begin{aligned}
 B_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| < d\} \cup \{(0, y) \in E_1^2 : y \in E_1\}, \\
 B_2 &= \{(x, y) \in E_1^2 : \|x + y\| < d\}.
 \end{aligned}
 \tag{2.18}$$

Since

$$\begin{aligned}
 E_1^2 \setminus B_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| \geq d\}, \\
 E_1^2 \setminus B_2 &= \{(x, y) \in E_1^2 : \|x + y\| \geq d\},
 \end{aligned}
 \tag{2.19}$$

we have

$$E_1^2 \setminus (B_1 \cup B_2) = \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| \geq d \text{ and } \|x + y\| \geq d\}.
 \tag{2.20}$$

Thus, it follows from our hypothesis that f satisfies inequality (2.3) for all $(x, y) \in E_1^2 \setminus (B_1 \cup B_2)$.

According to Theorem 2.1, there exists a unique additive mapping $A : E_1 \rightarrow E_2$ that satisfies inequality (2.4) for all $x \in E_1$. □

In 1983, Skof [12] presented an interesting asymptotic behavior of the additive mappings: a mapping $f : \mathbb{R} \rightarrow \mathbb{R}$ is additive if and only if $|f(x + y) - f(x) - f(y)| \rightarrow 0$ as $|x| + |y| \rightarrow \infty$.

Without difficulty, the above theorem of Skof can be extended to mappings from a real normed space to a Banach space. We now apply Corollary 2.2 to a generalization of Skof theorem.

COROLLARY 2.3. *A mapping $f : E_1 \rightarrow E_2$ is additive if and only if*

$$\|f(x + y) - f(x) - f(y)\| \rightarrow 0
 \tag{2.21}$$

as $\|x + y\| \rightarrow \infty$.

PROOF. On account of the hypothesis, there exists a decreasing sequence (ε_n) with $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ and

$$\|f(x + y) - f(x) - f(y)\| \leq \varepsilon_n,
 \tag{2.22}$$

for all $(x, y) \in E_1^2$ with $\|x + y\| \geq n$. With $y = 0$ and $\|x\| \rightarrow \infty$, our hypothesis implies that $f(0) = 0$.

By Corollary 2.2, there exists a unique additive mapping $A_n : E_1 \rightarrow E_2$ such that

$$\|f(x) - A_n(x)\| \leq 39\varepsilon_n, \quad \forall x \in E_1.
 \tag{2.23}$$

Now, let l and m be integers with $m > l > 0$. Then, inequality (2.23) implies that

$$\|f(x) - A_m(x)\| \leq 39\varepsilon_m \leq 39\varepsilon_l,
 \tag{2.24}$$

for $x \in E_1$, and further, the uniqueness of A_n implies that $A_m = A_l$ for all integers $l, m > 0$, that is, $A_n = A_1$ for any $n \in \mathbb{N}$. By letting $m \rightarrow \infty$ in the last inequality, we get

$$\|f(x) - A_1(x)\| = 0, \quad (2.25)$$

for any $x \in E_1$, which means that f is additive. The reverse assertion is trivial. \square

3. Stability of Jensen's equation on restricted domains. Kominek investigated in [8] the Hyers-Ulam stability of the Jensen's functional equation

$$2f\left(\frac{x+y}{2}\right) = f(x) + f(y), \quad (3.1)$$

for the class of mappings defined on a bounded subset of \mathbb{R}^N . On the other hand, the author proved in [7] the Hyers-Ulam stability of that equation on unbounded domains.

In this section, we use [Theorem 2.1](#) to generalize the theorems of the author and of Kominek.

Let $\varphi_1 : [0, \infty) \rightarrow [0, \infty)$ be a decreasing mapping that satisfies $\varphi_1(0) = d_0 > 0$. Define

$$\begin{aligned} B_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| < \varphi_1(\|x\|)\} \cup \{(0, y) \in E_1^2 : y \in E_1\}, \\ B_2 &= \{(x, y) \in E_1^2 : \|x + y\| < d_0\}, \\ D &= \{(0, y) \in E_1^2 : \|y\| \geq d_0\}. \end{aligned} \quad (3.2)$$

THEOREM 3.1. *If a mapping $f : E_1 \rightarrow E_2$ satisfies the inequality*

$$\left\| 2f\left(\frac{x+y}{2}\right) - f(x) - f(y) \right\| \leq \varepsilon, \quad (3.3)$$

for some $\varepsilon \geq 0$ and all $(x, y) \in E_1^2 \setminus (B_1 \cup B_2) \cup D$, then there exists a unique additive mapping $A : E_1 \rightarrow E_2$ such that

$$\|f(x) - A(x) - f(0)\| \leq 78\varepsilon, \quad (3.4)$$

for any $x \in E_1$.

PROOF. If we substitute $g(x)$ for $f(x) - f(0)$ in (3.3), then

$$\left\| 2g\left(\frac{x+y}{2}\right) - g(x) - g(y) \right\| \leq \varepsilon, \quad (3.5)$$

for any $(x, y) \in E_1^2 \setminus (B_1 \cup B_2) \cup D$. With $x = 0$ and $\|y\| \geq d_0$, inequality (3.5) yields

$$\left\| 2g\left(\frac{y}{2}\right) - g(y) \right\| \leq \varepsilon, \tag{3.6}$$

for each $y \in E_1$ with $\|y\| \geq d_0$. Replace y by $x + y$ ($\|x + y\| \geq d_0$) in inequality (3.6) to get

$$\left\| 2g\left(\frac{x+y}{2}\right) - g(x+y) \right\| \leq \varepsilon, \tag{3.7}$$

for all $x, y \in E_1$ with $\|x + y\| \geq d_0$.

It follows from (3.5) and (3.7) that

$$\begin{aligned} & \|g(x+y) - g(x) - g(y)\| \\ & \leq \left\| g(x+y) - 2g\left(\frac{x+y}{2}\right) \right\| + \left\| 2g\left(\frac{x+y}{2}\right) - g(x) - g(y) \right\| \\ & \leq 2\varepsilon, \end{aligned} \tag{3.8}$$

for every $(x, y) \in E_1^2 \setminus (B_1 \cup B_2) \cup D$ with $\|x + y\| \geq d_0$. Since $(x, y) \in E_1^2 \setminus (B_1 \cup B_2)$ implies that $\|x + y\| \geq d_0$, the mapping g surely satisfies

$$\|g(x+y) - g(x) - g(y)\| \leq 2\varepsilon, \tag{3.9}$$

for all $(x, y) \in E_1^2 \setminus (B_1 \cup B_2)$.

It trivially holds that $\varphi_1(s) \leq s$ for all $s \geq d_0$. On account of [Theorem 2.1](#), there exists a unique additive mapping $A : E_1 \rightarrow E_2$ such that

$$\|g(x) - A(x)\| \leq 78\varepsilon, \tag{3.10}$$

for each x in E_1 . □

Let $\varphi_2 : (0, \infty) \rightarrow [0, \infty)$ be a continuous and decreasing mapping that satisfies

$$0 < d = \inf \{s > 0 : \varphi_2(s) = 0\} < \infty. \tag{3.11}$$

Furthermore, assume that the restriction $\varphi_2|_{(0,d]}$ is strictly decreasing.

Now, we define

$$\begin{aligned} B_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| < \varphi_2(\|x\|)\} \cup \{(0, y) \in E_1^2 : y \in E_1\}, \\ B_2 &= \{(x, y) \in E_1^2 : \|x + y\| < d_0\}, \\ D &= \{(0, y) \in E_1^2 : \|y\| \geq d_0\}, \end{aligned} \quad (3.12)$$

where we set $d_0 = \inf\{d, \lim_{s \rightarrow 0^+} \varphi_2(s)\}$.

COROLLARY 3.2. *If a mapping $f : E_1 \rightarrow E_2$ satisfies inequality (3.3) for some $\varepsilon \geq 0$ and all $(x, y) \in E_1^2 \setminus (B_1 \cup B_2) \cup D$, then there exists a unique additive mapping $A : E_1 \rightarrow E_2$ satisfying inequality (3.4) for all $x \in E_1$.*

PROOF. First, we define a mapping $\varphi_0 : [0, \infty) \rightarrow [0, \infty)$ by

$$\varphi_0(s) = \begin{cases} d_0, & \text{for } s = 0, \\ \inf\{\varphi_2(s), \inf \varphi_2^{-1}(s)\}, & \text{for } s > 0, \end{cases} \quad (3.13)$$

where we set $\varphi_2^{-1}(t) = \{s > 0 : \varphi_2(s) = t\}$ and $\inf \emptyset = \infty$. (We cannot exclude the case $\varphi_2^{-1}(s) = \emptyset$ from the above definition.) We define

$$\begin{aligned} \tilde{B}_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| < \varphi_0(\|x\|)\} \cup \{(0, y) \in E_1^2 : y \in E_1\}, \\ \tilde{B}_2 &= \{(x, y) \in E_1^2 : \|x + y\| < d_0\}, \\ \tilde{D} &= \{(0, y) \in E_1^2 : \|y\| \geq d_0\}. \end{aligned} \quad (3.14)$$

The fact that $\varphi_0(s) \leq \varphi_2(s)$ for all $s > 0$ implies that $\tilde{B}_1 \subset B_1$. Since $B_2 = \tilde{B}_2$ and $D = \tilde{D}$, we get

$$E_1^2 \setminus (B_1 \cup B_2) \cup D \subset E_1^2 \setminus (\tilde{B}_1 \cup \tilde{B}_2) \cup \tilde{D}. \quad (3.15)$$

Now, assume that $(x, y) \in E_1^2 \setminus (\tilde{B}_1 \cup \tilde{B}_2) \cup \tilde{D}$ but $(x, y) \notin E_1^2 \setminus (B_1 \cup B_2) \cup D$. Because $(x, y) \notin D$ and $(x, y) \notin B_2$, we have

$$x \neq 0, \quad \|x + y\| \geq d_0. \quad (3.16)$$

Moreover, (x, y) should belong to $B_1 \setminus \tilde{B}_1$, that is,

$$0 < \inf \varphi_2^{-1}(\|x\|) \leq \|y\| < \varphi_2(\|x\|). \quad (3.17)$$

(Since $\|x\| > 0$ and $\varphi_2|_{(0,d]}$ is strictly decreasing, we have $\inf \varphi_2^{-1}(\|x\|) > 0$.) If we assume that $(y, x) \in B_1$, then we get $\|x\| < \varphi_2(\|y\|)$. This fact implies that $\|y\| < \inf \varphi_2^{-1}(\|x\|)$, which is contrary to (3.17). Hence, by (3.16), we conclude that $(y, x) \notin B_1 \cup B_2$. This fact together with (3.3), yields

$$\left\| 2f\left(\frac{y+x}{2}\right) - f(y) - f(x) \right\| \leq \varepsilon, \tag{3.18}$$

for all $(x, y) \in E_1^2 \setminus (\tilde{B}_1 \cup \tilde{B}_2) \cup \tilde{D}$.

We now define another mapping $\varphi : [0, \infty) \rightarrow [0, \infty)$ by

$$\varphi(s) = \begin{cases} d_0, & \text{for } s = 0, \\ \inf \{ \varphi_2(s), \inf \varphi_2^{-1}(s) \}, & \text{for } 0 < s \leq d_1, \\ \sup \{ \varphi_2(s), \sup \varphi_2^{-1}(s) \}, & \text{for } s > d_1, \end{cases} \tag{3.19}$$

where $d_1 > 0$ is the unique fixed point of φ_2 , that is, $d_1 = \varphi_2(d_1)$, and we set $\inf \emptyset = \infty$ and $\sup \emptyset = 0$.

Let $s_i > 0$ ($i = 1, 2, 3, 4$) be arbitrarily given with $0 < s_1 < s_2 \leq d_1 < s_3 < s_4$. Since φ_2 is decreasing, we have

$$\begin{aligned} \lim_{s \rightarrow 0^+} \varphi_2(s) &\geq \varphi_2(s_1) \geq \varphi_2(s_2) \geq d_1 \geq \varphi_2(s_3) \geq \varphi_2(s_4), \\ d &\geq \inf \varphi_2^{-1}(s_1) \geq \inf \varphi_2^{-1}(s_2) \geq d_1 \geq \sup \varphi_2^{-1}(s_3) \geq \sup \varphi_2^{-1}(s_4). \end{aligned} \tag{3.20}$$

Hence, we get

$$\varphi(0) \geq \varphi(s_1) \geq \varphi(s_2) \geq \varphi(s_3) \geq \varphi(s_4) \tag{3.21}$$

which implies that φ is decreasing.

Similarly as before, we define

$$\begin{aligned} \hat{B}_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| < \varphi(\|x\|)\} \cup \{(0, y) \in E_1^2 : y \in E_1\}, \\ \hat{B}_2 &= \{(x, y) \in E_1^2 : \|x + y\| < d_0\}, \\ \hat{D} &= \{(0, y) \in E_1^2 : \|y\| \geq d_0\}. \end{aligned} \tag{3.22}$$

Since $\hat{B}_1 \supset \tilde{B}_1$, $\hat{B}_2 = \tilde{B}_2$, and $\hat{D} = \tilde{D}$, we may conclude that inequality (3.3) holds true for all $(x, y) \in E_1^2 \setminus (\hat{B}_1 \cup \hat{B}_2) \cup \hat{D}$.

According to Theorem 3.1, there exists a unique additive mapping $A : E_1 \rightarrow E_2$ such that inequality (3.4) is true for any $x \in E_1$. □

The author in [7] proved that it needs only to show an asymptotic property of the Jensen difference to identify a given mapping with an additive one.

Let X and Y be a real normed space and a real Banach space, respectively. A mapping $f : X \rightarrow Y$ with $f(0) = 0$ is additive if and only if

$$\left\| 2f\left(\frac{x+y}{2}\right) - f(x) - f(y) \right\| \rightarrow 0 \quad (3.23)$$

as $\|x\| + \|y\| \rightarrow \infty$.

By using [Theorem 3.1](#), we now prove an asymptotic behavior of additive mappings which generalizes the above result.

COROLLARY 3.3. *A mapping $f : E_1 \rightarrow E_2$ with $f(0) = 0$ is additive if and only if*

$$\left\| 2f\left(\frac{x+y}{2}\right) - f(x) - f(y) \right\| \rightarrow 0 \quad (3.24)$$

as $\|x+y\| \rightarrow \infty$.

PROOF. According to our hypothesis, there exists a decreasing sequence (ε_n) with $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ and

$$\left\| 2f\left(\frac{x+y}{2}\right) - f(x) - f(y) \right\| \leq \varepsilon_n, \quad (3.25)$$

for all $(x, y) \in E_1^2$ with $\|x+y\| \geq n$.

The mapping $\varphi_1 : [0, \infty) \rightarrow [0, \infty)$ defined by $\varphi_1(s) = -s + n$ ($s \geq 0$) is decreasing. Moreover, it holds that $\varphi_1(0) = n$. We define

$$\begin{aligned} B_1 &= \{(x, y) \in E_1 \setminus \{0\} \times E_1 : \|y\| < -\|x\| + n\} \cup \{(0, y) \in E_1^2 : y \in E_1\}, \\ B_2 &= \{(x, y) \in E_1^2 : \|x+y\| < n\}, \\ D &= \{(0, y) \in E_1^2 : \|y\| \geq n\}. \end{aligned} \quad (3.26)$$

Since $B_1 \cup B_2 = \{(x, y) \in E_1^2 : x = 0 \text{ or } \|x+y\| < n\}$ and $D = \{(x, y) \in E_1^2 : x = 0 \text{ and } \|x+y\| \geq n\}$, we have

$$E_1^2 \setminus (B_1 \cup B_2) = \{(x, y) \in E_1^2 : x \neq 0 \text{ and } \|x+y\| \geq n\}, \quad (3.27)$$

and hence

$$E_1^2 \setminus (B_1 \cup B_2) \cup D = \{(x, y) \in E_1^2 : \|x+y\| \geq n\}. \quad (3.28)$$

Therefore, inequality [\(3.25\)](#) holds true for all $(x, y) \in E_1^2 \setminus (B_1 \cup B_2) \cup D$.

According to [Theorem 3.1](#), there exists a unique additive mapping $A_n : E_1 \rightarrow E_2$ such that

$$\|f(x) - A_n(x)\| \leq 78\varepsilon_n, \quad \forall x \in E_1. \quad (3.29)$$

Now, let l and m be positive integers with $m > l$. Then, it follows from [\(3.29\)](#) that

$$\|f(x) - A_m(x)\| \leq 78\varepsilon_m \leq 78\varepsilon_l, \quad (3.30)$$

for $x \in E_1$. However, the uniqueness of A_n implies that $A_m = A_l$ for all positive integers l and m , that is, $A_n = A_1$ for any $n \in \mathbb{N}$. By letting $m \rightarrow \infty$ in the last inequality, we get

$$\|f(x) - A_1(x)\| = 0, \quad (3.31)$$

for each $x \in E_1$, which implies that f is an additive mapping.

The reverse assertion is trivial because every additive mapping $f : E_1 \rightarrow E_2$ is a solution of the Jensen functional equation (see [\[10\]](#)). \square

ACKNOWLEDGMENT. The first author was supported by Korea Research Foundation grant KRF-DP0031.

REFERENCES

- [1] G. L. Forti, *The stability of homomorphisms and amenability, with applications to functional equations*, Abh. Math. Sem. Univ. Hamburg **57** (1987), 215–226.
- [2] ———, *Hyers-Ulam stability of functional equations in several variables*, Aequationes Math. **50** (1995), no. 1-2, 143–190.
- [3] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [4] D. H. Hyers, G. Isac, and T. M. Rassias, *Stability of Functional Equations in Several Variables*, Progress in Nonlinear Differential Equations and Their Applications, vol. 34, Birkhäuser Boston, Massachusetts, 1998.
- [5] D. H. Hyers and T. M. Rassias, *Approximate homomorphisms*, Aequationes Math. **44** (1992), no. 2-3, 125–153.
- [6] S.-M. Jung, *Hyers-Ulam-Rassias stability of functional equations*, Dynam. Systems Appl. **6** (1997), no. 4, 541–565.
- [7] ———, *Hyers-Ulam-Rassias stability of Jensen's equation and its application*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3137–3143.
- [8] Z. Kominek, *On a local stability of the Jensen functional equation*, Demonstratio Math. **22** (1989), no. 2, 499–507.
- [9] L. Losonczi, *On the stability of Hosszú's functional equation*, Results Math. **29** (1996), no. 3-4, 305–310.
- [10] J. C. Parnami and H. L. Vasudeva, *On Jensen's functional equation*, Aequationes Math. **43** (1992), no. 2-3, 211–218.
- [11] T. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), no. 2, 297–300.
- [12] F. Skof, *Local properties and approximation of operators*, Rend. Sem. Mat. Fis. Milano **53** (1983), 113–129 (Italian).

- [13] ———, *On the approximation of locally δ -additive mappings*, Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **117** (1983), no. 4-6, 377-389 (Italian).
- [14] S. M. Ulam, *Problems in Modern Mathematics*, John Wiley & Sons, New York, 1964.

Soon-Mo Jung: Mathematics Section, College of Science and Technology, Hong-Ik University, 339-701 Chochiwon, Korea

E-mail address: smjung@wow.hongik.ac.kr

Byungbae Kim: Mathematics Section, College of Science and Technology, Hong-Ik University, 339-701 Chochiwon, Korea

E-mail address: bkim@wow.hongik.ac.kr

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be