

THE DILATATION INVARIANT IN THE HOMOTOPY OF SPHERES

SENLIN XU and JINCHI LV

Received 15 May 2002

We obtain some results on the filtration invariant in the homotopy of spheres. Then by the relation between the dilatation invariant and filtration invariant, we get the corresponding results on the dilatation invariant in the homotopy of spheres.

2000 Mathematics Subject Classification: 55Q40, 57R35.

1. Introduction. Let S^m , respectively S^n , denote the standard unit sphere in Euclidean space \mathbb{R}^{m+1} , respectively \mathbb{R}^{n+1} , and let $f : S^m \rightarrow S^n$ be a smooth map. Following Olivier [5], we define $\delta(f)$, the dilatation of f , by

$$\delta(f) = \sup_{|X|=1} |df(X)|, \quad (1.1)$$

where X ranges over all unit tangent vectors and $|\cdot|$ is a Euclidean metric. Moreover, if $\alpha \in \pi_m(S^n)$, we define $\delta(\alpha)$, the dilatation invariant of α , to be the real number

$$\delta(\alpha) = \inf_f \delta(f), \quad (1.2)$$

where $f : S^m \rightarrow S^n$ ranges all smooth representatives f of α . It has been conjectured by Olivier [5] that $\delta(\alpha)$ is always an integer, which is still an open problem.

The filtration invariant of $\alpha \in \pi_m(S^n)$, $\text{filt}(\alpha)$, is the integer originally defined by James [2] in connection with his study of the reduced product construction. The definition is as follows [6]. For $p, q \in S^n$, $n \geq 2$, let $\Omega^* = \Omega^*(S^n; p, q)$ be the space of all continuous paths $\omega : [0, 1] \rightarrow S^n$ with $\omega(0) = p$, $\omega(1) = q$, equipped with the compact-open topology. As well known, if p and q are not conjugate on S^n , that is, $q \neq p, p'$ where p' denotes the antipode of p , then Ω^* admits a CW-decomposition of the form

$$\Omega^* \simeq S^{n-1} \cup e^{2(n-1)} \cup \dots \cup e^{r(n-1)} \cup \dots. \quad (1.3)$$

Let $\Omega_{r(n-1)}^* \simeq S^{n-1} \cup e^{2(n-1)} \cup \dots \cup e^{r(n-1)}$ be its $r(n-1)$ -skeleton. This allows us to define a filtration on the homotopy group $\pi_m(S^n)$ as follows. For any

$\alpha \in \pi_m(S^n)$, $\text{filt}(\alpha)$ is the least integer $r \geq 0$ for which the adjoint of α , $\text{ad}(\alpha) \in \pi_{m-1}(\Omega^*)$, lies in the image of the naturally induced map

$$\pi_{m-1}(\Omega_{r(n-1)}^*) \rightarrow \pi_{m-1}(\Omega^*). \quad (1.4)$$

One important fact is that $\text{filt}(\alpha) = 0$ if and only if $\alpha = 0$ and $\text{filt}(\alpha) \leq 1$ if and only if α is a suspension element.

The composition problem is very interesting. As we all know, $\text{filt}(\cdot)$ and $\delta(\cdot)$ are “submultiplicative,” that is,

$$\text{filt}(\alpha \circ \beta) \leq \text{filt}(\alpha) \cdot \text{filt}(\beta), \quad \delta(\alpha \circ \beta) \leq \delta(\alpha) \cdot \delta(\beta). \quad (1.5)$$

The former is verified in [2] and the latter is obvious. If $\alpha \in \pi_{2n-1}(S^n)$ is an element of Hopf invariant unity, then we have $\text{filt}(\alpha \circ \beta) = 2 \text{filt}(\beta) = \text{filt}(\alpha) \cdot \text{filt}(\beta)$ from [2]. With the same assumption, it has been conjectured by Roitberg [6] that $\delta(\alpha \circ \beta) = \delta(\alpha) \cdot \delta(\beta)$. But, unfortunately, this conjecture is not true even for the Hopf homotopy class.

In [6], Roitberg got a relation between the dilatation and filtration invariants in the homotopy of spheres by the fundamental theorem of Morse theory. In this note, we want to exploit more of the dilatation invariant. First, we get the following theorem, which is a generalization of [7, Theorem 1].

THEOREM 1.1. *Let $m > n \geq 2$, $f : S^m \rightarrow S^n$ be a smooth map and $\lambda = \max_{x \in S^n} d(f(x), f(-x))$, where d is the usual unit sphere metric. If $\text{filt}([f]) = r$, then $\delta(f) \geq 2[(r+1)/2] + (1+(-1)^r)\lambda/2\pi$.*

2. Proof of Theorem 1.1. Let $\Omega = \Omega(S^n; p, q)$ be the space of all piecewise smooth paths $\omega : [0, 1] \rightarrow S^n$ with $\omega(0) = p$, $\omega(1) = q$, equipped with the topology induced by the metric d^* defined in Milnor [4], let $E : \Omega \rightarrow \mathbb{R}$ be the energy function defined by $E(\omega) = \int_0^1 |d\omega/dt|^2 dt$, and set

$$\Omega^c = \Omega^c(S^n; p, q) = E^{-1}([0, c^2]), \quad c \geq 0. \quad (2.1)$$

It is well known [4, Theorem 17.1] that the inclusion $i : \Omega \rightarrow \Omega^*$ is a homotopy equivalence. Naturally, f induces a map

$$\tilde{f} : S^{m-1} = \Omega^\pi(S^m; x, -x) \rightarrow \Omega^{\delta(f)\pi} = \Omega^{\delta(f)\pi}(S^n; f(x), f(-x)) \quad (2.2)$$

for any $x \in S^m$, and the composition

$$i \circ j \circ \tilde{f} : S^{m-1} \rightarrow \Omega^*, \quad (2.3)$$

$j : \Omega^{\delta(f)\pi} \rightarrow \Omega$ the inclusion, is precisely the adjoint of f . The proof then follows from a study of the homotopy type of $\Omega^{\delta(f)\pi}$ for suitable choice of $x \in S^m$.

The Borsuk-Ulam theorem guarantees the existence of a point $x \in S^m$ such that $f(x) = f(-x)$. So, $\min_{x \in S^m} d(f(x), f(-x)) = 0$. We first assume that $\lambda > 0$. Suppose that $\delta(f) < 2\lceil (r+1)/2 \rceil + (-1)^r \lambda / \pi$. Let μ be a number such that $0 < \mu < \lambda \leq \pi$ and

$$\delta(f) < 2\left\lceil \frac{r+1}{2} \right\rceil + (-1)^r \mu / \pi. \tag{2.4}$$

Thus, by the continuity of d , there is a point $x \in S^m$ such that

$$d(f(x), f(-x)) = \mu. \tag{2.5}$$

From (2.4), we have $\delta(f)\pi < 2\lceil (r+1)/2 \rceil \pi + (-1)^r \mu$. Hence, by (2.5) and the fundamental theorem of Morse theory [4, Theorem 17.3 and Corollary 17.4], we deduce the following commutative diagram

$$\begin{array}{ccc} \Omega^{\delta(f)\pi} & \xrightarrow{j} & \Omega \\ \downarrow i' & & \downarrow i \\ \Omega_{(r-1)(n-1)}^* & \xrightarrow{j'} & \Omega^* \end{array} \tag{2.6}$$

where j' is the inclusion and i' is a homotopy equivalence. The conclusion $\text{filt}([f]) \leq r - 1$ then follows from (2.3), which is a contradiction. Furthermore, when r is odd, we can get $\delta(f) \geq r + 1$ by letting $\mu \rightarrow 0^+$. Then, we consider the case that $\lambda = 0$. For any $\epsilon > 0$, we can choose a suitable C^1 -approximation g of f such that $[g] = [f]$, $|\delta(g) - \delta(f)| < \epsilon$, and $\lambda(g) < \epsilon$. Letting $\epsilon \rightarrow 0^+$, we can get the conclusions.

REMARK 2.1. Noting that if $\text{filt}(\alpha) = r$ is even and $\inf_f \lambda(f) = \mu > 0$, then we have $\delta(\alpha) \geq r + \mu / \pi$. If $\delta(\alpha)$ is an integer, then we could get $\delta(\alpha) \geq r + 1$. The case that $\mu = \pi$ is very interesting since if true we can get $\delta(\alpha) \geq r + 1$. When m is even and $2\alpha \neq 0$, Lawson Jr. [3] showed that for any representative f of α , there exists a pair of antipodal points mapped by f into an antipodal pair, which had been contained in [7, Theorem 1].

COROLLARY 2.2. *If $\alpha \in \pi_m(S^2)$ with $m > 2$ and $2\alpha \neq 0$, then $\delta(\alpha) \geq 4$.*

PROOF. According to [1], $\text{filt}(\alpha) > 3$; so $\text{filt}(\alpha) = r \geq 4$. Applying Theorem 1.1, we have

$$\delta(f) \geq 2\left\lceil \frac{r+1}{2} \right\rceil + \frac{(1+(-1)^r)\lambda}{2\pi} \geq 4 \tag{2.7}$$

for any smooth representative f of α . Thus, $\delta(\alpha) \geq 4$. □

3. The case of S^3 . In this section, we consider the filtration of any element in $\pi_{3+k}(S^3; p)$ for $k = 2p - 3, 2(i+1)(p-1) - 2$, and $2(i+1)(p-1) - 1$,

$i = 1, \dots, p-1$, where p is an odd prime and $\pi_m(S^n; p)$ represents the p -primary component of the Abelian group $\pi_m(S^n)$. From [8, Theorem 13.4], we have the following two propositions.

PROPOSITION 3.1. *Let p be an odd prime, then $\pi_{3+k}(S^3; p)$ is \mathbb{Z}_p for $k = 2p-3, 2(i+1)(p-1)-2$, and $2(i+1)(p-1)-1, i = 1, \dots, p-1$, and is 0 otherwise for $k < 2(p+1)(p-1)-3$.*

PROPOSITION 3.2. *Let p be an odd prime, then $\pi_{2mp-1+k}(S^{2mp-1}; p)$ is \mathbb{Z}_p for $k = 2i(p-1)-1, 1 \leq i < p, m \geq 1$, and is 0 otherwise for $k < 2p(p-1)-2, m \geq 1$.*

We identify S^m with the suspension of S^{m-1} in the usual way. Then, we have an isomorphism $\phi : \pi_m(S^n) \rightarrow \pi_{m-1}(S_\infty^{n-1})$ (see [2]). For any element α in $\pi_m(S^n)$ of order q which is not necessarily a prime, $\phi(\alpha)$ is also an element of order q in $\pi_{m-1}(S_\infty^{n-1})$. Let $\text{filt}(\alpha) = r$, then $\phi(\alpha)$ lies in the image of the natural homomorphism $\pi_{m-1}(S_r^{n-1}) \rightarrow \pi_{m-1}(S_\infty^{n-1})$. When we consider the natural homomorphisms

$$\pi_{m-1}(S_r^{n-1}) \rightarrow \pi_m(S_k^{n-1}) \rightarrow \pi_{m-1}(S_\infty^{n-1}), \quad (3.1)$$

we know that $\phi(\alpha)$ is also an element of order q in $\pi_{m-1}(S_k^{n-1})$ for $k \geq r$. So, we have the following proposition.

PROPOSITION 3.3. *If $\pi_{m-1}(S_k^{n-1}; p) = 0$, then $\text{filt}(\alpha) \geq k+1$ for any nonzero element $\alpha \in \pi_m(S^n; p)$.*

As we all know, $\pi_n(S^n) \cong \mathbb{Z}$, so $\pi_{2+k}(S^{2p-1}; p) = 0$ for $k = 2p-3$. By [Proposition 3.2](#), we have

$$\pi_{2+k}(S^{2p-1}; p) = 0 \quad \text{for } k = 2(i+1)(p-1)-1, i = 1, \dots, p-1. \quad (3.2)$$

Thus, by the isomorphism $\pi_i(S_{p-1}^2; p) \rightarrow \pi_i(S^{2p-1}; p)$ for $i \geq 2$ and [Proposition 3.3](#), we get the following theorem.

THEOREM 3.4. *The elements of order p in $\pi_{3+k}(S^3)$ are of filtration at least p for $k = 2p-3$ and $2(i+1)(p-1)-1, i = 1, \dots, p-1$.*

By [Theorems 1.1](#) and [3.4](#), we can easily get the following corollary.

COROLLARY 3.5. *The elements of order p in $\pi_{3+k}(S^3)$ are of dilatation at least $p+1$ for $k = 2p-3$ and $2(i+1)(p-1)-1, i = 1, \dots, p-1$.*

COROLLARY 3.6. *If α is an element of order p in $\pi_{3+k}(S^3)$ as in [Corollary 3.5](#) and $\gamma \in \pi_3(S^2)$ is the Hopf class, then $\delta(\gamma \circ \alpha) \geq 2p+1$.*

PROOF. According to [2], $\text{filt}(\gamma \circ \alpha) = 2 \text{filt}(\alpha)$ since γ has Hopf invariant one. Noting that the composition is a homomorphism, the order of $\gamma \circ \alpha$ is

then 1 or p as p is prime. As $\text{filt}(\alpha) \geq p$ by [Theorem 3.4](#), we have $\text{filt}(\gamma \circ \alpha) = 2\text{filt}(\alpha) \geq 2p > 0$, so $\gamma \circ \alpha \neq 0$. Thus, the order of $\gamma \circ \alpha$ is p . It follows immediately from [Remark 2.1](#) that $\delta(\gamma \circ \alpha) \geq 2p + 1$ since $3 + k$ is even and p is an odd prime. □

From [[8](#), Theorem 13.4], we know that $\pi_{2p-3+k}(S^{2p-3}; p)$ is \mathbb{Z}_p for $k = 2(p - 1)^2 - 2$ and $2i(p - 1) - 1, i = 1, \dots, p - 1$, and is 0 otherwise for $k < 2p(p - 1) - 2$. So we have

$$\pi_{2+k}(S^{2p-3}; p) = 0 \quad \text{for } k = 2(i + 1)(p - 1) - 2, i = 1, \dots, p - 1. \tag{3.3}$$

Thus, by the isomorphism $\pi_i(S_{p-2}^2; p) \rightarrow \pi_i(S^{2p-3}; p)$ and [Proposition 3.3](#), we get the following proposition.

PROPOSITION 3.7. *The elements of order p in $\pi_{3+k}(S^3)$ are of filtration at least $p - 1$ for $k = 2(i + 1)(p - 1) - 2, i = 1, \dots, p - 1$.*

By [Theorem 1.1](#) and [Proposition 3.7](#), we can easily get the following corollary.

COROLLARY 3.8. *The elements of order p in $\pi_{3+k}(S^3)$ are of dilatation at least $p - 1$ for $k = 2(i + 1)(p - 1) - 2, i = 1, \dots, p - 1$.*

COROLLARY 3.9. *If α is an element of order p in $\pi_{3+k}(S^3)$ as in [Corollary 3.8](#) and $\gamma \in \pi_3(S^2)$ is the Hopf class, then $\delta(\gamma \circ \alpha) \geq 2(p - 1)$.*

PROOF. According to [[2](#)], $\text{filt}(\gamma \circ \alpha) = 2\text{filt}(\alpha)$ since γ has Hopf invariant one. As $\text{filt}(\alpha) \geq p - 1$ by [Proposition 3.7](#), it follows immediately from [Theorem 1.1](#) that $\delta(\gamma \circ \alpha) \geq 2(p - 1)$. □

4. Further results on the case that $\pi_m(S^n; p) = \mathbb{Z}_p$. Let p be a prime. We consider the case that $\pi_m(S^n; p) = \mathbb{Z}_p$. For any element α of order p in $\pi_m(S^n)$, we denote by $\lambda(\alpha)$ the least integer k such that $\pi_{m-1}(S_k^{n-1}; p) \neq 0$, then we have

$$\pi_{m-1}(S_{\lambda(\alpha)-1}^{n-1}; p) = 0, \quad \pi_{m-1}(S_{\lambda(\alpha)}^{n-1}; p) = \mathbb{Z}_p. \tag{4.1}$$

Thus by [Proposition 3.3](#), we have

$$\text{filt}(\alpha) \geq \lambda(\alpha). \tag{4.2}$$

Let β be an element of order p in $\pi_{m-1}(S_{\lambda(\alpha)}^{n-1})$, then β can be naturally regarded as an element in $\pi_{m-1}(S_{\infty}^{n-1})$ with order p . Then, $\phi^{-1}(\beta)$ is an element of order p in $\pi_m(S^n)$. Since $\pi_m(S^n; p) = \mathbb{Z}_p$, we have

$$\pi_m(S^n; p) = \langle \phi^{-1}(\beta) \rangle. \tag{4.3}$$

So, $\alpha = k\phi^{-1}(\beta)$ for some $1 \leq k \leq p-1$. Thus, $\alpha = k\phi^{-1}(\beta) = \phi^{-1}(k\beta)$ and $\phi(\alpha) = k\beta \in \pi_{m-1}(S_{\lambda(\alpha)}^{n-1})$. So we have

$$\text{filt}(\alpha) \leq \lambda(\alpha). \quad (4.4)$$

By (4.2) and (4.4), we get the following theorem.

THEOREM 4.1. *If $\pi_m(S^n; p) = \mathbb{Z}_p$, then $\text{filt}(\alpha) = \lambda(\alpha)$ for any element α of order p in $\pi_m(S^n)$.*

By Proposition 3.2, we have

$$\pi_{2+k}(S^{2p-1}; p) = \mathbb{Z}_p \quad \text{for } k = 2(i+1)(p-1) - 2, \quad i = 1, \dots, p-1. \quad (4.5)$$

So we have $\lambda(\alpha) = p-1$ for any element α of order p in $\pi_{3+k}(S^3)$ where $k = 2(i+1)(p-1) - 2, i = 1, \dots, p-1$. Thus by Theorem 4.1, we get the following theorem.

THEOREM 4.2. *The elements of order p in $\pi_{3+k}(S^3)$ are of filtration $p-1$ for $k = 2(i+1)(p-1) - 2, i = 1, \dots, p-1$.*

REFERENCES

- [1] I. M. James, *The suspension triad of a sphere*, Ann. of Math. (2) **63** (1956), 407-429.
- [2] ———, *Filtration of the homotopy groups of spheres*, Quart. J. Math. Oxford Ser. (2) **9** (1958), 301-309.
- [3] H. B. Lawson Jr., *Stretching phenomena in mappings of spheres*, Proc. Amer. Math. Soc. **19** (1968), 433-435.
- [4] J. W. Milnor, *Morse Theory*, Annals of Mathematics Studies, no. 51, Princeton University Press, New Jersey, 1963.
- [5] R. Olivier, *Über die Dehnung von Sphärenabbildungen*, Invent. Math. **1** (1966), 380-390 (German).
- [6] J. Roitberg, *Dilatation phenomena in the homotopy groups of spheres*, Advances in Math. **15** (1975), 198-206.
- [7] ———, *The dilatation and filtration invariants in the homotopy of spheres*, Advances in Math. **20** (1976), no. 2, 280-284.
- [8] H. Toda, *Composition Methods in Homotopy Groups of Spheres*, Annals of Mathematics Studies, no. 49, Princeton University Press, New Jersey, 1962.

Senlin Xu: Department of Mathematics, The University of Science and Technology of China, Hefei 230026, China

Jinchi Lv: Department of Mathematics, The University of Science and Technology of China, Hefei 230026, China

E-mail address: lvxi@mail.ustc.edu.cn

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from “Qualitative Theory of Differential Equations,” allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk