Internat. J. Math. & Math. Sci.
VOL. 21 NO. 1 (1998) 159-164

L_1 SPACES FAIL A CERTAIN APPROXIMATIVE PROPERTY

AREF KAMAL
Department of Mathematics and Computer Sciences
U A.E University
P.O Box 17551
Al-Ain, UNITED ARAB EMIRATES

(Received May 4, 1994)

ABSTRACT. In this paper the author studies some cases of Banach space that does not have the property P_1. He shows that if $X = \ell_1$ or $L_1(\mu)$ for some non-purely atomic measure μ, then X does not have the property P_1. He also shows that if $X = \ell_\infty$ or $C(Q)$ for some infinite compact Hausdorff space Q, then X^* does not have the property P_1.

KEY WORDS AND PHRASES: Property P_1, classical Banach spaces ℓ_1, $L_1(\mu)$ ℓ_1^n, compact width
1991 AMS SUBJECT CLASSIFICATION CODES: 41A65.

1. INTRODUCTION

The Banach space X is said to have the property P_1, if for each $\in > 0$ and each $r > 0$, there is $\delta > 0$, such that for each x and y in X, there is $z \in \overline{B(x, \in)}$ satisfying that for each θ with $0 < \theta < \delta$

$$B(x, r+\delta) \cap B(y, r+\theta) \subseteq B(z, r+\theta)$$

where $B(x,r)$ is the open ball of radius r and centered at x, and $\overline{B(x,r)}$ is its clouser u

The property P_1 plays an important role in approximation theory, and many authors used it This property appears in approximation by compact operators, simultaneous approximation and other areas (see for example Roversi [1], Lau [2], Mach [3] and Kamal [4]). Mach [3] showed that if X is uniformly convex then it has the property P_1 [3], and that if $X = C(Q)$, or $X = B(Q)$ then X has the property P_1 [4]. Mach [4, page 259] asked if the space $L_1(\mu)$ has the property P_1

In this paper the author studies some cases of normed linear space X, for which X does not have the property P_1 In section 2, it is shown that if $X = \ell_1$ then X does not have the property P_1, and in section 3, it is shown that if μ is a non-purely atomic measure, then $L_1(\mu)$ does not have the property P_1 These two results give a negative answer for the question of Mach [4] In section 3, it will be shown also that if $X = (\ell_\infty)^*$, or $X = (C(Q))^*$, where Q is an infinite compact Hausdorff space, then X does not have the property P_1

In this paper ℓ_1 is the Banach space of all real sequences $x = \{x_i\}$ satisfying that $\Sigma |x_i| < \infty$, together with the norm $\|x\| = \Sigma |x_i|$ Also ℓ_1^n is the Banach space of all real n-tuples $x = (x_1, x_2, ..., x_n)$ together with the norm $\|x\| = \sum_{i=1}^{n} |x_i|$.

2. I_1 DOES NOT HAVE THE PROPERTY P_1

The proof of the fact that ℓ_1 does not have the property P_1 depends on the behavior of the property P_1 in ℓ_1^n In Lemma 2.3, it will be shown that if $\epsilon > 0$ is fixed, and δ_n corresponds to ϵ for $X = \ell_1^n$ in Lemma 2 1, then $\delta_n \to 0$ when $n \to \infty$, so using the fact that ℓ_1^n is a norm-one-complemented subspace of ℓ_1, it will be shown in Theorem 2.4, that ℓ_1 does not have the property P_1.

LEMMA 2.1. If the Banach space X has the property P_1 then for each $\epsilon > 0$, there is $\delta > 0$ such that for each $y \in X$, there is $z \in \overline{B(0,\epsilon)}$ such that if $0 < \theta < \delta$ then

$$B(0, 1+\delta+\theta) \cap B(y, 1+\theta) \subseteq B(z, 1+\theta).$$

PROOF. Let $r = 1$ and let $\epsilon > 0$ be given By the definition of the property P_1 there is $\delta > 0$ such that for each x and y in X, there is $z \in B(x,\epsilon)$ satisfying the following; for each θ' such that $0 < \theta' < \delta'$

$$B(x, 1+\delta') \cap B(y, 1+\theta') \subseteq B(z, 1+\theta').$$

Let $x = 0$ and $\delta = 1/2\,\delta'$, then for all θ satisfying $0 < \theta < \delta$;

$$B(0, 1+\delta+\theta) \cap B(y, 1+\theta) \subseteq B(0, 1+\delta') \cap B(y, 1+\theta) \subseteq B(z, 1+\theta).$$

LEMMA 2.2. Let $n \geq 3$ be a positive integer, let $\delta > 0$ be given and let $(z_1, ..., z_n)$ be an n-tuple of real numbers

If $\sum_{i=1}^{n} z_i \geq \delta$, and for each $i \leq n-1$

$$z_1 + ... + z_{i-1} - z_i + z_{i+1} + ... + z_n \leq -\delta,$$

then $\sum_{i=1}^{n} |z_i| \geq (2n-3)\delta$.

PROOF. For each $i = 1, 2, ..., n$, let $y_i = z_1 + ... + z_{i-1} - z_i + z_{i+1} + ... + z_n$, then $\sum_{i=1}^{n} y_i = (n-2)\sum_{i=1}^{n} z_i$, thus

$$y_n = (n-2)\sum_{i=1}^{n} z_i - \sum_{i=1}^{n-1} y_i.$$

Therefore

$$\sum_{i=1}^{n} |z_i| \geq |y_n| \geq (n-2)\delta + (n-1)\delta = (2n-3)\delta.$$

LEMMA 2.3. Let $n \geq 3$ be a positive integer and let $\delta > 0$ be a given real number such that $(2n-5)\delta \leq 1$ Then the element $x_n = (\delta, \delta, ..., \delta, -(n-2)\delta)$ in ℓ_1^n satisfies the following conditions

(1) $\forall\theta$ such that $\theta > 0$

$$B(0, 1+\delta+\theta) \cap B(x_n, 1+\theta) \neq \phi.$$

(2) If $z \in \ell_1^n$ and for each θ with $0 < \theta < \delta$;

$$B(0, 1+\delta+\theta) \cap B(x_n, 1+\theta) \subseteq B(z, 1+\theta),$$

then $\|z\| \geq (2n-3)\delta$

PROOF. Let $\{e_i'\}_{i=1}^{n}$ be the standard basis in ℓ_1^n, that is $e_i' = (x_1^i, x_2^i, ..., x_n^i)$, where $x_j^i = 1$ if $i = j$ and $x_j^i = 0$ if $i \neq j$ and let

$$x_n = \delta\sum_{i=1}^{n-1} e_i' - (n-2)\delta e_n' = (\delta, \delta, ..., \delta, -(n-2)\delta) \in \ell_1^n.$$

Then $\|x_n\| = (n-1)\delta + (n-2)\delta = (2n-3)\delta \leq 1 + 2\delta \leq 2 + \delta$, therefore for each θ such that $0 < \theta < \delta$

$$B(0, 1+\delta+\theta) \cap B(x_n, 1+\theta) \neq \phi.$$

Assume that $z = (z_1, z_2, ..., z_n) \in \ell_1^n$ is such that for each θ with $0 < \theta < \delta$

$$B(0, 1+\delta+\theta) \cap B(x_n, 1+\theta) \subseteq B(z, 1+\theta).$$

It will be shown that

(1) $\sum_{i=1}^{n} z_i \geq \delta$, and

(2) for each $i \leq n-1$

$$z_1 + \ldots + z_{i-1} - z_i + z_{i+1} + \ldots + z_n \leq -\delta.$$

If these are true then by Lemma 2.2, $\|z\| = \sum_{i=1}^{n} |z_i| \geq (2n-3)\delta$

(1) Assume that $z_1 + \ldots\ldots + z_n < \delta$. Let

$$\begin{aligned} y &= \delta \sum_{i=1}^{n-2} e_i' + \frac{1+\delta}{2} e_{n-1}' + \left[\frac{1-(2n-5)}{2}\right] e_n' \\ &= \left(\delta, \ldots, \delta, \frac{1+\delta}{2}, \frac{1-(2n-5)}{2}\right) \in \ell_1^n. \end{aligned}$$

Then

$$\|y\| = (n-2)\delta + \frac{1+\delta}{2} + \frac{1-(2n-5)\delta}{2} = 1+\delta.$$

On the other hand

$$\begin{aligned} \|y - x_n\| &= \left|\frac{1+\delta}{2} - \delta\right| + \left|\frac{1-(2n-5)\delta}{2} + (n-2)\delta\right| \\ &= 1. \end{aligned}$$

Thus, for each θ such that $0 < \theta < \delta$,

$$y \in B(0, 1+\delta+\theta) \cap B(x_n, 1+\theta).$$

But

$$\begin{aligned} \|y - z\| = \sum_{i=1}^{n} |y_i - z_i| \geq \left|\sum_{i=1}^{n} y_i - \sum_{i=1}^{n} z_i\right| &= \left|1 + \delta - \sum_{i=1}^{n} z_i\right| \\ &= 1 + \left(\delta - \sum_{i=1}^{n} z_i\right) \\ &> 1, \end{aligned}$$

so for any $\theta < \left(\delta - \sum_{i=1}^{n} z_i\right)$, $y \notin B(z, 1+\theta)$.

(2) Assume that for a certain $i_0 \leq n-1$

$$z_1 + \ldots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \ldots + z_n > -\delta.$$

Let

$$y = \left(\frac{1+\delta}{2}\right) e_{i_0}' - \left(\frac{1+\delta}{2}\right) e_n' = \left(0, 0, \ldots, 0, \underset{i_0\text{-th term}}{\frac{1+\delta}{2}}, 0, \ldots, 0, -\left(\frac{1+\delta}{2}\right)\right) \in \ell_1^n.$$

Then

$$\|y\| = 1+\delta,$$

and

$$\begin{aligned}\|y - x_n\| &= (n-2)\delta + \left|\frac{1+\delta}{2} - \delta\right| + \left|\frac{1+\delta}{2} + (n-2)\delta\right| \\ &= (n-2)\delta + \frac{1-\delta}{2} + \frac{1-(2n-5)\delta}{2} \\ &= 1.\end{aligned}$$

Thus, for each θ such that $0 < \theta < \delta$, $y \in B(0, 1+\delta+\theta) \cap B(x_n, 1+\theta)$. But

$$\begin{aligned}\|y - z\| &= \sum_{i=1}^{n} |y_i - z_i| \\ &\geq |(y_1 + \ldots + y_{i_0-1} - y_{i_0} + y_{i_0+1} + \ldots + y_n) - (z_1 + \ldots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \ldots + z_n)| \\ &= |-1-\delta-(z_1 + \ldots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \ldots + z_n)| \\ &= |1 + [\delta + (z_1 + \ldots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \ldots + z_n)]| \\ &> 1.\end{aligned}$$

Thus, for some $\theta > 0$, $y \notin B(z, 1+\theta)$.

THEOREM 2.4. ℓ_1 does not have the property P_1

PROOF. It will be shown that for each $\delta > 0$, there is $x_\delta \in \ell_1$, such that if $z \in \ell_1$ and for all θ with $0 < \theta < \delta$ it is true that $B(0, 1+\delta+\theta) \cap B(x_\delta, 1+\theta) \subseteq B(z, 1+\theta)$, then $\|z\| > \frac{1}{2}$. Let $\{e_i\}_{i=1}^{\infty}$ be the standard basis in ℓ_1, and let $\delta > 0$ be given If $\delta > 1$ then for each $\theta > 0$

$$B(0, 1+1+\theta) \cap B(x_1, 1+\theta) \subseteq B(0, 1+\delta+\theta) \cap B(x_1, 1+\theta).$$

Thus one can take x_1 to be x_δ So without loss of generality one may assume that $\delta \leq 1$

Let $n \geq 3$ be a positive integer satisfying $(2n-5)\delta \leq 1$ and $(2n-3)\delta > \frac{1}{2}$, and let x_n be as in Lemma 2.3 Define

$$x_\delta = \delta \sum_{i=1}^{n-1} e_i - (n-2)\delta e_n = (\delta, \delta, \ldots, \delta, -(n-2)\delta, 0, 0, \ldots) \in \ell_1.$$

Then $\|x_\delta\| = \|x_n\| \leq 2 + \delta$, thus

$$B(0, 1+\delta+\theta) \cap B(x_\delta, 1+\theta) \neq \phi \quad \text{for} \quad 0 < \theta < \delta.$$

Let $P_n : \ell_1 \to \ell_1^n$ be the mapping defined by $P_n(\{x_i\}_{i=1}^{\infty}) = \{x_i\}_{i=1}^{n}$ By the construction of x_δ its image under P_n is the element x_n

Assume that for some $z \in \ell_1$

$$B(0, 1+\delta+\theta) \cap B(x_\delta, 1+\theta) \subseteq B(z, 1+\theta) \quad 0 < \theta < \delta,$$

then in ℓ_1^n

$$B(0, 1+\delta+\theta) \cap B(x_n, 1+\theta) \subseteq B(P_n(z), 1+\theta) \quad 0 < \theta < \delta,$$

Thus by Lemma 2.3 $\|P_n(z)\| \geq (2n-3)\delta > \frac{1}{2}$. Therefore

$$\|z\| \geq \|P_n(z)\| > \frac{1}{2}.$$

3. OTHER SPACES THAT DO NOT HAVE THE PROPERTY P_1

The subspace Y of X is called a norm-one-complemented subspace of X if there is a linear projection $P : X \to Y$ satisfying that $\|P\| = 1$. If A is a subset of X, and $x \in X$ then

$$d(x, A) = \inf\{\|x - y\|; y \in A\},$$

and if B is another subset of X, then the deviation of A from B is defined by

$$\delta(A, B) = \sup\{d(x, B); x \in A\}.$$

The compact width of A in X is defined by

$$a(A, X) = \inf\{\delta(A, K); K \text{ is a compact subset of } X\}.$$

The compact width is said to be attained if there is a compact subset K of X satisfying that $a(A, X) = \delta(A, K)$

In this section it will be shown that if $X = (C(Q))^*$, where Q is an infinite compact Hausdorff space, $X = (\ell_\infty)^*$, or $X = L_1(\mu)$ where μ is non-purely atomic measure, then X does not have the property P_1.

The proof of the following proposition is elementary

PROPOSITION 3.1. Let X be a Banach space that has the property P_1, and let Y be a closed subspace of X If Y is a norm-one-complemented subspace of X, then Y has the property P_1

COROLLARY 3.2. If μ is non-purely atomic measure then $L_1(\mu)$ does not have the property P_1

PROOF. By Feder [5, Theorem 2], $L_1[0,1]$ has a subset A for which the compact width $a(A, L_1[0,1])$ is not attained, thus by Kamal [6, Theorem 4.3] $L_1[0,1]$ does not have the property P_1, but by Lacy [7, sec 8], $L_1[0,1]$ is a norm-one-complemented subspace of $L_1(\mu)$, therefore by Proposition 3 1, $L_1(\mu)$ does not have the property P_1.

NOTE 3.3. Theorem 2.4 together with Corollary 3.2 give a negative answer to the question of Mach [4, page 259].

COROLLARY 3.4. If $X = \ell_\infty$ or $X = C(Q)$ for some compact infinite Hausdorff space Q Then X^* does not have the property P_1

PROOF. If $X = \ell_\infty$ then ℓ_1 is a norm-one-complemented subspace of X^*, and if $X = C(Q)$ then by Kamal [8, Lemma 3.2], ℓ_1 is a norm-one-complemented subspace of X^*, in both cases one concludes by Proposition 3.1 that X^* does not have the property P_1.

REFERENCES

[1] ROVERSI, M., Best approximation of bounded functions by continuous functions, *J. Approx. Theory* **41** (1984), 135-148

[2] LAU, K., Approximation by continuous vector valued functions, *Studia Math.* **68** (1980), 291-298

[3] MACH, J., Best simultaneous approximation of bounded functions with values in certain Banach spaces, *Math. Ann.* **240** (1979), 157-164.

[4] MACH, J., On the existence of best simultaneous approximation, *J. Approx. Theory* **25** (1979), 258-265.

[5] FERDER, M., On certain subset of $L_1[0,1]$ and non-existence of best approximation in some spaces of operators, *J. Approx. Theory* **29** (1980), 170-177.

[6] KAMAL, A., On certain diameters of bounded sets, *J. Port. Math.* **51** (1994), 321-333

[7] LACY, H.E., *The Isometric Theory of Classical Banach Spaces*, Springer-Verlag, 1974

[8] KAMAL, A., On proximinality and sets of operators III, Approximation by finite rank operators on spaces of continuous functions, *J. Approx. Theory* **47** (1986), 156-171.

Mathematical Problems in Engineering

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at http://www.hindawi.com/journals/mpe/guidelines.html. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at http://mts.hindawi.com/ according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br