

RESEARCH NOTES

THE SOLUTION OF A SINGULAR INTEGRAL EQUATION WITH SOME APPLICATIONS  
IN POTENTIAL THEORY

N.T. SHAWAGFEH

Department of mathematics  
University of Jordan  
Amman, JORDAN

(Received January 6, 1995 and in revised form December 9, 1996)

**ABSTRACT.** An analytical solution is derived for a singular integral equation which governs some two-dimensional potential boundary value problems in a region exterior to  $n$ -infinite co-axial circular strips. An application in electrostatics is discussed.

**KEY WORDS AND PHRASES:** Singular integral equation, electrostatics, Laplace's equation

**1991 AMS SUBJECT CLASSIFICATION CODES:** 31B10, 45B05, 78A30

1 INTRODUCTION

In this paper we derive a solution to the Fredholm singular integral equation

$$\sum_{k=0}^{n-1} \int_a^{\alpha} l(\phi) \left[ q + \log \left| \sin \left( \frac{1}{2} (\theta - \phi - \frac{2k\pi}{n}) \right) \right| \right] d\phi = f(\theta), \quad (1.1)$$

where  $q$  and  $\alpha$  are constants and  $f(\theta)$  is a differentiable function for  $|\theta - (2k\pi/n)| < \alpha, k = 0, \dots, n-1$ . This integral equation governs the solution of various two-dimensional Dirichlet and Neumann potential boundary value problems for the region consisting of the whole  $(r, \theta)$  plane outside the circular strips

$$r = a, \quad \left| \theta - \frac{2k\pi}{n} \right| < \alpha, k = 0, \dots, n-1, \quad |\alpha| < \frac{\pi}{n} \quad (1.2)$$

The previous investigations in potential problems of circular strips [1-8] were concerned mainly with the case of two strips, where Green's function approach leads to a singular integral equation with kernel

$q + \log \left| \sin \frac{1}{2} (\theta - \phi) \right|$ . Shail [3] transformed this equation into a well known Carleman type

A different technique has been used by Sampath and Jain [4] based on decoupling the equation into two singular equations which can be solved using eigenfunctions expansion method. The same technique has been used to solve various boundary value problems involving two circular strips [5-8].

In section 2 we use the approach of [4] to solve equation(1.1),and in section 3 we apply the results to solve laplace equation associated with Dirichlit and Neumann boundary conditions As an illustrative example we consider in section 4 the electrostatics problem in a region external to perfectly conducting n- circular strips Formulae for the surface charge density and concentration factor are derived,and these are believed to be new

**2.SOLUTION OF THE INTEGRAL EQUATION**

To obtain the solution of the integral equation (1 1) we first reduce it to simpler singular integral equations, so interchanging the sum and integral signs and using some properties of the function in the kernel we obtain

$$\int_{\alpha}^{\alpha} I(\phi) \left[ qn - \frac{n}{2} \log 2 + \frac{1}{2} \log \prod_{k=0}^{n-1} \left( 1 - \cos\left(\theta - \phi - \frac{2k\pi}{n}\right) \right) \right] d\phi = F(\theta), \tag{2 1}$$

Using the identity [9],

$$\prod_{k=0}^{n-1} \left( 1 - \cos\left(\theta - \phi - \frac{2k\pi}{n}\right) \right) = \frac{1}{2^{n-1}} (1 - \cos(n(\theta - \phi))), \tag{2 2}$$

in the equation (2 1) we get

$$\int_{\alpha}^{\alpha} I(\phi) \left[ Q + \log(1 - \cos n(\theta - \phi)) \right] d\phi = 2F(\theta), \quad |\theta| < \alpha \tag{2 3}$$

where  $Q = 2qn - (2n - 1) \log 2$

Now we write each of the functions  $I(\theta)$  and  $F(\theta)$  as

$$I(\theta) = I_e(\theta) + I_o(\theta) \tag{2 4a}$$

$$F(\theta) = F_e(\theta) + F_o(\theta) \tag{2 4b}$$

where the subscripts  $e$  and  $o$  stand for the even and odd parts respectively . Substiting (2 4) into (2 3) we obtain the following decoupled singular integral equations

$$\int_0^{\alpha} I_e(\phi) \left[ Q + \log \left\{ \cos(n\theta) - \cos(n\phi) \right\} \right] d\phi = F_e(\theta), \quad 0 < \theta < \alpha \tag{2 5}$$

$$\int_0^{\alpha} I_o(\phi) \log \left| \frac{\sin \frac{n}{2}(\phi - \theta)}{\sin \frac{n}{2}(\phi + \theta)} \right| d\phi = F_o(\theta) \tag{2 6}$$

In order to solve eq (2 5) we introduce the transformation

$$\cos(n\theta) = \sin^2\left(\frac{n\alpha}{2}\right) \cos x + \cos^2\left(\frac{n\alpha}{2}\right), \quad 0 < x < \pi \tag{2 7a}$$

$$\cos(n\phi) = \sin^2\left(\frac{n\alpha}{2}\right) \cos y + \cos^2\left(\frac{n\alpha}{2}\right), \quad 0 < y < \pi \tag{2 7b}$$

which transforms the equation into the form

$$\int_0^{\pi} H(y) \left[ R + \log(2|\cos x - \cos y|) \right] dy = h(x), \quad 0 < x < \pi \tag{2 8}$$

where

$$R = Q + \log \left[ \frac{1}{2} \sin^2 (n\alpha / 2) \right] \tag{2 9}$$

$$h(x) = F'_e(\theta) \tag{2 10}$$

$$\text{and } H(y) = \frac{\sin^2 \left( \frac{n\alpha}{2} \right) \sin y}{n \sin(n\phi)} I(\phi) \tag{2 11}$$

Expanding  $h(x)$  as Fourier cosine series

$$h(x) = \frac{1}{2} a_0 + \sum_{m=1}^{\infty} a_m \cos(mx) , \quad 0 < x < \pi \tag{2 12}$$

$$\text{where } a_m = \frac{2}{\pi} \int_0^{\pi} h(x) \cos(mx) dx , \quad m \geq 0 \tag{2 13}$$

and making use of the bilinear form [10]

$$\log(2|\cos x - \cos y|) = -2 \sum_{m=1}^{\infty} \frac{\cos(mx) \cos(my)}{m} , \quad 0 < x, y < \pi \tag{2 14}$$

, we find that the solution of (2 8) can be written in the form

$$H(y) = A_0 + \sum_{m=1}^{\infty} A_m \cos(my) , \tag{2 15}$$

where

$$A_0 = \frac{a_0}{2\pi R} , \quad A_m = -\frac{1}{\pi} a_m , \quad m \geq 1 \tag{2 16}$$

thus using (2 11) we find that the solution of (2 5) is given by

$$I_e(\phi) = \frac{\sqrt{2n} \cos(n\phi / 2)}{\sqrt{\cos(n\phi) - \cos(n\alpha)}} \left[ A_0 + \sum_{m=1}^{\infty} A_m \cos(my) \right] , \quad 0 < \phi < \alpha \tag{2 17}$$

where we have used the relation

$$\sin(y) = \sqrt{2} \csc^2 \left( \frac{n\alpha}{2} \right) \sin \left( \frac{n\phi}{2} \right) \sqrt{\cos(n\phi) - \cos(n\alpha)} , \tag{2 18}$$

Now to solve eq (2 6) we first differentiate both sides with respect to  $\theta$

$$\int_0^{\alpha} \frac{n I'_e(\phi) \sin(n\phi)}{\cos(n\phi) - \cos(n\theta)} d\phi = F'(\theta) , \tag{2 19}$$

Applying the transformation (2 7) into (2 19) reduces it to

$$\int_0^{\pi} \frac{G(y)}{\cos y - \cos x} dy = g(x) , \quad 0 < x < \pi \tag{2 20}$$

where

$$g(x) = F'(\theta) \tag{2 21}$$

$$G(y) = I'_e(\phi) \sin y \tag{2 22}$$

Expanding  $G(y)$  in Fourier cosine series

$$G(y) = B_0 + \sum_{m=1}^{\infty} B_m \cos(my) , \quad 0 < y < \pi \tag{2 23}$$

and noting the integral

$$\int_0^\pi \frac{\cos(my)}{\cos y - \cos x} dy = \pi \sin(mx) \csc(x), \quad 0 < x < \pi \tag{2.24}$$

which follows by putting  $w = e^i$  and integrating around the circle  $|w| = 1$  Then

$$\int_0^\pi \frac{G(y)}{\cos y - \cos x} dy = \pi \sum_{m=1}^\infty B_m \sin(mx) \csc(x), \tag{2.25}$$

Thus if we expand

$$g(x) \sin x = \sum_{m=1}^\infty C_m \sin(mx) \tag{2.26}$$

$$C_m = \frac{2}{\pi} \int_0^\pi g(x) \sin x \sin(mx) dx \tag{2.27}$$

it follows that

$$B_m = \frac{1}{\pi} C_m, \quad m \geq 1 \tag{2.28}$$

and hence

$$I_\alpha(\phi) = \frac{1}{\sin y} \left[ B_0 + \frac{1}{\pi} \sum_{m=1}^\infty C_m \cos(my) \right] \tag{2.29}$$

To evaluate the constant  $B_0$  we use the finiteness condition that  $G(y) \rightarrow 0$  as  $y \rightarrow 0$  this gives

$$B_0 = -\frac{1}{\pi} \sum_{m=1}^\infty C_m \tag{2.30}$$

Substituting (2.30) into (2.29) we find from (2.22) and the relation (2.18) that

$$I_\alpha(\phi) = \frac{1}{\pi \sin y} \sum_{m=1}^\infty C_m (1 - \cos my) \quad -\alpha < \phi < \alpha \tag{2.31}$$

Finally summing up the results we write the solution of eq (1.1) as

$$I(\phi) = \frac{\sqrt{2n \cos(n\phi/2)}}{\sqrt{\cos(n\phi) - \cos(n\alpha)}} \left[ A_0 + \sum_{m=1}^\infty A_m \cos(my) \right] - \frac{1}{\pi \sin y} \sum_{m=1}^\infty C_m (1 - \cos my), \quad -\alpha < \phi < \alpha \tag{2.32}$$

where

$$A_0 = \frac{a_0}{4\pi [qn + \log(2^{-n} |\sin(n\alpha/2)|)]} \tag{2.33}$$

$$A_m = -\frac{1}{\pi} a_m, \quad m \geq 1 \tag{2.34}$$

$$a_m = \frac{2}{\pi} \int_0^\pi h(x) \cos(mx) dx, \quad m \geq 0 \tag{2.35}$$

$$C_m = \frac{2}{\pi} \int_0^\pi g(x) \sin x \sin(mx) dx \tag{2.36}$$

$$h(x) = I'_\alpha(\theta), \quad g(x) = I'_\alpha(\theta) \tag{2.37}$$

**3.APPLICATION IN POTENTIAL THEORY**

The singular integral eq (1 1) governs the solutions of various Dirichlit and Neumann problems for two-dimensional Laplace equation in the domain D consisting of all the points (r, θ) lying ooutside the n-circular infinite strips

$$C_k \quad r = a \quad , \quad \left| \theta - \frac{2k\pi}{n} \right| < \alpha \quad , \quad k = 0, \dots, n-1 \quad , \quad (3 1)$$

These types of problems appear in different physical fields such as electrostatics, magnetostatics, steady state heat flow , and others

A Dirichlit problem We seek a function Φ(r, θ) that satisfies the follwing boundary value problem

$$\nabla^2 \Phi = 0 \quad (3 2)$$

$$\Phi(a, \theta) = f(\theta) \quad , \quad \left| \theta - \frac{2k\pi}{n} \right| < \alpha \quad , \quad k = 0, \dots, n-1 \quad (3 3)$$

$$\Phi(r, \theta) \sim -\frac{\Gamma}{2\pi} \log r \quad \text{as } r \rightarrow \infty \quad (3 4)$$

and are continuous across the arcs

$$C'_k \quad r = a \quad , \quad \frac{2k\pi}{n} + \alpha < \theta < 2(k+1)\frac{\pi}{n} - \alpha \quad (3 5)$$

The function f(θ) satisfies the symmetry condition

$$f\left(\theta + \frac{2k\pi}{n}\right) = f(\theta) \quad , \quad -\alpha < \theta < \alpha \quad (3 6)$$

$$\text{and } \Gamma = \sum_{k=0}^{n-1} \int_{C_k} \left[ \frac{\partial \Phi}{\partial r} \right]_{r=a} ds \quad (3 7)$$

Using Green's function approach and the symmetry in the problem the solution can be represented as

$$\Phi(r, \theta) = -\frac{a}{2\pi} \sum_{k=0}^{n-1} \int_{-\alpha}^{\alpha} I(\phi) \log \sqrt{r^2 + a^2 - 2ra \cos(\theta - \phi - \frac{2k\pi}{n})} d\phi \quad (3 8)$$

where the density function I(φ) is defined as

$$I(\phi) = \left[ \frac{\partial \Phi}{\partial r} \right]_{r=a}^{\phi, \phi'} \quad (3 9)$$

The boundary condition (3 3) leads to the following integral equation

$$(3 10)$$

which is eq (1 1) with  $q = \log(2a)$  ,  $F(\theta) = -\frac{2\pi}{a} f(\theta)$  , and hence it has the solution (2 32) with the

substitution of these values

B. Neumann problem We seek a function Ψ(r, θ) satisfying the boundary value problem

$$\nabla^2 \Psi = 0 \quad (3 11)$$

$$\Psi(a, \theta) = p(\theta) \quad , \quad \left| \theta - \frac{2k\pi}{n} \right| < \alpha \quad , \quad k = 0, \dots, n-1 \quad (3 12)$$

$$\Psi(r, \theta) \rightarrow 0, \text{ as } r \rightarrow \infty \tag{3 13}$$

$\Psi$ , and  $\frac{\partial \Psi}{\partial r}$  are continuous across the arcs  $C_k'$

The function  $P(\theta)$  satisfies the symmetry condition

$$P\left(\theta + \frac{2k\pi}{n}\right) = P(\theta), \quad -\alpha < \theta < \alpha \tag{3 14}$$

and the consistency condition

$$\int_{-\alpha}^{\alpha} P(\theta) d\theta = 0 \tag{3 15}$$

Green's function approach leads to the following representation

$$\Psi(r, \theta) = \frac{-a}{2\pi} \sum_{k=0}^{n-1} \int_{-\alpha}^{\alpha} J(\phi) \left[ \frac{\partial}{\partial \rho} \log \sqrt{r^2 + \rho^2 - 2r\rho \cos(\theta - \phi - \frac{2k\pi}{n})} \right]_{r', a} d\phi \tag{3 16}$$

$$\text{where } J(\phi) = [\Psi(r, \phi)]_{r', a}^{\alpha, -\alpha}, \quad -\alpha < \phi < \alpha. \tag{3 17}$$

Furthermore  $J(\theta)$  satisfies the edge condition

$$J(\pm\alpha) = 0 \tag{3 18}$$

Using the boundary condition (3 12) we obtain

$$\sum_{k=0}^{n-1} \int_{-\alpha}^{\alpha} J(\phi) \csc^2 \frac{1}{2} \left(\theta - \phi - \frac{2k\pi}{n}\right) d\phi = 8\pi a P(\theta) \tag{3 19}$$

Integrating by parts using the edge condition (3 18) we obtain

$$\sum_{k=0}^{n-1} \int_{-\alpha}^{\alpha} J'(\phi) \cot \frac{1}{2} \left(\theta - \phi - \frac{2k\pi}{n}\right) d\phi = -4\pi a P(\theta) \tag{3 20}$$

Upon integration with respect to  $\theta$  we get

$$\sum_{k=0}^{n-1} \int_{-\alpha}^{\alpha} J'(\phi) \log \left| \sin \frac{1}{2} \left(\theta - \phi - \frac{2k\pi}{n}\right) \right| d\phi = -2\pi a P'(\theta) + C' \tag{3 21}$$

where  $P'(\theta) = \int P(\theta) d\theta$  and  $C'$  is an arbitrary constant Eq (3 21) is special case of eq (1 1) with  $q = 0$

$P'(\theta) = C' - 2\pi a P(\theta)$ , and  $I(\phi) = J'(\phi)$  To determine the constant  $C'$  we use the edge condition (3 18)

Finally the density  $J(\theta)$  is given by

$$J(\theta) = \int_{-\alpha}^{\alpha} J'(\phi) d\phi, \quad -\alpha < \theta < \alpha \tag{3 22}$$

#### 4. ILLUSTRATIVE EXAMPLE

As an example to illustrate our results we consider the electrostatics potential problem of the n-equal infinite co-axial perfectly conducting strips in free space charged so that the total charge per height on each strip is unity. In cylindrical coordinates  $(r, \theta, z)$  let the strips be defined by

$$C_k: r = a, \quad \left| \theta - \frac{2k\pi}{n} \right| < \alpha, \quad k = 0, \dots, n-1, \quad -\infty \leq z \leq \infty \tag{4 1}$$

then the electrostatics potential satisfies the Dirichlet problem A, where on the boundary we assume that  $\Phi$  has the constant potential  $K$ . Thus

$$\Phi(r, \theta) = -a \sum_{k=0}^{n-1} \int_{-\alpha}^{\alpha} I(\phi) \log \sqrt{r^2 + a^2 - 2ra \cos(\theta - \phi - \frac{2k\pi}{n})} d\phi \quad (4.2)$$

and  $I(\phi)$  satisfy the integral equation (3.10) with  $f(\theta) = K$ , and hence

$$h(x) = I_c(\theta) = \frac{-K}{a}, \quad g(x) = I_o(\theta) = 0 \quad (4.3)$$

Substituting these values in (2.32) we find that

$$I(\phi) = \frac{-Kn \cos(n\phi/2)}{\sqrt{2\pi a} \log(a^n |\sin(n\alpha/2)| \sqrt{\cos(n\phi) - \cos(n\alpha)}), \quad -\alpha < \phi < \alpha \quad (4.4)$$

To find the value of  $K$  we use the condition that the total charge per unit height on each strip is unity, that is,

$$a \sum_{k=0}^{n-1} \int_{-\alpha}^{\alpha} I(\phi) d\phi = n, \quad \text{or} \quad \int_{-\alpha}^{\alpha} I(\phi) d\phi = \frac{1}{2a} \quad (4.5)$$

Inserting (4.4) into (4.5) and evaluating the resulting integral we find that

$$K = -\log(a^n |\sin(n\alpha/2)|) \quad (4.7)$$

and the charge density in this case is

$$I(\phi) = \frac{n \cos(n\phi/2)}{\sqrt{2\pi a} \sqrt{\cos(n\phi) - \cos(n\alpha)}}, \quad -\alpha < \phi < \alpha \quad (4.8)$$

while the charge concentration factor is given by

$$N = \lim_{\phi \rightarrow \alpha} (\alpha - \phi)^{-1/2} I(\phi) = \frac{1}{2a\pi} \sqrt{n \cot(n\alpha/2)} \quad (4.9)$$

All the above formulae are believed to be new. For  $n=4$  the formulae (4.6), (4.7), and (4.8) agree with those in [5].

and when  $\alpha \rightarrow \pi/n$  we recover the known results for a charged infinite cylinder when the total charge per unit height is  $n$ .

**ACKNOWLEDGMENT.** The present work is supported by the university of Jordan under project #356

#### REFERENCES

- [1] GAUTESSEN, A. K. & OLMESTEAD, W. E., On the solution of the integral equation for potential of two strips *SIAM J. Math. Anal.* **2** (1971), 293-306
- [2] GOEL, G. C. & JAIN, D. L., A note on electrostatic problem involving two strips, *J. Pure Appl. Math.* **7** (1976), 751-756
- [3] SHAIL, R., A class of singular integral equation with some application, *Int. J. Math. Edu. Tech.* **15** (1984), 359-374
- [4] SAMPATH, C. & JAIN, D. L., On solution of the integral equations for the potential problems of two

- circular strips, *Internat. J. Math. & Math.Sci.* **11** (1988) 751-762
- [5] SAMPATH,C & JAIN,D L , Some boundary value problems in electrostatics, *J. Math. Phy.Sci* **22** (1988)
- [6] JAIN,S & JAIN,D L , Diffraction of an H-polarized electromagnetic wave by two equal infinite circular strips, *Radio Sci* **24**(1989),443-454
- [7] SAMPATH,C & JAIN,D L , Some electrostatic problems of two equal co-axial circular strips, *J. Math.Phy. Sci* **25** (1991), 217-230
- [8] VARMA,S K & JAIN,D L , Diffraction of elastic P waves by two equal co-axial circular strips  
Eur J Mech A **11**(1992),157-168
- [9] GRADSHTEYN ,I S & RYZHIK,I M , *Tables of Integrals Series and Products*,Academic Press  
NewYork ,1965
- [10] KANWAL,R P , *Linear Integral Equations, Theory and Techniques* , Academic Press,NewYork  
(1971)

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

### Guest Editors

**Edson Denis Leonel**, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)