

ON COMMON FIXED AND PERIODIC POINTS OF COMMUTING FUNCTIONS

ALIASGHAR ALIKHANI-KOOPAEI

Mathematics Department
University of Isfahan
Isfahan, Iran

(Received March 8, 1996 and in revised form March 30, 1996)

ABSTRACT. It is known that two commuting continuous functions on an interval need not have a common fixed point. It is not known if such two functions have a common periodic point. In this paper we first give some results in this direction. We then define a new contractive condition, under which two continuous functions must have a unique common fixed point.

KEY WORDS AND PHRASES: FIXED POINTS, PERIODIC POINTS, COMMUTING FUNCTIONS, MINIMAL SETS.

1992 AMS SUBJECT CLASSIFICATION CODES: 26A16, 47H10.

1. INTRODUCTION.

For some time there was a rather well known conjecture that if f and g are continuous commuting selfmaps of $[0, 1]$ (i.e. $f(g(x)) = g(f(x))$), then they have a common fixed point. W.M. Boyce [1] and J.P. Huneke [2] answered this conjecture in the negative by constructing a pair of commuting continuous functions which have no common fixed point. It is easy to see that their pair of commuting continuous functions have a common periodic point. In fact for the Boyce's example zero is a common periodic point. Thus one may conjecture the following:

CONJECTURE 1.1. *If f and g are commuting continuous selfmaps of $[0, 1]$, then they must have a common periodic point.*

Even though we believe that the answer to the above conjecture is also negative, at present we are not able to construct a counterexample.

A.J. Schwartz [3] proved that if f and g are continuous functions of $[0, 1]$ into itself that

commute, then there always exists a point x such that $x = f(x) = g^n(x)$ for some positive integer n , under the additional assumption that f has a continuous derivative. In this paper we first give some results in this direction. Then we define a property P_1 and show that under this condition, two commuting continuous functions must have a unique common fixed point.

We begin with some preliminaries. The orbit of x under g (i.e. the set $\{g^k(x) : k \geq 0\}$) and its closure are denoted by $O(g, x)$ and $\overline{O(g, x)}$, respectively. The set of cluster points of $O(g, x)$ is denoted by $\omega(g, x)$. A subset Y of I is called invariant under g if $g(Y) \subseteq Y$. A closed, invariant, nonempty subset of I is called minimal if it contains no proper subset that is also closed, invariant and nonempty. A point x is called a recurrent point of g if x belongs to $\omega(g, x)$. Throughout f^n denotes the n fold composition of f with itself. The sets $P(f)$, $R(f)$ and $F(f)$ are the sets of periodic points of f , recurrent points of f , and the fixed points of f , respectively. We state some known facts concerning minimal sets. Every closed, invariant, nonempty subset of I contains a minimal set. If Y is a minimal set, then $Y \subseteq R(f)$. If Y is a minimal set, which is not the orbit of a periodic point, then Y is perfect. A minimal set is nowhere dense.

2. THE MAIN RESULTS.

THEOREM 2.1. *Let f and g be two commuting continuous selfmaps of the unit interval. If f and g have no common periodic points, then for any two positive integers m and n the set $A_{m,n} = \{x : f^n(x) = g^m(x)\}$ is uncountable.*

PROOF. First we show that $A_{m,n}$ is not empty. If for some m and n , $A_{m,n}$ is empty, the continuity of f and g permits us to assume without loss of generality, that

$$f^n(x) < g^m(x) \tag{2.1}$$

for all $x \in I$. Since $g^m(1) \leq 1$, the set $S = \{x \in I : g^m(x) \leq x\}$ is not empty. Thus, since S is closed, it has a minimum element c . Clearly, $c = g^m(c)$. Hence $f^n(c) = f^n(g^m(c)) = g^m(f^n(c))$, so that $f^n(c) \in S$. Consequently $f^n(c) \geq c = g^m(c)$. Since $f^n(c) \geq g^m(c)$ contradicts (2.1), the assertion that $A_{m,n}$ is empty is false. Now suppose that $x \in A_{m,n}$. Then $f^n(x) = g^m(x)$ hence, $f^n(f(x)) = f(g^m(x)) = g^m(f(x))$ and $f^n(g(x)) = g(f^n(x)) = g(g^m(x)) = g^m(g(x))$. Thus if $x \in A_{m,n}$, $\{f(x), g(x)\} \subseteq A_{m,n}$ and $f^{pn}(x) = g^{pm}(x)$ for each positive integer p . From this it follows that $O(g, x)$ is contained in $A_{m,n}$ whenever $x \in A_{m,n}$. The set $\overline{O(g, x)}$ is an invariant set under g . Suppose A_1 is a minimal set contained in $\overline{O(g, x)}$. Since f and g have no common periodic points, the set A_1 cannot be the orbit of a periodic point. Hence A_1 is a perfect set contained in $A_{m,n}$, implying that $A_{m,n}$ is uncountable.

COROLLARY 2.1. *Let f and g be two commuting continuous selfmaps of the unit interval. If for some positive integers m and n , the set $\{x : f^n(x) = g^m(x)\}$ is countable, then f and g have a common periodic point.*

Fixed points of contractive type mappings have been studied by a number of authors. B.E. Rhoades [4] has investigated a comparison of different kinds of contractive definitions in the literature. The following theorem is in this direction.

DEFINITION 2.1. *Let X be a compact metric space. The function $h : X \times X \rightarrow [0, \infty)$ is said to have property P_1 if it satisfies the following conditions:*

- (i): $h(x, y) = 0$ if and only if $x = y$,
- (ii): if $\lim_{n \rightarrow \infty} x_n = x_0$, $\lim_{n \rightarrow \infty} y_n = y_0$, and $\lim_{n \rightarrow \infty} h(x_n, y_n) = 0$,

then $x_0 = y_0$.

THEOREM 2.2. *Let f and g be selfmaps of the unit interval and let $h : I \times I \rightarrow [0, \infty)$ be a function having property P_1 . Suppose g is continuous on I and A is a nonempty closed g -invariant subset of $F(f)$. If there exists a real number a , $0 \leq a < 1$ such that for all x and y in $F(f)$, f and g satisfy the following inequality:*

$$h(fx, fy) \leq a \cdot \max\{h(gx, gy), h(gx, fx), h(gy, fy), h(gy, fx), h(fx, gy)\}, \tag{2.2}$$

then f and g have a unique common fixed point.

PROOF. First we show that any two functions f and g satisfying inequality (2.2), have at most one common fixed point. To see this, on the contrary, suppose q_1 and q_2 are two different common fixed points of f and g . Then using (2.2) we get

$$\begin{aligned} h(q_1, q_2) &= h(fq_1, fq_2) \\ &\leq a \cdot \max\{h(gq_1, gq_2), h(gq_1, fq_1), h(gq_2, fq_2), h(gq_2, fq_1), h(fq_1, gq_2)\}, \end{aligned}$$

implying

$$\begin{aligned} h(q_1, q_2) &\leq a \cdot \max\{h(q_1, q_2), h(q_1, q_1), h(q_2, q_2), h(q_2, q_1)\}, \\ &= a \cdot \max\{h(q_1, q_2), h(q_2, q_1)\}. \end{aligned}$$

In a similar manner we may show

$$h(q_2, q_1) \leq a \cdot \max\{h(q_1, q_2), h(q_2, q_1)\},$$

hence we shall have

$$\max\{h(q_1, q_2), h(q_2, q_1)\} \leq a \cdot \max\{h(q_1, q_2), h(q_2, q_1)\}.$$

This is impossible unless $\max\{h(q_1, q_2), h(q_2, q_1)\} = 0$, which implies that $h(q_1, q_2) = 0$, and hence $q_1 = q_2$, a contradiction. Now we show that f and g have a common fixed point. Without loss of generality we may assume that $A = F(f)$. It is clear that the set $B = \bigcap_{n=1}^{\infty} g^n(A)$ is a closed

nonempty subset of A , and that $g^2(B) = B$. If for some $y_0 \in B$, $f(y_0) = g(y_0)$, then y_0 is a common fixed point of f and g , so we may assume that $f(y) \neq g(y)$ for all $y \in B$. Let y_1 be an arbitrary point of B , and $x_2 \in B$ be such that $g^2(x_2) = y_1$. Since $g(y) \neq f(y)$ for all $y \in B$, if we let $x = y_1$ and $y_2 = g(x_2)$, then $g(y_2) = y_1 = f(y_1)$. Since $h(gy_2, fy_2) = h(fy_1, fy_2)$, from inequality (2.2) we have

$$\begin{aligned} h(fy_1, fy_2) &\leq a \cdot \max\{h(gy_1, gy_2), h(gy_1, fy_1), h(gy_2, fy_2), \\ &\quad h(gy_2, fy_1), h(fy_1, gy_2)\} \\ &= a \cdot \max\{h(gy_1, gy_2), h(gy_1, fy_1), h(gy_2, fy_2), 0\} \\ &= a \cdot \max\{h(gy_1, fy_1), h(gy_2, fy_2)\}, \end{aligned}$$

implying $h(gy_2, fy_2) \leq a \cdot h(gy_1, fy_1)$. If we define $\phi : B \rightarrow [0, \infty)$ by $\phi(x) = h(gx, fx)$, then we have $\phi(y_2) \leq a \cdot \phi(y_1)$. Suppose $x_3 \in B$ be such that $g^2(x_3) = y_2$. By taking $x = y_2$ and $y_3 = g(x_3)$, similarly we can show that $\phi(y_3) \leq a \cdot \phi(y_2) \leq a^2 \cdot \phi(y_1)$. By repeating this process we obtain a bounded sequence $\{y_n\}_{n=1}^{\infty}$ in B . Suppose $\{y_{n_k}\}_{k=1}^{\infty}$ is a convergent subsequence of $\{y_n\}_{n=1}^{\infty}$ and $\lim_{k \rightarrow \infty} y_{n_k} = y_0$. Since g is continuous on I ,

$$\begin{aligned} \lim_{k \rightarrow \infty} h(gy_{n_k}, y_{n_k}) &= \lim_{k \rightarrow \infty} h(gy_{n_k}, fy_{n_k}) \\ &= \lim_{k \rightarrow \infty} \phi(y_{n_k}) \\ &\leq \lim_{k \rightarrow \infty} a^{n_k-1} \cdot \phi(y_1) = 0. \end{aligned}$$

So $g(y_0) = y_0$, and $y_0 \in B \subseteq F(f)$, hence y_0 is a common fixed point of f and g .

REMARK 2.1. There is nothing in the proof of Theorem 2.2 that requires the underlying space to be the unit interval. However in our proof the compactness of the unit interval is used. Therefore one can let the underlying space to be a compact metric space. Also condition P_1 is not a necessary condition and it may be replaced with weaker conditions when f and g are nice. For example if h, f are nice enough that the function ϕ defined in Theorem 2.2 attains its minimum at some point $y_1 \in B$, then we can find a point $y_2 \in B$ such that $\phi(y_2) < \phi(y_1)$ a contradiction, so f and g must have a common fixed point. If we are not concerned about the uniqueness of the common fixed point, similar to the proof of Theorem 2.2 we have the following.

THEOREM 2.3. *Suppose f and g are two selfmaps of a compact metric space X and let $h : X \times X \rightarrow [0, \infty)$ be a function having property P_1 . Suppose also that g is continuous on X and A is a nonempty closed g -invariant subset of $F(f)$. If there exists a real number a , $0 \leq a < 1$ such that for all x and y in A , f and g satisfy the following inequality:*

$$\begin{aligned} h(fx, fy) &\leq a \cdot \max\{h(gx, gy), h(gx, fx), h(gy, fy), \\ &\quad h(gy, fx), h(fx, gy)\}, \end{aligned} \tag{2.3}$$

then f and g have a common fixed point.

The conclusion of Theorem 2.3 is not valid if inequality (2.3) is replaced with

$$h(fx, fy) \leq \max\{h(gx, gy), h(gx, fx), h(gy, fy), h(gy, fx), h(fx, gy)\}.$$

The following example illustrates this.

EXAMPLE 2.1. Let f and g be defined on the unit interval as follows:

$$g(x) = \begin{cases} (8x + 15)/24 & 0 \leq x \leq 3/4, \\ (-32x + 31)/8 & 3/4 < x \leq 13/16, \\ 2x - 1 & 13/16 < x \leq 7/8, \\ 3/4 & 7/8 < x \leq 1, \end{cases}$$

$$f(x) = \begin{cases} (8x + 39)/48 & 0 \leq x < 3/4, \\ 3/4 & x = 3/4, \\ (-32x + 39)/16 & 3/4 < x \leq 13/16, \\ x & 13/16 < x \leq 7/8, \\ 7/8 & 7/8 < x \leq 1. \end{cases}$$

Define $h(x, y) : I \times I \rightarrow [0, \infty)$ as $h(x, y) = \begin{cases} 0 & \text{if } x = y, \\ x + y & \text{if } x \neq y. \end{cases}$

One can easily check that h has property P_1 , $f(x) = \begin{cases} (1 + g(x))/2 & \text{if } x \neq 3/4, \\ 3/4 & \text{if } x = 3/4, \end{cases}$ and $F(f) = \{3/4\} \cup [13/16, 7/8]$. Let $A = \{3/4, 7/8\}$, then A is a closed g -invariant subset of $F(f)$ and for every $x \neq y$ in A we have

$$h(fx, fy) = \max\{h(gx, gy), h(gx, fx), h(gy, fy), h(gy, fx), h(fx, gy)\} \leq 13/8.$$

It is easy to see that f and g do not have a common fixed point.

THEOREM 2.4. Suppose f and g are two selfmaps of a compact metric space X with g continuous, and let $h : X \times X \rightarrow [0, \infty)$ is a function having property P_1 . If for all $x \neq y$ in X , f and g satisfy the following inequality:

$$h(fx, fy) < \max\{h(gx, gy), h(gx, fx), h(gy, fy), h(gy, fx), h(fx, gy)\}, \tag{2.4}$$

then one of the following holds:

- (i) either f and g have a common fixed point,

(ii) or every nonempty closed g -invariant subset of $F(f)$ contains a perfect minimal set B such that the functions $\phi_1(x) = h(gx, x)$ and $\phi_2(x) = h(x, gx)$, do not attain their minimum or maximum on B .

PROOF. From inequality (2.4) it follows that for each $x \in F(f)$ with $x \neq g(x)$, we have $h(gx, x) < h(g^2x, gx)$. This implies that $F(f)$ cannot contain any periodic point of g with period greater than 1. Let A be a nonempty closed g -invariant subset of $F(f)$, then it should contain a minimal set B . If this minimal set is finite, f and g have a common fixed point. Otherwise, B is a perfect set with $g(B) = B$. Since for each $x \in B$, $h(gx, x) < h(g^2x, gx)$, the functions ϕ_1 and ϕ_2 cannot attain their maximum or minimum on B .

LEMMA 2.1. Suppose $h : [0, 1] \times [0, 1] \rightarrow [0, \infty)$ is a lower semicontinuous function such that $h(x, y) = 0$ if and only if $x = y$. Then h satisfies property P_1 .

PROOF. Suppose $\lim_{n \rightarrow \infty} x_n = x_0$, $\lim_{n \rightarrow \infty} y_n = y_0$, and $\lim_{n \rightarrow \infty} h(x_n, y_n) = 0$. We need to show that $x_0 = y_0$. On the contrary suppose $x_0 \neq y_0$. Let $\delta = \text{dist}\{(x_0, y_0), \{(x, x) : x \in [0, 1]\}\}$, and $B_1 = \overline{B(x_0, \frac{\delta}{2})} = B[x_0, \frac{\delta}{2}]$, where $x_0 = (x_0, y_0)$. Let ϵ be an arbitrary positive number. Then there exists a positive integer \mathcal{N} such that for all $n \geq \mathcal{N}$, $(x_n, y_n) \in B_1$, and $|h(x_n, y_n)| < \epsilon$. Since h is lower semicontinuous and B_1 is compact, it attains its minimum at some point (s, t) in B_1 . Hence $|h(s, t)| = \text{minimum of } h(x, y) \text{ on } B_1 \leq |h(x_{\mathcal{N}}, y_{\mathcal{N}})| < \epsilon$, implying $h(s, t) = 0$. Thus $s = t$, a contradiction.

THEOREM 2.5. Let f and g be commuting selfmaps of $[0, 1]$ with g continuous. If A is a nonempty closed g -invariant subset of $F(f)$, then one of the following holds:

- (i) either $P(f) \cap P(g) \neq \emptyset$,
- (ii) or there is a perfect set $A_1 \subseteq (A \cap R(g) \cap P(f))$.

PROOF. Take $x_0 \in A$. For every positive integer n , $g^n(x_0) \in A$. Thus $O(g, x_0) \subseteq A$ implying $\omega(g, x_0) \subseteq \bar{A} = A$. Clearly $\omega(g, x_0)$ is nonempty and is invariant under g . Thus it contains a minimal g -invariant subset A_1 . Obviously $A_1 \subseteq (A \cap R(g))$. If A_1 is a finite set, then it is the orbit of a periodic point of g , implying $\emptyset \neq (P(g) \cap A) \subseteq P(f) \cap P(g)$. Otherwise A_1 is a perfect set contained in $A \cap R(g)$.

REMARK 2.2. As in the proof of Theorem 2.5, if f and g have no common periodic point then there exists a perfect set A_1 contained in $A \cap R(g) \cap P(f)$. Since f and g do not have common periodic points and, for a continuous function on a compact interval $\overline{P(f)} = \overline{P(g)}$ (see Coven and Hedlund [5]), we have $A_1 \subseteq A \cap \overline{R(g)} = A \cap \overline{P(g)} = A \cap [\overline{P(g)} \setminus P(g)]$. From this it follows that if $\overline{P(g)} \setminus P(g)$ does not contain a perfect set, then f and g must have a common periodic point. In particular, for two commuting continuous functions if either of $\overline{P(g)} \setminus P(g)$ or $\overline{P(f)} \setminus P(f)$ does not contain a perfect set, then f and g must have a common periodic point.

Consider the following ordering of positive integers:

1, 2, 4, 8, ..., $7 \cdot 8, 5 \cdot 8, 3 \cdot 8, \dots, 7 \cdot 4, 5 \cdot 4, 3 \cdot 4, \dots, 7 \cdot 2, 5 \cdot 2, 3 \cdot 2, \dots, 7, 5, 3$.

A.N. Sarkovskii has proven that if m is to the left of n in the above ordering and f has a periodic point of period n , then f must have a periodic point of period m (see Stefan [6]). Let $P_n(f) = \{x \in [0, 1] : f^n(x) = x\}$. The Sarkovskii's theorem immediately implies that if $P_1(f) = P_2(f)$, then $P_1(f) = P_n(f)$ for every $n > 1$. Suppose f and g are two commuting continuous selfmaps of the unit interval which do not have a common periodic point. We claim that f and g should have periodic points of all even orders. To see this, on the contrary suppose that $n = 2^k \cdot r$ (r odd, $k \geq 1$) and f has no periodic point of order n . By Sarkovskii's theorem $P_{2 \cdot r}(f) = P_r(f)$ implying $P_1(f^r) = P_2(f^r)$, hence $P_n(f^r) = P_1(f^r)$ for any $n > 1$. Thus we have $P(f^r) = \bigcup_{n=1}^{\infty} P_n(f^r) = P_1(f^r)$ which implies $\overline{P(f^r)} \setminus P(f^r) = \emptyset$. Since g commutes with f , it also commutes with f^r . Thus g and f^r should have a common periodic point which is also a common periodic point of f and g . We may interchange the roles of f and g . This implies that either f and g have a common periodic point or both have a rich orbit structure. By a rich orbit structure we mean that they have a homoclinic point and positive topological entropy (see Block [7]). Thus we have the following:

THEOREM 2.6. *Suppose f and g are two commuting continuous selfmaps of the unit interval. Then one of the following holds:*

- (i) *either $P(f) \cap P(g) \neq \emptyset$,*
- (ii) *or both f and g have periodic points of all even orders.*

ACKNOWLEDGMENT: I would like to express my sincere thanks to professor B.E. Rhoades for his valuable suggestions in improving the paper, especially on the comments that improved Theorems 2.2 and 2.3. This work also was completed while the author was spending his sabbatical at the University of North Carolina at Wilmington. The hospitality of its Department of Mathematics as well as the financial support of Isfahan University are greatly acknowledged.

AUTHOR'S PRESENT ADDRESS: Mathematical Sciences Department, The University of North Carolina at Wilmington, Wilmington, NC 28403-3297.

REFERENCES

1. BOYCE, W.M., Commuting functions with no common fixed point, Trans. Amer. Math. Soc. 137(1969), 77-92.
2. HUNEKE, J.P., On common fixed points of commuting functions on an interval, Trans. Amer. Math. Soc. 139(1969), 371-381.
3. SCHWARTZ, A.J., Common periodic points of commuting functions, Michigan Math. J. 12(1965), 353-355.

4. RHOADES, B.E., A comparison of various definitions of contractive mappings, *Trans. Amer. Math. Soc.* 226(1977), 257-290.
5. COVEN, E.M., and HEDLUND, G. A., $\overline{P} = \overline{R}$ for maps of the interval, *Proc. Amer. Math. Soc.* 79(1980), 316-318.
6. STEFAN, P., A theorem of Sarkowskii on the existence of periodic orbits of continuous endomorphisms of the real line, *Comm. Math. Phys.* 54(1977), 237-248.
7. BLOCK, L., Homoclinic points of mappings of the interval, *Proc. Amer. Math. Soc.* 72 (1978), 576-580.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleone@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru