

ON REAL ZEROS OF RANDOM POLYNOMIALS WITH HYPERBOLIC ELEMENTS

K. FARAHMAND

Department of Mathematics
University of Ulster
Jordanstown
Co. Antrim BT37 0QH, U.K.

M. JAHANGIRI

Department of Mathematics
Kent State University
14111 Claridon Troy Road
Burton, OH 44021-9500, U.S.A.

(Received May 1, 1996 and in revised form December 30, 1996)

ABSTRACT. This paper provides the asymptotic estimate for the expected number of real zeros of a random hyperbolic polynomial $g_1 \cosh x + 2g_2 \cosh 2x + \dots + ng_n \cosh nx$ where g_j , ($j = 1, 2, \dots, n$) are independent normally distributed random variables with mean zero and variance one. It is shown that for sufficiently large n this asymptotic value is $(1/\pi) \log n$.

KEY WORDS AND PHRASES: Number of real zeros, Kac-Rice formula, random algebraic polynomial, random trigonometric polynomial.

1991 AMS SUBJECT CLASSIFICATION CODES: Primary 60G99; Secondary 42BXX.

1. INTRODUCTION

Let $(\Omega, \mathcal{A}, \Pr)$ be a fixed probability space and $\{g_j(\omega)\}_{j=1}^n$ be a sequence of independent identically distributed random variables defined on Ω , each normally distributed with mean zero and variance one. Denote by $M_n(\alpha, \beta)$ the number of real zeros of $P_n(x)$ in the interval (α, β) , where,

$$P_n(x) \equiv P_n(x, \omega) = \sum_{j=1}^n j g_j(\omega) \cosh jx, \quad (1.1)$$

and by $EM_n(\alpha, \beta)$ its expected value. A similar hyperbolic polynomial to (1.1) has been studied in an unpublished work of Das [4] reported by Bharucha-Reid and Sambandham [1, page 110]. As oscillatory behavior of hyperbolic functions is well known it would be interesting to know how far this oscillatory behavior is transferred to the random hyperbolic polynomial. Since $EM_n(-\infty, \infty)$ could serve as the expected number of maxima and minima of a random hyperbolic polynomial, its value together with the expected number of real zeros and K -level crossings gives a better picture of such oscillations. This expected number of maxima and minima for other type of random polynomials, such as the random algebraic polynomial

$$F(x) \equiv F_n(x, \omega) = \sum_{j=1}^n g_j(\omega) x^j,$$

and the random trigonometric polynomial

$$T(x) \equiv T_n(x, \omega) = \sum_{j=1}^n g_j(\omega) \cos jx,$$

is obtained by Das [3] and [2], respectively. It is shown that in the algebraic case the expected number of maxima and minima is asymptotic to $\{(\sqrt{3} + 1)/\pi\} \log n$, that is about 1.366 times $(2/\pi) \log n$, the expected number of real zeros, see also Wilkins [9]. Therefore in the algebraic case there are about $(0.366) \log n$ oscillations which do not occur between two axis crossings, while for the trigonometric polynomial the expected number of maxima and minima in the interval $(0, 2\pi)$ is $2\sqrt{3}n/\sqrt{5}$, which is about 1.342 times $2n/\sqrt{3}$, the expected number of real zeros, see also Farahmand [6] and Wilkins [10]. Hence in the latter case, asymptotically, there are about 0.197 oscillations which do not occur between two axis crossings. It is, therefore, of special interest to have equivalent results for the hyperbolic case. We prove the following:

THEOREM 1 *The expected number of real zeros of $P_n(x)$ as $n \rightarrow \infty$, satisfies*

$$EM_n(-\infty, \infty) \sim (1/\pi) \log n.$$

From the proof of the theorem we will see that the above expected number arises from interval $(-1, 1)$ and in $(-\infty, -1) \cup (1, \infty)$ there is no significant number of zeros. This is also the case for the expected number of real zeros of $\sum_{j=1}^n g_j(\omega) \cosh jx$. The latter expected number is given in [4], or [5], as $(1/\pi) \log n$. Therefore, our theorem leads to the conjecture that for the hyperbolic case the number of oscillations are the same as the number of zero crossings. That is, for the hyperbolic case almost all the oscillations occur between two zero crossings. Indeed, unlike the algebraic case, these oscillations are all in the interval $(-1, 1)$ and outside this interval the polynomial does not oscillate significantly.

2. PRIMARY ANALYSIS

The Kac-Rice formula, [7] and [8], gives the expected number of real zeros of $P_n(x)$ in the interval (α, β) as

$$EN(\alpha, \beta) = \int_{\alpha}^{\beta} (\Delta/\pi A^2) dx, \tag{2.1}$$

where

$$A^2 = \text{var}\{P_n(x)\} = \sum_{j=1}^n j^2 \cosh^2 jx, \tag{2.2}$$

$$B^2 = \text{var}\{P'_n(x)\} = \sum_{j=1}^n j^4 \sinh^2 jx, \tag{2.3}$$

$$C = \text{cov}\{P_n(x), P'_n(x)\} = \sum_{j=1}^n j^3 \sinh jx \cosh jx \tag{2.4}$$

and

$$\Delta^2 = A^2 B^2 - C^2. \tag{2.5}$$

Since for each zero of $P_n(x)$ in the positive x -axis there corresponds a zero of $P_n(x)$ in the negative x -axis, it is sufficient to restrict ourselves to $(0, \infty)$. Successive differentiation of both sides of

$$\sum_{j=1}^n \cosh 2jx = \cosh nx \sinh(n+1)x / \sinh x - 1,$$

which is easy to prove, together with (2.2)-(2.4) and a little algebra yield,

$$A^2 = n(n+1)(2n+1)/12 + \sinh(2n+1)x \{n(n+1)$$

$$+ \cosh^2 x / 2 \sinh^2 x \} / 4 \sinh x - (2n + 1) \cosh x \cosh(2n + 1)x / 8 \sinh^2 x, \tag{2.6}$$

$$\begin{aligned} B^2 = & -n(n + 1)(2n + 1)(3n^2 + 3n - 1) / 60 + \{n(n^3 + 2n^2 - 1) \\ & + 3 \cosh^4 x / 2 \sinh^4 x\} \sinh(2n + 1)x / 4 \sinh x - \{2(2n^3 + 3n^2 - n - 1) \\ & + (6n + 3) \cosh^2 x / \sinh^2 x\} \cosh x \cosh(2n + 1)x / 8 \sinh^2 x \\ & + \{(3n^2 + 3n - 1) \cosh^2 x \sinh(2n + 1)x / 4 \sinh^3 x, \end{aligned} \tag{2.7}$$

and

$$\begin{aligned} C = & (16 \sinh x)^{-1} \left[\{(4n^3 + 6n^2 - 1) \right. \\ & + 3(2n + 1) \cosh^2 x / \sinh^2 x\} \cosh(2n + 1)x \\ & \left. - \{(6n^2 + 6n - 1) + 3 \cosh^2 x / \sinh^2 x\} \cosh x \sinh(2n + 1)x / \sinh x \right]. \end{aligned} \tag{2.8}$$

In order to be able to carry out the integration required in (2.1) we seek to obtain the dominant terms in (2.6)-(2.8). To this end we first let $x \in (\epsilon, 1)$ where $\epsilon = (\log n)^{1/2} / n$. Since in this interval, for all sufficiently large n , $2n + 1 > p \coth x \tanh(2n + 1)x$, for $p = 1, 2, \dots, 5$ then $f_{n,p}(x) = \sinh(2n + 1)x / \sinh^p x$ is an increasing function of x . Therefore, since $\sinh x < 4x$ in $(0, 1)$ for all sufficiently large n ,

$$f_{n,p} \geq (n/4)^p (\log n)^{-p/2} \exp\{2(\log n)^{1/2}\}.$$

Hence, those terms in (2.6)-(2.8) that are appearing in the form of $f_{n,p}$ for the largest value of p are dominant. Hence from (2.5) and (2.6)-(2.8) we can easily obtain,

$$A^2 = \{n^2 \sinh(2n + 1)x / 4 \sinh x\} \{1 + O(\log n)^{-1/2}\} \tag{2.9}$$

$$B^2 = \{n^4 \sinh(2n + 1)x / 4 \sinh x\} \{1 + O(\log n)^{-1/2}\} \tag{2.10}$$

$$C = \{n^3 \cosh(2n + 1)x / 4 \sinh x\} \{1 + O(\log n)^{-1/2}\}. \tag{2.11}$$

In order to evaluate $\Delta^2 = A^2 B^2 - C^2$ the dominant terms obtained in (2.9)-(2.11) somehow cancel out. Therefore we need to use (2.6)-(2.8). Some algebra, together with the same application of the upper limit of $f_{n,p}$ mentioned above yields,

$$\Delta^2 = \{n^4 \sinh(2n + 1)x \cosh(2n + 1)x / 64 \sinh^4 x\} \{1 + O(\log n)^{-1/2}\}. \tag{2.12}$$

These estimations are sufficient to enable us to prove the theorem.

3. PROOF OF THE THEOREM

We first avoid getting too close to zero by confining ourselves to the interval $(\epsilon, 1)$ where as above $\epsilon = (\log n)^{1/2} / n$. We can, therefore, use the estimates obtained in (2.9)-(2.12). Thus using (2.1), since for all $x \in (\epsilon, 1)$ and n sufficiently large $\coth(2n + 1)x = 1 + o(\log n)^{-1/2}$, some algebra yields,

$$\begin{aligned} EM_n(\epsilon, 1) &= (1/2\pi) \int_{\epsilon}^1 \operatorname{csch} x \, dx \{1 + O(\log n)^{-1/2}\} \\ &= (1/2\pi) \log n \{1 + O(\log n)^{-1/2}\}. \end{aligned} \tag{3.1}$$

Now we show that the expected number of maximum and minimum in the interval $(0, \epsilon)$ is small. To this end from (2.1) and the definition of A^2 and B^2 in (2.2) and (2.3) we can write,

$$\begin{aligned} EM_n(0, \epsilon) &< \int_0^{\epsilon} (B/\pi A) \, dx \\ &= \int_0^{\epsilon} \left\{ \sum_{j=1}^n j^4 \sinh^2 jx / \pi^2 \sum_{j=1}^n j^2 \cosh^2 jx \right\}^{1/2} \, dx \\ &< n\epsilon / \pi = O(\log n)^{1/2}. \end{aligned} \tag{3.2}$$

Therefore (3.1) and (3.2) gives $(1/2\pi) \log n$ as the asymptotic value for $EM_n(0, 1)$ and so.

$$EM_n(-1, 1) \sim (1/\pi) \log n. \quad (3.3)$$

Now it only remains to consider the case of $x \in (-\infty, -1) \cup (1, \infty)$. As we mentioned above it suffices to confine ourselves to the interval $(1, \infty)$. In this interval also the first leading terms of A^2 , B^2 and C which appeared in (2.6)-(2.8) in the evaluation of Δ^2 , cancel out. Therefore we need to use the following estimates, valid for sufficiently large n ,

$$A^2 \sim n\{(n+1) \sinh(2n+1)x - \coth x \cosh(2n+1)x\}/4 \sinh x, \quad (3.4)$$

$$B^2 \sim n^3\{(2n+4 - \coth x) \sinh(2n+1)x - 3 \coth x \cosh(2n+1)x\}/8 \sinh x, \quad (3.5)$$

$$C \sim n^2\{(2n+3) \cosh(2n+1)x - 3 \coth x \sinh(2n+1)x\}/8 \sinh x. \quad (3.6)$$

Therefore from (2.5) and (3.4)-(3.6) for all sufficiently large n we obtain

$$\Delta^2 \sim n^5 \cosh x \sin(2n+1)x \{\cosh(2n+1)x - \sinh(2n+1)x\}/32 \sinh^3 x. \quad (3.7)$$

Finally from (2.1), (3.7) and (3.4) we have,

$$\begin{aligned} EM_n(1, \infty) &\sim (\sqrt{n}/\pi\sqrt{2}) \int_1^\infty \coth^{1/2} x \sinh^{-1/2}(2n+1)x \{\cosh(2n+1)x \\ &\quad - \sinh(2n+1)x\}^{1/2} dx \\ &< (\sqrt{2n}/\pi) \int_1^\infty \coth^{1/2} x \exp(-nx) dx \\ &< o(n^{-1/2}). \end{aligned} \quad (3.8)$$

Since (3.8) can also serve as an upper limit for $EM_n(-\infty, -1)$ from (3.3) we have proof of the theorem.

ACKNOWLEDGMENT: The authors wish to thank the referee for useful comments on the earlier version of this paper.

References

- [1] A.T. Bharucha-Reid and M. Sambandham. *Random Polynomials*. Academic Press, N.Y., 1986.
- [2] M. Das. The average number of real zeros of a random trigonometric polynomial. *Proc. Camb. Phil. Soc.*, 64:721-729, 1968.
- [3] M. Das. The average number of maxima of a random algebraic curve. *Proc. Camb. Phil. Soc.*, 65:741-753, 1969.
- [4] M. Das. On the real zeros of a random polynomial with hyperbolic elements. *Unpublished Ph.D Dissertation, India*, 1971.
- [5] K. Farahmand. Level crossings of a random polynomial with hyperbolic elements. *Proc. Amer. Math. Soc.*, In Press.
- [6] K. Farahmand. The level crossings of random polynomials. *Appl. Math. Lett.*, 9:19-25, 1996.
- [7] M. Kac. On the average number of real roots of a random algebraic equation. *Bull. Amer. Math. Soc.*, 49:314-320, 1943.
- [8] S.O. Rice. Mathematical theory of random noise. *Bell. System Tech. J.*, 25:46-156, 1945.
- [9] J.E. Wilkins. An asymptotic expansion for the expected number of real zeros of a random polynomial. *Proc. Amer. Math. Soc.*, 103:1249-1258, 1988.
- [10] J.E. Wilkins. Mean number of real zeros of a random trigonometric polynomial. *Proc. Amer. Math. Soc.*, 111:851-863, 1991.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from “Qualitative Theory of Differential Equations,” allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk