

ON THE REFLECTION OF ALFVÉN WAVES IN AN IDEAL MAGNETOSPHERE

HADI YAHYA ALKAHBY

Department of Mathematics
Dillard University
New Orleans, LA 70122, U.S.A.

and

M.A. MAHROUS

Department of Mathematics,
University of New Orleans,
New Orleans, LA , 70148, U.S.A.

(Received March 20, 1996)

ABSTRACT: A linearized theory of magnetoatmospheric waves is developed where the restoring forces are those of compressibility and magnetic pressure. An equation for resonance is derived. Reflection and tunneling of upward propagating Alfvén waves in an ideal Magnetoatmosphere are considered. It is shown that the magnetic field produces a reflecting nonabsorbing critical layer. Below the critical layer, the solution of the problem can be written as a linear combination of an upward and a downward propagating wave and above it the solution decays exponentially with the altitude. The location of the critical layer and the magnitude of the reflection coefficient are determined and the conclusions are discussed in connection with the heating mechanism of the solar atmosphere.

KEY WORDS: Magnetoatmosphere, critical layer, wave propagation

1991 AMS SUBJECT CLASSIFICATION CODES: 76N,76Q

1 INTRODUCTION

The dynamics of the solar atmosphere is complicated by the fact that not only is it strongly stratified, in both gas density and temperature, but it is also permeated by a non-uniform magnetic field. The solar atmosphere is an example of a plasma that is both structured and stratified. More specifically, the sun is a compressible plasma and able to support sound waves. The presence of a strong magnetic field indicates that the solar atmosphere is an elastic medium. Thus, wave motions of various types will occur and become a source of energy in the solar atmosphere.

In this paper a linearized theory of magnetoatmospheric waves, involving the combined restoring forces due to compressibility and magnetic pressure, is developed for the case of a uniform horizontal magnetic field. A general propagation equation is derived for adiabatic perturbations with arbitrary vertical distribution of Alfvén and sound speeds. An exact analytical solution of the propagation equation is obtained for the case of an isothermal atmosphere permeated by a uniform horizontal magnetic field. We examine the propagation of Alfvén waves in two distinct regions in the solar atmosphere, which is taken to be an ideal one. It is shown that in the first region, where the strength of the magnetic field is weak, the solution can be written as a linear combination of upward and downward propagating waves

with equal wavelengths. In the first region the motion is approximately acoustic because the motion is dominated by the restoring force of compressibility. In the second region, where the magnetic field strength dominates the motion, the solution decays exponentially with altitude. The first and second regions are connected by a critical layer in which the reflection takes place and the motion is influenced greatly by the effect of the magnetic field. Also, the behavior of the solution, in both regions, indicates that the tunneling is weak as the waves, of all kinds, propagate between the two regions and the reflection is very strong. We expect the motion, in both regions to continue in its prescribed form because there is no physical mechanism for dissipation. The reflection coefficient, location of the transition region, critical layer, and the conclusions are presented in connection with the heating mechanism of the solar atmosphere.

Finally, in the formulation of the problem we will be able to introduce and justify the so-called "magnetic energy condition" in an ideal Magnetoatmosphere as an upper boundary condition to ensure a unique solution.

2 PROBLEM FORMULATION

We consider an ideal magnetoatmosphere which is inviscid and thermally non-conducting, and occupies the upper half-space $z > 0$. It is assumed that the gas is under the influence of a uniform horizontal magnetic field and that it has an infinite electrical conductivity. We investigate the problem of small vertical oscillations about equilibrium.

Let the equilibrium pressure, density, magnetic field intensity and external potential be denoted by $P_0(z)$, $\rho_0(z)$, $B(z)$ and $\psi_0(z)$. Let P , ρ , B , ψ and \mathbf{V} be the perturbations in the pressure, density, magnetic field intensity, external potential and velocity. The equations of ideal magnetohydrodynamics are those of momentum, induction, isotropy, and continuity:

$$\rho \left[\frac{\partial \mathbf{V}}{\partial t} + \mathbf{V} \cdot \nabla \mathbf{V} \right] = -\nabla p - \mathbf{B} \times (\nabla \times \mathbf{B}) - \rho \nabla \psi, \quad (2.1)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{V} \times \mathbf{B}), \quad (2.2)$$

$$\left[\frac{\partial}{\partial t} + \mathbf{V} \cdot \nabla \right] (P \rho^{-\gamma}) = 0, \quad (2.3)$$

$$\left[\frac{\partial}{\partial t} + \mathbf{V} \cdot \nabla \right] \rho = -\rho \nabla \cdot \mathbf{V}. \quad (2.4)$$

Here γ denotes the ratio of the specific heats. These equations are linearized about a static equilibrium defined by

$$\nabla P_0 + \mathbf{B}_0 \times (\nabla \times \mathbf{B}_0) = -\rho_0 \nabla \psi_0. \quad (2.5)$$

In this problem ψ_0 is considered to represent a uniform gravitational field. Thus, $\nabla \psi_0 = -\mathbf{g} = (0, 0, g)$ and the last term in equations (2.1) and (2.5) is replaced by $\rho_0 \mathbf{g}$. In terms of the linear Lagrangian displacement field $\boldsymbol{\xi}(\mathbf{r}, t)$, the velocity perturbation is defined by:

$$\mathbf{V} = \frac{\partial \boldsymbol{\xi}}{\partial t}, \quad (2.6)$$

and the equation of motion of ideal magnetohydrodynamic is obtained by integrating and eliminating all perturbation quantities except $\boldsymbol{\xi}$. As a result, we have

$$\rho_0 \frac{\partial^2 \boldsymbol{\xi}}{\partial t^2} = \mathbf{F}(\boldsymbol{\xi}), \quad (2.7)$$

where the force operator $\mathbf{F}(\boldsymbol{\xi})$ is defined by

$$\mathbf{F}(\boldsymbol{\xi}) = \nabla(\gamma P_0 \nabla \cdot \boldsymbol{\xi} + \boldsymbol{\xi} \cdot \nabla P_0) - \mathbf{B}_0 \times (\nabla \times \mathbf{R}) - \mathbf{R} \times (\nabla \times \mathbf{B}_0) + \nabla \cdot (\rho_0 \boldsymbol{\xi}) \nabla \psi_0, \quad (2.8)$$

where

$$\mathbf{R} = \nabla \times (\boldsymbol{\xi} \times \mathbf{B}_0). \quad (2.9)$$

The linear Lagrangian displacement field is assumed to be of the form $\boldsymbol{\xi}(x, y, z, t)$ and as an integral superposition of harmonic terms, with $\boldsymbol{\xi}$ and its first x derivative vanishing at infinity. As a result,

$$\boldsymbol{\xi} = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-\omega t} \boldsymbol{\phi}(z, k) e^{ikx} dk, \quad (2.10)$$

where the wave number is assumed to be $\mathbf{k} = (k, 0, 0)$. Consequently, the horizontal magnetic field can be written as

$$\mathbf{B}_0 = (B_0(z), B_y(z), 0). \quad (2.11)$$

Using equation (2.10), equation (2.7) becomes

$$-\rho_0 \omega^2 \boldsymbol{\phi} = \mathbf{F}(\boldsymbol{\phi}, k). \quad (2.12)$$

Eliminating all variables except ϕ_z , we obtain the following differential equation:

$$\frac{d}{dz} \left[A(z, \omega) \frac{d\phi_z}{dz} \right] + C(z, \omega) \phi = 0, \quad (2.13)$$

where the coefficient $A(z, \omega)$ and $C(z, \omega)$ have the following form:

$$A(z, \omega) = \frac{\rho_0 A_{11} A_{12} A_{13}}{A_{14} A_{15}}, \quad (2.14)$$

$$C(z, \omega) = \rho_0 \left[A_{12} - \frac{k^2 g^2 A_{12}}{A_{14} A_{15}} - \frac{k^2 g}{\rho_0} \left(\frac{d}{dz} \left[\rho_0 \left(\frac{A_{11} - a_x}{A_{14} A_{15}} \right) A_{16} \right] \right) \right], \quad (2.15)$$

where

$$\begin{aligned} A_{11} &= a_x^2 + a_y^2 + c_0^2, & A_{12} &= \omega^2 - a_x^2 k^2, \\ A_{13} &= \omega^2 - \frac{a_x^2 k^2 c_0^2}{A_{11} - a_y}, & A_{14} &= \omega^2 - \frac{1}{2} [k^2 A_{11} - A_{17}], \\ A_{15} &= \omega - \frac{1}{2} [k^2 A_{11} + A_{17}], & A_{16} &= \omega - \left[\frac{a_x c_0^2 k^2}{A_{11} - a_x} \right], \\ A_{17} &= k^4 [A_{11}^2 - 4a_x^2 c_0^2]. \end{aligned} \quad (2.16)$$

The parameters a_x and a_y are the x and y components of Alfvén speed, while c_0^2 denotes the speed of sound.

3 SIMPLIFICATION OF THE PROBLEM AND BOUNDARY CONDITIONS

For this problem the atmosphere is assumed to be isothermal ($c_0 = c(z) = \text{constant}$) and permeated by a uniform horizontal magnetic field $\mathbf{B} = (B, 0, 0)$. The equilibrium pressure P_0 and constant temperature T_0 satisfy the gas law $P_0 = RT_0 \rho_0$ and the hydrostatic equation $P_0'(z) + g \rho_0 = 0$. Here, R denotes the gas constant and the prime denotes differentiation with respect with respect to z . As a result, the equilibrium pressure and density can be written as:

$$P_0(z) = P_0(0) \exp(-z/H), \quad \rho_0(z) = \rho_0(0) \exp(-z/H), \quad (3.1)$$

where $H = -\rho'_0(z)/\rho_0(z) = \frac{RT_0}{g}$ is the density scale height. Consequently, the differential equation (2.13) becomes

$$\begin{aligned} & [\omega^2 c^2 + (\omega^2 - c^2 k^2) \alpha^2(0) \exp\left(\frac{z}{H}\right)] \phi''(z) - \frac{c^2 \omega^2}{H} \phi'(z) + \\ & \left[(\omega^2 - c^2 k^2) (\omega^2 - k^2 \alpha_0^2(0) \exp\left(\frac{z}{H}\right) - g \left(g - \frac{c^2}{H}\right) k^2 \right] \phi(z) = 0, \end{aligned} \tag{3.2}$$

where

$$\alpha_0^2(0) = \frac{B^2}{4\pi\rho_0(0)},$$

denotes Alfvén speed at $z = 0$. Furthermore, we introduce the following dimensionless quantities and variables:

$$\phi(z) = W(x) \exp -kz, \quad x = x_0 \exp\left(-\frac{z}{H}\right), \tag{3.3}$$

$$\dot{z} = \frac{z}{H}, \quad k = Hk, \quad x_0 = \frac{\beta^2 \sigma^2}{k^2 - \sigma^2}, \quad \beta^2 = \frac{\alpha_0^2(0)}{c^2}, \quad \sigma = \frac{H\omega}{c}, \tag{3.4}$$

where the prime on z is eliminated for simplicity. Thus, the differential equation (3.2) becomes,

$$x(1-x) \frac{d^2W(x)}{dx^2} + [c - (a+b+1)x] \frac{dW(x)}{dx} - abW(x) = 0. \tag{3.5}$$

It is clear that the differential equation (3.5) is a special case of the hypergeometric differential equation with

$$a + b = c = 2k + 1, \quad ab = \sigma^2 + k + \left(\frac{\gamma - 1}{\gamma}\right) \frac{k^2}{\sigma^2}, \tag{3.6}$$

Boundary Conditions:

To complete the formulation of the problem, certain conditions must be imposed to ensure a unique solution. If the atmosphere is viscous, an appropriate condition would be the dissipation condition, which requires the finiteness of the rate of the energy dissipation in an infinite column of fluid of unit cross-section. Since the dissipation function depends on the squares of the velocity gradients, this implies

$$\int_0^\infty |W'|^2 dx < \infty. \tag{3.7}$$

In our problem the atmosphere is not viscous, but the integral in (3.7) is proportional to the magnetic energy in an infinite column of fluid. This condition is a reasonable one to apply so long as there is no energy radiation to infinity, which is true in our case, and we shall call this condition “magnetic energy condition.” Thus we will require (3.7) even when the atmosphere is inviscid. A boundary condition is also required at $x = 0$, and we shall set

$$W(0) = 1, \tag{3.8}$$

by suitably normalizing $W(x)$. It will be seen that the boundary conditions (3.7) and (3.8) will determine a unique solution to within a multiplicative constant.

4 SOLUTION OF THE PROBLEM

In this section we investigate solutions of the following differential equation,

$$x(1-x) \frac{d^2W(x)}{dx^2} + [c - (a+b+1)x] \frac{dW(x)}{dx} - abW(x) = 0, \tag{4.1}$$

where

$$a + b = c = 2k + 1, \quad ab = \sigma^2 + k + \left(\frac{\gamma - 1}{\gamma}\right) \frac{k^2}{\sigma^2}, \tag{4.2}$$

subject to the prescribed boundary conditions. Solving for the parameters a and b , we obtain

$$a = \frac{1}{2} + k - r, \quad b = \frac{1}{2} + k + r, \quad (4.3)$$

where

$$r = \sqrt{r_1 - r_2}, \quad r_1 = \frac{1}{4} + k^2 \quad \text{and} \quad r_2 = \sigma^2 + \frac{\gamma - 1}{\gamma} \frac{k^2}{\sigma^2}.$$

It is clear that the parameter r is a real number, for $r_1 > r_2$, $r = 0$ for $r_1 = r_2$ and $r = i\sqrt{r_2 - r_1}$ for $r_1 < r_2$. The differential equation (4.1) is a special case of the hypergeometric equation which has three regular singular points at $x = 0$, $x = 1$ and $x = \infty$. The intermediate regular singular point, $x = 1$, corresponds to the existence of the critical layer, which has a great importance for understanding the heating mechanism of the solar atmosphere. As a result, the differential equation (4.1) has two linearly independent solutions which can be written in the following form for $|x| < 1$.

$$W_1(x) = F(a, b; c; x), \quad (4.4)$$

$$W_2(x) = x^{1-c} F(a - c + 1, b - c + 1; 2 - c; x), \quad (4.5)$$

$$F(a, b; c; x) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(b)} \sum_{n=0}^{\infty} \frac{\Gamma(a+n)\Gamma(b+n)}{\Gamma(c+n)} \frac{x^n}{n!}. \quad (4.6)$$

Since $k > 0$, then $1 - c = -2k < 0$, and

$$x^{1-c} = [x_0 \exp(-z)]^{1-c} \rightarrow \infty, \text{ as } z \rightarrow \infty. \quad (4.7)$$

Thus, the solution of the differential equation (4.1) as defined by equation (4.5) will be eliminated by the magnetic energy condition. Consequently, the general solution of the differential equation (4.1) is

$$W_1(x) = AF(a, b; c; x), \quad (4.8)$$

where A is an arbitrary constant and can be determined by the boundary condition (3.8). Using the asymptotic behavior of $F(a, b; c; x)$ for $|x| > 1$ and reintroducing the variable z , the solution of (4.4) can be written as

$$W(z) = \text{Cons} \cdot \left[\left[\exp\left(\frac{1}{2} + k + ir\right) z \right] + R \exp\left(\frac{1}{2} + k - ir\right) z \right], \quad (4.9)$$

where R denotes the reflection coefficient and defined by

$$R = \exp[i(\theta + 2r \log x_0)],$$

and

$$\theta = \arg(R) = \arg \left[\frac{\Gamma(a)\Gamma(b-a)\Gamma(c-b)}{\Gamma(a-b)\Gamma(b)\Gamma(c-a)} \right].$$

5 GENERAL DISCUSSION

The structured nature of the solar magnetic field means that magnetism is of greater importance in some regions of the sun than in others. A simple guide to the relative importance of magnetic effects is provided by the plasma beta, β , defined by

$$\beta = \frac{P_0}{P_m} = \frac{2}{\gamma} \left(\frac{c}{\alpha} \right)^2, \quad (5.1)$$

where P_m denotes the magnetic pressure. A low- β plasma, such as the corona, is thus one for which the Alfvén speed greatly exceeds the sound speed, $\alpha \gg c$. Wave propagation, then, involves the two speeds c and α . In fact, the sound speed exists and inter into a description of propagation speeds only in the combination with Alfvén speed. As a result, the wave speed W_s is such, $S_s < W_s < F_s$, where F_s and S_s are the fast and slow speeds and defined by

$$F_s^2 = c^2 + \alpha^2, \quad S_s^2 = c^{-2} + \alpha^{-2}. \quad (5.2)$$

Moreover, F_s will be referred to as magnetoacoustic speed; it is super-sonic and super-Alfvénic. By contrast, S_s is both sub-sonic and sub-Alfvénic.

It is easy to see that the maximum of the kinetic energy $\text{Max}(K) \propto \left(\frac{1}{|1+R|}\right)$. As a result, when the reflection coefficient $R \rightarrow -1$, we have $\text{Max}(K) \rightarrow \infty$. This occurs when

$$\theta + 2r \log x_0 \rightarrow \pm (2n + 1) \pi. \quad (5.3)$$

We call this equation, resonance equation.

From the above discussion and the asymptotic behavior of the solution, expressed in (4.9), we have the following observations:

(A) It is clear that $|R| = 1$. As a result, the magnetic field produces a nonabsorbing critical layer, below it the solution can be written as a linear combination of an upward and a downward propagating wave with the same wavelengths. Above the critical layer the solution decays exponentially with altitude. Thus, the critical layer separates two distinct regions with different physical properties.

(B) In the critical layer the reflection and the wave modification take place. Since $|R| = 1$, the tunneling is very weak while the reflection is very strong. In this case the total energy of the wave is divided equally between the incident and reflected waves.

(C) Since the Magnetoatmosphere is ideal, there is a physical mechanism for dissipation. Thus, we expect the motion to continue in this form and become one of the main sources of energy in the solar atmosphere.

(D) As a result of (B) and (C) we see that the heating process is an acoustic one below the critical layer because the compressibility force dominates the oscillatory process. As we move from a region of weak magnetic field to another one with a strong magnetic field, the heating process becomes magnetoacoustic.

References

- [1] ALKAHBY, H., "Reflection and dissipation of hydromagnetic waves in a viscous and thermally conducting isothermal atmosphere," *Geophys. Astroph. Fluid. Dynam.*, **72**, (1993), 197-207.
- [2] ALKAHBY, H., "The dissipation of magneto-acoustic waves," *Computer Math. Applic.*, Vol. **27**, (1994), 9-15.
- [3] ALKAHBY, H., and YANOWITCH, M., "Reflection of vertically propagating waves in a thermally conducting isothermal atmosphere with a horizontal magnetic field," *Geophys. Astroph. Fluid. Dynam.* Vol. **56**, (1991), 227-235.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br