

NONLINEAR OSCILLATIONS IN DISCONJUGATE FORCED FUNCTIONAL EQUATIONS WITH DEVIATING ARGUMENTS

BHAGAT SINGH

Department of Mathematics
University of Wisconsin Center
Manitowoc, Wisconsin 54220 U.S.A.

(Received July 28, 1982)

ABSTRACT. For the equation

$$L_n v(t) + a(t)h(y(\sigma(t))) = f(t)$$

where

$$L_n y(t) = p_n(t) (p_{n-1}(t) (\dots (p_1(t) (p_0(t)y(t)))' \dots)')'$$

sufficient conditions have been found for all of its solutions to be oscillatory. The conditions found also lead to growth estimates for the nonoscillatory solutions.

KEY WORDS AND PHRASES. Oscillatory, nonoscillatory, disconjugate, nonlinear oscillations.

AMS MOS (1980) SUBJECT CLASSIFICATION CODE. 34K15

1. INTRODUCTION.

Our main purpose in this paper is to study the oscillatory phenomenon associated with the equation

$$L_n y(t) + a(t)h(y(g(t))) = f(t) \tag{1.1}$$

Where $n \geq 2$ and L_n is a disconjugate differential operator defined by

$$L_n y(t) = p_n(t) (p_{n-1}(t) (\dots (p_1(t) (p_0(t)(y(t)))' \dots)')' \dots)')' \tag{1.2}$$

Following our work (Singh and Kusano [6]), it is assumed that:

(i) $p_i \in C([\alpha, \infty), (0, \infty))$, $0 \leq i \leq n$

(ii) $a, f, g \in C([\alpha, \infty), \mathbb{R})$, there exists a $t_0 > \alpha$ such that

$$0 < g(t) \leq t \text{ for } t \geq t_0, \text{ and } g(t) \rightarrow \infty;$$

(iii) $h \in C(\mathbb{R}, \mathbb{R})$, h is nondecreasing and $\text{sign } h(y) = \text{sign } y$.

We introduce the notation:

$$L_0 y(t) = p_0 y(t), L_i y(t) = p_i(t)(L_{i-1}(y(t)))', \quad 1 \leq i \leq n. \quad (1.3)$$

The domain $D(L_n)$ of L_n is defined to be the set of all functions $y : [T_y, \infty) \rightarrow \mathbb{P}$ such that $L_i y(t)$, $0 \leq i \leq n$, exist and are continuous on $[T_y, \infty)$. In what follows by a "solution" of equation (1.1) we mean a function $y \in D(L_n)$ which is nontrivial in any neighborhood of ∞ and satisfies (1.1) for all sufficiently large t . A solution of (1.1) is called oscillatory if it has arbitrarily large zeros; otherwise the solution is called nonoscillatory.

A great many oscillation criteria are known for an equation of the form

$$(r(t)y'(t))^{(n-1)} + a(t)h(y(g(t))) = 0 \quad (1.4)$$

For this we refer the reader to Onose [2], Singh [3] and Kusano and Onose [1]. A recently published Russian book by Sivelevov [7] gives a detailed list of references on the subject. Obtaining an oscillation criterion for the forced equation

$$(r(t)y'(t))^{(n-1)} + a(t)h(y(g(t))) = f(t) \quad (1.5)$$

is not so simple. To the best of this author's knowledge, the first attempt to obtain conditions for the oscillation of the equation

$$(r(t)y'(t))' + a(t)h(y(g(t))) = f(t) \quad (1.6)$$

was made by Kusano and Onose [1], and later by other authors including this one [4].

The main technique employed rendered the forced equation into an almost homogeneous equation, i.e. a function $\lambda(t)$ was sought such that $\lambda^{(i)}(t) \rightarrow 0$ as $t \rightarrow \infty$ for $i = 0, 1, \dots, n$; and $(r(t)\lambda'(t))^{(n-1)} = f(t)$.

In this work, we shall present a new but elementary technique to obtain an oscillation criterion for the equation (1.1).

In order to shorten long expressions we introduce the following notations:

For any function $Q(t) \in C[\alpha, \infty)$ and $t, s \in [\alpha, \infty)$, define

$$I_0(Q(r), r) = p_n^{-1}(r)Q(r) \quad (1.7)$$

and

$$I_k(Q(t), t, s; p_n, p_{n-1}, \dots, p_{n-k}) = \int_s^t p_{n-k}^{-1}(r) I_{k-1}(Q, t, r; p_n, \dots, p_{n-k+1}) dr \quad (1.8)$$

for $1 \leq k \leq n-1$.

Any solution $y(t)$ of equation (1.1) which is continuous in a finite interval can be indefinitely extended to the right of α provided the coefficients are continuous.

In fact following our proof of Theorem 3.1 in [5] we can state the following theorem:

THEOREM 1.1 The continuity of p_1, p_2, \dots, p_n ; and that of h, g, a and f guarantee that any solution of equation (1.1) continuous in a finite interval $[\alpha, \tau]$ can be continuously extended to all of $[\alpha, \infty)$.

PROOF. Same as that of Theorem 3.1 in [5] with minor changes.

2. MAIN RESULTS.

THEOREM 2.1 Suppose there exists a function $\psi(t)$ such that

$$\psi(t) \in C^{(n)} [R], \psi(t) > 0 \text{ and } (\psi' p_1(t))' \geq 0 \text{ for } t \geq \alpha. \tag{2.1}$$

Further suppose that $\psi(t)$ satisfies

$$\int_{\alpha}^{\infty} 1/\psi^2(t) dt < \infty \tag{2.2}$$

$$\limsup_{t \rightarrow \infty} \int_{\alpha}^t 1/\psi^2(s) \int_{\alpha}^s \psi(x) I_{n-3}(f(x), x, \alpha; p_n, p_{n-1}, \dots, p_3) p_2^{-1}(x) dx ds = \infty \tag{2.3}$$

$$\liminf_{t \rightarrow \infty} \int_{\alpha}^t 1/\psi^2(s) \int_{\alpha}^s \psi(x) I_{n-3}(f(x), x, \alpha; p_n, p_{n-1}, \dots, p_3) p_2^{-1}(x) dx ds = -\infty \tag{2.4}$$

$$\int_{\alpha}^{\infty} \frac{1}{\psi^2(s)} \int_{\alpha}^s \psi(x) p_2^{-1}(x) \int_{\alpha}^x p_3^{-1}(x_3) \int_{\alpha}^{x_3} p_4^{-1}(x_4) \dots \int_{\alpha}^{x_{n-2}} p_{n-1}^{-1}(x_{n-1}) \tag{2.5}$$

$$dx_{n-1} dx_{n-2}, dx_{n-3} \dots dx < \infty$$

$$a(t) \geq 0 \text{ and } p_1'(t) \leq 0 \text{ for } t \geq \alpha. \tag{2.6}$$

Then all solutions of equation (1.1) are oscillatory.

PROOF. Suppose to the contrary that $y(t)$ is a nonoscillatory solution of equation (1.1). Let $T \geq \alpha$ be large enough so that $y(t)$ and $y(g(t))$ are of the same sign for $t \geq T$. Without any loss of generality, let $y(t) > 0$ and $y(g(t)) > 0$ for $t \geq T$. On repeated integration for $t \geq T$, we obtain from equation (1.1)

$$\begin{aligned}
 & (p_1(t)(p_0(t)y(t))')' - p_2^{-1}(t)L_2y(T) - (p_2^{-1}(t) \int_T^t p_3^{-1}(x)dx)L_3y(T) \\
 & - (p_2^{-1}(t) \int_T^t p_3^{-1}(s) \int_T^s p_4^{-1}(x)dx ds)L_4y(T) - \dots \\
 & -L_{n-1}y(t) p_2^{-1}(t) \int_T^t p_3^{-1}(x_3) \int_T^{x_3} p_4^{-1}(x_4) \dots \int_T^{x_{n-2}} p_{n-1}^{-1}(x_{n-1})dx_{n-1} dx_{n-2} dx_{n-3} \dots dx_3 \\
 & + p_2^{-1}(t) I_{n-3} (a(t)h(y(g(t))),t,T;p_n,\dots,p_3) \\
 & = p_2^{-1}(t) I_{n-3}(f(t),t,T;p_n,p_{n-1},\dots,p_3) \tag{2.7}
 \end{aligned}$$

Multiplying (15) by $\psi(x)$ and rearranging terms we have

$$\begin{aligned}
 & (p_1(t)\psi(t)(p_0(t)y(t))')' - \psi'(t)p_1(t)(p_0(t)y(t))' - \psi(t)p_2^{-1}(t)L_2y(T) \\
 & -\psi(t)((p_2^{-1}(t) \int_T^t p_3^{-1}(x)dx)L_3y(T) - \dots - \psi(t)L_{n-1}y(T)p_2^{-1}(t) \int_T^t p_3^{-1}(x_3) \dots \\
 & \dots \int_T^{x_{n-2}} p_{n-1}^{-1}(x_{n-1})dx_{n-1} \dots dx_3 + \psi(t)p_2^{-1}(t) I_{n-3}(ah,t,T;p_n,\dots,p_3) \\
 & = \psi(t)p_2^{-1}(t) I_{n-3}(f(t),t,T;p_n,p_{n-1},\dots,p_3) \tag{2.8}
 \end{aligned}$$

Integrating (2.8) and dividing by $\psi^2(t)$ we get

$$\begin{aligned}
 & \frac{p_1(t)(p_0y)'}{\psi(t)} - \frac{p_1(T)\psi(T)(p_0y)'(T)}{\psi^2(t)} - \frac{1}{\psi^2(t)} \int_T^t \psi'(x)p_1(x)(p_0y)' dx \\
 & - \left[\frac{1}{\psi^2(t)} \int_T^t \psi(x)p_2^{-1}(x)dx \right] L_2y(T) - \left[\frac{1}{\psi^2} \int_T^t \psi(x)p_2^{-1}(x) \int_T^x p_3^{-1}(s) ds dx \right] L_3y(T) \\
 & - \dots \frac{1}{\psi^2(t)} \left[\int_T^t \psi(x)p_2^{-1}(x) \int_T^x p_3^{-1}(x_3) dx_3 \dots \int_T^{x_{n-2}} p_{n-1}^{-1}(x_{n-1}) dx_{n-1} \dots dx_3 dx \right] L_{n-1}(T) \\
 & + \frac{1}{\psi^2(t)} \int_T^t \psi(x)p_2^{-1}(x) I_{n-3}(a(x)h(y(f(x))),x,T;p_n,\dots,p_3) dx \\
 & = \frac{1}{\psi^2(t)} \int_T^t \psi(x)p_2^{-1}(x) I_{n-3}(f(x),x,T;p_n,p_{n-1},\dots,p_3) dx. \tag{2.9}
 \end{aligned}$$

In (2.9), we integrate the third term by parts to obtain

$$\frac{p_1(t)(p_0y)'}{\psi(t)} - \frac{K_0}{\psi^2(t)} - \frac{p_1(t)p_0y\psi'}{\psi^2} + \frac{1}{\psi^2(t)} \int_T^t (\psi'(x)p_1(x))' p_0y(x) dx$$

$$\begin{aligned}
 & - \left[\frac{1}{\psi^2(t)} \int_T^t \psi(x) p_2^{-1}(x) dx \right] L_2 y(T) - \dots \\
 & - \dots - \left[\frac{1}{\psi^2(t)} \int_T^t \psi(x) p_2^{-1}(x) \int_T^x p_3^{-1}(x_3) \dots \int_T^{x_{n-2}} p_{n-1}^{-1}(x_{n-1}) dx_{n-1} \dots dx_3 dx \right] \\
 & + \frac{1}{\psi^2(t)} \int_T^t \psi(x) p_2^{-1}(x) I_{n-3}(a(x)h(y(g(x))), x, T; p_n, \dots, p_3) dx \\
 & = \frac{1}{\psi^2(t)} \int_T^t \psi(x) p_2^{-1}(x) I_{n-3}(f(x), x, T; p_n, \dots, p_3) dx \tag{2.10}
 \end{aligned}$$

where

$$K_0 = p_1(T)\psi(T)(p_0 y)'(T) + p_1(T)p_0(T)y(T)\psi'(T)$$

Integrating first term in (2.10) again by parts and observing that

$$\begin{aligned}
 & \int_T^t \frac{p_1(x)p_0(x)y(x)\psi'(x)}{\psi^2(x)} dx \\
 & = - \int_T^t \left(\frac{p_1(x)}{\psi(x)} \right)' p_0(x)y(x) dx + \int_T^t \frac{p_1'(x)p_0(x)y(x)}{\psi(x)} dx \tag{2.11}
 \end{aligned}$$

We get

$$\begin{aligned}
 & \frac{p_1(t)p_0(t)y(t)}{\psi(t)} - K_1 - \int_T^t \frac{p_1'(x)p_0(x)y(x)}{\psi(x)} dx + \int_T^t \frac{1}{\psi^2(x)} \int_T^x (\psi' p_1)' p_0(s)y(s) ds dx \\
 & - L_2 y(T) \int_T^t \frac{1}{\psi^2(x)} \int_T^t \psi(s) p_2^{-1}(s) ds - \dots \\
 & \dots - \int_T^t \frac{1}{\psi^2(s)} \int_T^s \psi(x) p_2^{-1}(x) \int_T^x p_3^{-1}(x_3) \dots \int_T^{x_{n-2}} p_{n-1}^{-1}(x_{n-1}) dx_{n-1} \dots dx_3 dx ds \\
 & + \int_T^t \frac{1}{\psi^2(s)} \int_T^s \psi(x) p_2^{-1}(x) I_{n-3}(a(x)h(v(g(x))), x, T, p_n, \dots, p_3) dx ds \\
 & = \int_T^t \frac{1}{\psi^2(s)} \int_T^s \psi(x) p_2^{-1}(x) I_{n-3}(f(x), x, T; p_n, p_{n-1}, \dots, p_3) dx ds \tag{2.12}
 \end{aligned}$$

where

$$K_1 = \frac{p_1(T)p_0(T)y(T)}{\psi(T)}$$

The terms on the left hand side of (2.12) are either positive or finite. Since the right hand side oscillates between $-\infty$ and ∞ , a contradiction is apparent. The theorem is proved.

EXAMPLE 1. Consider the equation

$$(t \frac{1}{t} v')' + y(t-3\pi/2) = t^8 \sin t, \quad t \geq \frac{3\pi}{2} \quad (2.13)$$

Here $p_0(t) \equiv 1$, $p_1(t) \equiv 1/t$, $p_2(t) \equiv t$ and $p_3(t) \equiv 1$.

Choosing $\psi(t) = t^3$ for $t \geq \alpha$, it is easily verified that all conditions of the theorem are satisfied. Hence all solutions of equation (1.1) are oscillatory.

THEOREM 2.2 Suppose conditions (2.2) - (2.6) of theorem 2.1 hold, let $\psi(t)$ be a nonnegative solution of the equation

$$(p_1(t)y'(t))' - a(t)h(v(g(t))) = 0. \quad (2.14)$$

Then all solutions of equation (1.1) are oscillatory.

PROOF. Since $(p_1(t)\psi(t))' \geq 0$, all conditions of theorem 2.1 are satisfied.

The proof is complete.

COROLLARY 2.1 For the equation

$$y^{(n)}(t) + a(t)h(y(g(t))) = f(t) \quad (2.15)$$

Conditions (2.3) - (2.5) respectively reduce to

$$\liminf_{t \rightarrow \infty} \int_{\alpha}^t 1/\psi^2(s) \int_{\alpha}^s \psi(x) \int_{\alpha}^x (x-u)^{n-3} f(u) du dx ds = -\infty \quad (2.16)$$

$$\limsup_{t \rightarrow \infty} \int_{\alpha}^t 1/\psi^2(s) \int_{\alpha}^s \psi(s) \int_{\alpha}^x (x-u)^{n-3} f(u) du dx ds = \infty \quad (2.17)$$

and

$$\lim_{t \rightarrow \infty} \int_{\alpha}^t 1/\psi^2(s) \int_{\alpha}^s \psi(x) x^{n-4} dx ds < \infty. \quad (2.18)$$

Thus subject to conditions (2.2), (2.6) and (2.15) - (2.18) all solutions of equation (2.15) are oscillatory.

EXAMPLE 2. Consider the equation

$$y^{(iv)}(t) + e^{2\pi} y(t-\pi) = -63 e^{2t} \sin(2t), \quad t > \pi \quad (2.19)$$

If we choose $\psi(t) = e^t$, then conditions of corollary 2.1 can be easily verified.

Hence all solutions of equation (2.19) are oscillatory. In fact $y(t) = e^{2t} \sin(\lambda t)$ is one such solution.

For the bounded solutions of equation (1.1), the condition $a(t) \geq 0$ can be improved as the following theorem shows:

THEOREM 2.3 Suppose there exists a function $\psi(t)$ such that

$\psi(t) \in C^{(n)}[\mathbb{R}]$, $\psi(t) > 0$, $(\psi' p_1)' \geq 0$ and $p_1'(t) \leq 0$ for $t \geq \alpha$. Further suppose that $\psi(t)$ satisfies conditions (2.2) - (2.5) of Theorem 2.1 and the condition

$$\int_{\alpha}^{\infty} 1/\psi^2(t) \int_{\alpha}^t \psi(s) I_{n-3}(|a(s)|, s, a; p_n, p_{n-1}, \dots, p_3) p_2^{-1}(s) ds dt < \infty \tag{2.20}$$

Then all bounded solutions of equation (1.1) are oscillatory.

PROOF. We proceed as in Theorem (2.1) and arrive at (2.12). If $y(t)$ is bounded then there exists a constant $C_y > 0$ such that

$$|h(y(g(t)))| \leq C_y. \tag{2.21}$$

In equation (2.12) the last term on the left hand side

$$\begin{aligned} & \left| \int_T^t 1/\psi^2(s) \int_T^s \psi(x) p_2^{-1}(x) I_{n-3}(a(x)h(y(g(x))), x, T; p_n, \dots, p_3) dx ds \right| \\ & \leq C_y \int_T^{\infty} 1/\psi^2(t) \int_T^t \psi(x) p_2^{-1}(x) I_{n-3}(|a(x)|, x, T; p_n, \dots, p_3) dx dt < \infty. \end{aligned} \tag{2.22}$$

Hence all terms on the left hand side of equation (2.12) are either finite or non-negative. The proof now follows as in theorem 2.1.

REMARK. When $n=2,3$, condition (2.18) reduces to condition (2.2) with obvious changes in (2.16) and (2.17).

Under the conditions of theorem 2.3, equation (1.1) may possess unbounded nonoscillatory solutions as the following example shows:

EXAMPLE 3. The equation

$$y''(t) + 50 \sin t y(t) = (4 + 50 \sin t) e^{2t} \tag{2.23}$$

has $v(t) = e^{2t}$ as a nonoscillatory solution.

If we choose $\psi(t) = e^t$ we have from (2.20)

$$\left| \int_{\pi}^{\infty} e^{-2t} \int_{\pi}^t e^s \sin s ds dt \right| < \infty \tag{2.24}$$

Conditions (2.16) and (2.17) yield

$$\liminf_{t \rightarrow \infty} \int_{\pi}^t e^{-2t} \int_{\pi}^s e^{3s} (4 + 50 \sin s) ds dt = -\infty$$

and

$$\limsup_{t \rightarrow \infty} \int_{\pi}^t e^{-2t} \int_{\pi}^s e^{3s} (4 + 50 \sin s) ds dt = \infty$$

Thus all conditions of theorem 2.3 are satisfied. Hence all bounded solutions of equation (2.23) are oscillatory even though it has an unbounded nonoscillatory solution.

COROLLARY 2.2. Subject to the conditions of theorem 2.3 all nonoscillatory solutions of equation (1.1) are unbounded.

Corollary 2.2 leads to the following theorem which gives a growth condition for the nonoscillatory solutions of equation (1.1).

THEOREM 2.4 Suppose conditions of theorem 2.3 hold. Let $v(t)$ be a nonoscillatory solution of equation (1.1). Then

$$\limsup_{t \rightarrow \infty} (|y(t)| p_n^{-1}(t)) = \infty \quad (2.25)$$

PROOF. Suppose to the contrary that there exist constant $D_y > 0$ and $T > \alpha$ such that

$$|y(g(t))| p_n^{-1}(t) \leq D_y \quad (2.26)$$

for $t \geq T$. Condition on h implies that there exists a $C_y > 0$ such that

$$|h(v(g(t)))| \leq C_y |y(g(t))| \leq C_y D_y \equiv K_y \quad (2.27)$$

Following the proof of theorem 2.3 we see that the constant K_y replaces C_y in equality (2.22) and the proof is complete.

REMARK. It is a matter of general interest to obtain theorems involving operator L_n in its most general form; i.e. when L_n is not necessarily in canonical form. L_n is said to be in canonical form if

$$\int_{\alpha}^{\infty} p_i^{-1}(t) dt = \infty, \quad 1 \leq i \leq n-1. \quad (2.28)$$

Recently Trench [8] has shown that when L_n is not in canonical form, i.e. when (2.28) doesn't hold, it can be put in a canonical form in a unique way with a different set of p s satisfying (2.28). More precisely the operator L_n can be rewritten as

$$L_n y = b_n (b_{n-1}(t) (\dots (b_1(t) (b_0(t) y(t))')' \dots))'$$

where

$$\int_a^\infty b_i^{-1}(t) dt = \infty, \quad 1 \leq i \leq n-1 \quad (2.29)$$

each b_i is obtained from $p_1, p_2, \dots, p_n, i = 0, 1, \dots, n$. The actual computations leading to b s are tedious.

It is interesting and important to note that the results in this work do not depend upon condition (2.28) or (2.29).

REFERENCES

1. KUSANO, T. and ONOSE, H. Oscillations of Functional Differential Equations With Retarded Arguments, J. Differential Equations, 15 (1974), 269-277.
2. ONOSE, H. Oscillatory Properties of Ordinary Differential Equations of Arbitrary Order, J. Differential Equations, 7 (1970), 454-458.
3. SINGH, B. A Necessary and Sufficient Condition for the Oscillation of an Even Order Nonlinear Delay Differential Equation, Canad. J. Math., 25 (1973), 1078-1089.
4. SINGH, B. Impact of Delays on Oscillations in General Functional Equations, Hiroshima Math. J., 5 (1975), 351-361.
5. SINGH, B. Forced Nonoscillations in Second Order Functional Equations, Hiroshima Math. J., 7 (1977), 657-665.
6. SINGH, B. and KUSANO, T. On Asymptotic Limits of Nonoscillations in Functional Equations with Retarded Arguments, J. Math. Anal. Appl., 10 (1980), 557-565.
7. SHEVELOV, V. N. "Oscillation Theory in Differential Equations with Deviating Arguments," (Russian) Academy of Sciences of Ukrainian CCP (1978).
8. TRENCH W. F. Canonical Forms and Principal Systems for General Disconjugate Equations, Trans. Amer. Math. Soc., 189 (1974), 319-327.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleone@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru