

ON THE RITT ORDER OF A CERTAIN CLASS OF FUNCTIONS

R. PARVATHAM and J. THANGAMANI

The Ramanujan Institute
University of Madras
Madras-600 005, INDIA

(Received August 14, 1981)

ABSTRACT. The authors introduce the notions of Ritt order and lower order to functions defined by the series $\sum_1^{\infty} f_n(s) \exp(-\lambda_n s)$ where (λ_n) is a D-sequence and $f_n(s)$ are entire functions of bounded index.

KEY WORDS AND PHRASES. Ritt order, entire functions of bounded index, Dirichlet series.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. 30A16, 30A64.

1. INTRODUCTION.

Let us consider an M-dirichletian element:

$$\{\phi\}: \sum_{n=1}^{\infty} f_n(s) \exp(-\lambda_n s), \quad s = \sigma + i\tau, \quad (\sigma, \tau) \in \mathbb{R}^2 \quad (1.1)$$

where (λ_n) is a D-sequence (a strictly increasing unbounded sequence of positive numbers) and $f_n(s)$ are entire functions of bounded index (defined below). Convergence properties of such elements were discussed by J.S.J. Mac Donnell in his doctoral dissertation [1] under the conditions $\lim_{n \rightarrow \infty} \frac{\log n}{\lambda_n} = 0$ and $\lim_{n \rightarrow \infty} \frac{m_n}{\lambda_n} = 0$

where m_n is the index of f_n . In this paper, we first study the convergence properties of these elements with less restrictions, namely,

$$L = \limsup_{n \rightarrow \infty} \frac{\log n}{\lambda_n} < \infty \text{ and} \quad (1.2)$$

$$\beta = \limsup_{n \rightarrow \infty} \frac{m_n}{\lambda_n} < \infty \quad (1.3)$$

As the functions defined by these series are unbounded in the half-plane, it is not possible to define Ritt order directly. However, by making use of functions

defined by associated intermediate series, we introduce the notions of Ritt order and lower order to these functions.

2. MAIN RESULTS.

DEFINITION 2.1. [2]. An entire function f is said to be of bounded index if there exists a non-negative integer N such that

$$\max_{0 \leq k \leq N} \left\{ \frac{|f^{(k)}(s)|}{k!} \right\} \geq \frac{|f^{(j)}(s)|}{j!} \quad (f^{(0)}(s) = f(s))$$

for all j and for all s . The least such integer N is called the index of f .

We require the following lemma which shows that an entire function of bounded index is of exponential type.

LEMMA 2.2. [3], [2]. Let f be an entire function of bounded index N . Then

$$|f(s)| \leq \left\{ \max_{0 \leq k \leq N} \frac{|f^{(k)}(0)|}{(N+1)^k} \right\} \exp(N+1) |s|. \tag{2.1}$$

Let $f_n(s) = \sum_{j=0}^{\infty} a_{nj} s^j$ be an entire function of bounded index m_n ;

$$A_n = \max \{ |a_{nj}| / j! : j = 0, 1, \dots, m_n \} = \max \frac{|f_n^{(j)}(0)|}{j!}, \quad j = 0, 1, \dots, m_n \tag{2.2}$$

$\{X\}$: $\sum_1^{\infty} A_n \exp(-\lambda_n s)$ the associated dirichletian element whose abscissa of convergence is denoted by σ_c^X ; $k = \lim_{n \rightarrow \infty} \frac{\log A_n}{\lambda_n}$.

REMARK 2.3. It can be easily seen from Lemma 2.2 that

$$\begin{aligned} |f_n(s)| &\leq \max_{0 \leq j \leq m_n} \left\{ \frac{|f_n^{(j)}(0)|}{j!} \right\} \exp(m_n + 1) |s| \\ &= A_n \exp(m_n + 1) |s|. \end{aligned}$$

THEOREM 2.4. If $0 < \beta < 1$, the region of absolute convergence of (1.1) is the exterior of the hyperbola centre $(k(1 - \beta^2)^{-1}, 0)$ and eccentricity β^{-1} contained in the half-plane $\sigma > k$.

PROOF. Using Remark 2.3, we have

$$|f_n(s) \exp(-\lambda_n s)| \leq A_n \exp(m_n + 1) |s| \exp(-\lambda_n \sigma). \tag{2.3}$$

From the definitions of k and β it follows that for $\epsilon > 0$

$$\begin{aligned} \exists_{n'} \quad \forall_{n \geq n'} \quad A_n < \exp(k + \epsilon)\lambda_n \quad \text{and} \\ \exists_{n''} \quad \forall_{n \geq n''} \quad m_n + 1 < (\beta + \epsilon)\lambda_n. \end{aligned}$$

Hence

$$\forall_{n(\epsilon) = \max(n', n'')} \quad |f_n(s) \exp(-\lambda_n s)| \leq \exp(-\lambda_n(\sigma - k - \epsilon - (\beta + \epsilon)|s|))$$

and

$$\sum_{n(\epsilon)}^{\infty} f_n(s) \exp(-\lambda_n s) \leq \sum_{n(\epsilon)}^{\infty} \exp(-\lambda_n(\sigma - k - \epsilon - (\beta + \epsilon)|s|)).$$

The series in the right hand side converges provided

$$\sigma - k - \beta |s| > 0 \tag{2.4}$$

which is valid only if $\sigma > k$ and $0 \leq \beta < 1$.

Thus any point in the region of convergence of (1.1) must satisfy

$$(\sigma - k)^2 - \beta^2(\sigma^2 + \tau^2) > 0$$

which reduces to

$$\left(\sigma - \frac{k}{1 - \beta^2}\right)^2 - \frac{\beta^2 \tau^2}{1 - \beta^2} > \frac{k^2 \beta^2}{(1 - \beta^2)^2} \tag{2.5}$$

from which the theorem follows.

REMARK 2.5. If $\beta = 0$ the M-dirichletian element converges in the half-plane $\sigma > k$ (which coincides with the half-plane of convergence of the associated series $\{\chi\}$) thus giving the result of MacDonnell [1] as a particular case.

Next we proceed to introduce the notions of Ritt order and lower order for functions defined by (1.1). We need the following lemmas.

Let the M-dirichletian element given by (1.1) converge absolutely on E_a^ϕ and $D_0 = \{s \in \mathbb{C} \mid \sigma = 0, \tau \in \mathbb{R}\}$ denote the imaginary axis.

LEMMA 2.6. Under the conditions $\sigma_c^\chi = -\infty$ and $0 \leq \beta < \infty$, we have $E_a^\phi = \phi$ and ϕ is holomorphic on ϕ .

PROOF. Using (2.3) we have

$$\forall_{n \in \mathbb{N} - \{0\}} \quad \forall_{s \in \phi - D_0} \quad |f_n(s) \exp(-\lambda_n s)| \leq A_n \exp[-\sigma \lambda_n (1 - \frac{m_n + 1}{\lambda_n} \theta_\sigma \frac{|s|}{|\sigma|})]$$

where $\theta_\sigma = 1$ if $\sigma > 0$ and $\theta_\sigma = -1$ if $\sigma < 0$.

Since $0 \leq \beta < \infty$ given $\varepsilon > 0$

$$\begin{aligned} \sum_{n' (=n_\varepsilon)} \forall n \geq n' \quad \forall s \in \mathbb{C} - D_0 \quad & |f_n(s) \exp(-\lambda_n s)| \\ & \leq A_n \exp[-\sigma \lambda_n (1 - (\beta + \varepsilon) \frac{|s|}{|\sigma|} \theta_\sigma)]. \end{aligned} \tag{2.6}$$

For any point (on the imaginary axis) $s_0 = i\tau_0$ of D_0 we extend $\frac{|s|}{|\sigma|} \theta_\sigma$ by its limiting value as $s \rightarrow s_0$.

The function $\mathbb{C} \ni s \rightarrow \sigma[1 - (\beta + \varepsilon) \frac{|s|}{|\sigma|} \theta_\sigma]$ is continuous on \mathbb{C} . Let $E_\mu = \{s \in \mathbb{C} / \sigma[1 - (\beta + \varepsilon) \frac{|s|}{|\sigma|} \theta_\sigma] \geq \mu\}$ indexed by μ on \mathbb{R} . Then $\{\phi_n\}$ converges uniformly on each E_μ as $\mu > \sigma_c^X$, where

$$\{\phi_n\}: \sum_{n=n'}^{\infty} f_n(s) \exp(-\lambda_n s).$$

Let G be any open subset of \mathbb{C} . We put

$$\mu_G = \inf\{\sigma[1 - (\beta + \varepsilon) \frac{|s|}{|\sigma|} \theta_\sigma] \mid s \in G\}.$$

The number μ_G is finite and $\{\phi_n\}$ converges absolutely in G if $\sigma_c^X = -\infty$; further, $\phi_G: G \ni s \rightarrow \phi(s)$ is holomorphic on G . Since G is arbitrary on \mathbb{C} , $\{\phi_n\}$ converges absolutely on each point of \mathbb{C} and ϕ_G can be continued analytically on the totality of \mathbb{C} . Let ϕ denote its analytic continuation. Now we put

$$\forall \sigma \in \mathbb{R} \quad M^\phi(\sigma, B) = \sup\{|\phi(s')| / \sigma' \geq \sigma \quad s \in B(\tau_1, \ell)\}$$

where $B(\tau_1, \ell) = \{s \in \mathbb{C} / |\tau - \tau_1| \leq \ell\}$ is the horizontal strip with $\tau = \tau_1$ as axis and of width 2ℓ . Then

LEMMA 2.7. Under the conditions $\sigma_c^X = -\infty$ and $0 \leq \beta < 1$ we have

$$\forall (\tau_1, \ell) \in \mathbb{R} \times \mathbb{R}_0^+ \quad M^\phi(\sigma, B) \text{ is bounded on each point } \sigma \in \mathbb{R} \text{ and } \lim M^\phi(\sigma, B) = 0$$

as $\sigma \rightarrow +\infty$.

PROOF. Let (τ_1, ℓ) be fixed arbitrarily on $\mathbb{R} \times \mathbb{R}_0^+$. Then given $\varepsilon' > 0$,

$$\exists \sigma_{\varepsilon'} \quad \forall s \in \{\sigma \geq \sigma_{\varepsilon'}, / s \in B(\tau_1, \ell)\} \quad \frac{|s|}{|\sigma|} < 1 + \varepsilon' \quad \text{and} \quad \text{with } 0 \leq \beta < 1. \text{ We have}$$

$$\forall \sigma \geq \sigma_\epsilon, \quad \sigma(1 - (\beta + \epsilon)) \frac{|s|}{|\sigma|} \theta_\sigma > \sigma(1 - (\beta + \epsilon)(1 + \epsilon')) \tag{2.7}$$

$$\geq \sigma_\epsilon (1 - (\beta + \epsilon)(1 + \epsilon'));$$

as a result of(2.6)and (2.7)

$$M^{\phi_{n'}}(\sigma, B) < A_n, \exp(-\sigma_\epsilon, (1 - (\beta + \epsilon)(1 + \epsilon')))$$

$$= \chi_n(\sigma_\epsilon, (1 - (\beta + \epsilon)(1 + \epsilon')))$$

and hence

$$\lim M^{\phi_{n'}}(\sigma, B) = 0 \text{ as } \sigma \rightarrow \infty;$$

finally,

$$\lim M^\phi(\sigma, B) = 0 \text{ as } \sigma \rightarrow \infty .$$

Further, we have

$$\forall \sigma \geq -\sigma_\epsilon, \quad \sigma(1 - (\beta + \epsilon)) \frac{|s|}{|\sigma|} \theta_\sigma > \sigma(1 + (\beta + \epsilon)(1 + \epsilon'));$$

we put $M(\sigma_\epsilon) = \text{Max} \{ |\phi_n(s)| / \sigma \leq \sigma_\epsilon, \wedge s \in B(\tau_1, \ell) \}$. As ϕ_n is holomorphic on the compact set, $\{s \in B(\tau_1, \ell) / |\sigma| \leq \sigma_\epsilon\}$, $M(\sigma_\epsilon)$ is finite. Then we get

$$\forall \sigma \in \mathbb{R} \quad M^{\phi_{n'}}(\sigma, B) \leq \text{Max} \{ \chi_n(\sigma(1 + (\beta + \epsilon)(1 + \epsilon'))), M(\sigma_\epsilon), \chi_n(\sigma_\epsilon, (1 - (\beta + \epsilon)(1 + \epsilon')))\};$$

χ_n is a strictly decreasing function in \mathbb{R} and hence

$$\chi_n(\sigma[1 + \beta + \epsilon)(1 + \epsilon')) > \chi_n(\sigma(1 - (\beta + \epsilon)(1 + \epsilon'))) \quad \text{if } \sigma < 0 \quad \text{and}$$

$$\chi_n(\sigma(1 + (\beta + \epsilon)(1 + \epsilon'))) < \chi_n(\sigma(1 - (\beta + \epsilon)(1 + \epsilon'))) \quad \text{if } \sigma > 0,$$

(equality holds for $\sigma = 0$) with $\chi_n(\sigma) = 0$ as $\sigma \rightarrow \infty$.

As all M-dirichletian polynomials satisfy the two properties of the lemma, in each horizontal strip $B(\tau, \ell)$ $M^\phi(\sigma, B)$ is bounded for each $\sigma \in \mathbb{R}$ and hence the function $\sigma \rightarrow M^\phi(\sigma, B)$ is decreasing on \mathbb{R} with $\lim_{\sigma \rightarrow \infty} M^\phi(\sigma, B) = 0$.

DEFINITION 2.8. We put

$$\rho_B^\phi = \lim_{\sigma \rightarrow -\infty} \sup \left\{ \frac{\log \log^+ M^\phi(\sigma, B)}{-\sigma} \right\} .$$

Then ρ_B^ϕ is called the Ritt order of ϕ on B . Let $\lambda_B^\phi = \lim_{\sigma \rightarrow \infty} \inf \left\{ \frac{\log \log^+ M^\phi(\sigma, B)}{-\sigma} \right\}$.

Then λ_B^ϕ is called the lower order of ϕ on B .

THEOREM 2.9. Under the conditions $\sigma_c^\chi = -\infty$ and $0 \leq \beta < 1$, we have

$$\forall (\tau_1, \ell) \in \mathbb{R} \times \mathbb{R}_0^+ \quad \rho_B^\phi \leq \rho_R^\chi \text{ and } \lambda_B^\phi \leq \lambda_R^\chi$$

where ρ_R^χ and λ_R^χ are respectively the Ritt order and lower order of χ in the whole plane.

PROOF. Proceeding as in Lemma 2.6 for $0 \leq \beta < 1$, we have by (2.6)

$$\forall \varepsilon \in \mathbb{R}_0^+ \exists n' (=n_\varepsilon) \forall s \in B(\tau_1, \ell) |\phi_{n'}(s)| < \chi_n, [\sigma(1 - (\beta + \varepsilon) \frac{|s|}{|\sigma|} \theta_\sigma)] .$$

Now denoting by ϕ_n , and χ_n , the holomorphic functions on \mathbb{C} defined by the elements $\{\phi_n\}$ and $\{\chi_n\}$, we have for σ negative with $|\sigma|$ sufficiently large:

$$M^{\phi_{n'}}(\sigma, B) \leq \chi_n, [\sigma(1 - (\beta + \varepsilon) \frac{|\sigma|}{|\sigma|} \theta_\sigma)]$$

which gives

$$\forall \varepsilon' \in \mathbb{R}_0^+ \rho_B^{\phi_{n'}} \leq \rho_R^{\chi_{n'}} [1 + (\beta + \varepsilon) \varepsilon']$$

and hence $\rho_B^{\phi_{n'}} \leq \rho_R^{\chi_{n'}}$.

As adding finite number of terms to a holomorphic function defined by a classical Dirichlet series does not affect its Ritt order [4], we add $\sum_{n=0}^{n'-1} A_n \exp(-\lambda_n s)$ to $\{\chi_n\}$ and then $\rho_R^{\chi_{n'}} = \rho_R^\chi$.

Now

$$\forall \sigma \in \mathbb{R} M^\phi(\sigma, B) \leq M^{\phi_{n'}}(\sigma, B) + M^{\phi_{n'}^0}(\sigma, B)$$

where $\{\phi_{n'}^0\}$: $\sum_{n=1}^{n'-1} f_n(s) \exp(-\lambda_n s)$; then

$$\begin{aligned} \forall (\tau, \ell) \in \mathbb{R} \times \mathbb{R}_0^+ \rho_B^\phi &\leq \max(\rho_B^{\phi_{n'}}, \rho_B^{\phi_{n'}^0}) \\ &= \rho_B^{\phi_{n'}} \text{ since } \rho_B^{\phi_{n'}^0} = 0. \end{aligned}$$

Finally we have

$$\forall (\tau, \ell) \in \mathbb{R} \times \mathbb{R}_0^+ \rho_B^\phi \leq \rho_R^\chi$$

and similarly we can show:

$$\forall (\tau, \ell) \in \mathbb{R} \times \mathbb{R}_0^+ \lambda_B^\phi \leq \lambda_R^\chi .$$

Now we are in a position to define the Ritt order and lower order of ϕ in the whole plane \mathbb{C} .

DEFINITION 2.10. We put

$$\rho_R^\phi = \sup \{ \rho_B^\phi / (\tau_1, \ell) \in \mathbb{R} \times \mathbb{R}_0^+ \}$$

and

$$\lambda_R^\phi = \sup \{ \lambda_B^\phi / (\tau_1, \ell) \in \mathbb{R} \times \mathbb{R}_0^+ \}$$

Then ρ_R^ϕ is called the Ritt order of ϕ on \mathcal{C} and λ_R^ϕ is called the lower order of ϕ on \mathcal{C} .

REFERENCES

1. MAC DONNELL, J.S.J. Doctoral dissertation, The Catholic University of America, (1957).
2. SHAH, S.M. Entire functions of bounded index, Proc. Amer. Math. Soc. 19, (1968) 1017-1022.
3. HAYMAN, W.K. Differential inequalities and local valency, Pacific J. Math. 44, (1973) 117-137.
4. BLAMBERT, M. Sur la comparaison des ordres boreliens et (R) des fonctions entieres d'une certaene classe, Bull. Sc. Math. 89, (1965) 103-125.

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be