

ASYMPTOTIC RELATIONSHIPS BETWEEN TWO HIGHER ORDER ORDINARY DIFFERENTIAL EQUATIONS

TAKASI KUSANO

Department of Mathematics
Faculty of Science
Hiroshima University
Hiroshima, Japan

(Received April 6, 1982)

ABSTRACT. Some asymptotic relationships between the two ordinary differential equations

$$(1) \quad x^{(n)} + p_1(t)x^{(n-1)} + \dots + p_n(t)x = 0,$$

$$(2) \quad y^{(n)} + p_1(t)y^{(n-1)} + \dots + p_n(t)y = f(t,y),$$

are studied. Conditions are given that lead to an asymptotic equivalence between certain of the solutions of (1) and certain of the solutions of (2). The case where the perturbation $f(t,y)$ depends on a functional argument is also discussed.

KEY WORDS AND PHRASES. *Ordinary differential equations, asymptotic relations, functional differential equations.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 34A.

1. INTRODUCTION.

We are concerned with some asymptotic relationships between the following two differential equations

$$x^{(n)} + p_1(t)x^{(n-1)} + \dots + p_n(t)x = 0, \quad (1.1)$$

$$y^{(n)} + p_1(t)y^{(n-1)} + \dots + p_n(t)y = f(t,y), \quad (1.2)$$

where $p_i: [t_0, \infty) \rightarrow \mathbb{R}$, $1 \leq i \leq n$, and $f: [t_0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions.

We obtain conditions that lead to an asymptotic equivalence between certain of the solutions of the linear equation (1.1) and certain of the solutions of the perturbed linear equation (1.2). No restriction is placed on the behavior of solutions of (1.1) which may be oscillatory or nonoscillatory. The example given at the end of this note deals with the case where (1.1) has both oscillatory and nonoscillatory solutions. Our results generalize those of Hallam [1] for second order equations.

2. MAIN RESULTS.

In what follows we denote by $W(\phi_1, \dots, \phi_m)(t)$ the Wronskian of $\phi_1(t), \dots, \phi_m(t)$.

Let a fundamental system of solutions of (1.1), $\{x_1(t), \dots, x_n(t)\}$, be fixed and suppose that there exist positive continuous functions $x_i^*(t)$, $\xi_i^*(t)$, $1 \leq i \leq n$, which satisfy the inequalities

$$|x_i(t)| \leq x_i^*(t) \quad \text{on} \quad [t_0, \infty), \quad 1 \leq i \leq n, \quad (2.1)$$

$$\left| \frac{W(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(t)}{W(x_1, \dots, x_n)(t)} \right| \leq \xi_i^*(t) \quad \text{on} \quad [t_0, \infty), \quad 1 \leq i \leq n. \quad (2.2)$$

Note that $W(x_1, \dots, x_n)(t)$ is a constant multiple of $\exp\left(-\int_{t_0}^t p_1(s) ds\right)$.

With regard to (1.2) we assume that $f(t, y)$ satisfies the inequality

$$|f(t, y)| \leq \phi(t, |y|) \quad \text{on} \quad [t_0, \infty) \times \mathbb{R}, \quad (2.3)$$

where $\phi : [t_0, \infty) \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$, $\mathbb{R}_+ = [0, \infty)$, is continuous and nondecreasing in the second variable for each fixed t .

THEOREM 1. Suppose conditions (2.1), (2.2), and (2.3) are satisfied. Also suppose that, for some k , $1 \leq k \leq n$, and some constant $c > 1$,

$$\int_{t_0}^{\infty} \xi_i^*(s) \phi(s, cx_k^*(s)) ds < \infty, \quad 1 \leq i \leq n, \quad (2.4)$$

and

$$\frac{x_i^*(t)}{x_k^*(t)} \int_{t_0}^{\infty} \xi_i^*(s) \phi(s, cx_k^*(s)) ds = o(1) \quad \text{as } t \rightarrow \infty \quad (2.5)$$

for $1 \leq i \leq n$ with $i \neq k$.

Then there exists a solution $y(t)$ of equation (1.2) such that

$$y(t) = x_k(t) + o(x_k^*(t)) \quad \text{as } t \rightarrow \infty. \quad (2.6)$$

In addition, for any solution $y_k(t)$ of (1.2) satisfying the inequality

$|y_k(t)| \leq cx_k^*(t)$ on $[t_0, \infty)$, there exists a solution $x(t)$ of (1.1) such that

$$x(t) = y_k(t) + o(x_k^*(t)) \quad \text{as } t \rightarrow \infty. \quad (2.7)$$

PROOF. In view of (2.4) and (2.5) we can choose $t_1 > t_0$ so large that

$$\int_{t_1}^{\infty} \xi_k^*(s) \Phi(s, cx_k^*(s)) ds < \frac{c-1}{2} \quad (2.8)$$

and

$$\frac{x_i^*(t)}{x_k^*(t)} \int_t^{\infty} \xi_i^*(s) \Phi(s, cx_k^*(s)) ds < \frac{c-1}{2(n-1)}, \quad t \geq t_1, \quad (2.9)$$

for $1 \leq i \leq n$ with $i \neq k$. Let $C[t_1, \infty)$ be the locally convex space of continuous functions on $[t, \infty)$ with the compact open topology and consider the closed convex subset F of $C[t_1, \infty)$ defined by

$$F = \{y \in C[t_1, \infty) : |y(t)| \leq cx_k^*(t), \quad t \geq t_1\}.$$

Define the operator $\mathcal{J} : F \rightarrow C[t_1, \infty)$ by the formula

$$y(t) = x_k(t) + \sum_{i=1}^n (-1)^{n-i-1} x_i(t) \int_t^{\infty} \mathfrak{w}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(s) f(s, y(s)) ds, \quad (2.10)$$

where

$$\mathfrak{w}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(t) = \frac{W(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(t)}{W(x_1, \dots, x_n)(t)}. \quad (2.11)$$

We seek for a fixed point $y = y(t)$ of \mathcal{J} in F . Using the identities

$$\sum_{i=0}^n (-1)^{n-i} x_i^{(j)}(t) W(x_1, \dots, x_{i+1}, \dots, x_n)(t) = 0, \quad 0 \leq j \leq n-2, \quad (2.12)$$

$$\sum_{i=0}^n (-1)^{n-i} x_i^{(n-1)}(t) W(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(t) = \mathfrak{w}(x_1, \dots, x_n)(t), \quad (2.13)$$

we easily see that a fixed point of \mathcal{J} is a solution of equation (1.2).

i) \mathcal{J} maps F into F. This follows immediately from (2.1), (2.2), (2.3), (2.8), (2.9) and (2.10).

ii) \mathcal{J} is continuous. Let $\{y_\nu\}$ be a sequence in F converging uniformly to $y \in F$

on every compact subinterval of $[t_1, \infty)$. Consider any compact subinterval of the form $[t_1, t_2]$. Let $\epsilon > 0$ be given and choose $t_3 > t_2$ so that

$$\int_{t_3}^{\infty} \xi_i^*(s) \phi(s, cx_k^*(s)) ds < \frac{\epsilon}{4nM}, \quad 1 \leq i \leq n, \tag{2.14}$$

where $M = \max_{1 \leq i \leq n} \max_{t \in [t_1, t_2]} x_i^*(t)$. Take an integer N such that

$$\xi_i^*(s) |f(s, y_\nu(s)) - f(s, y(s))| < \frac{\epsilon}{2nM(t_3 - t_1)}, \quad s \in [t_1, t_3], \quad 1 \leq i \leq n, \tag{2.15}$$

for all $\nu \geq N$; this is possible since $f(t, y)$ is continuous and $\{y_\nu\}$ converges uniformly to y on $[t_1, t_3]$.

Using (2.14) and (2.15), we have

$$\begin{aligned} |\mathcal{J}y_\nu(t) - \mathcal{J}y(t)| &\leq \sum_{i=1}^n x_i^*(t) \int_{t_1}^{t_3} \xi_i^*(s) |f(s, y_\nu(s)) - f(s, y(s))| ds \\ &\quad + 2 \sum_{i=1}^n x_i^*(t) \int_{t_3}^{\infty} \xi_i^*(s) \phi(s, cx_k^*(s)) ds < \epsilon \end{aligned}$$

for all $t \in [t_1, t_2]$ and all $\nu \geq N$. This shows that \mathcal{J} is continuous on F .

iii) $\mathcal{J}F$ is relatively compact. From (2.10), (2.11) and (2.12) (with $j = 0$) we obtain

$$|(\mathcal{J}y)'(t)| \leq |x_k'(t)| + \sum_{i=1}^n |x_i'(t)| \int_t^{\infty} \xi_i^*(s) \phi(s, cx_k^*(s)) ds.$$

On any compact subinterval of $[t_0, \infty)$ the right-hand side of the above inequality is bounded by a constant independent of $y \in F$. Therefore $\mathcal{J}F$ is equicontinuous on every compact subinterval of $[t_1, \infty)$ and its relative compactness follows from the Ascoli-Arzelà theorem.

Applying now the Schauder-Tychonoff fixed-point theorem, we see that the operator \mathcal{J} has a fixed point $y = y(t)$ in F . As remarked earlier, $y(t)$ is a solution of (1.2).

That $y(t)$ satisfies the order relation (2.6) follows from the inequality

$$|y(t) - x_k(t)| \leq \sum_{i=1}^n x_i^*(t) \int_t^{\infty} \xi_i^*(s) \phi(s, cx_k^*(s)) ds$$

with the help of hypotheses (2.4) and (2.5).

To prove the opposite relationship between the solutions of (1.1) and (1.2), let $y_k(t)$ be a solution of (2.2) satisfying the inequality $|y_k(t)| \leq cx_k^*(t)$ on $[t_0, \infty)$. Then it is easy to verify that the function $x(t)$ defined by

$$x(t) = y_k(t) + \sum_{i=1}^n (-1)^{n-i} x_i^*(t) \int_t^{\infty} \omega(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(s) f(s, y_k(s)) ds \quad (2.16)$$

is a solution of (1.1) which satisfies the order relations (2.7). Thus the proof of Theorem 1 is complete.

THEOREM 2. Let conditions (2.1), (2.2) and (2.3) be satisfied. Suppose that, for some integer k , $1 \leq k \leq n$, and some constant $c > 1$,

$$\int_{t_0}^{\infty} \xi_k^*(s) \phi(s, cx_k^*(s)) ds < \infty \quad (2.17)$$

and

$$\frac{x_i^*(t)}{x_k^*(t)} \int_{t_0}^t \xi_i^*(s) \phi(s, cx_k^*(s)) ds = o(1) \quad \text{as } t \rightarrow \infty \quad (2.18)$$

for $1 \leq i \leq n$ with $i \neq k$.

Then there exists a solution $y(t)$ of (1.2) such that (2.6) is satisfied. In addition, if $y_k(t)$ is any solution of (1.2) satisfying the inequality $|y_k(t)| \leq cx_k^*(t)$, then there exists a solution $x(t)$ of (1) such that (2.7) is satisfied.

PROOF. It suffices to proceed as in the proof of Theorem 1 by replacing (2.10) and (2.16), respectively, by

$$\begin{aligned} \mathcal{J}y(t) = & x_k(t) + (-1)^{n-k-1} x_k(t) \int_t^{\infty} \omega(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n)(s) f(s, y(s)) ds \\ & + \sum_{\substack{i=1 \\ i \neq k}}^n (-1)^{n-i} x_i^*(t) \int_{t_1}^t \omega(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(s) f(s, y(s)) ds \end{aligned} \quad (2.19)$$

and

$$\begin{aligned}
 x(t) = & y_k(t) + (-1)^{n-k} x_k(t) \int_t^\infty \ln(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n)(s) f(s, y_k(s)) ds \\
 & + \sum_{\substack{i=1 \\ i \neq k}}^n (-1)^{n-i-1} x_i(t) \int_{t_1}^t \ln(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)(s) f(s, y_k(s)) ds. \quad (2.20)
 \end{aligned}$$

The details will be left to the reader.

It is not hard to see that the above arguments can be applied to establish similar asymptotic relationship between (1.1) and the functional differential equation

$$y^{(n)}(t) + p_1(t)y^{(n-1)}(t) + \dots + p_n(t)y(t) = f(t, y(g(t))), \quad (2.21)$$

where $p_i(t)$ and $f(t, y)$ are as before and $g : [t_0, \infty) \rightarrow \mathbb{R}$ is a continuous function such that $\lim_{t \rightarrow \infty} g(t) = \infty$. For example, we have the following analogue of Theorem 1.

THEOREM 3. Let conditions (2.1), (2.2), and (2.3) be satisfied. Suppose that, for some $k, 1 \leq k \leq n$, and some constant $c > 1$,

$$\int_{t_0}^\infty \xi_i^*(s) \phi(s, cx_k^*(g(s))) ds < \infty, \quad 1 \leq i \leq n, \quad (2.22)$$

$$\frac{x_i^*(t)}{x_k^*(t)} \int_t^\infty \xi_i^*(s) \phi(s, cx_k^*(g(s))) ds = o(1) \quad \text{as } t \rightarrow \infty \quad (2.23)$$

for $1 \leq i \leq n$ with $i \neq k$.

Then (2.21) has a solution $y(t)$ which satisfies (2.6). In addition, if $y_k(t)$ is a solution of (2.21) satisfying $|y_k(t)| \leq cx_k^*(t)$, then (1.1) has a solution $x(t)$ such that (2.7) is satisfied.

3. EXAMPLE.

Consider the third order differential equations

$$x''' + x = 0, \quad (3.1)$$

$$y''' + y = b(t) |y|^\gamma \text{sgn } y, \quad (3.2)$$

where $\gamma \geq 0$ is a constant and $b : [0, \infty) \rightarrow \mathbb{R}$ is a continuous function. The functions $x_1(t) = (2/\sqrt{3})e^{-t}$, $x_2(t) = e^{t/2} \cos(\sqrt{3}/2)t$, $x_3(t) = e^{t/2} \sin(\sqrt{3}/2)t$ form a fundamental system of solutions of (3.1) such that $W(x_1, x_2, x_3)(t) = 1$. We can take

$$x_1^*(t) = (2/3\sqrt{3})e^{-t}, \quad x_2^*(t) = x_3^*(t) = e^{t/2}, \quad \xi_1^*(t) = (\sqrt{3}/2)e^t, \quad \xi_2^*(t) = \xi_3^*(t) = (2/3)e^{-t/2}.$$

Suppose that

$$\int_0^{\infty} e^{(1-\gamma)t} |b(t)| dt < \infty. \quad (3.3)$$

Theorem 1 (with $k = 1$) then ensures that (3.2) has a solution $y_1(t)$ such that

$$y_1(t) = x_1(t) + o(e^{-t}) \quad \text{as } t \rightarrow \infty. \quad (3.4)$$

Suppose next that

$$\int_0^{\infty} e^{(1+(\gamma/2))t} |b(t)| dt < \infty. \quad (3.5)$$

Then, applying Theorem 1 (with $k = 2$ and $k = 3$), we see that (3.2) has solutions $y_2(t)$ and $y_3(t)$ such that

$$y_2(t) = x_2(t) + o(e^{t/2}) \quad \text{as } t \rightarrow \infty, \quad (3.6)$$

and

$$y_3(t) = x_3(t) + o(e^{t/2}) \quad \text{as } t \rightarrow \infty. \quad (3.7)$$

Obviously, $y_1(t)$ is nonoscillatory, whereas $y_2(t)$ and $y_3(t)$ are oscillatory.

Since (3.5) is stronger than (3.3), (3.5) guarantees the existence of all the three solutions $y_1(t)$, $y_2(t)$, and $y_3(t)$ listed above.

From Theorem 1 it also follows that, in case (3.5) holds, if $y(t)$ is a solution of (3.2) satisfying

$$|y(t)| \leq c e^{t/2} \quad (3.8)$$

for some constant $c > 1$, then there exists a solution $x(t)$ of (3.1) such that $x(t) = y(t) + o(e^{t/2})$ as $t \rightarrow \infty$. Note that not all solutions of (3.2) are subject to this estimate. In fact, equation (3.2) with $\gamma = 3$ and $b(t) = 28e^{-6t}$ has a solution $y(t) = e^{3t}$, even though (3.5) is satisfied.

Finally, consider the functional differential equation

$$y'''(t) + y(t) = b(t)|y(t + \sin t)|^\gamma \operatorname{sgn} y(t + \sin t), \quad (3.9)$$

where γ and $b(t)$ are as above. Appealing to Theorem 1, we conclude that if (3.5) holds, (3.9) has three solutions $y_1(t)$, $y_2(t)$, and $y_3(t)$ with properties (3.4), (3.6), and (3.7), respectively.

ACKNOWLEDGEMENT. This work was done while the author was visiting the Iowa State University. The author wishes to express his sincere thanks to Professor R. S. Dahiya for this invitation and hospitality.

REFERENCE

1. HALLAM, T.G. Asymptotic relationships between the solutions of two second order differential equations, Ann. Polon. Math. 24 (1971), 295-300.

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.otero@usc.es