

SOME INVARIANT THEOREMS ON GEOMETRY OF EINSTEIN NON-SYMMETRIC FIELD THEORY

LIU SHU-LIN

Institute of Mathematics
The Academy of Sciences of China

XU SEN-LIN

Department of Mathematics
University of Science and Technology of China
and

Department of Mathematics
Princeton University, New Jersey 08544

(Received August 30, 1982)

ABSTRACT. This paper generalizes Einstein's theorem. It is shown that under the transformation

$$T_{\Lambda} : U_{ik}^{\ell} \rightarrow \bar{U}_{ik}^{\ell} \equiv U_{ik}^{\ell} + \delta_i^{\ell} \Lambda_k - \delta_k^{\ell} \Lambda_i,$$
curvature tensor $S_{k\ell m}^i(U)$, Ricci tensor $S_{ik}(U)$, and scalar curvature $S(U)$ are all invariant, where $\Lambda = \Lambda_j dx^j$ is a closed 1-differential form on an n-dimensional manifold M.

It is still shown that for arbitrary U, the transformation that makes curvature tensor $S_{k\ell m}^i(U)$ (or Ricci tensor $S_{ik}(U)$) invariant

$$T_v : U_{ik}^{\ell} \rightarrow \bar{U}_{ik}^{\ell} \equiv U_{ik}^{\ell} + V_{ik}^{\ell}$$

must be T_{Λ} transformation, where V (its components are V_{ik}^{ℓ}) is a second order differentiable covariant tensor field with vector value.

KEY WORDS AND PHRASES. Einstein non-symmetric field, Einstein theorem, curvature tensor, Ricci tensor, scalar curvature, T_{Λ} transformation, T_v transformation.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. 53B20, 53B05.

1. INTRODUCTION.

When A. Einstein devoted himself to research on relativism in his symmetric field [1], he regarded non-symmetric g_{ij} (or g^{ij}) and non-symmetric affine connection D (its coefficients are Γ_{ik}^{ℓ} in local coordinates $\{x^i\}$) as independent variables such that the number of independent variables increased from 50 (g_{ij} and Γ_{ik}^{ℓ} are all symmetric for lower coordinates) to 80 (16 g_{ij} or g^{ij} and 64 Γ_{ik}^{ℓ}). With so many covariant variables, it was impossible to choose them according to the

principle of relativism alone. To overcome this difficulty, Einstein introduced a very important concept, transposition invariance. This "transposition invariance" (or transposition symmetry) meant that when all A_{ik} were transposed ($A_{ik}^T = A_{ki}$), all equations were still applicable [2]. Einstein supposed that field equations were transposition invariant. He thought that in physics this hypothesis was equivalent to the law that positive and negative electricity occurred symmetrically.

As the Ricci tensor $R_{ik}(\Gamma)$ represented by connected coefficients Γ_{ik}^ℓ was not transposition invariant, Einstein introduced a "pseudo-tensor" U_{ik}^ℓ instead [3]; its definition was

$$U_{ik}^\ell \equiv \Gamma_{ik}^\ell - \Gamma_{it}^\ell \delta_k^t, \text{ where } \Gamma_{it}^t = \sum_{t=1}^4 \Gamma_{it}^t. \tag{1.1}$$

Denoting Γ_{ik}^ℓ by U_{ik}^ℓ , we obtained

$$\Gamma_{ik}^\ell = U_{ik}^\ell - \frac{1}{3} U_{it}^t \delta_k^\ell \quad (i, k, \ell = 1, \dots, 4). \tag{1.2}$$

Then the Ricci curvature denoted by U was

$$R_{ik} = U_{ik,s}^s - U_{it}^s U_{sk}^t + \frac{1}{3} U_{is}^s U_{tk}^t = S_{ik}(U) = S_{ik}. \tag{1.3}$$

Einstein proved that S_{ik} were transposition invariant and the following.

THEOREM (EINSTEIN). [1] Under the transformation

$$T_\lambda : U_{ik}^\ell \rightarrow \bar{U}_{ik}^\ell \equiv U_{ik}^\ell + \delta_{i,\lambda}^\ell - \delta_{k,\lambda}^\ell, \tag{1.4}$$

the Ricci tensor S_{ik} of U is invariant; i.e., under the transformation (4), there are $\bar{S}_{ik} = S_{ik}$ for arbitrary U , where $\bar{S}_{ik} \equiv S_{ik}(\bar{U})$. In (4), $\lambda, j = \frac{\partial \lambda}{\partial x^j}$ and λ is a differentiable function on a manifold M .

REMARK. Einstein gave transformation (1.4) for $n=4$, but we will still call transformation (1.4) the Einstein transformation for general $n (\geq 2)$.

One asks naturally, how about the converse of Einstein's theorem? A. Einstein and B. Kaufman did not solve the problem. It has remained unsolved.

In this paper, we generalize Einstein and Kaufman's results to an arbitrary $n (\geq 2)$ dimensional manifold M . Objects which we discuss are not limited to the Ricci tensor S_{ik} of U . Besides S_{ik} , we discuss curvature tensor $S_{k\ell m}^i$ and scalar curvature S .

Then, for general $n (\geq 2)$, we give some invariant theorems on curvature tensor $S_{k\ell m}^i$, Ricci tensor S_{ik} and scalar curvature S of U . For this, first we generalize Einstein's transformation. Finally, we give converse theorems of theorems for arbitrary $n (\geq 2)$. These are the main results of this paper. In the special case $n=4$, we answer the problem above mentioned; that is, a converse to Einstein's theorem.

2. DEFINITION AND MAIN RESULTS.

To give the definitions for curvature tensor $S_{k\ell m}^i$ and Ricci tensor S_{ik} of U , first let us give reasonable definitions for curvature tensor $R_{k\ell m}^i$ and Ricci tensor R_{ik} of connection $D (\Gamma_{ik}^\ell)$ (order of lower coordinates is very important; what we give here differs by a minus sign from what is sometimes used, for example, in Pauli's relativism).

$$\begin{aligned} R_{k\ell m}^i &\equiv \Gamma_{k\ell, m}^i - \Gamma_{km, \ell}^i - \Gamma_{s\ell}^i \Gamma_{km}^s + \Gamma_{sm}^i \Gamma_{k\ell}^s \\ &= (\Gamma_{k\ell, m}^i - \Gamma_{s\ell}^i \Gamma_{km}^s) - (\Gamma_{km, \ell}^i - \Gamma_{sm}^i \Gamma_{k\ell}^s) \\ &\equiv [\ell, m] - [m, \ell], \end{aligned} \tag{2.1}$$

In (2.1), let $i=m$. Adding from 1 to n the curvature tensor $R_{k\ell m}^i$ is contracted to obtain the Ricci tensor

$$R_{k\ell} \equiv R_{k\ell s}^s = \Gamma_{k\ell, s}^s - \Gamma_{ks, \ell}^s - \Gamma_{t\ell}^s \Gamma_{ks}^t + \Gamma_{ts}^s \Gamma_{k\ell}^t. \tag{2.2}$$

To establish expressions for the curvature tensor $S_{k\ell m}^i$, Ricci tensor S_{ik} , and scalar curvature S of U for arbitrary n (≥ 2), it is necessary to give transformation between U and Γ for arbitrary n (≥ 2).

In (1.1), let $\ell=k$, add from 1 to n obtaining

$$U_{it}^t = \Gamma_{it}^t - n\Gamma_{it}^t = -(n-1)\Gamma_{it}^t. \tag{2.3}$$

Substituting (2.3) into (1.1), we can solve

$$\Gamma_{ik}^\ell = U_{ik}^\ell - \frac{1}{n-1} \delta_k^\ell U_{it}^t \quad (i, k, \ell = 1, \dots, n) \tag{2.4}$$

From (2.1) - (2.3), and definitions we obtain immediately

PROPOSITION 1. Curvature tensor $S_{k\ell m}^i$, Ricci tensor S_{ik} and scalar curvature S of U are respectively

$$\begin{aligned} (1) \quad R_{k\ell m}^i &= U_{k\ell, m}^i - \frac{1}{n-1} \delta_\ell^i U_{kt, m}^t - U_{s\ell}^i U_{km}^s + \frac{1}{n-1} \delta_\ell^i U_{st}^t U_{km}^s \\ &+ \frac{1}{n-1} U_{kt}^t U_{m\ell}^i - \frac{1}{(n-1)^2} \delta_\ell^i U_{mt}^t U_{ks}^s - U_{km, \ell}^i + \frac{1}{n-1} \delta_m^i U_{kt, \ell}^t + U_{sm}^i U_{k\ell}^s \\ &- \frac{1}{n-1} \delta_m^i U_{st}^t U_{k\ell}^s - \frac{1}{n-1} U_{kt}^t U_{\ell m}^i + \frac{1}{(n-1)^2} U_{\ell t}^t U_{ks}^s \equiv S_{k\ell m}^i(U) \equiv S_{k\ell m}^i. \end{aligned} \tag{2.5}$$

$$(2) \quad R_{ik} = U_{ik, s}^s - U_{it}^s U_{sk}^t + \frac{1}{n-1} U_{is}^s U_{tk}^t \equiv S_{ik}(U) \equiv S_{ik}. \tag{2.6}$$

$$(3) \quad S \equiv g^{ik} R_{ik} = g^{ik} U_{ik, s}^s - g^{ik} U_{it}^s U_{sk}^t + \frac{1}{n-1} g^{ik} U_{is}^s U_{tk}^t \equiv S(U) \tag{2.7}$$

When $n \geq 2$, it is not difficult to verify that Ricci tensor S_{ik} and scalar curvature S of U are transposition invariant.

THEOREM 1. Curvature tensor $S_{k\ell m}^i$, Ricci tensor S_{ik} and scalar curvature S of U are all invariant under the following transformation

$$T_\Lambda : U_{ik}^\ell \rightarrow \bar{U}_{ik}^\ell \equiv U_{ik}^\ell + \delta_i^\ell \Lambda_k - \delta_k^\ell \Lambda_i, \tag{2.8}$$

where $\Lambda = \Lambda_j dx^j$ is a closed 1-differential form on a manifold M ; i.e., $d\Lambda = 0$.

REMARK. The transformation (2.8) is a generalization of Einstein's transformation (1.4). In fact, as Λ is a closed 1-differential form on a manifold M ($d\Lambda = 0$), then by the Poincaré Lemma, there exists a coordinate neighborhood $M_1 \subset M$ and a differentiable function λ such that $\Lambda_k = \frac{\partial \lambda}{\partial x^k} = \lambda_{,k}$ ($k=1, \dots, n$). Therefore, in a local neighborhood, for example M_1 , the transformation (2.8) conforms with Einstein's transformation (1.4).

Because an exact differential form $d\lambda$ is a closed differential form ($d^2\lambda = 0$), T_λ is a transformation which makes $S_{k\ell m}^i$, S_{ik} and S invariant. When $n=4$, Einstein's theorem is a special case of the above theorem 1.

PROOF. In local coordinates $\{x^i\}$, let $\Lambda_i = \lambda_{,i}$, then

$$\begin{aligned} \bar{S}_{k\ell m}^i &= S_{k\ell m}^i(\bar{U}) \\ &= (U_{k\ell, m}^i + \delta_{k^\lambda, \ell m}^i - \delta_{\ell^\lambda, km}^i) - \frac{1}{n-1} \delta_\ell^i (U_{kt, m}^t + \delta_{k^\lambda, tm}^t - \delta_{t^\lambda, km}^t) \\ &\quad - (U_{s\ell}^i + \delta_{s^\lambda, \ell}^i - \delta_{\ell^\lambda, s}^i) (U_{km}^s + \delta_{k^\lambda, m}^s - \delta_{m^\lambda, k}^s) \\ &\quad + \frac{1}{n-1} \delta_\ell^i (U_{st}^t + \delta_{s^\lambda, t}^t - \delta_{t^\lambda, s}^t) (U_{km}^s + \delta_{k^\lambda, m}^s - \delta_{m^\lambda, k}^s) \\ &\quad + \frac{1}{n-1} (U_{kt}^t + \delta_{k^\lambda, t}^t - \delta_{t^\lambda, k}^t) (U_{m\ell}^i + \delta_{m^\lambda, \ell}^i - \delta_{\ell^\lambda, m}^i) \\ &\quad - \frac{1}{(n-1)^2} \delta_\ell^i (U_{mt}^t + \delta_{m^\lambda, t}^t - \delta_{t^\lambda, m}^t) (U_{ks}^s + \delta_{k^\lambda, s}^s - \delta_{s^\lambda, k}^s) \\ &\quad - (U_{km, \ell}^i + \delta_{k^\lambda, m\ell}^i - \delta_{m^\lambda, k\ell}^i) + \frac{1}{n-1} \delta_m^i (U_{kt, \ell}^t + \delta_{k^\lambda, t\ell}^t - \delta_{t^\lambda, k\ell}^t) \\ &\quad + (U_{sm}^i + \delta_{s^\lambda, m}^i - \delta_{m^\lambda, s}^i) (U_{k\ell}^s + \delta_{k^\lambda, \ell}^s - \delta_{\ell^\lambda, k}^s) \\ &\quad - \frac{1}{n-1} \delta_m^i (U_{st}^t + \delta_{s^\lambda, t}^t - \delta_{t^\lambda, s}^t) (U_{k\ell}^s + \delta_{k^\lambda, \ell}^s - \delta_{\ell^\lambda, k}^s) \\ &\quad - \frac{1}{n-1} (U_{kt}^t + \delta_{k^\lambda, t}^t - \delta_{t^\lambda, k}^t) (U_{\ell m}^i + \delta_{\ell^\lambda, m}^i - \delta_{m^\lambda, \ell}^i) \\ &\quad + \frac{1}{(n-1)^2} \delta_m^i (U_{\ell t}^t + \delta_{\ell^\lambda, t}^t - \delta_{t^\lambda, \ell}^t) (U_{ks}^s + \delta_{k^\lambda, s}^s - \delta_{s^\lambda, k}^s) \\ &= S_{k\ell m}^i(U) - U_{k\ell}^i \lambda_{, m} + U_{m\ell}^i \lambda_{, k} - U_{km}^i \lambda_{, \ell} - \delta_{k^\lambda, \ell}^i \lambda_{, m} \\ &\quad + \delta_{m^\lambda, \ell}^i \lambda_{, k} + \delta_{\ell^\lambda, s}^i U_{km}^s + \delta_{\ell^\lambda, k}^i \lambda_{, m} - \delta_{\ell^\lambda, m}^i \lambda_{, k} \\ &\quad + \frac{1}{n-1} \delta_{\ell^\lambda, m}^i U_{kt}^t - \frac{1}{n-1} \delta_{\ell^\lambda, k}^i U_{mt}^t + \frac{1}{n-1} \delta_{\ell^\lambda, s}^i U_{km}^s + \frac{1}{n-1} \delta_{\ell^\lambda, k}^i \lambda_{, m} \\ &\quad - \frac{1}{n-1} \delta_{\ell^\lambda, m}^i \lambda_{, k} - \frac{n}{n-1} \delta_{\ell^\lambda, s}^i U_{km}^s - \frac{n}{n-1} \delta_{\ell^\lambda, k}^i \lambda_{, m} \\ &\quad + \frac{n}{n-1} \delta_{\ell^\lambda, m}^i \lambda_{, k} + \frac{1}{n-1} \delta_{m^\lambda, \ell}^i U_{kt}^t - \frac{1}{n-1} \delta_{\ell^\lambda, m}^i U_{kt}^t \\ &\quad + \frac{1}{n-1} \lambda_{, k} U_{m\ell}^i + \frac{1}{n-1} \delta_{m^\lambda, k}^i \lambda_{, \ell} - \frac{1}{n-1} \delta_{\ell^\lambda, k}^i \lambda_{, m} \\ &\quad - \frac{n}{n-1} \lambda_{, k} U_{m\ell}^i - \frac{n}{n-1} \delta_{m^\lambda, k}^i \lambda_{, \ell} + \frac{n}{n-1} \delta_{\ell^\lambda, k}^i \lambda_{, m} \\ &\quad - \frac{1}{(n-1)^2} \delta_{\ell^\lambda, k}^i U_{mt}^t + \frac{n}{(n-1)^2} \delta_{\ell^\lambda, k}^i U_{mt}^t - \frac{1}{(n-1)^2} \delta_{\ell^\lambda, m}^i U_{ks}^s \\ &\quad - \frac{1}{(n-1)^2} \delta_{\ell^\lambda, m}^i \lambda_{, k} + \frac{n}{(n-1)^2} \delta_{\ell^\lambda, m}^i \lambda_{, k} + \frac{n}{(n-1)^2} \delta_{\ell^\lambda, m}^i U_{ks}^s \\ &\quad + \frac{n}{(n-1)^2} \delta_{\ell^\lambda, m}^i \lambda_{, k} - \frac{n^2}{(n-1)^2} \delta_{\ell^\lambda, m}^i \lambda_{, k} - [m, \ell] = S_{k\ell m}^i, \end{aligned}$$

where $[m, \ell] = -\lambda_{, \ell} U_{km}^i + \dots - \frac{n^2}{(n-1)^2} \delta_{m^\lambda, \ell}^i \lambda_{, k}$.

Consider the transformation

$$T_{\Omega} : U_{ik}^{\ell} \rightarrow \bar{U}_{ik}^{\ell} \equiv U_{ik}^{\ell} + \delta_{i\Omega_k}^{\ell} - \delta_k^{\ell}\Omega_i, \tag{2.9}$$

where Ω is a 1-differential form, in local coordinates $\{x^i\}$, $\Omega = \Omega_j dx^j$.

Having the above result for theorem 1, we ask naturally if the transformation which makes curvature tensor $S_{k\ell m}^i$ (or Ricci tensor S_{ik}) invariant is the transformation T_{Ω} ? For this, although we cannot give the complete answer - it is a very difficult problem - we have the following results.

THEOREM 2. The transformation that makes curvature tensor $S_{k\ell m}^i$ (or Ricci tensor S_{ik}) of some U invariant

$$T_{\Omega} : U_{ik}^{\ell} \rightarrow \bar{U}_{ik}^{\ell} \equiv U_{ik}^{\ell} + \delta_{i\Omega_k}^{\ell} - \delta_k^{\ell}\Omega_i$$

must be T_{Ω} , where $\Omega = \Omega_j dx^j$ is a 1-differential form.

PROOF. Similar to the proof of theorem 1, we obtain $\bar{S}_{k\ell m}^i = S_{k\ell m}^i(\bar{U})$

$$\begin{aligned} &= (U_{k\ell, m}^i + \delta_{k\Omega_{\ell, m}}^i - \delta_{\ell\Omega_{k, m}}^i) - \frac{1}{n-1} \delta_{\ell}^i (U_{kt, m}^t + \delta_{k\Omega_t, m}^t - \delta_t^t\Omega_{k, m}) \\ &- (U_{s\ell}^i + \delta_{s\Omega_{\ell}}^i - \delta_{\ell\Omega_s}^i) (U_{km}^s + \delta_{k\Omega_m}^s - \delta_m^s\Omega_k) \\ &+ \frac{1}{n-1} \delta_{\ell}^i (U_{st}^t + \delta_{s\Omega_t}^t - \delta_t^t\Omega_s) (U_{km}^s + \delta_{k\Omega_m}^s - \delta_m^s\Omega_k) \\ &+ \frac{1}{n-1} (U_{kt}^t + \delta_{k\Omega_t}^t - \delta_t^t\Omega_k) (U_{m\ell}^i + \delta_{m\Omega_{\ell}}^i - \delta_{\ell\Omega_m}^i) \\ &- \frac{1}{(n-1)^2} \delta_{\ell}^i (U_{mt}^t + \delta_{m\Omega_t}^t - \delta_t^t\Omega_m) (U_{ks}^s + \delta_{k\Omega_s}^s - \delta_s^s\Omega_k) - [m, \ell] \\ &= S_{k\ell m}^i(U) + (\delta_{k\Omega_{\ell, m}}^i - \delta_{\ell\Omega_{k, m}}^i) - \frac{1}{n-1} \delta_{\ell}^i (\delta_{k\Omega_t, m}^t - \delta_t^t\Omega_{k, m}) \\ &- U_{k\ell\Omega_m}^i + U_{m\ell\Omega_k}^i - U_{km\Omega_{\ell}}^i - \delta_{k\Omega_{\ell}\Omega_m}^i + \delta_{m\Omega_{\ell}\Omega_k}^i + \delta_{\ell\Omega_s}^i U_{ksm}^s \\ &+ \delta_{\ell\Omega_k\Omega_m}^i - \delta_{\ell\Omega_m\Omega_k}^i + \frac{1}{n-1} \delta_{\ell}^i U_{kt\Omega_m}^t - \frac{1}{n-1} \delta_{\ell}^i U_{mt\Omega_k}^t + \frac{1}{n-1} \delta_{\ell}^i U_{s\Omega_{km}}^s \\ &+ \frac{1}{n-1} \delta_{\ell}^i \Omega_{k\Omega_m} - \frac{1}{n-1} \delta_{\ell}^i \Omega_{m\Omega_k} - \frac{n}{n-1} \delta_{\ell}^i U_{ksm}^s - \frac{n}{n-1} \delta_{\ell}^i \Omega_{k\Omega_m} \\ &+ \frac{n}{n-1} \delta_{\ell}^i \Omega_{m\Omega_k} + \frac{1}{n-1} \delta_{m\Omega_{kt}\Omega_{\ell}}^i - \frac{1}{n-1} \delta_{\ell}^i U_{kt\Omega_m}^t + \frac{1}{n-1} \Omega_k U_{m\ell}^i \\ &+ \frac{1}{n-1} \delta_{m\Omega_k\Omega_{\ell}}^i - \frac{1}{n-1} \delta_{\ell}^i \Omega_{k\Omega_m} - \frac{n}{n-1} \Omega_k U_{m\ell}^i - \frac{n}{n-1} \delta_{m\Omega_k\Omega_{\ell}}^i \\ &+ \frac{n}{n-1} \delta_{\ell}^i \Omega_{k\Omega_m} - \frac{1}{(n-1)^2} \delta_{\ell}^i U_{mt\Omega_k}^t + \frac{n}{(n-1)^2} \delta_{\ell}^i U_{mt\Omega_k}^t - \frac{1}{(n-1)^2} \delta_{\ell}^i U_{ms}^s \\ &- \frac{1}{(n-1)^2} \delta_{\ell}^i \Omega_{m\Omega_k} + \frac{n}{(n-1)^2} \delta_{\ell}^i \Omega_{m\Omega_k} + \frac{n}{(n-1)^2} \delta_{\ell}^i U_{ksm}^s + \frac{n}{(n-1)^2} \delta_{\ell}^i \Omega_{m\Omega_k} \\ &- \frac{n^2}{(n-1)^2} \delta_{\ell}^i \Omega_{m\Omega_k} - [m, \ell] = S_{k\ell m}^i + \delta_k^i (\Omega_{\ell, m} - \Omega_{m, \ell}). \end{aligned}$$

Therefore, $\bar{S}_{k\ell m}^i = S_{k\ell m}^i$ if and only if $\Omega_{\ell, m} = \Omega_{m, \ell}$, i.e., $d\Omega = 0$.

From $\bar{S}_{k\ell} = \bar{S}_{k\ell i}^i = S_{k\ell i}^i + \delta_k^i (\Omega_{\ell, i} - \Omega_{i, \ell}) = S_{k\ell} + (\Omega_{\ell, k} - \Omega_{k, \ell})$ it follows that $\bar{S}_{k\ell} = S_{k\ell}$ if and only if

$$\Omega_{\ell, k} = \Omega_{k, \ell},$$

i.e., $d\Omega = 0$.

THEOREM 3. A necessary and sufficient condition that transformation T_Ω makes scalar curvature S of some U invariant is

$$g^{ik} \Omega_{k, i} - g^{ik} \Omega_{i, k} = 0.$$

where $\Omega_{k, i} = \frac{\partial \Omega_k}{\partial x^i}$.

PROOF. From $\bar{S} = g^{ik} \bar{S}_{ik} = g^{ik} S_{ik} + g^{ik} (\Omega_{k, i} - \Omega_{i, k}) = S + (g^{ik} \Omega_{k, i} - g^{ik} \Omega_{i, k})$ it follows that $\bar{S} = S$ if and only if

$$g^{ik} \Omega_{k, i} - g^{ik} \Omega_{i, k} = 0.$$

REMARK. If g^{ik} are symmetric, then $g^{ik} \Omega_{k, i} - g^{ik} \Omega_{i, k} = 0$. If g^{ik} are not symmetric, for example $g^{12} \neq g^{21}$, let

$$\Omega_k = \begin{cases} x^2, & k = 1, \\ 0, & k = 2, \dots, n, \end{cases}$$

then $g^{ik} \Omega_{k, i} - g^{ik} \Omega_{i, k} = g^{21} - g^{12} \neq 0$.

Now we give the converse of theorem 1. For this, what we must emphasize is that because of theorem 1, the transformation T_Λ makes curvature tensor $S_{k\ell m}^i$ and Ricci tensor S_{ik} of every U invariant.

The following theorems, 4 and 5 respectively, are the converses of theorem 1 on curvature tensor $S_{k\ell m}^i$ and Ricci tensor S_{ik} .

THEOREM 4. Let V be a second order differentiable covariant tensor field with vector value and its components be V_{ik}^ℓ in local coordinates $\{x^i\}$. If the transformation

$$T_V : U_{ik}^\ell \rightarrow \bar{U}_{ik}^\ell \equiv U_{ik}^\ell + V_{ik}^\ell \quad (i, k, \ell = 1, \dots, n) \tag{2.10}$$

makes curvature tensor $S_{k\ell m}^i$ of every U invariant, then it implies

$$V_{ik}^\ell = \delta_i^\ell \Lambda_k - \delta_k^\ell \Lambda_i,$$

where $\Lambda = \Lambda_j dx^j$ is a closed 1-differential form; i.e., T_V must be T_Λ .

PROOF. By (2.10),

$$\begin{aligned} \bar{S}_{k\ell m}^i &= S_{k\ell m}^i(\bar{U}) = (U_{k\ell, m}^i + V_{k\ell, m}^i) - \frac{1}{n-1} \delta_\ell^i (U_{kt, m}^t + V_{kt, m}^t) \\ &- (U_{s\ell}^i + V_{s\ell}^i) (U_{km}^s + V_{km}^s) + \frac{1}{n-1} \delta_\ell^i (U_{st}^t + V_{st}^t) (U_{km}^s + V_{km}^s) \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{n-1} (U_{kt}^t + V_{kt}^t) (U_{m\ell}^i + V_{m\ell}^i) - \frac{1}{(n-1)^2} \delta_\ell^i (U_{mt}^t + V_{mt}^t) (U_{ks}^s + V_{ks}^s) \\
 - [m, \ell] & = S_{k\ell m}^i + V_{k\ell, m}^i - \frac{1}{n-1} V_{kt, m}^t \delta_\ell^i - V_{s\ell}^i U_{km}^s - U_{s\ell}^i V_{km}^s \\
 & - V_{s\ell}^i V_{km}^s + \frac{1}{n-1} \delta_\ell^i V_{st}^t U_{km}^s + \frac{1}{n-1} \delta_\ell^i V_{km}^s U_{st}^t + \frac{1}{n-1} \delta_\ell^i V_{st}^t V_{km}^s + \frac{1}{n-1} U_{kt}^t V_{m\ell}^i \\
 & + \frac{1}{n-1} V_{kt}^t U_{m\ell}^i + \frac{1}{n-1} V_{m\ell}^i V_{kt}^t - \frac{1}{(n-1)^2} \delta_\ell^i U_{mt}^t V_{ks}^s - \frac{1}{(n-1)^2} \delta_\ell^i U_{ks}^s V_{mt}^t - \frac{1}{(n-1)^2} \delta_\ell^i V_{mt}^t V_{ks}^s \\
 - [m, \ell] & = S_{k\ell m}^i + F_{k\ell m}^i.
 \end{aligned}$$

Since $\bar{S}_{k\ell m}^i = S_{k\ell m}^i$ (for every U), now $F_{k\ell m}^i(U) = 0$ ($i, k, \ell, m = 1, \dots, n$ and for every U). Therefore,

$$\begin{aligned}
 0 & = \frac{\partial F_{k\ell m}^i}{\partial U_{\alpha\beta}^\gamma} = -V_{\gamma\ell}^i \delta_k^\alpha \delta_m^\beta - V_{km}^\alpha \delta_\gamma^\beta + \frac{1}{n-1} \delta_\ell^i V_{\gamma t}^t \delta_k^\alpha \delta_m^\beta \\
 & + \frac{1}{n-1} \delta_\ell^i V_{km}^\alpha \delta_\gamma^\beta + \frac{1}{n-1} V_{m\ell}^i \delta_k^\alpha \delta_\gamma^\beta + \frac{1}{n-1} V_{kt}^t \delta_\gamma^i \delta_m^\alpha \delta_\ell^\beta - \frac{1}{(n-1)^2} \delta_\ell^i V_{ks}^s \delta_m^\alpha \delta_\gamma^\beta \\
 & - \frac{1}{(n-1)^2} \delta_\ell^i V_{mt}^t \delta_k^\alpha \delta_\gamma^\beta + V_{\gamma m}^i \delta_k^\alpha \delta_\ell^\beta + \delta_\gamma^i \delta_\ell^\beta V_{km}^\alpha - \frac{1}{n-1} \delta_m^i V_{\gamma t}^t \delta_k^\alpha \delta_\ell^\beta \\
 & - \frac{1}{n-1} \delta_m^i V_{k\ell}^\alpha \delta_\gamma^\beta - \frac{1}{n-1} V_{\ell m}^i \delta_k^\alpha \delta_\gamma^\beta - \frac{1}{n-1} V_{kt}^t \delta_\gamma^i \delta_\ell^\alpha \delta_m^\beta + \frac{1}{(n-1)^2} \delta_m^i V_{ks}^s \delta_\ell^\alpha \delta_\gamma^\beta \\
 & + \frac{1}{(n-1)^2} \delta_m^i V_{\ell t}^t \delta_k^\alpha \delta_\gamma^\beta.
 \end{aligned}$$

In the above formula, let $i = \ell$. Adding from 1 to n we obtain

$$\begin{aligned}
 0 & = \frac{\partial F_{k\ell m}^\ell}{\partial U_{\alpha\beta}^\gamma} = -V_{\gamma\ell}^\ell \delta_k^\alpha \delta_m^\beta - \delta_\gamma^\beta V_{km}^\alpha + \frac{n}{n-1} V_{\gamma t}^t \delta_k^\alpha \delta_m^\beta + \frac{n}{n-1} V_{km}^\alpha \delta_\gamma^\beta \\
 & + \frac{1}{n-1} V_{m\ell}^\ell \delta_k^\alpha \delta_\gamma^\beta + \frac{1}{n-1} V_{kt}^t \delta_\gamma^\beta \delta_m^\alpha - \frac{n}{(n-1)^2} V_{ks}^s \delta_m^\alpha \delta_\gamma^\beta - \frac{n}{(n-1)^2} V_{mt}^t \delta_k^\alpha \delta_\gamma^\beta \\
 & + V_{\gamma m}^\beta \delta_k^\alpha + \delta_m^\beta V_{k\gamma}^\alpha - \frac{1}{n-1} V_{\gamma t}^t \delta_k^\alpha \delta_m^\beta - \frac{1}{n-1} V_{km}^\alpha \delta_\gamma^\beta - \frac{1}{n-1} V_{\ell m}^\ell \delta_k^\alpha \delta_\gamma^\beta \\
 & - \frac{1}{n-1} V_{kt}^t \delta_\gamma^\alpha \delta_m^\beta + \frac{1}{(n-1)^2} V_{ks}^s \delta_m^\alpha \delta_\gamma^\beta + \frac{1}{(n-1)^2} V_{mt}^t \delta_k^\alpha \delta_\gamma^\beta \\
 & = \delta_k^\alpha V_{\gamma m}^\beta + \delta_m^\beta V_{k\gamma}^\alpha - \frac{1}{n-1} V_{\ell m}^\ell \delta_k^\alpha \delta_\gamma^\beta - \frac{1}{n-1} V_{kt}^t \delta_\gamma^\alpha \delta_m^\beta.
 \end{aligned}$$

Let $\alpha = k$. Adding from 1 to n we obtain

$$0 = nV_{\gamma m}^\beta + \delta_m^\beta V_{k\gamma}^\alpha - \frac{n}{n-1} V_{\ell m}^\ell \delta_\gamma^\beta - \frac{1}{n-1} V_{\gamma t}^t \delta_\gamma^\beta. \tag{2.11}$$

In (2.11) let $\beta = \gamma$ again. Adding from 1 to n we obtain

$$nV_{\beta m}^{\beta} + V_{\alpha m}^{\alpha} - \frac{n^2}{n-1} V_{\ell m}^{\ell} - \frac{1}{n-1} V_{mt}^t = -\frac{V_{\beta m}^{\beta}}{n-1} - \frac{V_{mt}^t}{n-1} = 0,$$

$$-V_{mt}^t = V_{tm}^t \quad (= V_m^t).$$

Substituting the above formula into (2.11) to obtain

$$V_{\gamma m}^{\beta} = \delta_{\gamma}^{\beta} \frac{V_m}{n-1} - \delta_m^{\beta} \frac{V_{\gamma}}{n-1} = \delta_{\gamma}^{\beta} \Omega_m - \delta_m^{\beta} \Omega_{\gamma},$$

where $\Omega_{\gamma} = \frac{V_{\gamma}}{n-1} = \frac{V_{t\gamma}^t}{n-1}.$

This proves that the transformation T_V must be T_{Ω} ; then, according to theorem 2, $\Omega = \Omega_j dx^j$ must be a closed differential form; i.e., $\Omega = \Lambda.$

The analogous result for S_{ik} is stronger.

THEOREM 5. The transformation T_V that makes the Ricci tensor S_{ik} of arbitrary U invariant must be the transformation $T_{\Lambda}.$

PROOF. In fact,

$$\begin{aligned} \bar{S}_{ik} &= S_{ik}(\bar{U}) \equiv \bar{U}_{ik,s}^s - \bar{U}_{it}^s \bar{U}_{sk}^t + \frac{1}{n-1} \bar{U}_{is}^s \bar{U}_{tk}^t \\ &= (U_{ik,s}^s + V_{ik,s}^s) - (U_{it}^s + V_{it}^s) (U_{sk}^t + V_{sk}^t) + \frac{1}{n-1} (U_{is}^s + V_{is}^s) (U_{tk}^t + V_{tk}^t) \\ &= S_{ik} + F_{ik}, \end{aligned}$$

where,

$$\begin{aligned} F_{ik} &= V_{ik,s}^s - U_{it}^s V_{sk}^t - V_{it}^s U_{sk}^t - V_{it}^s V_{sk}^t + \frac{1}{n-1} U_{is}^s V_{tk}^t \\ &+ \frac{1}{n-1} V_{is}^s U_{tk}^t + \frac{1}{n-1} V_{is}^s V_{tk}^t. \end{aligned}$$

According to the condition of the theorem, $\bar{S}_{ik} = S_{ik},$ and we obtain $F_{ik} = 0$ (for arbitrary U). Therefore,

$$\begin{aligned} 0 &= \frac{\partial F_{ik}}{\partial U_{\alpha\beta}^{\gamma}} = -\delta_{\gamma}^s \delta_i^{\alpha} \delta_t^{\beta} V_{sk}^t - \delta_{\gamma}^t \delta_s^{\alpha} \delta_k^{\beta} V_{it}^s + \frac{1}{n-1} \delta_{\gamma}^s \delta_i^{\alpha} \delta_s^{\beta} V_{tk}^t \\ &+ \frac{1}{n-1} \delta_{\gamma}^t \delta_t^{\alpha} \delta_k^{\beta} V_{is}^s = -\delta_i^{\alpha} V_{\gamma k}^{\beta} - \delta_k^{\beta} V_{i\gamma}^{\alpha} + \frac{1}{n-1} \delta_i^{\alpha} \delta_{\gamma}^{\beta} V_{tk}^t + \frac{1}{n-1} \delta_{\gamma}^{\alpha} \delta_k^{\beta} V_{is}^s. \end{aligned}$$

In the above formula, let $\alpha = \gamma.$ Adding from 1 to n we obtain

$$\begin{aligned} -V_{ik}^{\beta} - \delta_k^{\beta} V_{it}^t + \frac{1}{n-1} \delta_i^{\beta} V_{tk}^t + \frac{n}{n-1} \delta_k^{\beta} V_{is}^s &= 0, \\ -V_{ik}^{\beta} + \frac{1}{n-1} \delta_i^{\beta} V_{tk}^t + \frac{1}{n-1} \delta_k^{\beta} V_{it}^t &= 0, \\ V_{ik}^{\beta} = \delta_i^{\beta} \left(\frac{1}{n-1} V_{tk}^t \right) + \delta_k^{\beta} \left(\frac{1}{n-1} V_{it}^t \right). \end{aligned} \tag{2.12}$$

In the above formula, let $\beta = i;$ again, adding from 1 to n we obtain

$$V_k = V_{tk}^t = \frac{n}{n-1} V_{tk}^t + \frac{1}{n-1} V_{kt}^t = \frac{n}{n-1} V_k + \frac{1}{n-1} V_{kt}^t,$$

$$V_k = -V_{kt}^t = V_{tk}^t.$$

Substituting the above formula into (2.12) we obtain (note $\Omega_k = \frac{1}{n-1} V_k$)

$$V_{ik} = \delta_i^\beta \Omega_k - \delta_i^\beta \Omega_i.$$

From theorem 2, it follows that $\Omega = \Omega_j dx^j$ is a closed differential form; namely T_V must be T_Λ .

Moreover, we have the following

THEOREM 6. The transformation T_V that makes scalar curvature $S = g^{ik} S_{ik}$ of arbitrary U invariant must satisfy the following system of equations

$$g^{\alpha k} (V_{\gamma k}^\beta - \frac{1}{n-1} \delta_\gamma^\beta V_k) + g^{i\beta} (V_{i\gamma}^\alpha + \frac{1}{n-1} \delta_\gamma^\alpha V_i) = 0, \tag{2.13}$$

where $V_k = V_{tk}^t = -V_{kt}^t$.

PROOF. From (2.7), we have

$$\begin{aligned} \bar{S} &= g^{ik} \bar{S}_{ik} = g^{ik} U_{ik,s}^s - g^{ik-s} U_{it}^t U_{sk}^s + \frac{1}{n-1} g^{ik-s} U_{is}^s U_{tk}^t \\ &= g^{ik} (U_{ik,s}^s + V_{ik,s}^s) - g^{ik} (U_{it}^s + V_{it}^s) (U_{sk}^t + V_{sk}^t) \\ &+ \frac{1}{n-1} g^{ik} (U_{is}^s + V_{is}^s) (U_{tk}^t + V_{tk}^t) = S + F, \end{aligned}$$

where,

$$\begin{aligned} F &= g^{ik} V_{ik,s}^s - g^{ik} U_{it}^s V_{sk}^t - g^{ik} V_{it}^s U_{sk}^t - g^{ik} V_{it}^s V_{sk}^t \\ &+ \frac{1}{n-1} g^{ik} U_{is}^s V_{tk}^t + \frac{1}{n-1} g^{ik} V_{is}^s U_{tk}^t + \frac{1}{n-1} g^{ik} V_{is}^s V_{tk}^t. \end{aligned}$$

From the condition of the theorem, $\bar{S} = S$, it follows that $F = 0$ (for arbitrary U). Therefore,

$$\begin{aligned} 0 &= \frac{\partial F}{\partial U_{\alpha\beta}^\gamma} = -g^{ik} \delta_i^\alpha V_{\gamma k}^\beta - g^{ik} V_{i\gamma}^\alpha \delta_k^\beta + \frac{1}{n-1} g^{ik} \delta_\gamma^\beta \delta_i^\alpha V_{tk}^t \\ &+ \frac{1}{n-1} g^{ik} V_{is}^s \delta_\gamma^\alpha \delta_k^\beta = -g^{\alpha k} V_{\gamma k}^\beta - g^{i\beta} V_{i\gamma}^\alpha + \frac{1}{n-1} g^{\alpha k} \delta_\gamma^\beta V_{tk}^t + \frac{1}{n-1} g^{i\beta} \delta_\gamma^\alpha V_{is}^s \tag{2.14} \end{aligned}$$

Multiply the above formula by $g_{\alpha\beta}$, adding from 1 to n for α and β to obtain

$$\begin{aligned} -g_{\alpha\beta} g^{\alpha k} V_{\gamma k}^\beta - g_{\alpha\beta} g^{i\beta} V_{i\gamma}^\alpha + \frac{1}{n-1} g_{\alpha\beta} g^{\alpha k} \delta_\gamma^\beta V_{tk}^t + \frac{1}{n-1} g_{\alpha\beta} g^{i\beta} V_{is}^s \delta_\gamma^\alpha &= 0, \\ -V_{\gamma k}^k - V_{\alpha\gamma}^\alpha + \frac{1}{n-1} V_{t\gamma}^t + \frac{1}{n-1} V_{\gamma s}^s &= 0. \end{aligned}$$

From this we obtain,

$$V_{\gamma} = V_{t\gamma}^t = -V_{\gamma t}^t .$$

Substituting the above formula into (2.14), we have

$$g^{\alpha k} (V_{\gamma k}^{\beta} - \frac{1}{n-1} \delta_{\gamma}^{\beta} V_k) + g^{i\beta} (V_{i\gamma}^{\alpha} + \frac{1}{n-1} \delta_{\gamma}^{\alpha} V_i) = 0 .$$

REMARK. Let $\Omega = \Omega_j dx^j$ be a 1-differential form. It is easy to prove that $V_{ik}^l = \delta_i^l \Omega_k - \delta_k^l \Omega_i$ is a second order differentiable covariant tensor field with vector value. By the following computation, we know it satisfies (2.13).

$$V_{\gamma k}^{\beta} - \frac{1}{n-1} \delta_{\gamma}^{\beta} V_k = (\delta_{\gamma}^{\beta} \Omega_k - \delta_k^{\beta} \Omega_{\gamma}) - \frac{1}{n-1} \delta_{\gamma}^{\beta} (\delta_t^t \Omega_k - \delta_k^t \Omega_t)$$

$$= \delta_{\gamma}^{\beta} \Omega_k - \delta_k^{\beta} \Omega_{\gamma} - \frac{n}{n-1} \delta_{\gamma}^{\beta} \Omega_k + \frac{1}{n-1} \delta_{\gamma}^{\beta} \Omega_k = -\delta_k^{\beta} \Omega_{\gamma} ,$$

$$V_{i\gamma}^{\alpha} + \frac{1}{n-1} \delta_{\gamma}^{\alpha} V_i = (\delta_i^{\alpha} \Omega_{\gamma} - \delta_{\gamma}^{\alpha} \Omega_i) + \frac{1}{n-1} \delta_{\gamma}^{\alpha} (\delta_t^t \Omega_i - \delta_i^t \Omega_t)$$

$$= \delta_i^{\alpha} \Omega_{\gamma} - \delta_{\gamma}^{\alpha} \Omega_i + \frac{n}{n-1} \delta_{\gamma}^{\alpha} \Omega_i - \frac{1}{n-1} \delta_{\gamma}^{\alpha} \Omega_i = \delta_i^{\alpha} \Omega_{\gamma} ,$$

$$g^{\alpha k} (V_{\gamma k}^{\beta} - \frac{1}{n-1} \delta_{\gamma}^{\beta} V_k) + g^{i\beta} (V_{i\gamma}^{\alpha} + \frac{1}{n-1} \delta_{\gamma}^{\alpha} V_i)$$

$$= g^{\alpha k} (-\delta_k^{\beta} \Omega_{\gamma}) + g^{i\beta} (\delta_i^{\alpha} \Omega_{\gamma}) = -g^{\alpha\beta} \Omega_{\gamma} + g^{\alpha\beta} \Omega_{\gamma} = 0 .$$

From the remark of theorem 3, it follows that although T_{Ω} satisfies (2.13), perhaps it does not make scalar curvature S invariant. Therefore, (2.13) is only a necessary condition under which the transformation T_V makes scalar curvature S invariant.

REFERENCES

1. EINSTEIN, A., The Meaning of Relativity. 5th ed. (1955). Appendix II.
2. EINSTEIN, A. & KAUFMAN, B., Louis de Broglie, Physicien et Penseur. Armand Colin. Paris (1953).
or
KAUFMAN, B., Fünfzig Jahre Relativitätstheorie. Helvetia Physica Acta. Supplementum IV (1956), pp. 227-238.
3. EINSTEIN, A. & KAUFMAN, B., Annals of Math. 62 (1955) 1., 128-138.
4. BERGER, M. & GOSTIAUX, B., Géométrie différentielle. Armand Colin. Paris (1972).

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be