

FINITE DIFFERENCE METHODS FOR COMPUTING EIGENVALUES OF FOURTH ORDER BOUNDARY VALUE PROBLEMS

RIAZ A. USMANI

Department of Applied Mathematics
University of Manitoba
Winnipeg, Manitoba, Canada
R3T 2N2

(Received September 7, 1984)

ABSTRACT. This brief report describes some new finite difference methods of order 2 and 4 for computing eigenvalues of a two point boundary value problem associated with a fourth order linear differential equation $y^{(4)} + (p(x) - \lambda q(x))y = 0$. These methods are derived from the formula

$$h^4 y_1^{(4)} = (\delta^4 - \frac{1}{6} \delta^6 + \frac{7}{240} \delta^8 - \dots) y_1 \quad .$$

Numerical results are included to demonstrate practical usefulness of our methods.

KEY WORDS AND PHRASES. Central difference formula, Finite difference methods, Generalized symmetric eigenvalue problem, Positive definite matrix, Two-point boundary value problem.

1980 AMS SUBJECT CLASSIFICATION CODE: 65L15.

1. INTRODUCTION

We shall consider the fourth order linear differential equation

$$y^{(4)} + [p(x) - \lambda q(x)]y = 0 \tag{1.1}$$

associated with one of the following pairs of homogeneous boundary conditions:

$$(a) \quad y(a) = y'(a) = y(b) = y'(b) = 0$$

$$(b) \quad y(a) = y''(a) = y(b) = y''(b) = 0 \tag{1.2}$$

$$(c) \quad y(a) = y'(a) = y''(b) = y'''(b) = 0$$

Such boundary value problems occur frequently in applied mathematics, engineering and modern physics, see [1,2,3]. In (1.1), the functions $p(x)$, $q(x) \in C[a,b]$ and they satisfy the conditions

$$p(x) \geq 0, \quad q(x) > 0, \quad x \in [a,b] \quad . \tag{1.3}$$

Recently, Chawla and Katti [4] have developed a finite difference method of order 2 for computing approximate values of λ for a boundary value problem (1.1)-(1.2a). For the same problem, a fourth order method was developed by Chawla [5] which leads to a generalized seven-band symmetric matrix eigenvalue problem.

The purpose of this note is to present some new finite difference methods for computing approximate values of λ for the boundary value problems (1.1)-(1.2b) and (1.1)-(1.2c). These methods lead to generalized five-band and seven-band symmetric matrix eigenvalue problem and provide $O(h^2)$ and $O(h^4)$ -convergent

approximations for the eigenvalues.

2. A SECOND ORDER METHOD FOR COMPUTING λ FOR (1.1) and (1.2b)

For a positive integer $N \geq 4$, let $h = (b - a)/(N + 1)$ and $x_i = a + ih$, $i = 0(1)N + 1$. We shall designate $y_i = y(x_i)$, $p_i = p(x_i)$ and $q_i = q(x_i)$. The boundary value problem (1.1) and (1.2b) is discretized by the difference equations

$$\begin{aligned} (a) \quad & -2y_0 + 5y_1 - 4y_2 + y_3 = -h^2 y_0'' + h^4 \left[-\frac{1}{12} y^{(4)} + y_1^{(4)} \right] + t_1, \\ (b) \quad & \delta^4 y_i = h^4 y_1^{(4)} + t_i, \quad i = 2(1)N - 1, \\ (c) \quad & y_{N-2} - 4y_{N-1} + 5y_N - 2y_{N+1} = -h^2 y_{N+1}'' + h^4 \left[y_N^{(4)} - \frac{1}{12} y_{N+1}^{(4)} \right] + t_N. \end{aligned} \tag{2.1}$$

Note that the truncation errors t_i , $i = 1(1)N$, are

$$t_i = \begin{cases} \frac{59}{360} h^6 y^{(6)}(\xi_1), & \xi_1 \in (a, x_3) \\ \frac{1}{6} h^6 y^{(6)}(\xi_i), & \xi_i \in (x_{i-2}, x_{i+2}), \quad i = 2(1)N - 1 \\ \frac{59}{360} h^6 y^{(6)}(\xi_N), & \xi_N \in (x_{N-2}, b). \end{cases} \tag{2.2}$$

The formula 2.1(b) is obtained from the well-known central difference formula

$$\begin{aligned} h^4 y_i^{(4)} &= \left[\delta^4 - \frac{1}{6} \delta^6 + \frac{7}{240} \delta^8 - \frac{41}{7560} \delta^{10} + \dots \right] y_i, \\ (i = 2, 3, 4, \dots). \end{aligned} \tag{2.3}$$

The discretizations 2.1(a) and 2.1(c) are introduced so that the resulting coefficient matrix in (2.1) is a five-band symmetric matrix. The system of linear equations (2.1) can be written in matrix form

$$(J^2 + h^4 P)Y = \lambda h^4 QY + t, \tag{2.4}$$

where J^2 is a symmetric five-band matrix and $J = (j_{rs})$ is a tridiagonal matrix such that

$$J_{rs} = \begin{cases} 2, & r = s \\ -1, & |r - s| = 1 \\ 0, & \text{otherwise.} \end{cases} \tag{2.5}$$

The matrices P and Q are diagonal matrices

$$P = \text{diag}[p_1 \ p_2 \ \dots \ p_N], \quad Q = \text{diag}[q_1 \ q_2 \ \dots \ q_N]$$

and

$$Y = [y_1 \ y_2 \ \dots \ y_N]^T, \quad t = [t_1 \ t_2 \ \dots \ t_N]^T.$$

Thus, our method for computing approximations Λ for λ of (1.1)-(1.2b) can be expressed as a generalized five-band symmetric matrix eigenvalue problem

$$(J^2 + h^4 P)\tilde{Y} = \Lambda h^4 Q\tilde{Y}. \tag{2.6}$$

In fact, the matrix J^2 is a positive definite matrix and hence for any step-size $h > 0$, the approximations Λ for λ by (2.6) are real and positive for all $p(x) \geq 0$ and $q(x) > 0$. That our method provides $O(h^2)$ convergent approximations Λ for λ can be established following Grigorieff [6]. We omit the details of

convergence proof for brevity.

3. A FOURTH ORDER METHOD

The boundary value problem (1.1)-(1.2b) is now discretized by the following difference equations

$$\begin{aligned}
 \text{(a)} \quad & -17y_0 + 44y_1 - 38y_2 + 12y_3 - y_4 = -8h^2 y_0'' h^4 \left[\frac{1}{3} y_0^{(4)} + 6y_1^{(4)} \right] + \frac{13}{90} h^6 y_0^{(6)} + \dots \\
 \text{(b)} \quad & 10y_0 - 38y_1 + 56y_2 - 39y_3 + 12y_4 - y_5 = h^2 y_0'' \\
 & \quad \quad \quad + h^4 \left[\frac{1}{12} y_0^{(4)} + 6y_2^{(4)} \right] + \frac{1}{360} h^6 y_0^{(6)} + \dots, \\
 \text{(c)} \quad & (6\delta^4 - \delta^6) y_i = 6h^4 y_i^{(4)} - \frac{7}{40} h^8 y_i^{(8)} + \dots, \quad i = 3(1)N-2; \quad (3.1) \\
 \text{(d)} \quad & -y_{N-4} + 12y_{N-3} - 39y_{N-2} + 56y_{N-1} - 38y_N + 10y_{N+1} \\
 & \quad \quad \quad = h^2 y_{N+1}'' + h^4 \left[6y_{N-1}^{(4)} + \frac{1}{12} y_{N+1}^{(4)} \right] + \frac{1}{360} h^6 y_N^{(6)} + \dots, \\
 \text{(e)} \quad & -y_{N-3} + 12y_{N-2} - 38y_{N-1} + 44y_N - 17y_{N+1} = -8h^2 y_{N+1}'' \\
 & \quad \quad \quad + h^4 \left[6y_N^{(4)} + \frac{1}{3} y_{N+1}^{(4)} \right] + \frac{13}{90} h^6 y_N^{(6)} + \dots
 \end{aligned}$$

As in [7], the derivation of (3.1c) is immediate from (2.3), on truncating the infinite series on the right of equality sign after the two terms. The additional difference equations (3.1a, b, d, e) are chosen so that the resulting matrix associated with the system of linear equations in (3.1) is a seven-band symmetric matrix. It turns out that our method for computing approximations Λ for λ of (1.1)-(1.2b) can be expressed as a generalised seven-band symmetric matrix eigenvalue problem

$$[(6J^2 + J^3) + 6h^4 P] \tilde{Y} = 6\Lambda h^4 Q \tilde{Y}. \quad (3.2)$$

The matrix $6J^2 + J^3$ is a positive definite matrix and hence for any step-size $h > 0$, the approximations Λ for λ by (3.2) are real and positive for all $p(x) \geq 0$ and $q(x) > 0$. As before, it follows from the results of Grigorieff (1975) that our present method provides $O(h^4)$ -convergent approximations Λ for λ .

4. METHODS FOR COMPUTING λ FOR (1.1)-(1.2c)

For this section, let $h = (b-a)/N$ and $x_i = a + ih$, $i = 0(1)N$. The boundary value problem (1.1)-(1.2c) is discretized by the following scheme

$$\begin{aligned}
 \text{(a)} \quad & -4y_0 + 7y_1 - 4y_2 + y_3 = 2hy_0' + h^4 y_1^{(4)} + \left(\frac{h^3}{3} y_0'' + \dots \right), \\
 \text{(b)} \quad & \delta^4 y_i = h^4 y_i^{(4)} + \frac{1}{6} h^4 y_i^{(6)}(\xi_i), \quad \xi_i \in (x_{i-2}, x_{i+2}), \\
 & \quad \quad \quad i = 2(1)N-2, \quad (4.1) \\
 \text{(c)} \quad & y_{N-3} - 4y_{N-2} + 5y_{N-1} - 2y_N = -h^2 y_N'' + \frac{11}{12} h^4 y_{N-1}^{(4)} \\
 & \quad \quad \quad + \left(\frac{1}{12} h^5 y_N^{(5)} + \dots \right), \\
 \text{(d)} \quad & y_{N-2} - 2y_{N-1} + y_N = h^2 y_N'' - h^3 y_N''' + \frac{7}{12} h^4 y_N^{(4)} + \left(\frac{1}{4} h^5 y_N^{(5)} + \dots \right).
 \end{aligned}$$

This system can be written in matrix form

5. CHAWLA, M.M. A new fourth-order finite difference method for computing eigenvalues of fourth-order two-point boundary value problem, IMA Journal of Numerical Analysis, 3 (1983), 291-293.
6. GRIGORIEFF, R.D. Diskrete Approximation von Eigenwertproblemen II, Number. Math., 24 (1975), 415-433.
7. USMANI, R.A. On a class of numerical integration methods for a problem of plate deflection theory, Internat. J. Comput. Math., 11 (1982), 305-318.
8. USMANI, R.A. Finite difference methods for a certain two point boundary value problem, Indian J. Pure Appl. Math., 14 (1983), 398-411.

Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Lead Guest Editor

Juan J. Nieto, Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;
juanjose.nieto.roig@usc.es

Guest Editor

Donal O'Regan, Department of Mathematics, National University of Ireland, Galway, Ireland;
donal.oregan@nuigalway.ie