

NOTES ON ALMOST-PERIODICITY IN TOPOLOGICAL VECTOR SPACES

GASTON MANDATA N'GUÉRÉKATA

Faculte des Sciences
Universite de Bangui
BF 1450 BANGUI
Republique Centrafricaine

(Received December 27, 1983)

ABSTRACT. A study is made of almost-periodic functions in topological vector spaces with applications to abstract differential equations.

KEY WORDS AND PHRASES. *Topological vector spaces, abstract differential equations.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 34G10

1. INTRODUCTION.

In our recent papers [1, 2], we extended the theory of almost-periodic functions from Banach spaces to topological vector spaces and gave a few results concerning its applications to abstract differential equations. The following results are the continuation of discussions begun there. Specifically Theorem 2 is a version of a theorem contained in [1, 2] (see Theorem 5.1 in [2]) which was originally inspired from a result due to A. I. Perov (cf. [3] Theorem 1.1).

Let us first recall some useful facts (see [1, 2] for more details). The reader can also find in [4] the elementary properties of linear topological spaces needed here.

DEFINITION 1. A continuous function $f: \mathbb{R} \rightarrow E$, where E is a complete locally convex space and \mathbb{R} is the set of real numbers, is called almost periodic (a.p.) if for each neighborhood (of the origin in E) U , there exists a real number $\lambda = \lambda(U) > 0$ such that every interval $[a, a + \lambda]$ contains at least a point τ such that

$$f(t + \tau) - f(t) \in U \quad \text{for every } t \in \mathbb{R}.$$

τ is then called a U -translation number of the function f .

REMARK: $U = U(\epsilon; p_i, 1 \leq i \leq n)$

$$= \{x \in E; p_i(x) < \epsilon, 1 \leq i \leq n\}$$

where each $p_i \in Q$, the set of semi-norms on E .

Finally we recall Bochner's criteria: If E is a Frechet space, then a function $f: \mathbb{R} \rightarrow E$ is a.p. iff for every real sequence $(s'_n)_{n=1}^{\infty}$ there exists a subsequence $(s_n)_{n=1}^{\infty}$ such that $(f(t + s_n))_{n=1}^{\infty}$ converges uniformly in $t \in \mathbb{R}$.

DEFINITION 2. A Frechet space E is called a perfect Frechet space if the following property is verified in E : every function $\phi: \mathbb{R} \rightarrow E$ such that

- (i) $\{\phi(t); t \in R\}$ is bounded in E
 (ii) the derivative $\phi'(t)$ is a.p. in E , is necessarily a.p. in E .

2. MAIN RESULTS.

Now let us state and prove:

THEOREM 1. If $f(t)$ is a.p. in a complete locally convex space E , then for every real sequence $(s_n)_{n=1}^{\infty}$ there exists a subsequence $(s'_n)_{n=1}^{\infty}$ such that for every neighborhood (of the origin in E) U ,

$$f(t + s'_n) - f(t + s'_m) \in U$$

for all $t \in R$, m and n .

PROOF. Let $U = U(\varepsilon; p_i, 1 \leq i \leq n)$ be a neighborhood and $V = V(\frac{\varepsilon}{4}; p_i, 1 \leq i \leq n)$ a symmetric neighborhood such that $V + V + V + V \subset U$. By the definition of almost-periodicity, there exists $\ell = \ell(V)$ (therefore ℓ depends on U) such that in every real interval of length ℓ , there exists τ such that

$$f(t + \tau) - f(t) \in V$$

for every $t \in R$.

Now for each s_n , we can find τ_n and σ_n such that $s_n = \tau_n + \sigma_n$ with τ_n a V -translation number of f and $\sigma_n \in [0, \ell]$ (it suffices to take $\tau_n \in [s_n - \ell, s_n]$ and then $\sigma_n = s_n - \tau_n$).

As f is uniformly continuous on R (cf. [1, 2]), there exists $\delta = \delta(\varepsilon)$ such that

$$f(t') - f(t'') \in V \quad (2.1)$$

for all $t', t'', |t' - t''| < 2\delta$.

Also $0 \leq \sigma_n \leq \ell$ for every n ; we can then subtract from $(\sigma_n)_{n=1}^{\infty}$, a convergent subsequence $(\sigma_{n_k})_{k=1}^{\infty}$, by the Bolzano-Weierstrass theorem.

Let $\sigma = \lim_{k \rightarrow \infty} \sigma_{n_k}$, with $0 \leq \sigma \leq \ell$.

Now consider the subsequence $(\sigma_{n_k})_{k=1}^{\infty}$ with

$$\sigma - \delta < \sigma_{n_k} < \sigma + \delta, k = 1, 2, \dots$$

and let $(s_{n_k})_{k=1}^{\infty}$ be the corresponding subsequence where

$$s_{n_k} = \tau_{n_k} + \sigma_{n_k}, k = 1, 2, \dots$$

Let us prove the relation

$$f(t + s_{n_k}) - f(t + s_{n_j}) \in U \quad (2.2)$$

for all $t \in R$.

For this, write

$$\begin{aligned} f(t + s_{n_k}) - f(t + s_{n_j}) &= f(t + \tau_{n_k} + \sigma_{n_k}) - f(t + \sigma_{n_k}) \\ &\quad + f(t + \sigma_{n_k}) - f(t + \sigma_{n_j}) \\ &\quad + f(t + \sigma_{n_j}) - f(t + \tau_{n_j} + \sigma_{n_j}). \end{aligned}$$

Because τ_{n_k} and τ_{n_j} are V -translation numbers of f , we shall get

$$\begin{aligned} f(t + \tau_{n_k} + \sigma_{n_k}) - f(t + \sigma_{n_k}) &\in V, \text{ for every } t \in \mathbb{R} \\ f(t + \tau_{n_j} + \sigma_{n_j}) - f(t + \sigma_{n_j}) &\in V, \text{ for every } t \in \mathbb{R}. \end{aligned} \quad (2.3)$$

On the other hand

$$|(t + \sigma_{n_k}) - (t + \sigma_{n_j})| = |\sigma_{n_k} - \sigma_{n_j}| < 2\delta;$$

therefore, by using relation (2.1), we get

$$f(t + \sigma_{n_k}) - f(t + \sigma_{n_j}) \in V, \text{ for every } t \in \mathbb{R}. \quad (2.4)$$

Finally we can deduce (2.2) from (2.3) and (2.4). The theorem is proved by taking

$$s'_n = s_{n_k}, \quad k = 1, 2, \dots \quad \square$$

3. APPLICATIONS

Let E be a perfect Fréchet space and A a closed linear operator with domain $D(A)$ dense in E . Suppose A generates a strongly continuous one-parameter group $T(t)$, $t \in \mathbb{R}$.

Consider in such E the differential equation

$$x'(t) = Ax(t), \quad t \in \mathbb{R}. \quad (3.1)$$

THEOREM 2. Assume for every semi-norm $p \in Q$, there exists a semi-norm $q \in Q$ such that

$$p(T(t)u) \leq q(u)$$

for every $u \in E$ and $t \in \mathbb{R}$.

Then every solution $x(t)$ of (3.1) such that $\{x'(t); t \in \mathbb{R}\}$ is relatively compact in E is a.p.

PROOF. Let $x(t)$ be such a solution; we can write $x(t) = T(t)x(0)$, $t \in \mathbb{R}$; by the property on $T(t)$, $x(t)$ is obviously bounded.

Consider a given real sequence $(s'_n)_{n=1}^\infty$; we can extract a subsequence $(s_n)_{n=1}^\infty$ such that $(x'(s_n))_{n=1}^\infty$ is a Cauchy sequence in E , for $\{x'(t); t \in \mathbb{R}\}$ is relatively compact in E . We have

$$\begin{aligned} x'(t + s_n) &= Ax(t + s_n) \\ &= AT(t + s_n)x(0) \\ &= AT(t)T(s_n)x(0) \\ &= AT(t)x(s_n) \\ &= T(t)Ax(s_n) \\ &= T(t)x'(s_n) \end{aligned}$$

for every n and every $t \in \mathbb{R}$. Therefore

$$x'(t + s_n) - x'(t + s_m) = T(t)[x'(s_n) - x'(s_m)]$$

for every n, m and $t \in \mathbb{R}$.

Take now any $p \in Q$; then there exists $q \in Q$ such that

$$p[x'(t + s_n) - x'(t + s_m)] \leq q[x'(s_n) - x'(s_m)]$$

for every $t \in \mathbb{R}$; which shows $x'(t)$ is a.p. by Bochner's criteria. As E is a perfect Fréchet space, the conclusion is immediate.

ACKNOWLEDGEMENT

The author was supported by a Fulbright grant at the University of California at Berkeley from June to October 1983.

REFERENCES

1. N'GUÉKATA, G. M. Remarques sur les Équations Différentielles Abstraites, Thèse de Ph.D., Université de Montréal, Juin 1980.
2. N'GUÉKATA, G. M. Almost-Periodicity in Linear Topological Spaces and Applications to Abstract Differential Equations, Internat. J. Math. Math. Sci. 7 (1984) 529-540.
3. ZAIDMAN, S. Solutions Presque-Periodiques des Équations Différentielles Abstraites, Enseign. Math. 24 (1978), 87-110.
4. ROBERTSON, A. P. and ROBERTSON, W. Topological Vector Spaces, Cambridge University Press, 1973.
5. AMERIO, L. and PROUSE, G. Almost-Periodic Functions and Functional Equations, Van Nostrand, 1971.
6. CORDUNEANU, C. Almost-Periodic Functions, Interscience, 1968.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from “Qualitative Theory of Differential Equations,” allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk