

ON COMMON FIXED POINTS OF WEAKLY COMMUTING MAPPINGS AND SET-VALUED MAPPINGS

S. SESSA

Istituto di Matematica
Facolta di Architettura
Universita di Napoli
Via Monteoliveto 3
80134 Napoli, Italy

and

B. FISHER

Department of Mathematics
The University
Leicester, ENGLAND

(Received September 7, 1984)

ABSTRACT. Our main theorem establishes the uniqueness of the common fixed point of two set-valued mappings and of two single-valued mappings defined on a complete metric space, under a contractive condition and a weak commutativity concept. This improves a theorem of the second author.

KEY WORDS AND PHRASES. Common fixed point, set-valued mapping, weak commutativity.

1980 AMS SUBJECT CLASSIFICATION CODES. 54H25, 47H10.

1. BASIC PRELIMINARIES.

Let (X, d) be a complete metric space and let $B(X)$ be the set of all nonempty, bounded subsets of X . As in [1], let $\delta(A, B)$ be the function defined by

$$\delta(A, B) = \sup \{d(a, b) : a \in A, b \in B\}$$

for all A, B in $B(X)$.

If A consists of a single point a we write

$$\delta(A, B) = \delta(a, B)$$

and if B also consists of a single point b we write

$$\delta(A, B) = d(a, b).$$

It follows immediately from the definition that

$$\delta(A, B) = \delta(B, A) \geq 0, \quad \delta(A, A) = \text{diam } A,$$

$$\delta(A, B) \leq \delta(A, C) + \delta(C, B)$$

for all A, B, C in $B(X)$.

We say that a subset A of X is the limit of a sequence $\{A_n\}$ of nonempty subsets of X if each point a in A is the limit of a convergent sequence $\{a_n\}$, where a_n is in A_n for $n = 1, 2, \dots$, and if for arbitrary $\epsilon > 0$, there exists an integer N such that $A_n \subseteq A_\epsilon$ for $n > N$, where A_ϵ is the union of all open spheres with

centres in A and radius ϵ .

LEMMA 1. If $\{A_n\}$ and $\{B_n\}$ are sequences of bounded subsets of (X, d) which converge to the bounded sets A and B respectively, then the sequence $\{\delta(A_n, B_n)\}$ converges to $\delta(A, B)$.

This lemma was proved in [2].

Now let F be a mapping of X into $B(X)$. We say that F is continuous at the point x in X if whenever $\{x_n\}$ is a sequence of points in X converging to x , the sequence $\{F x_n\}$ in $B(X)$ converges to $F x$ in $B(X)$. If F is continuous at each point x in X , we say that F is a continuous mapping of X into $B(X)$. A point z in X is said to be a fixed point of F if z is in $F z$.

For a selfmap I of (X, d) , the authors of [3], extending the results of [2] and [4], defined F and I to be weakly commuting on X if

$$\delta(FIx, IFx) \leq \max\{\delta(Ix, Fx), \text{diam } IFx\} \quad (1.1)$$

for all x in X . Two commuting mappings F and I clearly commute, but two weakly commuting mappings F and I do not necessarily commute as is shown in the following example.

EXAMPLE 1. Let $X = [0, 1]$, let δ be the function induced by the euclidean metric d and define

$$F x = [0, x/(x+a^h)], \quad I x = x/a$$

for all x in X , where $h \geq 1$ and $a \geq 2$. Then for any non-zero x in X we have

$$FIx = [0, x/(x+a^{h+1})] \neq [0, x/(ax+a^{h+1})] = IFx$$

but for any x in X we have

$$\delta(FIx, IFx) = x/(x+a^{h+1}) \leq x/a = \delta(Ix, Fx).$$

Note that if F is a single-valued mapping, then the set $\{IFx\}$ consists of a single point and therefore $\text{diam } \{IFx\} = 0$ for all x in X . Condition (1.1) therefore reduces to the condition given in [5], i.e.

$$d(FIx, IFx) \leq d(Ix, Fx) \quad (1.2)$$

for all x in X .

An extensive literature exists about (common) fixed points of set-valued mappings satisfying contractive conditions controlled from non-negative real functions f from $[0, \infty)$ into $[0, \infty)$. Suitable properties of f guarantee the convergence to the (common) fixed point of the sequence of successive approximations: see for example the papers of Barcz [6], Chen and Shih [7], Guay, Singh, and Whitfield [8], Miczko and Palezewski [9], Nhan [10], Papageorgiou [11], Popa [12], Sharma [13] and Wegrzyk [14]. In this paper we consider the family F of functions f from $[0, \infty)$ into $[0, \infty)$ such that

- (a) f is non-decreasing,
- (aa) f is continuous from the right,
- (aaa) $f(t) < t$ for all $t > 0$.

LEMMA 2. For any $t > 0$, $\lim_{n \rightarrow \infty} f^n(t) = 0$.

The proof of this lemma is obvious but see also [15].

Further details about the usage of functions with properties similar to (α) , $(\alpha\alpha)$, and $(\alpha\alpha\alpha)$ can be found in the papers of Benedykt and Matkowski [16], Browder [17], Conserva and Fedele [18], Hegedüs and Szilágyi [19], Hikida [20], Park and Rhoades [21], Rhoades [22], and Singh and Kasahara [23].

2. RESULTS IN COMPLETE METRIC SPACES.

Let F, G be two set-valued mappings of X into $B(X)$ and let I, J be two selfmaps of X such that

$$F(X) \subseteq I(X), \quad G(X) \subseteq J(X) . \tag{2.1}$$

Let x_0 (resp. y_0) be an arbitrary point in X and define inductively a sequence $\{x_n\}$ (resp. $\{y_n\}$) such that, having defined the point x_{n-1} (resp. y_{n-1}), choose a point x_n (resp. y_n) with Ix_n (resp. Jy_n) in Fx_{n-1} (resp. Gy_{n-1}) for $n = 1, 2, \dots$.

This can be done since the range of I (resp. J) contains the range of F (resp. G).

Further, assume that

$$\sup\{\delta(Fx_n, Gy_0), \delta(Gy_n, Fx_0) : n = 1, 2, \dots\} < \infty . \tag{2.2}$$

REMARK 1. IF X is bounded then (2.2) will always be satisfied for all x, y in X .

We consider the following conditions:

(γ_1) I continuous,

(γ_2) F continuous and $IFx \subseteq FIx$ for all x in X .

(λ_1) J continuous,

(λ_2) G continuous and $JGx \subseteq GJx$ for all x in X .

Modifying the proof of theorem 1 of [1] we are now able to prove the following:

THEOREM 1. Let F, G be two set-valued mappings of X into $B(X)$ and let I, J be two selfmaps of X satisfying (2.1) and

$$\delta(Fx, Gy) \leq f(\max\{d(Ix, Jy), \delta(Ix, Gy), \delta(Jy, Fx)\}) \tag{2.3}$$

for all x, y in X , where f is in F . Further let F and G weakly commute with I and J respectively. If there exist points x_0 and y_0 in X satisfying (2.2) and if the conditions (γ_i) and (λ_j) with $i, j = 1, 2$, hold, then F, G, I and J have a unique common fixed point z . Further, $Fz = Gz = \{z\}$ and z is the unique common fixed point of F and I and of G and J .

PROOF. Since

$$\delta(Fx_r, Gy_s) \leq \delta(Fx_r, Gy_0) + \delta(Gy_0, Fx_0) + \delta(Fx_0, Gy_s),$$

it follows from (2.2) that

$$M = \sup \{\delta(Fx_r, Gy_s) : r, s = 0, 1, 2, \dots\}$$

is finite.

If $M > 0$, then for arbitrary $\epsilon > 0$, we can choose an integer p such that $f^p(M) < \epsilon$ by lemma 2. If $M = 0$, then $f^p(M) = 0 < \epsilon$ for any integer p .

As in the proof of theorem 1 of [24], we have on using inequality (2.3) p times and property (α) :

$$\delta(Fx_m, Gy_n) \leq f^p(\max\{\delta(Fx_r, Gy_q) : m - p \leq r \leq m ; n - p \leq q \leq n\})$$

$< \epsilon$

for $m, n > p$. Thus

$$\delta(Fx_m, Fx_n) \leq \delta(Fx_m, Gy_s) + \delta(Gy_s, Fx_r) < 2\epsilon$$

for $m, n > p$. The sequence $\{z_n\}$ is therefore a Cauchy sequence in the complete metric space X and so has a limit z in X , where z is independent of the particular choice of each z_n . It follows in particular that the sequence $\{Ix_n\}$ converges to z and the sequence of sets $\{Fx_n\}$ converges to the set $\{z\}$.

Similarly, it can be proved that the sequence $\{Jy_n\}$ converges to a point w and the sequence of sets $\{Gy_n\}$ converges to the set $\{w\}$.

Using (2.3) we have

$$\delta(Fx_n, Gy_n) \leq f(\max\{d(Ix_n, Jy_n), \delta(Ix_n, Gy_n), \delta(Jy_n, Fx_n)\})$$

Letting n tend to infinity and using lemma 1 and properties $(\alpha\alpha)$ and $(\alpha\alpha\alpha)$, it is seen that $w = z$.

Now suppose that (γ_1) holds. Then the sequence $\{I^2x_n\}$ and $\{IFx_n\}$ converge to Iz and $\{Iz\}$ respectively. Let w_n be an arbitrary point in FIx_n for $n = 1, 2, \dots$. Then since I weakly commutes with F we have on using (1.1)

$$\begin{aligned} d(w_n, Iz) &\leq \delta(FIx_n, Iz) \\ &\leq \delta(FIx_n, IFx_n) + \delta(IFx_n, Iz) \\ &\leq \max\{\delta(Ix_n, Fx_n), 2\delta(I^2x_{n+1}, IFx_n)\} + \delta(IFx_n, Iz). \end{aligned}$$

Letting n tend to infinity and using lemma 1 we see that the sequence $\{w_n\}$ converges to Iz . But Iz is independent of the particular choice of w_n in FIx_n and this means that the sequence of sets $\{FIx_n\}$ converges to the set $\{Iz\}$.

Using inequality (2.3) we have

$$\delta(FIx_n, Gy_n) \leq f(\max\{d(I^2x_n, Jy_n), \delta(I^2x_n, Gy_n), \delta(Jy_n, FIx_n)\}).$$

Letting n tend to infinity and using lemma 1 and property $(\alpha\alpha)$, we have

$$d(Iz, z) \leq f(d(Iz, z))$$

which implies $Iz = z$ by $(\alpha\alpha\alpha)$.

Since

$$\delta(Fz, Gy_n) \leq f(\max\{d(Iz, Jy_n), \delta(Iz, Gy_n), \delta(Jy_n, Fz)\})$$

we have on letting n tend to infinity and using lemma 1 and property $(\alpha\alpha)$

$$\delta(Fz, z) \leq f(\delta(z, Fz))$$

which gives $Fz = \{z\}$ by $(\alpha\alpha\alpha)$.

Similarly, the weak commutativity of G and J and condition (λ_1) implies $Jz = z$ and $Gz = \{z\}$.

Now assume that (γ_2) holds. Then the sequence $\{FIx_n\}$ converges to Fz and using inequality (2.3) we have

$$\begin{aligned} \delta(FIx_n, Gy_n) &\leq f(\max\{d(I^2x_n, Jy_n), \delta(I^2x_n, Gy_n), \delta(Jy_n, FIx_n)\}) \\ &\leq f(\max\{\delta(FIx_n, Jy_n), \delta(FIx_n, Gy_n), \delta(Jy_n, FIx_n)\}) \end{aligned}$$

since f is non-decreasing and Ix_n is in Fx_{n-1} and so I^2x_n is in $IFx_{n-1} \subseteq FIx_{n-1}$.

Letting n tend to infinity and using lemma 1 and property $(\alpha\alpha)$, we have

$$\delta(Fz, z) \leq f(\delta(Fz, z))$$

which implies $Fz = \{z\}$ by $(\alpha\alpha\alpha)$. Thus by (2.1) there must exist a point u in X such that $Iu = z$.

Using inequality (2.3) we have

$$\delta(Fu, Gy_n) \leq f(\max\{d(Iu, Jy_n), \delta(Iu, Gy_n), \delta(Jy_n, Fu)\}).$$

Letting n tend to infinity and using lemma 1 and property $(\alpha\alpha)$, we obtain the inequality

$$\delta(Fu, z) \leq f(\max\{d(Iu, z), \delta(z, Fu)\}) = f(\delta(z, Fu)).$$

Thus $Fu = \{z\}$ by $(\alpha\alpha\alpha)$ and since F and I weakly commute, we have

$$\{z\} = Fz = FIu = IFu = \{Iz\}.$$

It follows that $Iz = z$.

Similarly property (λ_2) assures that $Gz = \{z\}$ and $Jz = z$.

We have therefore shown that if the conditions (γ_1) and (λ_j) , with $i, j = 1, 2$, hold then $Iz = Jz = z$ and $Fz = Gz = \{z\}$.

That z is the unique common fixed point of F and I and of G and J follows easily. This completes the proof of the theorem.

COROLLARY 1. Let F, G be two set-valued mappings of X into $B(X)$ and let I, J be two selfmaps of X satisfying (2.1) and

$$\delta(Fx, Gy) \leq c \cdot \max\{d(Ix, Jy), \delta(Ix, Gy), \delta(Jy, Fx)\} \tag{2.4}$$

for all x, y in X , where $0 \leq c < 1$. Further, let F and G commute with I and J respectively. If F or I and G or J are continuous, then F, G, I and J have a unique common fixed point z . Further, $Fz = Gz = \{z\}$ and z is the unique common fixed point of F and I and of G and J .

PROOF. As in the proof of theorem 1 of [1], it is proved that (2.2) holds for any x_0, y_0 in X . Since F and G commute with I and J respectively, we have $FIx = IFx$ and $GJx = JGx$ for all x in X . The thesis then follows from theorem 1 if we assume that $f(t) = ct$ for all $t \geq 0$.

The result of this corollary was given in [1].

We now give an example in which theorem 1 holds but corollary 1 is not applicable.

EXAMPLE 2. Let $X = [0, 1]$ with δ induced by the euclidean metric d and let $Fx = [0, x/(x + 4)]$, $Gx = [0, x/(x + 8)]$, $Ix = Jx = \frac{1}{2}x$ for all x in X .

By example 1, F and G weakly commute with I . Further, we have

$$\begin{aligned} F(X) &= [0, 1/5] \subset [0, \frac{1}{2}] = I(X), \\ G(X) &= [0, 1/9] \subset [0, \frac{1}{2}] = J(X), \\ IFx &= [0, x/(2x + 8)] \subset [0, x/(x + 8)] = FIx \\ JGx &= [0, x/(2x + 16)] \subset [0, x/(x + 16)] = GJx \end{aligned}$$

for all x in X .

Since

$$\begin{aligned} \delta(Fx, Gy) &= \max\{x/(x+4), y/(y+8)\} \\ &\leq \max\{x/(x+4), y/(y+4)\} \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{2} \max\{\frac{1}{2}x, \frac{1}{2}y\} \\ &= \begin{cases} \frac{1}{2} \delta(Ix, Gy), & \text{if } x \geq y, \\ \frac{1}{2} \delta(Jy, Fx), & \text{if } x < y \end{cases} \end{aligned}$$

and since X is bounded all the hypotheses of theorem 1 are satisfied if we assume $f(t) = \frac{1}{2}t$ for all $t \geq 0$. Clearly f is in F and 0 is the unique common fixed point of F, G and I .

Theorem 1 is a stronger result than corollary 1, even if the mappings under consideration are commutative, as is shown in the following example.

EXAMPLE 3. Let X be the reals with δ induced by the euclidean metric d , let

$$F_x = \begin{cases} \{0\}, & \text{if } x \leq 0, \\ [0, x/(1+3x)], & \text{if } 0 < x \leq 1, \\ [0, 1/4], & \text{if } x > 1 \end{cases}$$

$$G_x = \begin{cases} \{0\}, & \text{if } x \leq 0, \\ [0, x/(1+2x)], & \text{if } 0 < x \leq 1, \\ \{1/3\}, & \text{if } x > 1, \end{cases}$$

$$I_x = \begin{cases} 0, & \text{if } x \leq 0, \\ x, & \text{if } 0 < x \leq 1, \\ 1, & \text{if } x > 1, \end{cases} \quad J_x = \begin{cases} 0, & \text{if } x \leq 0, \\ x, & \text{if } x > 0, \end{cases}$$

for all x in X and let f in F be given by

$$f(t) = \begin{cases} t/(1+2t), & \text{if } 0 \leq t \leq 1, \\ t/3, & \text{if } t > 1. \end{cases}$$

We have

$$\begin{aligned} \delta(F_x, G_y) &= 0 = f(d(I_x, J_y)), \text{ if } x, y \leq 0, \\ \delta(F_x, G_y) &= y/(1+2y) = f(y) = f(d(I_x, J_y)), \text{ if } x \leq 0 \text{ and } 0 < y \leq 1, \\ \delta(F_x, G_y) &= 1/3 < y/3 = f(y) = f(d(I_x, J_y)), \text{ if } x \leq 0 \text{ and } y > 1, \\ \delta(F_x, G_y) &= x/(1+3x) < x/(1+2x) = f(x) = f(d(I_x, J_y)), \text{ if } 0 < x \leq 1 \text{ and } y \leq 0, \\ \delta(F_x, G_y) &= \max\{x/(1+3x), y/(1+2y)\} \\ &< \max\{x/(1+2x), y/(1+2y)\} \\ &= \begin{cases} f(y) = f(\delta(F_x, J_y)), & \text{if } x \leq y, \\ f(x) = f(\delta(I_x, G_y)), & \text{if } x > y, \text{ and if } 0 < x, y \leq 1, \end{cases} \\ \delta(F_x, G_y) &= 1/3 < y/3 = f(y) = f(\delta(J_y, F_x)), \text{ if } 0 < x \leq 1 \text{ and } y > 1, \\ \delta(F_x, G_y) &= 1/4 < 1/3 = f(1) = f(d(I_x, J_y)), \text{ if } x > 1 \text{ and } y \leq 0, \\ \delta(F_x, G_y) &= \max\{1/4, y/(1+2y)\} \leq 1/3 = f(1) = f(\delta(I_x, G_y)), \text{ if } x > 1 \text{ and } \\ &0 < y \leq 1, \\ \delta(F_x, G_y) &= 1/3 < y/3 = f(y) = f(\delta(J_y, F_x)), \text{ if } x, y > 1. \end{aligned}$$

Condition (2.3) therefore holds in every case since f is non-decreasing. Further

$$F(X) = [0, 1.4] \subset [0, 1] = I(X),$$

$$G(X) = [0, 1/3] \subset [0, \infty] = J(X)$$

and F and G commute with I and J respectively. Since $F_x \subseteq [0, 1/4]$ and $G_x \subseteq [0, 1/3]$ for all x in X , it is easily seen that $M \leq 1/3$ and so (2.2) holds for any x_0 and y_0 chosen in X . As I and J are continuous, theorem 1 is applicable. However, the conditions of the corollary are not satisfied. Otherwise for $x=0$ and $0 < y \leq 1$, condition (2.4) should imply

$$\delta(Fx, Gy) = \frac{y}{1+2y} \leq c \cdot \max\{y, \frac{y}{1+2y}, y\} = cy$$

and so $1/(1+2y) \leq c$ which as y tends to zero, gives $c \geq 1$, a contradiction.

REFERENCES

1. FISHER, B., "Common fixed point theorems for mappings and set-valued mappings", J. Univ. Kuwait Sci., to appear.
2. FISHER, B., "Common fixed points of mappings and set-valued mappings", Rostock. Math. Kolloq., 18(1981), 69-77.
3. SESSA, S., KHAN, M.S., and IMDAD, M., "Common fixed point theorems with a weak commutativity condition", submitted for publication.
4. FISHER, B., "Fixed points of mappings and set-valued mappings", J. Univ. Kuwait Sci., 9(1982), 175-180.
5. SESSA, S., "On a weak commutativity condition in fixed point considerations", Publ. Inst. Math., 32(46)(1982), 149-153.
6. BARCZ, E. "Some fixed point theorems for multi-valued mappings", Demonstratio Math., 16(1983), 735-744.
7. CHEN, M. P. and SHIN, M. H., "Fixed point theorems for point to point and point to set maps", J. Math. Anal. Appl., 71(1979), 515-524.
8. GUAY, M.D., SINGH, K. L., and WHITFIELD, J.H.M., "Common fixed points for set-valued mappings", Bull. Acad. Polon. Sci. Sér. Sci. Math., 30(1982), 545-551.
9. MICZKO, A. and PALCZEWSKI, B., "On convergence of successive approximations of some generalized contraction mappings", Ann. Polon. Math., 4(1983), 213-232.
10. DUANG TRONG NHAN, "Pair of nonlinear contraction mappings common fixed points", Studia Univ. Babes-Bolyai Math., 26(1981), 34-51.
11. PAPAGEORGIOU, N. S., "Fixed point theorems for multifunctions in metric and vector spaces", Nonlinear Anal. TMA 7(1983), 763-770.
12. POPA, V., "Fixed point theorems for multifunctions", Studia Univ. Babes-Bolyai Math., 27(1982), 21-27.
13. SHARMA, A. K., "Common fixed points of set-valued maps", Bull. Acad. Polon. Sci. Sér. Sci. Math., 27(1979), 407-412.
14. WEGRZYK, R., "Fixed-point theorems for multi-valued functions and their applications to functional equations", Dissertations Math., 201, Warszawa (1982), 1-28.
15. SINGH, S.P. and MEADE, B.A., "On common fixed point theorems", Bull. Austral. Math. Sco., 16(1977), 49-53.
16. BENEDYKT, Z. and MATKOWSKI, J., "Remarks on fixed point theorems", Demonstratio Math., 14(1981), 227-232.
17. BROWDER, F. E., "Remarks on fixed point theorems of contractive type", Nonlinear Anal. TMA, 3(1979), 657-661.
18. CONSERVA, V. and FEDELE, F., "Remark on a Browder's fixed point theorem", Math. Japon., 28(1983), 147-157.
19. HEGEDÜS, M. and SZILÁGYI, T., "Equivalent conditions and a new fixed point theorem in the theory of contractive type mappings", Math. Japon., 25(1980), 147-157.
20. HIKIDA, M., "A note on contractions in ranked space", Math. Japon., 26(1981), 131-137.
21. PARK, S. and RHOADES, B. E., "Extensions of some fixed point theorems of Hegedüs and Kasahara", Math. Sem. Notes, Kobe Univ., 9(1981), 113-118.
22. RHOADES, B. E., "Contractive definitions revisited", Topological Methods in Nonlinear Functional Analysis, Contemporary Math. AMS, 21(1983), 189-205.
23. SINGH, S. L. and KASAHARA, S., "On some recent results on common fixed points", Indian J. Pure Appl. Math., 13(7)(1982), 757-761.
24. FISHER, B., "Four mappings with a common fixed point", J. Univ. Kuwait Sci., 8(1981), 131-139.

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

| | |
|------------------------|-----------------|
| Manuscript Due | April 1, 2009 |
| First Round of Reviews | July 1, 2009 |
| Publication Date | October 1, 2009 |

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.otero@usc.es