

A DIOPHANTINE SYSTEM

ANDREW BREMNER

Department of Mathematics
Arizona State University
Tempe, Arizona 85287

(Received November 20, 1985)

ABSTRACT: It is shown how to find all integers a, b such that $a+b$, a^2+b^2 and a^3+b^3 are simultaneously perfect squares.

KEYWORDS AND PHRASES: Diophantine equation, elliptic curve.

1980 AMS SUBJECT CLASSIFICATION CODE. 11B10.

1. INTRODUCTION.

In the Gentleman's Diary of 1795, J. Saul found a solution in integers a, b of the Diophantine system

$$\begin{aligned} a + b &= \text{square} \\ a^2 + b^2 &= \text{square} \\ a^3 + b^3 &= \text{square} . \end{aligned} \tag{1.1}$$


It is the intention of this note to show that recurrence formulae may be developed giving all integral solutions of the system. This amounts to displaying the additive group structure on an elliptic curve, where the curve is given not in standard Weierstrass form, but as the intersection of two quadrics. Such explicit group structure occurs not too frequently in print. For an example, see Cassels [1], or Selmer [2].

2. RESULTS.

First observe that any rational solution of (1.1) gives rise to a rational point $(x, y, u, v) = (a, b, u, v)$ on the elliptic curve

$$E: \begin{cases} x^2 + y^2 = u^2 \\ x^2 - xy + y^2 = v^2 . \end{cases}$$

Conversely, any x, y corresponding to a rational point on E give rise to a solution of the original system (1.1) by multiplying by an appropriate scalar. Thus it suffices to determine all rational points on E . Now E is just a 2-covering of its Jacobian, and indeed we have the commutative diagram



$$J: Y^2 = 2X(X+1)(X+3) ,$$

the map $J \rightarrow J$ is multiplication by 2, and the diagonal map $E \rightarrow J$ is given by

$$X = -3u^2/2v^2 , \quad Y = 3(y^2-x^2)u/2v^3 . \tag{2.1}$$

The vertical map $J \rightarrow E$ is birational (but not, of course, necessarily defined over \mathbb{Q}).

Now from (2.1), rational points of E correspond precisely to points of J of which the x -coordinate is -6 modulo squares. It is a straightforward (albeit dull) calculation to show that the group of rational points on J has precisely one generator of infinite order, which may be taken as the point $P = (-3/2, 3/2)$. Then any rational point on J is of type $mP + P_0$, $m \in \mathbb{Z}$, where P_0 is one of the four 2-division points of J , namely: $(0,0)$, $(-1,0)$, $(-3,0)$, O (the point at infinity, being the zero of the group structure on J).

The addition formula on J is $(X_1, Y_1) + (X_2, Y_2) = (X_3, Y_3)$ with

$$X_3 = \frac{1}{2X_1X_2} \left(\frac{X_1Y_2 - X_2Y_1}{X_1 - X_2} \right)^2 ,$$

and so it is readily verified that the first coordinate of $mP + P_0$ is congruent to -6 modulo squares if and only if n is odd, and $P_0 = O$. Since $2P = (\frac{1}{8}, \frac{15}{16})$, it follows on writing $m = 2n + 1$ that the rational points of E are in 1-1 correspondence with the points of J

$$(X_n, Y_n) = n\left(\frac{1}{8}, \frac{15}{16}\right) + \left(-\frac{3}{2}, \frac{3}{2}\right) , \quad n \in \mathbb{Z} .$$

Using the group structure on J it is easy to deduce a recurrence relation for X_n :

$$X_{n+1} = \frac{1}{X_n} \left[\frac{15X_n - 2Y_n}{8X_n - 1} \right]^2 ;$$

and thus

$$\frac{-3u_{n+1}^2}{2v_{n+1}^2} = -\frac{3}{2} \left[\frac{15u_n v_n + 2(y_n^2 - x_n^2)}{12u_n^2 + v_n^2} \right]^2 .$$

Taking

$$u_{n+1} = 15u_n v_n + 2(y_n^2 - x_n^2) \tag{2.2}$$

$$v_{n+1} = 12u_n^2 + v_n^2 \tag{2.3}$$

one can now solve for x_{n+1}, y_{n+1} to get

$$x_{n+1} = -4x_n(u_n - v_n) + y_n(v_n + 14u_n) \quad (2.4)$$

$$y_{n+1} = x_n(v_n - 14u_n) + 4y_n(u_n + v_n). \quad (2.5)$$

Notice that $(x_{-n}, y_{-n}, u_{-n}, v_{-n}) = (x_{n-1}, y_{n-1}, -u_{n-1}, v_{n-1})$ and so it is only necessary to consider in the recurrences (2.2)-(2.5) non-negative values of n . The initial value $n = 0$ corresponding to $(a, b) = (0, 1)$. Then $(x_1, y_1, u_1, v_1) = (15, 8, 17, 13)$, corresponding to $(a, b) = (345, 184)$, $(x_2, y_2, u_2, v_2) = (1768, -2415, 2993, 3637)$, corresponding to $(a, b) = (-1143896, 1562505)$, etc.

REFERENCES

1. CASSELS, J.W.S., Arithmetic on an elliptic curve, Proc. Intern. Congress Math., Stockholm 1962, 234-246.
2. SELMER, E.S., The Diophantine equation $ax^3 + by^3 + cz^3 = 0$, Acta Math., 85, 1951, 203-362.

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.otero@usc.es