

A REGULAR GRAPH OF GIRTH 6 AND VALENCY 11

P.K. WONG

Department of Mathematics
Seton Hall University
South Orange, New Jersey 07079 U.S.A.

(Received November 20, 1985 and in revised form March 7, 1986)

ABSTRACT. Let $f(11,6)$ be the number of vertices of an $(11,6)$ -cage. By giving a regular graph of girth 6 and valency 11, we show that $f(11,6) \leq 240$. This is the best known upper bound for $f(11,6)$.

KEY WORDS AND PHRASES. *Regular graph, cage, latin square.*
1980 AMS SUBJECT CLASSIFICATION CODE. 05C35.

1. INTRODUCTION.

A graph is said to be regular of valency v if each of its vertices has valency v . A regular graph of valency v and girth g with the least possible number of vertices is called a (v,g) -cage. The number of vertices of a (v,g) -cage is denoted by $f(v,g)$. The existence of (v,g) -cages was proved by Erdős and Sachs [1].

When $g = 6$, it is easy to see that $f(v,6) \geq 2(v^2 - v + 1)$ ($v \geq 3$). Also, it is known that, when $v-1$ is a prime power, $f(v,6) = 2(v^2 - v + 1)$ and the graph can be constructed by using a complete set of orthogonal latin squares (see [2, p.7]). When $v=7$, O'Keefe and Wong showed that $f(7,6) = 90$ [3]. Thus far, no other $(v,6)$ -cages have been found.

The construction of an $(11,6)$ -cage is of great interest and this would settle an open problem in projective geometry: can a projective plane of order 10 exist? We now give a regular graph of girth 6 and valency 11 with 240 vertices. Therefore, $f(11,6) \leq 240$. This is the best known upper bound for $f(11,6)$. The graph is presented in the following way. Eighteen (18) "opposite" vertices A_n and B_n ($n = 1,2, \dots,9$) are added to Figure 8 in [2, p.7] with $k = 10$. B_n is adjacent to $A_n, 1n, 2n, \dots, 10n$ ($n=1,2,\dots,9$). The array of numbers in Table I shows how the vertices of set Y are joined to those of set X , row N standing for set N of Y ; the number M appearing in row N means that vertex NM is adjacent to the vertex of set X in the same column ($M, N = 1,2, \dots, 10$). We note that the sets $2,3, \dots, 10$ are obtained by permuting cyclically the rows of set 1. Because of this very unusual property of the graph, the author believes that this upper bound is not the best possible one.

Table I

(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(110)
A_1	8	7	1	4	9	5	3	10	6
3	A_2	9	8	2	5	10	6	4	7
2	4	A_3	10	9	3	6	1	7	8
6	3	5	A_4	1	10	4	7	2	9
9	7	4	6	A_5	2	1	5	8	10
4	10	8	5	7	A_6	3	2	6	1
10	5	1	9	6	8	A_7	4	3	2
8	1	6	2	10	7	9	A_8	5	3
5	9	2	7	3	1	8	10	A_9	4
7	6	10	3	8	4	2	9	1	5
(21)	(22)	(23)	(24)	(25)	(26)	(27)	(28)	(29)	(210)
3	A_2	9	8	2	5	10	6	4	7
2	4	A_3	10	9	3	6	1	7	8
6	3	5	A_4	1	10	4	7	2	9
9	7	4	6	A_5	2	1	5	8	10
4	10	8	5	7	A_6	3	2	6	1
10	5	1	9	6	8	A_7	4	3	2
8	1	6	2	10	7	9	A_8	5	3
5	9	2	7	3	1	8	10	A_9	4
7	6	10	3	8	4	2	9	1	5
A_1	8	7	1	4	9	5	3	10	6
(31)	(32)	(33)	(34)	(35)	(36)	(37)	(38)	(39)	(310)
2	4	A_3	10	9	3	6	1	7	8
6	3	5	A_4	1	10	4	7	2	9
9	7	4	6	A_5	2	1	5	8	10
4	10	8	5	7	A_6	3	2	6	1
10	5	1	9	6	8	A_7	4	3	2
8	1	6	2	10	7	9	A_8	5	3
5	9	2	7	3	1	8	10	A_9	4
7	6	10	3	8	4	2	9	1	5
A_1	8	7	1	4	9	5	3	10	6
3	A_2	9	8	2	5	10	6	4	7

(41)	(42)	(43)	(44)	(45)	(46)	(47)	(48)	(49)	(410)
6	3	5	A ₄	1	10	4	7	2	9
9	7	4	6	A ₅	2	1	5	8	10
4	10	8	5	7	A ₆	3	2	6	1
10	5	1	9	6	8	A ₇	4	3	2
8	1	6	2	10	7	9	A ₈	5	3
5	9	2	7	3	1	8	10	A ₉	4
7	6	10	3	8	4	2	9	1	5
A ₁	8	7	1	4	9	5	3	10	6
3	A ₂	9	8	2	5	10	6	4	7
2	4	A ₃	10	9	3	6	1	7	8
(51)	(52)	(53)	(54)	(55)	(56)	(57)	(58)	(59)	(510)
9	7	4	6	A ₅	2	1	5	8	10
4	10	8	5	7	A ₆	3	2	6	1
10	5	1	9	6	8	A ₇	4	3	2
8	1	6	2	10	7	9	A ₈	5	3
5	9	2	7	3	1	8	10	A ₉	4
7	6	10	3	8	4	2	9	1	5
A ₁	8	7	1	4	9	5	3	10	6
3	A ₂	9	8	2	5	10	6	4	7
2	4	A ₃	10	9	3	6	1	7	8
6	3	5	A ₄	1	10	4	7	2	9
(61)	(62)	(63)	(64)	(65)	(66)	(67)	(68)	(69)	(610)
4	10	8	5	7	A ₆	3	2	6	1
10	5	1	9	6	8	A ₇	4	3	2
8	1	6	2	10	7	9	A ₈	5	3
5	9	2	7	3	1	8	10	A ₉	4
7	6	10	3	8	4	2	9	1	5
A ₁	8	7	1	4	9	5	3	10	6
3	A ₂	9	8	2	5	10	6	4	7
2	4	A ₃	10	9	3	6	1	7	8
6	3	5	A ₄	1	10	4	7	2	9
9	7	4	6	A ₅	2	1	5	8	10

(71) (72) (73) (74) (75) (76) (77) (78) (79) (710)

10	5	1	9	6	8	A ₇	4	3	2
8	1	6	2	10	7	9	A ₈	5	3
5	9	2	7	3	1	8	10	A ₉	4
7	6	10	3	8	4	2	9	1	5
A ₁	8	7	1	4	9	5	3	10	6
3	A ₂	9	8	2	5	10	6	4	7
2	4	A ₃	10	9	3	6	1	7	8
6	3	5	A ₄	1	10	4	7	2	9
9	7	4	6	A ₅	2	1	5	8	10
4	10	8	5	7	A ₆	3	2	6	1

(81) (82) (83) (84) (85) (86) (87) (88) (89) (810)

8	1	6	2	10	7	9	A ₈	5	3
5	9	2	7	3	1	8	10	A ₉	4
7	6	10	3	8	4	2	9	1	5
A ₁	8	7	1	4	9	5	3	10	6
3	A ₂	9	8	2	5	10	6	4	7
2	4	A ₃	10	9	3	6	1	7	8
6	3	5	A ₄	1	10	4	7	2	9
9	7	4	6	A ₅	2	1	5	8	10
4	10	8	5	7	A ₆	3	2	6	1
10	5	1	9	6	8	A ₇	4	3	2

(91) (92) (93) (94) (95) (96) (97) (98) (99) (910)

5	9	2	7	3	1	8	10	A ₉	4
7	6	10	3	8	4	2	9	1	5
A ₁	8	7	1	4	9	5	3	10	6
3	A ₂	9	8	2	5	10	6	4	7
2	4	A ₃	10	9	3	6	1	7	8
6	3	5	A ₄	1	10	4	7	2	9
9	7	4	6	A ₅	2	1	5	8	10
4	10	8	5	7	A ₆	3	2	6	1
10	5	1	9	6	8	A ₇	4	3	2

(101)	(102)	(103)	(104)	(105)	(106)	(107)	(108)	(109)	(1010)
7	6	10	3	8	4	2	9	1	5
A_1	8	7	1	4	9	5	3	10	6
3	A_2	9	8	2	5	10	6	4	7
2	4	A_3	10	9	3	6	1	7	8
6	3	5	A_4	1	10	4	7	2	9
9	7	4	6	A_5	2	1	5	8	10
4	10	8	5	7	A_6	3	2	6	1
10	5	1	9	6	8	A_7	4	3	2
8	1	6	2	10	7	9	A_8	5	3
5	9	2	7	3	1	8	10	A_9	4

REFERENCES

1. ERDÖS, P. and SACHS, H., Reguläre Graphen Gegebener Tailenweite mit minimaler Knotenzahl, WISS.Z. Uni. Halle (Math. Nat.) 12(1963),251-257.
2. WONG, P. K., Cages-A survey. J. Graph Theory 6(1982), 1-22.
3. O'KEEFE, M. and WONG, P. K., The smallest graph of girth 6 and valency 7, J. Graph Theory 5(1981), 79-85.
4. DENES, J. and KEEDWELL, A. D., Latin squares and their applications, Academic, New York (1974).

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br