

RESEARCH NOTES

ON COLLINEATION GROUPS OF TRANSLATION PLANES OF ORDER q^4

V. JHA and N.L. JOHNSON

Department of Mathematics
The University of Iowa
Iowa City, IA 52242 U.S.A.

(Received April 9, 1984 and in revised form March 31, 1986)

ABSTRACT. Let P be an affine translation plane of order q^4 admitting a nonsolvable group G in its translation complement. If G fixes more than $q+1$ slopes, the structure of G is determined. In particular, if G is simple then q is even and $G = L_2(2^s)$ for some integer s at least 2.

KEY WORDS AND PHRASES. *Translation planes, nonsolvable groups.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. 50D05, 05B25

1. INTRODUCTION.

Let π be a translation plane of square order p^{2r} . If π admits a collineation group isomorphic to $SL(2, p^t)$ and the Sylow p -subgroups are planar, then usually (in the known cases) the group fixes p^r+1 components (or slopes). Generally, however, simply knowing that a group fixes a number of slopes says essentially nothing concerning the structure of the group. However, for planes of order q^4 , we can make some progress. That is, in this note our objective is to prove the following.

THEOREM A. Let π be an affine translation plane of order q^4 admitting a nonsolvable group G in its translation complement. Suppose G fixes more than $q+1$ slopes.

(1) If q is odd then $8 \mid |G|$ and the 2-rank of $G \leq 2$. Furthermore, G always contains the kern involution.

(2) If q is even then G contains a normal subgroup N such that $N \cong L_2(2^s)$, for some s , and G/N is of odd order. Now the Sylow 2-subgroups fix Baer subplanes elementwise. Furthermore, the Baer subplanes share the same points at infinity and so N fixes exactly q^2+1 slopes.

COROLLARY 1. If G is simple then q is even and $G = L_2(2^s)$ for some integer $s \geq 2$.

PROOF. When q is odd the kern involution is central in G and so G is not simple.

REMARKS. (1) When q is odd it is possible to find G 's with 2-rank one and 2-rank two such that they satisfy the hypothesis of Theorem A. For example, $G_1 = \text{SL}(2, q)$, acting on Hall planes of order q^4 , has 2-rank one; if we now choose $G_2 = \langle G_1, \alpha \rangle$, where α is any Baer involution, we get a ' G ' with 2-rank two.

(2) Both the theorem and its corollary cease to be unconditionally true if we allow G to fix "at least $q+1$ slopes", instead of "more than $q+1$ slopes": the only known counterexamples seem to be the Lorimer-Rahilly planes [1] and its transpose, the Johnson-Walker plane [2].

The following well known consequences of Foulser [3] will be used on several occasions in the proof of Theorem A.

RESULT 0. Suppose B is a collineation group of an affine translation plane of order p^{2r} that fixes a Baer subplane elementwise. Then

- (i) B is solvable;
- (ii) the Sylow p -subgroups of B are elementary abelian; and
- (iii) the Hall p' subgroups of B are cyclic.

2. PROOF OF THEOREM A.

We begin by dealing with the case when q is odd. The first step is folklore and corresponds to Ostrom's ideas in [4].

LEMMA 1. If q is odd then any Klein 4-group in G must contain the unique kern involution of π ; we shall always denote this involution by \hat{i} .

PROOF. Let $K = \{1, \alpha, \beta, \alpha\beta\}$ be a Klein group in G and suppose, if possible that $\hat{i} \notin K$. For any involution x in K write π_x for its fixed Baer subplane. Now we claim $\pi_x \cap \pi_y$ cannot be a subplane of π if x, y are distinct involutions in K . If $\pi_x = \pi_y$, we have a Klein group fixing elementwise a Baer subplane of π , contrary to Result 0. So $\pi_x \cap \pi_y$ is a fourth root subplane of π and now we contradict the assumption that G fixes more than $q+1$ slopes. Thus xy acts like -1 on the fixed components of G , in the spread associated with π ; i.e., xy is the required involution.

LEMMA 2. If q is odd then $|G|$ is divisible by 8.

PROOF. If 2 exactly divides $|G|$ then, by Burnside's theorem, G has a normal 2-complement [5, 6.2.11] and we contradict the assumption that G is nonsolvable. For the same reason the Sylow 2-subgroups of G cannot be cyclic of order 4. Hence $4 \parallel |G|$ only if the Sylow 2-subgroups are Klein groups. So by Lemma 1, G contains the kern involution \hat{i} and $G/\langle \hat{i} \rangle$ is solvable. Thus we contradict the nonsolvability of G when $8 \nmid |G|$. The result follows.

LEMMA 3. Suppose q is odd. Then G cannot contain an elementary abelian 2-group of order 8.

PROOF. If S is an elementary abelian subgroup of G , whose order is 8, we may write

$$S = \{1, \alpha, \beta, \alpha\beta, \gamma, \alpha\gamma, \beta\gamma, \alpha\beta\gamma\} \quad (2.1)$$

and assume that

$$L = \{1, \alpha, \beta, \alpha\beta\}, \quad M = \{1, \alpha, \gamma, \alpha\gamma\} \tag{2.2}$$

are distinct subgroups of order 4. But by Lemma 1, both L and M contain the kern involution \hat{i} . Hence $\alpha = \hat{i}$. Interchanging the role of α and β , we find that β is also \hat{i} . The lemma follows, since we have contradicted the assumption that $|S| = 8$.

LEMMA 4. G contains \hat{i} , the kern involution of π .

PROOF. Let S denote a Sylow 2-subgroup of G. So $|S| \geq 8$ (Lemma 2) and non-cyclic, because G is nonsolvable. Now Lemma 1 applies unless the 2-rank of S is one. Thus S is the generalized quaternion group

$$\langle x, y \mid x^{2^n} = 1, y^2 = x^{2^{n-1}}, y^{-1}xy = x^{-1} \text{ for } n \geq 2 \rangle \tag{2.3}$$

and so contains $Q = \langle x^{2^{n-2}}, y \rangle$, the quaternion group of order 8.

Now let α denote the unique involution in Q and, to get a contradiction, assume α is a Baer involution with fixed plane π_α . Now Q leaves π_α invariant but does not fix it elementwise because of Result 0. Moreover, no element of Q can induce a Baer involution on π_α because Q fixes $> q+1$ slopes of π_α . Thus the restriction map

$$\rho : Q \longrightarrow Q \mid \pi_\alpha \tag{2.4}$$

has as its image $\langle \beta \rangle$, where β is the kern involution of π_α . So $\ker \rho$ is clearly a noncyclic group Σ of order 4, contrary to Result 0.

The lemmas proved so far add up to Theorem A, Part (1). To deal with the case when q is even we need the following version of a theorem of Johnson [6], deduced from Hering [7].

RESULT 5. Suppose ψ is an affine translation plane of even order admitting a nonsolvable group H in its translation complement. Assume a Sylow 2-subgroup of H fixes a Baer subplane elementwise. Then H contains a normal subgroup N such that H/N is of odd order and $N \cong L_2(2^s)$ for some integer s.

PROOF. Use Johnson's argument in [6, Theorem 2.3].

LEMMA 6. If q is even then the Sylow 2-subgroups of G fix Baer subplanes of π elementwise.

PROOF. Let S be a Sylow 2-subgroup of G and note that elements of π fixed by S form a subplane π_S , because S fixes many slopes. To get a contradiction we assume π_S is not a Baer subplane of π . Now let α be any involution in the center of S and let π_α be its fixed Baer subplane. Then we have a chain of planes $\pi \supset \pi_\alpha \supset \pi_S$. Hence the order of $\pi_S \leq q$ and we contradict the assumption that G fixes more than $q+1$ slopes. The result follows.

Now by Result 5 and Lemma 6 we immediately have

LEMMA 7. Suppose q is even. Then the Sylow 2-subgroups of G fix Baer subplanes elementwise and generate of a subgroup $N \cong L_2(2^s)$ where $2^s \parallel |G|$.

To complete the proof of Theorem A we restrict ourselves from now on to the situation described in Lemma 7.

LEMMA 8. N fixes a unique affine point O .

PROOF. Suppose N , which is in the translation complement of π , fixes a second affine point of π . Then N is a planar group of π of order $> q$. Now by Lemma 7 we have a Baer chain $\pi \supset \pi_S \supset \pi_N$. If $\pi_N \neq \pi_S$ we have the same contradiction as in Lemma 6; otherwise we have a nonsolvable group fixing a Baer subplane element-wise, contrary to Result 0.

LEMMA 9. The only affine point fixed by distinct Sylow 2-subgroups of N is O , the unique point fixed by N .

PROOF. N is generated by any two of its Sylow 2-subgroups, so we contradict Lemma 8 unless the lemma is valid.

For $2^S \neq 4$, Lemma 9, when combined with Foulser and Johnson [8, Proposition 3.4], shows that N fixes q^2+1 slopes. From Johnson [9, Theorem 2.1], the same is true if $2^S = 4$ as N fixes at least 3 slopes. This proves Part (2) of Theorem A.

REFERENCES

1. LORIMER, P. A Projective Plane of Order 16, J. Combinatorial Theory A16 (1974), 334-347.
2. WALKER, M. A Note on Tangentially Transitive Affine Planes, Bull. London Math. Soc. 8 (1976), 273-277.
3. FOULSER, D.A. Subplanes of Partial Spreads in Translation Planes, Bull. London Math. Soc. 4 (1972), 32-38.
4. OSTROM, T.G. Elementary Abelian 2-Groups in Finite Translation Planes, Arch. Mat. 36 (1981).
5. SCOTT, W.R. Group Theory, Prentice Hall, New Jersey, 1964.
6. JOHNSON, N.L. The Geometry of $GL(2,q)$ in Translation Planes of Even Order q^2 , Internat. J. Math. & Math. Sci. 1 (1978), 447-458.
7. HERING, Ch. On Subgroups with Trivial Normalizer Intersection, J. Algebra 20 (1972), 622-629.
8. FOULSER, D.A. and JOHNSON, N.L. The Translation Planes of Order p^2 that Admit $SL(2,q)$ as a Collineation Group. II. Odd Order, J. Geometry 18 (1982), 122-139.
9. JOHNSON, N.L. Representation of $SL(2,4)$ on Translation Planes of Even Order, J. Geometry 21 (1983), 184-200.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br