

Betti numbers of lex ideals over some Macaulay-Lex rings

Jeff Mermin · Satoshi Murai

Received: 2 August 2008 / Accepted: 15 June 2009 / Published online: 27 June 2009
© Springer Science+Business Media, LLC 2009

Abstract Let $A = K[x_1, \dots, x_n]$ be a polynomial ring over a field K and M a monomial ideal of A . The quotient ring $R = A/M$ is said to be Macaulay-Lex if every Hilbert function of a homogeneous ideal of R is attained by a lex ideal. In this paper, we introduce some new Macaulay-Lex rings and study the Betti numbers of lex ideals of those rings. In particular, we prove a refinement of the Frankl–Füredi–Kalai Theorem which characterizes the face vectors of colored complexes. Additionally, we disprove a conjecture of Mermin and Peeva that lex-plus- M ideals have maximal Betti numbers when A/M is Macaulay-Lex.

Keywords Lex ideals · Graded Betti numbers · Hilbert functions · Colored simplicial complexes

1 Introduction

The Hilbert function is an important invariant of homogeneous ideals of a polynomial ring $A = K[x_1, \dots, x_n]$ over a field K , studied in commutative algebra, algebraic geometry and combinatorics. One of the central results in the study of Hilbert functions is Macaulay’s Theorem [14], which characterizes the Hilbert functions of homogeneous ideals of A in terms of lex ideals. In the 1990’s, a remarkable extension

The first author is supported by an NSF Postdoctoral fellowship (award No. DMS-0703625).
The second author is supported by JSPS Research Fellowships for Young Scientists.

J. Mermin

Department of Mathematics, University of Kansas, Lawrence, KS 66045, USA
e-mail: mermin@math.ku.edu

S. Murai (✉)

Department of Mathematics, Graduate School of Science, Kyoto University, Sakyou-ku,
Kyoto 606-8502, Japan
e-mail: murai@math.kyoto-u.ac.jp

of Macaulay's Theorem was proved by Bigatti [2], Hulett [12] and Pardue [23]. They proved that lex ideals have the greatest graded Betti numbers among all homogeneous ideals having the same Hilbert function. In this paper, we introduce a class of monomial ideals M such that Macaulay's Theorem holds for the quotient ring A/M , and study the graded Betti numbers of lex ideals of those rings.

Let M be a monomial ideal of A and set $R = A/M$. Recall that the *Hilbert function* $\text{Hilb}(N)(-): \mathbb{Z} \rightarrow \mathbb{Z}$ of a finitely generated graded R -module N is the function defined by

$$\text{Hilb}(N)(d) = \dim_K N_d,$$

where N_d is the homogeneous component of degree d of N . A set W of monomials of R is said to be a *lex-segment* if, for all monomials $u, v \in R$ of the same degree, $u \in W$ and $v >_{\text{lex}} u$ imply $v \in W$, where $>_{\text{lex}}$ is the degree lexicographic order. We say that a monomial ideal I of R is a *lex ideal* if the set of monomials in I is a lex-segment. The ring R is said to be *Macaulay-Lex* if, for any homogeneous ideal J of R , there exists a lex ideal of R having the same Hilbert function as J .

By Macaulay's Theorem [14], the polynomial ring A itself is Macaulay-Lex. A famous class of Macaulay-Lex rings is the Clements–Lindström rings [4] $R = A/(x_1^{a_1}, \dots, x_n^{a_n})$, where $1 \leq a_1 \leq \dots \leq a_n$ are integers or ∞ . The notion of Macaulay-Lex rings was introduced in [17], and basic properties of Macaulay-Lex rings were established in [17, 18]. A fundamental problem about Macaulay-Lex rings is the following.

Problem 1.1 (Mermin–Peeva) Find classes of monomial ideals M of A such that A/M is Macaulay-Lex.

A homogeneous ideal I of A is said to be *homogeneous-plus- M* (resp. *lex-plus- M*) if there exists a homogeneous (resp. lex) ideal J such that $I = J + M$. Clearly, A/M is Macaulay-Lex if and only if, for any homogeneous-plus- M ideal I , there exists a lex-plus- M ideal having the same Hilbert function as I . Inspired by the Bigatti–Hulett–Pardue Theorem as well as Evans' Lex-plus-powers Conjecture [7], Mermin and Peeva made the following conjecture in [18].

Conjecture 1.2 (Mermin–Peeva) Suppose that $R = A/M$ is Macaulay-Lex.

- (1) Every lex ideal L of R has the greatest graded Betti numbers among all homogeneous ideals of R having the same Hilbert function as L .
- (2) Every lex-plus- M ideal L of A has the greatest graded Betti numbers among all homogeneous-plus- M ideals of A having the same Hilbert function as L .

Note that (1) considers infinite free resolutions, while (2) considers finite free resolutions. Conjecture 1.2 has been well studied for Clements–Lindström rings $R = A/(x_1^{a_1}, \dots, x_n^{a_n})$. In this special case, Conjecture 1.2(2) was proved in a series of papers [16, 19, 20], and Conjecture 1.2(1) was proved in [22] when the characteristic of K is 0. On the other hand, little is known for other Macaulay-Lex rings. In this paper we consider the following rings.

Definition 1.3 Let $V = \bigcup_{j=1}^r V_j$ be a set of variables with $V_j = \{x_{j,1}, \dots, x_{j,n_j}\}$, where $n_1 \geq n_2 \geq \dots \geq n_r$. Denote by $S = K[V]$ the polynomial ring over K with the set of variables V . We will work with the lexicographic order $>_{\text{lex}}$ on S induced by the ordering of the variables defined by $x_{k,\ell} > x_{k',\ell'}$ if $\ell > \ell'$ or $\ell = \ell'$ and $k < k'$. Let $Q = \sum_{j=1}^r (x_{j,1}, \dots, x_{j,n_j})^2 \subset S$. We call the ring $R = S/Q$ an *r-colored squarefree ring of type (n_1, \dots, n_r)* .

We say that a quotient ring $R = A/M$ admits ideals with maximal Betti numbers over A if, whenever H is the Hilbert function of some homogeneous-plus- M ideal of A , there exists a homogeneous-plus- M ideal L with Hilbert function H such that $\beta_{i,j}(L) \geq \beta_{i,j}(I)$ for all i, j and for all homogeneous-plus- M ideals I with Hilbert function H , where $\beta_{i,j}(J)$ are the graded Betti numbers of an ideal J of A . Thus Conjecture 1.2(2) states that Macaulay-Lex rings admit ideals with maximal Betti numbers. The main results of this paper are the following.

- Colored squarefree rings are Macaulay-Lex.
- An r -colored squarefree ring of type (n_1, \dots, n_r) does not admit ideals with maximal Betti numbers over S if $r = 2$ and $n_2 \geq 4$ or if $r \geq 3$ and $n_r \geq 3$.
- A computation of the graded Betti numbers of Borel ideals over R .

In particular, the second result disproves Conjecture 1.2(2).

The first result is inspired by the Frankl–Füredi–Kalai Theorem [8], which characterizes face vectors of colored simplicial complexes. Indeed, if $n_1 = \dots = n_r$ then the result is equivalent to the Frankl–Füredi–Kalai Theorem (see Remark 2.12 for details). Our proof is different from the original proof of the Frankl–Füredi–Kalai Theorem, but similar to another proof of the theorem given by London [13]. Frankl–Füredi–Kalai used a combinatorial technique called *shifting*, while our proof is based on *compression*, a technique which was introduced by Macaulay [14] and used efficiently by Clements–Lindström [4].

This paper is organized as follows: In Section 2, we show that colored squarefree rings are Macaulay-Lex. In Section 3, we study the graded Betti numbers of lex-plus- Q ideals and disprove Conjecture 1.2(2). In Section 4, we study the graded Betti numbers of Borel ideals of r -colored squarefree rings. In Section 5, we discuss some related problems.

2 The Macaulay-Lex property

In studying Problem 1.1 and Conjecture 1.2, it is enough to consider monomial ideals since the initial ideal of a homogeneous-plus- M ideal I is a monomial ideal which contains M and which has the same Hilbert function as I (see e.g., [6, Chap. 15]). Thus, throughout this paper, we assume that all the ideals are monomial ideals. The main result of this section is the following.

Theorem 2.1 *Let R be an r -colored squarefree ring. For any monomial ideal I of R , there exists a unique lex ideal L of R having the same Hilbert function as I .*

In the rest of this section, $R = S/Q$ stands for an r -colored squarefree ring as defined in Definition 1.3. For a subset $\mathcal{A} \subset [r] = \{1, 2, \dots, r\}$, write $\bar{\mathcal{A}} = [r] \setminus \mathcal{A}$, $V_{\mathcal{A}} = \bigcup_{j \in \mathcal{A}} V_j$, $S_{\mathcal{A}} = K[V_{\mathcal{A}}]$, $Q_{\mathcal{A}} = \sum_{j \in \mathcal{A}} (x_{j,1}, \dots, x_{j,n_j})^2 \subset S_{\mathcal{A}}$ and $R_{\mathcal{A}} = S_{\mathcal{A}}/Q_{\mathcal{A}}$. For any set W of monomials of R , we write W_d for the set of monomials in W of degree d .

Definition 2.2 Let W be a set of monomials of R and $\mathcal{A} \subset [r]$. Then we may decompose W as the disjoint union

$$W = \bigsqcup_{f \in R_{\bar{\mathcal{A}}}} fW_f$$

where f ranges over the monomials in $R_{\bar{\mathcal{A}}}$ and each W_f is a set of monomials in $R_{\mathcal{A}}$. We say that W is \mathcal{A} -compressed if all the W_f are lex-segments of $R_{\mathcal{A}}$. Moreover, we say that W is compressed if W is \mathcal{A} -compressed for all $\mathcal{A} \subset [r]$ with $|\mathcal{A}| = r - 1$, where $|\mathcal{A}|$ is the cardinality of \mathcal{A} . Let $L_f = \bigsqcup_{d \geq 0} (L_f)_d$ be the lex-segment set of monomials in $R_{\mathcal{A}}$ such that $|(W_f)_d| = |(L_f)_d|$ for all d . The set of monomials $X = \bigsqcup_f fL_f$ is called the \mathcal{A} -compression of W .

Let I be a monomial ideal of R and M the set of monomials in I . The \mathcal{A} -compression of I is the K -vector space spanned by the \mathcal{A} -compression of M .

Notation 2.3 Let W be a set of monomials in R_d . Define

$$\text{Shad}(W) = \{ym \in R_{d+1} : y \in V, m \in W\}.$$

Write $\text{Lex}(W) \subset R_d$ for the lex-segment set of monomials with $|W| = |\text{Lex}(W)|$.

For any monomial $m \in R$, let $\text{first}(m)$ (resp. $\text{last}(m)$) be the greatest (resp. smallest) variable which divides m . Let

$$\text{color}(m) = \{j \in [r] : \text{there exists } y \in V_j \text{ such that } y \text{ divides } m\}.$$

The following facts are straightforward (see, e.g., [17]).

Lemma 2.4 *If $W \subset R$ is a lex-segment set of monomials of the same degree then $\text{Shad}(W)$ is also a lex-segment.*

Corollary 2.5 *Theorem 2.1 holds if and only if, for any set $W \subset R$ of monomials of the same degree, one has $|\text{Shad}(W)| \geq |\text{Shad}(\text{Lex}(W))|$.*

We will prove Theorem 2.1 by using Corollary 2.5 and induction on r .

Lemma 2.6 *Theorem 2.1 holds if $r \leq 2$.*

Proof The statement is obvious if $r = 1$. Suppose $r = 2$. Note that $R = R_0 \oplus R_1 \oplus R_2$. Then, by Corollary 2.5, it is enough to show that, for any set $W \subset R_1$ of monomials, one has $|\text{Shad}(W)| \geq |\text{Shad}(\text{Lex}(W))|$.

Let $a_k = |W \cap V_k|$ for $k = 1, 2$. A routine computation implies

$$|\text{Shad}(W)| = n_1a_2 + n_2a_1 - a_1a_2 = -(n_1 - a_1)(n_2 - a_2) + n_1n_2.$$

Then $|\text{Shad}(W)|$ is smallest when the difference between $(n_1 - a_1)$ and $(n_2 - a_2)$ is minimized. Hence $|\text{Shad}(W)| \geq |\text{Shad}(\text{Lex}(W))|$ by the definition of $>_{\text{lex}}$. \square

Definition 2.7 Let $W = \{u_1, \dots, u_t\}$ and $W' = \{u'_1, \dots, u'_t\}$ be sets of monomials of R with $u_1 >_{\text{lex}} \dots >_{\text{lex}} u_t$ and $u'_1 >_{\text{lex}} \dots >_{\text{lex}} u'_t$. We say that W is *lex-greater than* W' if there exists $1 \leq j \leq t$ such that $u_k = u'_k$ for $k < j$ and $u_j >_{\text{lex}} u'_j$.

Lemma 2.8 *Suppose that Theorem 2.1 holds for all $(r - 1)$ -colored squarefree rings. Let $\mathcal{A} \subset [r]$ with $|\mathcal{A}| = r - 1$.*

- (i) *For any monomial ideal I of R , the \mathcal{A} -compression of I is an ideal of R .*
- (ii) *Let $W \subset R_d$ be a set of monomials and W' the \mathcal{A} -compression of W . Then W' is lex-greater than or equal to W and $|\text{Shad}(W)| \geq |\text{Shad}(W')|$.*

Proof (i) Let $M = \bigoplus_f f M_f$ be the set of monomials in I , where $f \in R_{\bar{\mathcal{A}}}$ is a monomial and $M_f \subset R_{\mathcal{A}}$. Let $L = \bigoplus_f f L_f$ be the \mathcal{A} -compression of M . Note that the vector space spanned by M_f is a monomial ideal of $R_{\mathcal{A}}$. By the assumption, the vector space spanned by L_f is an ideal of $R_{\mathcal{A}}$. Hence, what we must prove is that, for any $fu \in f L_f$ and for any variable $y \in V_{\bar{\mathcal{A}}}$, one has $yfu \in L$ or $yfu = 0$.

Suppose $yf \neq 0$. Since I is an ideal, $W_f \subset W_{yf}$. Hence $L_f \subset L_{yf}$. This implies $u \in L_{yf}$ and $yfu \in yf L_{yf} \subset L$.

(ii) It is clear that W' is lex-greater than or equal to W . Let I be the monomial ideal generated by W and J the \mathcal{A} -compression of I . Then $|\text{Shad}(W)| = \dim_K I_{d+1}$ and $|\text{Shad}(W')| \leq \dim_K J_{d+1} = \dim_K I_{d+1}$ by (i). Hence the statement follows. \square

Lemma 2.9 *Let $W \subset R_d$ be a compressed set of monomials. Let $u \in W$ and $v \in R_d$. If $v >_{\text{lex}} u$ and if u and v are divisible by some variable y then $v \in W$.*

Proof Let $\mathcal{A} = [r] \setminus \text{color}(y)$. Since W is \mathcal{A} -compressed and $\frac{u}{y} \in W_y$, we have $\frac{v}{y} \in W_y$, i.e., $v \in W$. \square

For any monomial $m \in R$, write

$$\text{grow}(m) = |\{y \in V_{[r] \setminus \text{color}(m)} : y < \text{last}(m)\}|.$$

Note that if $W \subset R_d$ is a lex-segment, then $|\text{Shad}(W)| = \sum_{m \in W} \text{grow}(m)$. This definition is inspired by work of Bigatti [2], who used the analogous formula in a polynomial ring to study Borel ideals.

Lemma 2.10 *Let m and m' be monomials of degree d with $\text{last}(m) = x_{s,t}$ and $\text{last}(m') = x_{s',t'}$. If $t < t'$ then $\text{grow}(m) \leq \text{grow}(m')$.*

Proof We may assume $t' = t + 1$. For any $j \in \text{color}(m)$ one has

$$|\{y \in V_j : y \leq x_{1,t}\}| = t. \tag{1}$$

Indeed, if $|\{y \in V_j : y \leq x_{1,t}\}| \leq t - 1$ then $n_j \leq t - 1$. However, since $j \in \text{color}(m)$, some $x_{j,\ell}$ is greater than $x_{s,t}$. This means $n_j \geq \ell \geq t$, a contradiction. Then (1) implies $\text{grow}(m') \geq |\{y \in V : y \leq x_{1,t}\}| - dt \geq \text{grow}(m)$. \square

Lemma 2.11 *Suppose $r \geq 3$. Let $W \subset R_d$ be a compressed set of monomials which is not a lex-segment. There exists a set $W' \subset R_d$ of monomials such that $|W| = |W'|$, $|\text{Shad}(W)| \geq |\text{Shad}(W')|$, and W' is lex-greater than W .*

Proof If $r \geq 3$ then any compressed subset $W \subset R_1$ is a lex-segment. Suppose $d \geq 2$. Let $g \in R_d$ be the lex-greatest monomial which is not in W and b the lex-smallest monomial in W . Set $\tilde{W} = W \cup \{g\}$ and $W' = (W \setminus \{b\}) \cup \{g\}$. Then, by the choice of g and b , a straightforward computation implies

$$|\text{Shad}(\tilde{W})| \leq |\text{Shad}(W)| + \text{grow}(g) \quad \text{and} \quad |\text{Shad}(\tilde{W})| \geq |\text{Shad}(W')| + \text{grow}(b).$$

Hence, to prove the statement, it is enough to show

$$\text{grow}(g) \leq \text{grow}(b). \tag{2}$$

The statement is obvious if $d = r$. Hence we may assume $2 \leq d < r$. Let u be the lex-greatest monomial in W such that $u <_{\text{lex}} g$. Set $y_1 = \text{first}(u)$, $y_0 = \text{first}(g)$ and $z = \text{last}(b)$. Clearly, $y_0 \geq y_1$ since $g >_{\text{lex}} u$. Moreover, since W is compressed, Lemma 2.9 implies

$$y_0 > y_1.$$

[Case 1]: Suppose $\text{color}(y_0) \neq \text{color}(z)$. Since $u \geq_{\text{lex}} b$, we have $y_0 > y_1 \geq \text{first}(b)$. Let f be the lex-smallest monomial of degree d which is divisible by y_0z . Since $y_0 > \text{first}(b)$, $f >_{\text{lex}} b$. Since f and b are divisible by z , Lemma 2.9 implies $f \in W$. Also, since $f >_{\text{lex}} u$ and since u is the lex-greatest monomial in W with $u <_{\text{lex}} g$, we have $f >_{\text{lex}} g$.

Let $w = \text{first}(\frac{f}{y_0})$. Since f is the lex-smallest monomial of degree d which is divisible by y_0z , if $w \neq z$ then f is the lex-smallest monomial of degree d which is divisible by y_0 . However, this cannot happen since $f >_{\text{lex}} g$ and g is divisible by y_0 . Hence $z = w$. Write $g = y_0w_1 \cdots w_{d-1}$, where $y_0 > w_1 > \cdots > w_{d-1}$. Since $\text{first}(\frac{f}{y_0}) = z$, $\frac{f}{y_0} >_{\text{lex}} \frac{g}{y_0}$, and $\frac{f}{y_0}$ is the lex-smallest monomial of degree $d - 1$ which is divisible by z , we have $z > w_1 > \cdots > w_{d-1}$. Let $z = x_{p,q}$ and $w_{d-1} = x_{s,t}$. Note that $t \leq q$ since $z > w_{d-1}$. If $t < q$ then, by Lemma 2.10, we have $\text{grow}(g) \leq \text{grow}(b)$ as desired. If $t = q$ then, for each k , $w_k = x_{\ell_k,q}$ for some $p < \ell_k \leq s$. Hence, by (1),

$$\begin{aligned} \text{grow}(b) &\geq \left| \{y \in V_{[r] \setminus \text{color}(y_0)} : y < x_{p,q}\} \right| - (d - 1)q \\ &\geq \left| \{y \in V_{[r] \setminus \text{color}(y_0)} : y < x_{s,t}\} \right| + (d - 1) - (d - 1)q \\ &= \left| \{y \in V_{[r] \setminus \text{color}(y_0)} : y < x_{s,t}\} \right| - (d - 1)(q - 1) = \text{grow}(g). \end{aligned}$$

[Case 2]: Suppose $\text{color}(y_0) = \text{color}(z) = \{c\}$. Fix $a \in [r] \setminus \text{color}(b)$. Note that $\{a\} \neq \text{color}(y_0)$. Let $f = y_0f'$ be the lex-smallest monomial of degree d such that f is divisible by y_0 and $a \notin \text{color}(f)$. Since $y_0 > y_1$, $f >_{\text{lex}} u \geq_{\text{lex}} b$. Then, since $f, b \in R_{[r] \setminus \{a\}}$ and W is compressed, we have $f \in W$. In particular, $f >_{\text{lex}} g$ by the choice of u . Hence f' cannot be the lex-smallest monomial of degree $d - 1$ in $R_{[r] \setminus \{c\}}$.

Let m be the lex-smallest monomial of degree d in $R_{[r]\setminus\{c\}}$. Since f' is lex-smallest in $(R_{[r]\setminus\{a,c\}})_{d-1}$ but is not lex-smallest in $(R_{[r]\setminus\{c\}})_{d-1}$, m is divisible by $x_{a,1}$ and $f' = \frac{m}{x_{a,1}}$. Then $\frac{g}{y_0} <_{\text{lex}} f' \leq_{\text{lex}} \frac{m}{\text{last}(m)}$. Since m is lex-smallest in $(R_{[r]\setminus\{c\}})_d$,

$$\text{last}(g) = \text{last}(m) = \begin{cases} x_{r,1}, & \text{if } c \neq r, \\ x_{r-1,1}, & \text{if } c = r. \end{cases}$$

In both cases, we have $\text{grow}(g) = 0 \leq \text{grow}(b)$. □

Now we are in position to prove Theorem 2.1.

Proof of Theorem 2.1 We use induction on r . We may assume $r \geq 3$ by Lemma 2.6. Suppose that the statement holds for all $(r - 1)$ -colored squarefree rings.

Let W be a set of monomials of degree d . By Corollary 2.5, it is enough to show that $|\text{Shad}(W)| \geq |\text{Shad}(\text{Lex}(W))|$. By Lemmas 2.8 and 2.11, if W is not a lex-segment then there exists $W' \subset R_d$ such that $|W'| = |W|$, $|\text{Shad}(W')| \leq |\text{Shad}(W)|$ and W' is lex-greater than W . Arguing inductively, we have $|\text{Shad}(W)| \geq |\text{Shad}(\text{Lex}(W))|$. □

Remark 2.12 Here we note the relation between Theorem 2.1 and face vectors of colored simplicial complexes. A simplicial complex Δ on $[n] = \{1, 2, \dots, n\}$ is a collection of subsets of $[n]$ such that, if $F \in \Delta$ and $G \subset F$, then $G \in \Delta$. A simplicial complex Δ is said to be r -colored if there exists a partition of $[n]$, $[n] = C_1 \cup \dots \cup C_r$, such that for every $F \in \Delta$ and every $1 \leq i \leq r$, $|C_i \cap F| \leq 1$. In particular, if $r = \max\{|F| : F \in \Delta\}$ then Δ is called *completely balanced*.

Let $H_i = \{k \in [n] : k \equiv i \pmod{r}\}$ for $i = 1, 2, \dots, r$ and let \mathcal{C} be the collection of subsets $F \subset [n]$ satisfying $|F \cap H_i| \leq 1$ for all i . Let $>_{\text{rev}}$ be the reverse lexicographic order induced by $1 >_{\text{rev}} \dots >_{\text{rev}} n$. An r -colored *rev-lex complex* $\Gamma \subset \mathcal{C}$ is an r -colored simplicial complex such that, for any faces $F \in \Gamma$ and $G \in \mathcal{C}$ with $|F| = |G|$, if $G >_{\text{rev}} F$ then $G \in \Gamma$. Considering the partition $[n] = H_1 \cup \dots \cup H_r$, we see that r -colored rev-lex complexes are r -colored. While many r -colored complexes are not defined by the special partition above, Frankl, Füredi and Kalai [8] proved that they all share a face vector with an r -colored rev-lex complex. In particular, since r -colored rev-lex complexes are uniquely determined by their face vectors, this result characterizes the possible face vectors of colored complexes in terms of colored rev-lex complexes (a numerical characterization was also given in [8]). This result of Frankl, Füredi and Kalai can be recovered from Theorem 2.1.

Suppose that Δ is an r -colored simplicial complex on $[n]$. Then there exists a monomial ideal I of an r -colored squarefree ring R of type (n, \dots, n) such that the set of monomials of R which are not in I can be identified with Δ . (Since R has nr variables, the ideal I will contain at least $(nr - n)$ variables corresponding to vertices which do not appear in Δ .) Theorem 2.1 shows that there exists a lex ideal L of R having the same Hilbert function as I . The r -colored rev-lex complex having the same face vector as Δ corresponds to the set of monomials of R which are not in L .

Actually, Theorem 2.1 refines this result. Theorem 2.1 characterizes the face vectors of colored simplicial complexes on $[n]$ with a fixed partition $[n] = C_1 \cup \dots \cup C_r$.

For example, our result gives the complete description of face vectors of 2-colored complexes on $\{1, 2, 3, 4, 5, 6\}$ with the specific partition $\{1, 2, 3, 4\} \cup \{5, 6\}$, while the Frankl–Füredi–Kalai theorem does not guarantee this.

3 Betti numbers of lex-plus- Q ideals

In this section, we show that most colored squarefree rings do not admit ideals with maximal Betti numbers over S . As before, let $R = S/Q$ be an r -colored squarefree ring of type (n_1, \dots, n_r) as defined in Definition 1.3. For a finitely generated graded S -module M , the integers $\beta_{i,j}(M) = \dim_K \text{Tor}_i(M, K)_j$ and $\beta_i(M) = \dim_K \text{Tor}_i(M, K)$ are called the *graded Betti numbers* of M and the *total Betti numbers* of M respectively. A monomial ideal I of S is said to be *strongly color-stable* if $ux_{j,k} \in I$ implies $ux_{j,\ell} \in I$ for all $k < \ell$ and for all $j \in [r]$. The next fact easily follows from [1, Theorem 5.4] or [15, Theorem 5.9].

Lemma 3.1 *Let I be a homogeneous-plus- Q ideal. There exists a strongly color-stable ideal J with $J \supset Q$ and $\text{Hilb}(J) = \text{Hilb}(I)$ such that $\beta_{ij}(J) \geq \beta_{ij}(I)$ for all i and j .*

The above lemma shows that, to study Conjecture 1.2(2) for colored squarefree rings, it is enough to consider strongly color-stable ideals.

3.1 2-colored squarefree rings

We first consider 2-colored squarefree rings. Let $n_1 \geq n_2 \geq 4$, $S = K[x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}]$ and $Q = (x_1, \dots, x_{n_1})^2 + (y_1, \dots, y_{n_2})^2$.

Proposition 3.2 *A 2-colored squarefree ring $R = S/Q$ of type (n_1, n_2) with $n_2 \geq 4$ does not admit ideals with maximal Betti numbers over S .*

We first give an example.

Example 3.3 Let $A = K[x_1, \dots, x_4, y_1, \dots, y_4]$ and $P = (x_1, \dots, x_4)^2 + (y_1, \dots, y_4)^2$. Let

$$L = (x_4, x_3, y_4, x_2y_3, x_1y_3) + P,$$

$$I = (x_4, x_3, y_4, x_2y_3, x_2y_2) + P$$

and

$$J = (x_4, x_3, x_2) + P.$$

Then L is lex-plus- P and any strongly color-stable ideal B with $B \supset P$ and $\text{Hilb}(B) = \text{Hilb}(L)$ is isomorphic to L, I , or J . The following are Betti diagrams of these ideals computed by the computer algebra system Macaulay 2 [9]:

$$\begin{aligned} \text{betti}(\text{res}(S/L)) = \text{total: } & 1 \ 14 \ 65 \ 156 \ 224 \ 202 \ 113 \ 36 \ 5 \\ & 0: \ 1 \ 3 \ 3 \ 1 \ . \ . \ . \ . \\ & 1: \ . \ 11 \ 53 \ 107 \ 118 \ 78 \ 32 \ 8 \ 1 \\ & 2: \ . \ . \ 9 \ 48 \ 106 \ 124 \ 81 \ 28 \ 4 \end{aligned}$$

$$\begin{aligned} \text{betti}(\text{res}(S/I)) = \text{total: } & 1 \ 14 \ 66 \ 159 \ 225 \ 196 \ 104 \ 31 \ 4 \\ & 0: \ 1 \ 3 \ 3 \ 1 \ . \ . \ . \ . \\ & 1: \ . \ 11 \ 53 \ 108 \ 120 \ 77 \ 27 \ 4 \ . \\ & 2: \ . \ . \ 10 \ 50 \ 105 \ 119 \ 77 \ 27 \ 4 \end{aligned}$$

$$\begin{aligned} \text{betti}(\text{res}(S/J)) = \text{total: } & 1 \ 14 \ 64 \ 150 \ 209 \ 182 \ 98 \ 30 \ 4 \\ & 0: \ 1 \ 3 \ 3 \ 1 \ . \ . \ . \ . \\ & 1: \ . \ 11 \ 53 \ 106 \ 113 \ 68 \ 22 \ 3 \ . \\ & 2: \ . \ . \ 8 \ 43 \ 96 \ 114 \ 76 \ 27 \ 4 \end{aligned}$$

None of these ideals has maximal graded Betti numbers. Hence, by Lemma 3.1, the ring A/P does not admit ideals with maximal Betti numbers.

Proof of Proposition 3.2 We use the ideals given in Example 3.3. Let L' be the ideal of S defined by

$$L' = LS + (x_5, \dots, x_{n_1}, y_5, \dots, y_{n_2}).$$

Define I' and J' in the same way as L' . Then L' is lex-plus- Q , and every strongly color-stable ideal B with $B \supset Q$ and $\text{Hilb}(B) = \text{Hilb}(L')$ is isomorphic to L', I' , or J' . Recall that, if M is a homogeneous ideal of S and f is a non-zero divisor of S/M , then the tensor product of the minimal free resolutions of S/M and $S/(f)$ is a minimal free resolution of $S/(M + (f))$. By using this fact together with the computations given in Example 3.3, it follows that

$$\beta_1(I') > \beta_1(L') > \beta_1(J')$$

and

$$\beta_{n_1+n_2-1}(L') > \beta_{n_1+n_2-1}(I') = \beta_{n_1+n_2-1}(J').$$

Then, by Lemma 3.1, the ring S/Q does not admit ideals with maximal Betti numbers over S . □

Remark 3.4 While we used a computer system for the computations of Betti diagrams of L, I , and J , one can compute those Betti numbers by using Lemma 3.7 and Hochster’s formula.

3.2 General construction

In the rest of this section, $R = S/Q$ is an r -colored squarefree ring of type (n_1, \dots, n_r) with $r \geq 3$ and $n_r \geq 3$. The goal is to show the following.

Theorem 3.5 *With the same notation as above, R does not admit ideals with maximal Betti numbers over S .*

First, we give a formula to compute the graded Betti numbers of a strongly color-stable ideal I with $I \supset Q$ using simplicial complexes. Let $\tilde{V}_j = \{x_{j,0}\} \cup V_j$, $\tilde{V} = \bigcup_{j=1}^r \tilde{V}_j$ and $\tilde{S} = K[\tilde{V}]$. Let

$$\tilde{Q} = (\{x_{j,k}x_{j,\ell} \in \tilde{S} : k \neq \ell, j = 1, 2, \dots, r\}).$$

For any strongly color-stable ideal $I = I' + Q$ of S , where I' is generated by monomials not in Q , define

$$\tilde{I} = I'\tilde{S} + \tilde{Q}.$$

Thus the ideal \tilde{I} is obtained from $I = I' + Q$ by replacing Q by \tilde{Q} .

Example 3.6 Let $I = (x_{1,2}x_{2,2}, x_{1,2}x_{2,1}) + (x_{1,1}^2, x_{1,1}x_{1,2}, x_{1,2}^2, x_{2,1}^2, x_{2,1}x_{2,2}, x_{2,2}^2)$. Then

$$\tilde{I} = (x_{1,2}x_{2,2}, x_{1,2}x_{2,1}) + (x_{1,0}x_{1,1}, x_{1,0}x_{1,2}, x_{1,1}x_{1,2}, x_{2,0}x_{2,1}, x_{2,0}x_{2,2}, x_{2,1}x_{2,2}).$$

Actually, the map $I \rightarrow \tilde{I}$ is a special case of the colored squarefree operation introduced in [1]. Thus, by [21, Theorem 0.1], we obtain

Lemma 3.7 *Let I be a strongly color-stable ideal of S with $I \supset Q$. Then I and \tilde{I} have the same graded Betti numbers.*

Since \tilde{I} is a squarefree monomial ideal, we can compute its graded Betti numbers using Hochster’s formula. We recall Hochster’s formula. Let J be a squarefree monomial ideal of \tilde{S} , and let $\Delta(J)$ be its Stanley-Reisner complex,

$$\Delta(J) = \{u \in \tilde{S} \setminus J : u \text{ is a squarefree monomial}\}.$$

(We identify squarefree monomials of \tilde{S} with subsets of \tilde{V} and regard $\Delta(J)$ as a simplicial complex on the vertex set \tilde{V} .) Hochster’s formula [11] says

$$\beta_{i,j}(J) = \sum_{W \subset \tilde{V}, |W|=j} \dim_K \tilde{H}_{j-i-2}(\Delta(J)_W; K),$$

where $\Delta(J)_W = \Delta(J) \cap K[W]$ and where $\tilde{H}_i(\Delta; K)$ is the i -th reduced homology group of a simplicial complex Δ over a field K .

A simplicial complex Δ on the vertex set \tilde{V} is said to be *colored shifted* if $ux_{j,k} \in \Delta$ implies $ux_{j,\ell} \in \Delta$ for any $j \in [r]$ and for any $0 \leq \ell < k$ such that $x_{j,\ell}$ does not divide u . The top Betti numbers of colored shifted complexes can be computed as follows (see [1, Theorem 5.7] and [21, Proposition 4.2]).

Lemma 3.8 (Babson–Novik) *If Δ is a colored shifted simplicial complex on \tilde{V} then*

$$\dim_K \tilde{H}_{r-1}(\Delta; K) = |\{u \in \Delta : \deg u = r, u \text{ is not divisible by any of } x_{j,0}\}|.$$

Let I be a strongly color-stable ideal of S . Define

$$\Gamma(I) = \{u \in S \setminus I : u \text{ is a squarefree monomial with } \deg u = r\}.$$

Lemma 3.9 *If I is a strongly color-stable ideal of S with $I \supset Q$, then*

$$\beta_{r-1,2r}(I) = \sum_{x_{1,i_1}x_{2,i_2}\cdots x_{r,i_r} \in \Gamma(I)} \left(\prod_{k=1}^r i_k \right).$$

Proof Note that $\Gamma(I)$ is the set of monomials u in $\Delta(\tilde{I})$ of degree r which are not divisible by any of the $x_{j,0}$. Since I is strongly color-stable, the simplicial complex $\Delta(\tilde{I})$ is colored shifted. Note also that, for any $W \subset \tilde{V}$, $\Delta(\tilde{I})_W$ is colored shifted on W . Hence, for any $W \subset \tilde{V}$ with $|W| = 2r$, Lemma 3.8 implies

$$\dim_K \tilde{H}_{r-1}(\Delta(\tilde{I})_W; K) = \begin{cases} 1, & \text{if } |W \cap V_j| = 2 \text{ for all } j \text{ and } \Delta(\tilde{I})_W \text{ contains all} \\ & \text{monomials of the form } x_{1,i_1} \cdots x_{r,i_r} \in K[W], \\ 0, & \text{otherwise.} \end{cases}$$

(Observe that, if $W \cap V_j = \{x_{j,k}, x_{j,\ell}\}$ with $k < \ell$, then $x_{j,k}$ plays the role of $x_{j,0}$ in Lemma 3.8.)

By Hochster’s formula,

$$\begin{aligned} \beta_{r-1,2r}(\tilde{I}) &= \sum_{W \subset \tilde{V}, |W|=2r} \dim_K \tilde{H}_{r-1}(\Delta(\tilde{I})_W; K) \\ &= \left| \left\{ W = \bigcup_{j=1}^r \{x_{j,p_j}, x_{j,q_j}\} \subset \tilde{V} : p_j < q_j, \prod_{j=1}^r x_{j,q_j} \in \Delta(\tilde{I}) \right\} \right| \\ &= \sum_{x_{1,q_1}x_{2,q_2}\cdots x_{r,q_r} \in \Gamma(I)} \left(\prod_{k=1}^r q_k \right). \end{aligned}$$

The last equality follows since I is strongly color-stable. Then the statement follows from Lemma 3.7. □

We need the following fact.

Lemma 3.10 *Let $a_1 \geq \cdots \geq a_t$ and $b_1 \geq \cdots \geq b_t$ be two sequences of integers. If $a_k + \cdots + a_t \geq b_k + \cdots + b_t$ for all k then $a_1 \cdots a_t \geq b_1 \cdots b_t$.*

Proof We induct on t . If $t = 1$ then there is nothing to prove. Suppose $t > 1$. If $a_1 \geq b_1$ then the statement immediately follows by induction. Suppose $a_1 < b_1$. Note that $a_2 + \cdots + a_t > b_2 + \cdots + b_t$ by the assumption. Let $p \geq 2$ be the greatest integer such that $a_p = a_2$. We claim that $a_k + \cdots + a_t > b_k + \cdots + b_t$ for $2 \leq k \leq p$.

Suppose to the contrary that $a_k + \cdots + a_t = b_k + \cdots + b_t$ for some $k, 2 < k \leq p$. Since $a_{k+1} + \cdots + a_t \geq b_{k+1} + \cdots + b_t$, we have

$$a_2 = \cdots = a_k \leq b_k \leq \cdots \leq b_2.$$

Then $\sum_{j=2}^t a_j = \sum_{j=2}^{k-1} a_j + \sum_{j=k}^t a_j \leq \sum_{j=2}^t b_j$, a contradiction.

Thus, the sequence $(a_1 + 1, a_2, \dots, a_{p-1}, a_p - 1, a_{p+1}, \dots, a_r)$ also satisfies the assumption of the lemma, and, since $a_1 \geq a_p$,

$$a_1 a_p \left(\prod_{j \neq 1, p} a_j \right) \geq (a_1 + 1)(a_p - 1) \left(\prod_{j \neq 1, p} a_j \right).$$

By repeating this procedure, we may assume $a_1 = b_1$. The statement follows by induction. □

Proof of Theorem 3.5 Let

$$\alpha = x_{1,n_1} x_{2,n_2} \cdots x_{r-2,n_{r-2}}.$$

[Case 1]: Suppose $n_{r-1} = n_r$. To simplify, set $n = n_{r-1} = n_r$. Let

$$I = (\alpha x_{r-1,n}, \alpha x_{r-1,n-1}) + Q$$

and

$$L = (\alpha x_{r-1,n}, \alpha x_{r,n}, \alpha x_{r-1,n-1} x_{r,n-1}) + Q.$$

Note that I and L are strongly color-stable ideals having the same Hilbert function. First, we show that the graded Betti numbers of I and L are incomparable.

Clearly, $\beta_0(I) < \beta_0(L)$. It is enough to show that $\beta_{r-1,2r}(I) > \beta_{r-1,2r}(L)$. Since

$$\Gamma(I) \setminus \Gamma(L) = \{\alpha x_{r,n} x_{r-1,\ell} : \ell = 1, 2, \dots, n - 2\}$$

and

$$\Gamma(L) \setminus \Gamma(I) = \{\alpha x_{r-1,n-1} x_{r,\ell} : \ell = 1, 2, \dots, n - 2\},$$

it follows from Lemma 3.9 that

$$\begin{aligned} & \beta_{r-1,2r}(I) - \beta_{r-1,2r}(L) \\ &= \sum_{x_{1,i_1} x_{2,i_2} \cdots x_{r,i_r} \in \Gamma(I) \setminus \Gamma(L)} \binom{r}{k=1}^{i_k} - \sum_{x_{1,i_1} x_{2,i_2} \cdots x_{r,i_r} \in \Gamma(L) \setminus \Gamma(I)} \binom{r}{k=1}^{i_k} \\ &= \left(\prod_{j=1}^{r-2} n_j \right) \{n - (n - 1)\} (1 + 2 + \cdots + (n - 2)) > 0. \end{aligned}$$

Since L is lex-plus- Q , any monomial ideal $J \supset Q$ with $\text{Hilb}(J) = \text{Hilb}(L)$ satisfies $\beta_0(J) \leq \beta_0(L)$. Thus, to complete the proof, it is enough to show that, if J is any homogeneous-plus- Q ideal which has the same Hilbert function as L and satisfies $\beta_0(J) = \beta_0(L)$, then $\beta_{r-1,2r}(J) \leq \beta_{r-1,2r}(L)$.

We may assume that J is strongly color-stable. Since J has the same Hilbert function as L , J contains two monomials u_1 and u_2 of degree $r - 1$ which are not in Q . Let $\{c_k\} = [r] \setminus \text{color}(u_k)$. Then $|\text{Shad}(\{u_1, u_2\})| \geq n_{c_1} + n_{c_2} - 1$. Since J has a generator

of degree r and has the same Hilbert function as L , $|\text{Shad}(\{u_1, u_2\})|$ must be equal to $2n - 1$. It follows that $n_{c_1} = n_{c_2} = n$, and $c_1 \neq c_2$. Thus $(u_1, u_2) + Q$ is isomorphic to $(\alpha x_{r-1,n}, \alpha x_{r,n}) + Q$. Hence we may assume $u_1 = \alpha x_{r-1,n}$ and $u_2 = \alpha x_{r,n}$.

Let v be a generator of J of degree r . Then $\Gamma(J) \setminus \Gamma(L) = \{\alpha x_{r-1,n-1} x_{r,n-1}\}$ and $\Gamma(L) \setminus \Gamma(J) = \{v\}$. Also, since $J \neq L$ is strongly color-stable, v is a monomial of the form $v = x_{k,n_k-1} (\prod_{j \neq k} x_{j,n_j})$ for some $1 \leq k \leq r - 2$. Then, by Lemma 3.9, we have

$$\beta_{r-1,2r}(L) - \beta_{r-1,2r}(J) \geq \left(\prod_{j \neq k, r-1, r} n_j \right) \{ (n_k - 1)n^2 - n_k(n - 1)^2 \} \geq 0$$

as desired.

[Case 2]: Suppose $n_{r-1} > n_r$. Let

$$I = \alpha(x_{r-1,n_{r-1}}, \dots, x_{r-1,n_r+2}, x_{r,n_r}) + Q$$

and

$$L = \alpha(x_{r-1,n_{r-1}}, \dots, x_{r-1,n_r+2}, x_{r-1,n_r+1}, x_{r-1,n_r} x_{r,n_r}) + Q.$$

Then I and L are strongly color-stable ideals having the same Hilbert function. First, we show that the graded Betti numbers of I and those of L are incomparable. Clearly, $\beta_0(I) < \beta_0(L)$. On the other hand, in the same way as in [Case 1], we have $\beta_{r-1,2r}(I) > \beta_{r-1,2r}(L)$ since

$$\Gamma(I) \setminus \Gamma(L) = \{\alpha x_{r-1,n_r+1} x_{r,\ell} : \ell = 1, 2, \dots, n_r - 1\}$$

and

$$\Gamma(L) \setminus \Gamma(I) = \{\alpha x_{r,n_r} x_{r-1,\ell} : \ell = 1, 2, \dots, n_r - 1\}.$$

Since L is lex-plus- Q , to complete the proof, it is enough to show that, for any strongly color-stable ideal $J \supset Q$ having the same Hilbert function as L , if $\beta_0(J) = \beta_0(L)$ then $\beta_{r-1,2r}(J) \leq \beta_{r-1,2r}(L)$.

Let $t = n_{r-1} - n_r$. Then J has generators u_1, \dots, u_t of degree $r - 1$ which are not in Q . First, we claim that $\text{color}(u_k) = [r - 1]$ for all k .

Let $\{c_k\} = [r] \setminus \text{color}(u_k)$. Suppose $c_1 = \dots = c_s = r$ and $c_k \neq r$ for $k > s$. Then

$$|\text{Shad}(\{u_1, \dots, u_t\})| \geq sn_r + (n_{c_{s+1}} - s) + (n_{c_{s+2}} - s - 1) + \dots + (n_{c_t} - t + 1).$$

However, since J has a generator of degree r and has the same Hilbert function as L , $|\text{Shad}(\{u_1, \dots, u_t\})|$ must be equal to tn_r . Since $n_{c_k} - (k - 1) \geq n_{r-1} - t + 1 > n_r$ for $k = s + 1, \dots, t$, it follows that $s = t$ and $\text{color}(u_k) = [r - 1]$ for all k .

Let $v = x_{1,j_1} \cdots x_{r,j_r}$ be the generator of J with $\text{deg } v = r$. Write

$$u_k = x_{1,i_{1,k}} \cdots x_{r-1,i_{r-1,k}}.$$

Since J is strongly color-stable, we may assume that

$$i_{1,k} + \dots + i_{r-1,k} \geq n_1 + \dots + n_{r-1} - (k - 1) \quad \text{for } k = 1, 2, \dots, t \quad (3)$$

and

$$j_1 + \dots + j_{r-1} \geq n_1 + \dots + n_{r-1} - t. \tag{4}$$

Also, since J is strongly color-stable, we have

$$j_r = n_r.$$

Now, by Lemma 3.9, we have

$$\begin{aligned} &\beta_{r-1,2r}(L) - \beta_{r-1,2r}(J) \\ &= \left[\sum_{x_1, k_1 \dots x_r, k_r \in \Gamma(L)} k_1 \dots k_r \right] - \left[\sum_{x_1, k_1 \dots x_r, k_r \in \Gamma(J)} k_1 \dots k_r \right] \\ &= \left[\sum_{k=1}^t i_{1,k} \dots i_{r-1,k} (1 + \dots + n_r) \right] + j_1 \dots j_{r-1} n_r \\ &\quad - \left[\sum_{k=1}^t n_1 \dots n_{r-2} (n_{r-1} - k + 1) (1 + \dots + n_r) \right] - n_1 \dots n_{r-2} (n_{r-1} - t) n_r. \end{aligned}$$

By (3), the sequences $(i_{1,k}, \dots, i_{r-1,k})$ and $(n_1, \dots, n_{r-2}, n_{r-1} - k + 1)$ satisfy the conditions of Lemma 3.10, and, by (4), so do (j_1, \dots, j_{r-1}) and $(n_1, \dots, n_{r-2}, n_{r-1} - t)$. Then the desired inequality $\beta_{r-1,2r}(L) \geq \beta_{r-1,2r}(J)$ follows from Lemma 3.10. \square

Example 3.11 Let $A = K[x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3]$ and $Q = (x_1, x_2, x_3)^2 + (y_1, y_2, y_3)^2 + (z_1, z_2, z_3)^2$. Let

$$L = (x_3y_3, x_3z_3, x_3y_2z_2) + Q$$

and

$$I = (x_3y_3, x_3y_2) + Q.$$

Then, L is the lex-plus- Q ideal having the same Hilbert function as I . The following are betti diagrams of these ideals computed by Macaulay 2.

```
betti(res(S/L)) = total: 1 21 142 490 1004 1305 1090 566 166 21
0: 1 . . . . . . . . . .
1: . 20 37 24 7 1 . . . .
2: . 1 105 343 460 321 122 24 2 .
3: . . . 123 537 983 968 542 164 21
```

```
betti(res(S/I)) = total: 1 20 141 493 1016 1324 1105 572 167 21
0: 1 . . . . . . . . . .
1: . 20 36 24 7 1 . . . .
2: . . 105 343 463 330 132 29 3 .
3: . . . 126 546 993 973 543 164 21
```

4 Betti numbers of Borel ideals over a colored squarefree ring

In the previous section, we saw that most colored squarefree rings do not satisfy Conjecture 1.2(2). On the other hand, we are not sure whether Conjecture 1.2(1) fails for those rings. Indeed, the ideals used in the proof of Theorem 3.5 do not give a counterexample to Conjecture 1.2(1). The purpose of this section is to give a way to compute the graded Betti numbers of lex ideals of a colored squarefree ring, which will be the first step in studying Conjecture 1.2(1) for colored squarefree rings.

Let $R = S/Q$ be an r -colored squarefree ring as defined in Definition 1.3. For a finitely generated graded R -module M , let $\beta_{i,j}^R(M)$ and $\beta_i^R(M)$ be the graded Betti numbers of M and the total Betti numbers of M over R respectively. A homogeneous ideal I of R has a *linear resolution* if there exists an integer d such that $\beta_{i,i+j}^R(I) = 0$ for all i if $j \neq d$.

Lemma 4.1 *Let $A = K[x_1, \dots, x_n]$, $P = (x_1, \dots, x_n)^2$ and $B = A/P$. Then the ideal $I = (x_1, \dots, x_p)$ of B has a linear resolution and $\beta_i^B(I) = pn^i$ for all i .*

Proof It is clear that the first syzygy module of the ideal (x_1) of B is (x_1, \dots, x_n) . Since $I = (x_1) \oplus_B (x_2) \oplus_B \dots \oplus_B (x_p)$, the first syzygy module of I is isomorphic to $(x_1, \dots, x_n) \oplus_B \dots \oplus_B (x_1, \dots, x_n) \subset B^p$. Then I has a linear resolution and $\beta_{i+1}(I) = n\beta_i(I)$ for all $i \geq 0$. □

For a subset $W \subset V$, write I_W for the ideal of R generated by W .

Lemma 4.2 *Let $W \subset V$ and $p_j = |W \cap V_j|$. Then I_W has a linear resolution and*

$$\beta_i^R(I_W) = \sum_{i_1 + \dots + i_r = i} (n_1^{i_1} \dots n_r^{i_r}) \left(\prod_{i_k \neq 0} p_k \right) \text{ for all } i.$$

Proof Let $I_{\{k\}}$ be the ideal of $R_{\{k\}} = K[V_k]/(x_{k,1}, \dots, x_{k,n_k})^2$ generated by $W \cap V_k$. Let $\mathbf{F}_{\{k\}}$ be the minimal free resolution of $R_{\{k\}}/I_{\{k\}}$ over $R_{\{k\}}$. Since

$$R = R_{\{1\}} \otimes_K R_{\{2\}} \otimes_K \dots \otimes_K R_{\{r\}}$$

and

$$R/I_W = (R_{\{1\}}/I_{\{1\}}) \otimes_K (R_{\{2\}}/I_{\{2\}}) \otimes_K \dots \otimes_K (R_{\{r\}}/I_{\{r\}}),$$

by the Künneth tensor formula, $\mathbf{F}_{\{1\}} \otimes_K \dots \otimes_K \mathbf{F}_{\{r\}}$ is the minimal free resolution of R/I_W over R . This fact and Lemma 4.1 prove the desired formula. □

Definition 4.3 For monomials $u = y_1 \dots y_t$ and $v = z_1 \dots z_t$ of R of the same degree, where $y_1 > \dots > y_t$ and $z_1 > \dots > z_t$ are variables, write $u \geq_P v$ if $y_k \geq z_k$ for all k . A monomial ideal I of R is said to be *Borel* if, for all monomials $u, v \in R$, $u \in I$ and $v >_P u$ implies $v \in I$.

Note that lex ideals are Borel. For all monomials $u, v \in R$, write $u \succ v$ if $\deg u < \deg v$ or $\deg u = \deg v$ and $u \succ_{\text{lex}} v$. Let $G(I)$ be the set of minimal monomial generators of a monomial ideal $I \subset R$.

Lemma 4.4 *Let $I \subset R$ be a Borel ideal with $G(I) = \{u_1, \dots, u_t\}$, where $u_1 \succ \dots \succ u_t$. Then, for $k = 1, 2, \dots, t$,*

$$(u_1, \dots, u_{k-1} : u_k) = \{y \in V : y \succ \text{last}(u_k) \text{ or } \text{color}(y) \subset \text{color}(u_k)\},$$

where $(u_1, \dots, u_{k-1}) = 0$ if $k = 1$.

Proof Since I is Borel, it is clear that the left-hand side contains the right-hand side. We show that the right-hand side contains the left-hand side.

Let $fu_k \in (u_1, \dots, u_{k-1})$ be a monomial and $d = \deg u_k$. If $fu_k = 0$ then there exists a monomial $y \in V_a$ with $a \in \text{color}(u_k)$ such that y divides f . Suppose $fu_k \neq 0$. Write $fu_k = y_1 \cdots y_s$, where $y_1 \succ \dots \succ y_s$ are variables. Since (u_1, \dots, u_{k-1}) is Borel, there exists an integer $1 \leq \delta \leq d$ such that $y_1 \cdots y_\delta \in G(I)$. Let $u_p = y_1 \cdots y_\delta$. Then $\text{last}(u_p) \succ \text{last}(u_k)$ and u_p does not divide u_k . Hence there exists y_ℓ which divides u_p but does not divide u_k . This y_ℓ must divide f , and $y_\ell \succ \text{last}(u_k)$. \square

For any monomial $m \in R$, let

$$W(m) = \{y \in V : y \prec \text{last}(m) \text{ and } \text{color}(y) \not\subset \text{color}(m)\}$$

and

$$T(m) = V \setminus W(m) = \{y \in V : y \succ \text{last}(m) \text{ or } \text{color}(y) \subset \text{color}(m)\}.$$

Proposition 4.5 *Let $I \subset R$ be Borel. Then*

$$\beta_{i,i+j}^R(I) = \sum_{u \in G(I), \deg(u)=j} \beta_i^R(R/I_{T(u)}) \quad \text{for all } i, j.$$

Proof The proof is the same as that of [10, Lemma 1.5]. Let $G(I) = \{u_1, \dots, u_t\}$ with $u_1 \succ \dots \succ u_t$. We use induction on t . If $t = 1$ then the first syzygy module of $I = (u_1)$ is $(0 : u_1) = I_{T(u_1)}$, and therefore the statement follows from Lemma 4.2.

Suppose $t > 1$. Let $J = (u_1, \dots, u_{t-1})$ and $d = \deg u_t$. Note that J is also Borel. Consider the short exact sequence

$$0 \longrightarrow R/(J : u_t)(-d) \xrightarrow{\times u_t} R/J \longrightarrow R/I \longrightarrow 0. \tag{5}$$

By Lemma 4.4, $(J : u_t) = I_{T(u_t)}$. Let \mathbf{G} be the minimal free resolution of R/J over R and \mathbf{F} the minimal free resolution of $R/(J : u_t)(-d) = R/I_{T(u_t)}(-d)$. One has $\mathbf{G}_i = \bigoplus_{j \in \mathbb{N}} R(-j)^{\beta_{i,j}^R(R/J)}$. By induction, $\max\{j : \beta_{i,j}^R(R/J) \neq 0\} \leq i + d - 1$. On the other hand, by Lemma 4.2 $\mathbf{F}_i = R(-d - i)^{\beta_i^R(R/I_{T(u_t)})}$. Hence, R/I is minimally resolved by the mapping cone arising from the map $R/(J : u_t)(-d) \xrightarrow{\times u_t} R/J$. The statement follows. \square

Corollary 4.6 *The graded Betti numbers of Borel ideals of R do not depend on the characteristic of the base field K .*

Remark 4.7 The key idea in the proof of Proposition 4.5 is that the colon ideal $((u_1, \dots, u_{k-1}) : u_k)$ has a 1-linear resolution. This property also holds for Borel ideals in the quotient ring S/M , whenever M is an ideal generated by monomials of degree 2. Thus, the Betti numbers of Borel ideals of such rings can be computed in the same way as in the proof of Proposition 4.5.

One might think that the graded Betti numbers of a Borel ideal $I = (u_1, \dots, u_t)$ depend on the shape of $T(u_j)$. We will show that they only depend on $\text{grow}(u_j) = |W(u_j)|$. For an integer $0 \leq a \leq |\bigcup_{j=d+1}^r V_j|$, let $W(a, d)$ be the set of lex-smallest variables in $\bigcup_{j=d+1}^r V_j$ with $|W(a, d)| = a$. Set $T(a, d) = V \setminus W(a, d)$.

Lemma 4.8 *Let $m \in R$ be a monomial of degree d with $\text{grow}(m) = a$. Then $a \leq |\bigcup_{j=d+1}^r V_j|$ and $I_{T(m)}$ has the same graded Betti numbers as $I_{T(a,d)}$.*

Proof For any subset $W \subset V$, let $R[W] = K[W]/(Q \cap K[W]) \cong R/I_{V \setminus W}$. By Lemma 4.2, we must prove that there exists a subset \tilde{W} such that $R[\tilde{W}]$ has the same Hilbert function as $R[W(m)]$ and \tilde{W} is a set of lex-smallest variables in $\bigcup_{j=d+1}^r V_j$.

For a subset $W \subset V$, let $c_i(W) = |W \cap V_i|$ for $i = 1, 2, \dots, r$ and $\mathbf{c}(W) = (c_1(W), \dots, c_r(W))$. Note that the Hilbert function of $R[W]$ only depends on the vector $\mathbf{c}(W)$. Let $e_i = |W(m) \cap V_i|$ for $i = 1, 2, \dots, r$ and $\mathbf{e} = (e_1, \dots, e_r)$. Let $\text{last}(m) = x_{s,t}$ and $q = \max\{j : n_j \geq t\}$. Then, by (1) (in the proof of Lemma 2.10), the vector \mathbf{e} can be written in the form

$$\mathbf{e} = (e_1, \dots, e_q, n_{q+1}, \dots, n_r)$$

and each e_k is either 0, $t - 1$ or t for $k = 1, 2, \dots, q$. Consider the vector of the form

$$\tilde{\mathbf{e}} = (\tilde{e}_1, \dots, \tilde{e}_r) = (0, \dots, 0, t - 1, \dots, t - 1, t, \dots, t, n_{q+1}, \dots, n_r)$$

which is obtained by a permutation of entries of \mathbf{e} . Let $\tilde{W} = \bigcup_{i=1}^r \{x_{i,j} : 1 \leq j \leq \tilde{e}_i\}$. Then $\mathbf{c}(\tilde{W}) = \tilde{\mathbf{e}}$. By the construction of $\tilde{\mathbf{e}}$, it follows that $R[\tilde{W}]$ and $R[W(m)]$ have the same Hilbert function. Also, by the formula of $\tilde{\mathbf{e}}$, it is clear that \tilde{W} is a set of lex-smallest variables in $\bigcup_{j=d+1}^r V_j$. \square

By Proposition 4.5 and Lemma 4.8, we get

Theorem 4.9 *Let $I = (u_1, \dots, u_t)$ be a Borel ideal of R with $\text{grow}(u_k) = a_k$. Then*

$$\beta_{i,i+j}^R(I) = \sum_{u_k \in G(I), \text{deg } u_k = j} \beta_i^R(R/I_{T(a_k,j)}) \quad \text{for all } i, j.$$

Example 4.10 Let I, L and $R = S/Q$ be as in Example 3.11. Let $\tilde{I} = (x_3y_3, x_3y_2)$ and $\tilde{L} = (x_3y_3, x_3z_3, x_3y_2z_2)$ be ideals of R . Since L is lex-plus- Q , \tilde{L} is lex. Also $W(x_3y_3) = \{z_1, z_2, z_3\}$, $W(x_3z_3) = \{y_1, y_2\}$ and $W(x_3y_2z_2) = \emptyset$. Hence, by Proposition 4.5,

$$\begin{aligned} \beta_{i,i+2}^R(\tilde{L}) &= \beta_i^R(R/(x_1, x_2, x_3, y_1, y_2, y_3)) + \beta_i^R(R/(x_1, x_2, x_3, y_3, z_1, z_2, z_3)) \\ &= (i + 1)3^i + \left\{ (i + 1)3^i + \binom{i + 1}{2}3^{i-1} \right\} \end{aligned}$$

and

$$\beta_{i,i+3}^R(\tilde{L}) = \beta_i^R(R/(x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3)) = \binom{i + 2}{2}3^i.$$

Note that the ideal \tilde{I} has a linear resolution and $\beta_{i,i+2}^R(\tilde{I}) = 2(i + 1)3^i$ for all i .

5 Concluding remarks and open problems

5.1 Betti numbers of lex ideals

As we show in Section 3, Conjecture 1.2(2) is false. However Conjecture 1.2(1) is still open even for colored squarefree rings.

Problem 5.1 Do lex ideals of a colored squarefree ring R have the greatest graded Betti numbers among all homogeneous ideals of R having the same Hilbert function?

While we give a combinatorial way to compute the graded Betti numbers of Borel ideals of colored squarefree rings, we do not know whether lex ideals have the greatest graded Betti numbers among all Borel ideals having the same Hilbert function. The only known case is

Proposition 5.2 *Lex ideals of a 2-colored squarefree ring R have the greatest graded Betti numbers among all homogeneous ideals of R having the same Hilbert function.*

Proof Let $I = (u_1, \dots, u_s, v_1, \dots, v_t)$ be a monomial ideal of R with $\deg u_k = 1$ and $\deg v_\ell = 2$. Then $I = (u_1, \dots, u_s) \oplus_R (v_1) \oplus_R \dots \oplus_R (v_t)$. Then, by Lemma 4.2,

$$\beta_{i,i+2}^R(I) = \sum_{j=1}^t \beta_{i,i+2}^R((v_j)) = t\beta_i^R(R/\mathfrak{m}) \quad \text{for all } i,$$

where \mathfrak{m} is the maximal ideal of R , and $\beta_{i,i+j}^R(I) = 0$ if $j \neq 1, 2$.

Let $L = (u'_1, \dots, u'_s, v'_1, \dots, v'_t)$ be the lex ideal of R having the same Hilbert function as I , where $\deg u'_k = 1$ and $\deg v'_\ell = 2$. Since L is lex, $t' \geq t$. Hence $\beta_{i,i+2}^R(L) = t'\beta_i^R(R/\mathfrak{m}) \geq t\beta_i^R(R/\mathfrak{m}) = \beta_{i,i+2}^R(I)$ for all i . Since I and L have the same Hilbert

function, we have $\sum_i (-1)^i \beta_{i,j}^R(I) = \sum_i (-1)^i \beta_{i,j}^R(L)$ for all j . Since $\beta_{i,i+j}^R(I) = \beta_{i,i+j}^R(L) = 0$ if $j \neq 1, 2$, it follows that

$$\beta_{j-1,j}^R(L) - \beta_{j-2,j}^R(L) = \beta_{j-1,j}^R(I) - \beta_{j-2,j}^R(I).$$

Thus we have $\beta_{i,i+1}^R(L) \geq \beta_{i,i+1}^R(I)$ for all i as desired. □

5.2 Betti numbers of Stanley–Reisner ideals

One may ask the following question which is similar to Problem 5.1: Does there always exist an r -colored complex whose Stanley–Reisner ideal has the greatest graded Betti numbers among all r -colored complexes for a fixed face vector?

Unfortunately, the answer is no. Indeed, 2-colored complexes are bipartite graphs, and if we consider bipartite graphs with 10 edges and 7 vertices (there are only 4 such bipartite graphs up to a permutation of the vertices), then computer computations show that none of them have the greatest graded Betti numbers.

5.3 h -vectors of balanced Cohen–Macaulay complexes

A further generalization of Theorem 2.1 is the next problem.

Problem 5.3 Let $S = K[V]$ and $Q = \sum_{j=1}^r (x_{j,1}, \dots, x_{j,n_j})^{a_j}$, where a_1, \dots, a_r are positive integers. Characterize the Hilbert functions of homogeneous ideals of S/Q .

Unfortunately, the above ring S/Q is generally not Macaulay-Lex with respect to any order of the variables. However, Problem 5.3 would be interesting since it yields the complete description of h -vectors of balanced Cohen–Macaulay complexes (see [3, 24]).

Acknowledgements We would like to thank Ryota Okazaki for useful discussions regarding Section 4. The results and examples presented in this paper have been inspired by computations performed by the computer algebra system CoCoA [5] and Macaulay 2 [9].

References

1. Babson, E., Novik, I.: Face numbers and nongeneric initial ideals. *Electron. J. Combin.* **11** (2004/2006). Research Paper 25, 23 pp.
2. Bigatti, A.: Upper bounds for the Betti numbers of a given Hilbert function. *Comm. Algebra* **21**, 2317–2334 (1993)
3. Björner, A., Frankl, P., Stanley, R.P.: The number of faces of balanced Cohen–Macaulay complexes and a generalized Macaulay theorem. *Combinatorica* **7**, 23–34 (1987)
4. Clements, G.F., Lindström, B.: A generalization of a combinatorial theorem of Macaulay. *J. Combinatorial Theory* **7**, 230–238 (1969)
5. CoCoA Team: CoCoA: a system for doing Computations in Commutative Algebra. Available at <http://cocoa.dima.unige.it>
6. Eisenbud, D.: *Commutative algebra with a view toward algebraic geometry*. Grad. Texts in Math., vol. 150. Springer, New York (1995)
7. Francisco, C.A., Richert, B.P.: Lex-plus-powers ideals. In: Peeva, I. (ed.) *Syzygies and Hilbert functions*. Lect. Notes Pure Appl. Math., vol. 254, pp. 113–144. CRC Press, Boca Raton (2007)

8. Frankl, P., Füredi, Z., Kalai, G.: Shadows of colored complexes. *Math. Scand.* **63**, 169–178 (1988)
9. Grayson, D., Stillman, M.: Macaulay 2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>
10. Herzog, J., Takayama, Y.: Resolutions by mapping cones. *Homology Homotopy Appl.* **4**, 277–294 (2002)
11. Hochster, M.: Cohen-Macaulay rings, combinatorics, and simplicial complexes. In: McDonald, B.R., Morris, R. (eds.) *Ring theory, II, Proc. Second Conf., Univ. Oklahoma, Norman, Oklahoma. Lecture Notes in Pure and Appl. Math.*, vol. 26, pp. 171–223. Dekker, New York (1977)
12. Hulett, H.: Maximum Betti numbers of homogeneous ideals with a given Hilbert function. *Comm. Algebra* **21**, 2335–2350 (1993)
13. London, E.: A new proof of the colored Kruskal-Katona theorem. *Discrete Math.* **126**, 217–223 (1994)
14. Macaulay, F.S.: Some properties of enumeration in the theory of modular systems. *Proc. London Math. Soc.* **26**, 531–555 (1927)
15. Mermin, J.: Compressed ideals. *Bull. London Math. Soc.* **40**, 77–87 (2008)
16. Mermin, J., Murai, S.: The lex-plus-powers conjecture holds for pure powers (2008, submitted)
17. Mermin, J., Peeva, I.: Lexifying ideals. *Math. Res. Letters* **13**, 409–422 (2006)
18. Mermin, J., Peeva, I.: Hilbert functions and lex ideals. *J. Algebra* **313**, 642–656 (2007)
19. Mermin, J., Peeva, I., Stillman, M.: Ideals containing the squares of the variables. *Adv. Math.* **217**, 2206–2230 (2008)
20. Murai, S.: Borel-plus-powers monomial ideals. *J. Pure Appl. Algebra* **212**, 1321–1336 (2008)
21. Murai, S.: Betti numbers of strongly color-stable ideals and squarefree strongly color-stable ideals. *J. Algebraic Combin.* **27**, 383–398 (2008)
22. Murai, S., Peeva, I.: Hilbert schemes and Betti numbers over a Clements–Lindström ring (2008, submitted)
23. Pardue, K.: Deformation classes of graded modules and maximal Betti numbers. *Illinois J. Math.* **40**, 564–585 (1996)
24. Stanley, R.P.: Balanced Cohen–Macaulay complexes. *Trans. Amer. Math. Soc.* **249**, 139–157 (1979)