Journal of Convex Analysis
Volume 1 (1994), No. 2, 143–150

Structure of Efficient Sets for Strictly Quasi Convex Objectives

C. Malivert, N. Boissard

LACO (URA 1586) - Département de Mathématiques, Faculté des Sciences,
123 Avenue Albert Thomas, F-87060 Limoges Cedex, France.
e-mail: malivert@cict.fr, boissard@unilim.fr

Received 7 February 1994
Revised manuscript received 13 October 1994

This paper studies the weak efficient set (*WEff P*) of a minimization problem P with k objectives defined on a convex set X of $\mathbb{R}^n$. These objectives are continuous and belong to the class of so-called strictly quasiconvex functions, which contains, in particular, convex as well as linear fractional functions. When k is greater than n, it is of interest to replace the original problem by several subproblems, having at most n objectives. We show that if *WEff P* is bounded, the knowledge of the efficient sets of such subproblems, completely determines *WEff P*.

Keywords : Multiple objectives, Efficient sets, Quasiconvexity, Fractional programming.

1991 Mathematics Subject Classification: 90C31 90C32

1. Introduction

We consider the multiobjective problem

$$(P) \qquad \min_{x \in X} (f_1, \cdots, f_k)$$

where X is a closed convex set in $\mathbb{R}^n$, $f_i : X \longrightarrow \mathbb{R}$, $i = 1, \cdots, k$ and we focus our attention on the structure of the weak efficient set *WEff P*. In [7], it is shown that if the objectives are continuous and convex, the set *WEff P* is determinable from the spatial structure of the efficient sets of subproblems having at most n objectives. The purpose of this paper is to extend these results to a class of non convex objectives. Namely, we consider the class of strictly quasiconvex functions, which has been previously introduced in [2]. This class contains in particular convex functions as well as linear fractional functions, which gives a wide range of applications. See for instance [5][6] for many examples and an extensive bibliography on fractional programming in the scalar case. The paper is divided into two sections. In section 2 we state the problem and give definitions used in the sequel. In section 3 we extend several results, known in the convex case, to the strictly quasiconvex case. Then we deduce Theorem 3.7, which is the main result of this paper.

ISSN 0944-6532 / $ 2.50 © Heldermann Verlag

2. Basic concepts

Recall that X being a convex set in $\mathbb{R}^n$, a functional $f : X \longrightarrow \mathbb{R}$ is quasiconvex on X iff for all points x, y in X and $\lambda \in [0,1]$

$$f(\lambda x + (1-\lambda)y) \leq \max(f(x), f(y)).$$

This class has been extensively studied in the literature, see for instance [1] for many examples, properties and a bibliography.

We say that $f : X \longrightarrow \mathbb{R}$ is strictly quasiconvex [2] iff for every x, y in X and $\lambda \in]0,1[$ one has :

$$f(\lambda x + (1-\lambda)y) < max(f(x), f(y)) \text{ if } f(x) \neq f(y)$$

and

$$f(\lambda x + (1-\lambda)y) \leq f(x) \text{ if } f(x) = f(y)\cdot$$

In particular, linear fractional functions of the form $f(x) = \dfrac{a.x + s}{b.x + t}$ where $a, b \in \mathbb{R}^n$ and $s, t \in \mathbb{R}$, are strictly quasiconvex on every convex set X contained in their domain. (Here . stands for the scalar product in $\mathbb{R}^n$)

Convexity implies strict quasiconvexity and strict quasiconvexity implies quasiconvexity.

As an immediate consequence of the definition, it is easy to prove that if $x = \sum_{i=1}^{l} \lambda_i x_i,\ \lambda_i > 0,\ i = 1, \cdots, l,\ \sum_{i=1}^{l} \lambda_i = 1$, and $f(x_i) \leq z$ for $i = 1, \cdots, l$ then $f(x) < z$ whenever $\{f(x_i) : i = 1, \cdots, l\}$ is not a singleton.

Consider the problem

$$(P) \qquad \min_{x \in X} F(x)$$

with $F = (f_1, \cdots, f_k)$, each $f_i,\ i = 1, \cdots, k$ being strictly quasiconvex. We refer to a subproblem P_I of P when only a nonempty subset $I \subset \{1, \cdots, k\}$ of all objectives, is considered.

The notation $|I|$ stands for the cardinality of I.

Recall that the sets of minimal points and weakly minimal points of P are defined by

$$MinF(X) = \{z \in F(X) : (z - F(X)) \cap (\mathbb{R}^k_+ \setminus \{0\}) = \emptyset\}$$

$$WMinF(X) = \{z \in F(X) : (z - F(X)) \cap \operatorname{int} \mathbb{R}^k_+ = \emptyset\}\cdot$$

The corresponding efficient sets in the argument space are:

$$Eff\, P = \{x \in X : F(x) \in MinF(X)\}$$

$$WEff\, P = \{x \in X : F(x) \in WMinF(X)\}.$$

3. Determination of *WEff P*

In 1984, Lowe et al.[3] showed that, when all objectives are convex, *WEff P* is the union of efficient sets of all subproblems P_I , $I \subset \{1, \cdots, k\}, I \neq \emptyset$.

At first we give a similar result in the strictly quasiconvex case.

Theorem 3.1. *Suppose that all objectives f_i, $i = 1, \cdots, k$ are strictly quasiconvex and upper semicontinuous along line segments in X then*

$$WEff\ P = \cup\{Eff\ P_I : I \subset \{1, \cdots, k\}, I \neq \emptyset\}$$

Proof. Consider $x \in X$ such that $x \notin WEff\ P$. There exists $y \in X$ with $f_i(y) < f_i(x)$ for all i and then, for each $I \subset \{1, \cdots, k\}, x \notin Eff\ P_I$.

Now, suppose that $x \notin \cup\{Eff\ P_I : I \subset \{1, \cdots, k\}, I \neq \emptyset\}$. Taking $I = \{1, \cdots, k\}$, there must exist $i_1 \in I$ and $x_1 \in X$ such that

$$f_{i_1}(x_1) < f_{i_1}(x) \text{ and } f_i(x_1) \leq f_i(x), \quad i \in I. \tag{1}$$

Now if $I = I_1 = \{1, \cdots, k\}\backslash\{i_1\}$, there are $i_2 \in I_1$ and $x_2 \in X$ such that

$$f_{i_2}(x_2) < f_{i_2}(x) \quad \text{and} \quad f_i(x_2) \leq f_i(x), \quad i \in I_1. \tag{2}$$

Setting $y_2 = \lambda x_1 + (1-\lambda)x_2, \lambda \in]0,1[$, and using the upper semicontinuity of f_{i_1} on the segment $[x_1, x_2]$, we have by (1), for $(1-\lambda)$ small enough,

$$f_{i_1}(y_2) < f_{i_1}(x).$$

On the other hand, the strict quasiconvexity of f_{i_2} implies, for each $\lambda \in]0,1[$

$$f_{i_2}(y_2) < f_{i_2}(x).$$

Further from the quasiconvexity of f_i 's , we have for each $\lambda \in]0,1[$

$$f_i(y_2) \leq f_i(x), \quad i \in I_2 = \{1, \cdots, k\}\backslash\{i_1, i_2\}.$$

Suppose now, that we have obtained y_ℓ and $I_\ell = I\backslash\{i_1, \cdots, i_\ell\}$ such that $f_i(y_\ell) < f_i(x), \quad i \in \{i_1, \cdots, i_\ell\}$ and $f_i(y_\ell) \leq f_i(x), \quad i \in I_\ell$. Using $x \notin Eff\ P_{I_\ell}$ we get $i_{\ell+1} \in I_\ell$ and $x_{\ell+1} \in X$ satisfying

$$f_{i_{\ell+1}}(x_{\ell+1}) < f_{i_{\ell+1}}(x) \quad \text{and} \quad f_i(x_{\ell+1}) \leq f_i(x), \quad i \in I_\ell.$$

Then from the upper semicontinuity of $f_{i_1}, \cdots, f_{i_\ell}$ on the segment $[y_l, x_{\ell+1}]$, the strict quasiconvexity of $f_{i_{\ell+1}}$ and the quasiconvexity of other objectives, we obtain $y_{\ell+1}$, a convex combination of y_ℓ and $x_{\ell+1}$, such that

$$f_i(y_{\ell+1}) < f_i(x), \quad i \in \{i_1, \cdots, i_{\ell+1}\}$$

and

$$f_i(y_{\ell+1}) \leq f_i(x), \quad i \in I_{\ell+1} = I\backslash\{i_1, \cdots, i_{\ell+1}\}.$$

As the number of objectives is k, we obtain finally $y_k \in X$ such that
$f_i(y_k) < f_i(x), \quad i \in \{1, \cdots, k\}$, which means that $x \notin WEff\, P$. □

Note that convex functions are automatically upper semicontinuous along line segments and also that the previous theorem remains valid if $X \subset E$, where E is a linear space without topology.

Now, following an idea developped by Ward [7] in the convex case, which uses Helly's Theorem, we give an extension of Theorem 3.1 when $n < k$.

Helly's Theorem : Let C_i, $i = 1, \cdots, m$ be a collection of convex sets in $\mathbb{R}^n$. If every subcollection of $n+1$ or fewer of these C_i has a nonempty intersection, then the entire collection of the m sets has a nonempty intersection.

Theorem 3.2. *Suppose that f_i, $i = 1, \cdots, k$ are strictly quasiconvex and upper semicontinuous along line segments, then*

$$WEff\, P = \cup\{Eff\, P_I : I \subset \{1, \cdots, k\}, I \neq \emptyset, |I| \leq n+1\}$$

Proof. From Theorem 3.1 it is sufficient to consider the case $k > n+1$ and to prove the inclusion $\subset$.

Consider $x \notin \cup\{Eff\ P_I : I \subset \{1, \cdots, k\}, I \neq \emptyset, |I| \leq n+1\}$. Then for each $J \subset \{1, \cdots, k\}, J \neq \emptyset$ with $|J| \leq n+1$, we have $x \notin \cup\{Eff\ P_I : I \subset J, I \neq \emptyset\}$ and from Theorem 3.1, it follows that $x \notin WEff\, P_J$. Therefore there exists

$$x_J \in X \text{ such that } f_j(x_J) < f_j(x) \text{ for all } j \in J. \tag{3}$$

For each $i \in \{1, \cdots, k\}$ we define the closed convex set

$$C_i = \text{conv}\{x_J : J \subset \{1, \cdots, k\}, J \neq \emptyset, |J| \leq n+1, \ \ i \in J\}.$$

It is clear from (3) that, for all $J \subset \{1, \cdots, k\}$ with $J \neq \emptyset, |J| \leq n+1$, $i \in J$, we have $f_i(x_J) < f_i(x)$ and the quasiconvexity of f_i entails that for every $y \in C_i$

$$f_i(y) < f_i(x). \tag{4}$$

Now, for a fixed J with $|J| \leq n+1$, the collection $\{C_i\ ,\ i \in J\}$ has x_J in common and from Helly's Theorem, there exists some y^* belonging to $\bigcap_{i=1}^{k} C_i$. Thus from (4), for each $i \in \{1, \cdots, k\}$

$$f_i(y^*) < f_i(x)$$

and $x \notin WEff\, P$. □

Recall that if $C \subset \mathbb{R}^n$, $\dim C$ denotes the dimension of the affine space generated by C. The following lemma will be useful in the sequel.

Lemma 3.3. *Let $C = \text{conv}\{y_i \in \mathbb{R}^n : i = 1, \cdots, n+1\}$. Suppose that there exists $x \in C$ which cannot be written as a convex combination of fewer than $n+1$ points y_i, then $\dim C = n$ and $x \in \text{int}\, C$.*

Proof. Suppose that $\dim C \leq n-1$. By Carathéodory's Theorem [4] , every point of C can be expressed as a convex combination of n elements y_i. This is a contradiction with the assumption about x. It remains to prove that $x \in \operatorname{int} C$. Suppose that $x = \sum_{i=1}^{n+1} \lambda_i y_i$, with all $\lambda_i > 0$. As $\dim C = n$, the vectors $y_{n+1} - y_i, i = 1, \cdots, n$ are independent. We consider the neighborhood of 0 in $\mathbb{R}^n$ defined by

$$N = \{z : z = \sum_{i=1}^{n} \gamma_i (y_{n+1} - y_i) \ , \ |\gamma_i| < \lambda_i \quad i = 1, \cdots, n, \quad |\sum_{i=1}^{n} \gamma_i| < \lambda_{n+1}\}.$$

We have for every $z \in N$,

$$x + z = \sum_{i=1}^{n} (\lambda_i - \gamma_i) y_i + (\lambda_{n+1} + \sum_{i=1}^{n} \gamma_i) y_{n+1} \in C.$$

Thus $x + N \subset C$ and $x \in \operatorname{int} C$. □

Lemma 3.4. *Suppose that $x \in WEff\, P_J$ and that for every $j \in J$ there exists y_j such that*

$$f_i(y_j) < f_i(x), \quad i \in J \backslash \{j\}, \tag{5}$$

then

$$x \in \operatorname{int} C \text{ where } C = \operatorname{conv}\{y_j : j \in J\}.$$

Proof. For each $j \in J$, consider the closed, convex subset of X,

$$C_j = \operatorname{conv}\{y_k : k \in J \backslash \{j\}\}.$$

Obviously, by (5), $f_j(y_k) < f_j(x)$ for $k \in J \backslash \{j\}$ and the quasiconvexity of f_j implies

$$\forall y \in C_j \qquad f_j(y) < f_j(x). \tag{6}$$

From (6), for all $j \in J$, $x \notin C_j$ and since $x \in WEff\, P_J$ we have $\cap\{C_j : j \in J\} = \emptyset$.

Now let us define $C'_j = \operatorname{conv}(C_j, \{x\})$ for $j \in J$.

The intersection of $n+1$ of the sets $\{(C'_j)_{j \in J}, C\}$ is nonempty. Indeed, all C'_j contain x and if we take a collection of the form $\{(C'_j)_{j \in J \backslash \{j_0\}}, C\}$, all of these sets contain y_{j_0}.

Applying Helly's Theorem, there exists $z \in \left(\bigcap_{j \in J} C'_j \right) \bigcap C$. If $z \neq x$, then for each $j \in J$, $z = \lambda_j x + (1 - \lambda_j) y$, with $\lambda_j \in [0, 1[$, $y \in C_j$. From (6) and the strict quasiconvexity of f_j, for each $j \in J$, $f_j(z) < f_j(x)$ which contradicts $x \in WEff\, P_J$. Thus $z = x$ and $x \in C$. Thus we can write $x = \sum_{j \in J} \lambda_j y_j, \ \lambda_j \geq 0, \quad \sum_{j \in J} \lambda_j = 1$.

Suppose that $\lambda_{j_0} = 0$ for some $j_0 \in J$. By definition of y_j, $f_{j_0}(y_j) < f_{j_0}(x)$, for all $j \in J\backslash\{j_0\}$ and f_{j_0} being quasiconvex, $f_{j_0}(x) \leq \max(f_{j_0}(y_j), j \in J\backslash\{j_0\}) < f_{j_0}(x)$, a contradiction. Thus every λ_j is strictly positive and using Lemma 3.3 we conclude $x \in \operatorname{int} C$. □

It is well known that the continuity of f_i's implies that *WEff P* is a closed set. Let us denote by $\operatorname{bd}(WEff\ P)$ the boundary of *WEff P*.

Theorem 3.5. *Suppose that f_i, $i = 1, \cdots, k$ are strictly quasiconvex and continuous. Then*

$$\operatorname{bd}(WEff\ P) \subset \cup\{Eff\ P_I : I \subset \{1, \cdots, k\}, I \neq \emptyset, |I| \leq n\}.$$

Proof. By Theorem 3.1 it is sufficient to consider the case $k > n$.

Suppose that $x \in \operatorname{bd}(WEff\ P) \setminus \cup\{Eff\ P_I : I \subset \{1, \cdots, k\}, I \neq \emptyset, |I| \leq n\}$. Using Theorem 3.2 we have $x \in Eff\ P_J$ for some $J \subset \{1, \cdots, k\}$, with $|J| = n+1$ and $x \notin Eff\ P_{J'}$ for all $J' \subset J, J' \neq \emptyset, J' \neq J$. In particular for every $j \in J$, $x \notin \cup\{Eff\ P_{J'} : J' \subset J\backslash\{j\}, J' \neq \emptyset\}$. But from Theorem 3.1 , this latter set is $WEff\ P_{J\backslash\{j\}}$. Thus

$$\forall j \in J \quad \exists y_j \in X \quad f_i(y_j) < f_i(x), i \in J\backslash\{j\}.$$

It follows from Lemma 3.4 that $x \in \operatorname{int}\ C$, where $C = \operatorname{conv}\{y_j : j \in J\}$. From Lemma 3.3, each $y \in \operatorname{bd}\ C$ can be written as a convex combination of only n points y_j, say $\{y_j : j \in J\backslash\{i\}\}$ and by the quasiconvexity of $f_i, f_i(y) < f_i(x)$. It follows that

$$\forall y \in \operatorname{bd}\ C \quad \max_{j\in J}(f_j(x) - f_j(y)) > 0\cdot \tag{7}$$

As $\max_{j\in J}(f_j(x) - f_j(y))$ is continuous with respect to y, it achieves its minimum on the compact set $\operatorname{bd}\ C$.

Denote by $m = \min_{y\in \operatorname{bd}\ C} \max_{j\in J}(f_j(x) - f_j(y))$, we have $m > 0$.

Now consider the neighborhood of x in $\mathbb{R}^n$ defined by

$$V = \{y \in C : \max_{j\in J}(f_j(x) - f_j(y)) < m\}\cdot$$

Note that by the definition of m, $V \cap \operatorname{bd}\ C = \emptyset$ and then $V \subset \operatorname{int} C$. We show that $V \subset Eff\ P_J$.

Suppose on the contrary that $y \in V\backslash Eff\ P_J$. Then

$$\exists u \in X \quad (f_i(y) - f_i(u))_{i\in J} \in \mathbb{R}^n_+\backslash\{0\}\cdot \tag{8}$$

As $y \in \operatorname{int} C$ and C being a compact set, we can find $t_0 = \max\{t > 0 : y + t(y-u) \in C\}$. We have $z = y + t_0(y-u) \in \operatorname{bd}\ C$ and by (7), there exists $i_0 \in J$ such that

$$f_{i_0}(x) - f_{i_0}(z) \geq m > 0\cdot \tag{9}$$

As $y \in V$ we have also $f_{i_0}(x) - f_{i_0}(y) < m$ and with (9)

$$f_{i_0}(y) - f_{i_0}(z) > 0\cdot \tag{10}$$

Now using the fact that y is a convex combination of u and z, and the strict quasiconvexity of f_{i_0}, we get either $f_{i_0}(u) = f_{i_0}(z) \geq f_{i_0}(y)$, which contradicts (10), or

$$f_{i_0}(u) \neq f_{i_0}(z) \text{ and } f_{i_0}(y) < \max(f_{i_0}(u), f_{i_0}(z)). \tag{11}$$

But $f_{i_0}(y) > f_{i_0}(z)$ by (10) and then (11) entails $f_{i_0}(y) < f_{i_0}(u)$, a contradiction with (8).

Thus we have proved that $V \subset \mathit{Eff}\ P_J$, which means that $x \in \text{int}(\mathit{Eff}\ P_J)$. As $\mathit{Eff}\ P_J \subset \mathit{WEff}\ P_J$, we have got a contradiction with the assumption $x \in \text{bd}(\mathit{WEff}\ P_J)$ and the proof is complete. □

In the following we adopt the notations of Ward [7] :

$$r[x,v] = \{x + tv : t \geq 0\} \quad x \in X, v \in \mathbb{R}^n,$$
$$U(P,n) = \cup\{\mathit{Eff}\ P_I : I \subset \{1,\cdots,k\}, I \neq \emptyset, |I| \leq n\}$$
$$S(P,n) = \{x \in X \backslash U(P,n) : \forall v \neq 0,\ r[x,v] \cap U(P,n) \neq \emptyset\}.$$

The result of (lemma 4, [7]) remains true in the strictly quasiconvex case.

Lemma 3.6. *Suppose that f_i, $i = 1,\cdots,k$ are strictly quasiconvex, then $S(P,n) \subset \mathit{Eff}\ P$.*

Proof. Assume that $x \in S(P,n) \backslash \mathit{Eff}\ P$. Then, there exists $y \in X$ such that

$$(f_i(x) - f_i(y))_{i=1,\cdots,k} \in \mathbb{R}^k_+ \backslash \{0\}$$

and we can find $z \in r[x, x-y] \cap U(P,n)$. As $x \neq z$ and $x \neq y$, $x = \lambda z + (1-\lambda)y$ with $\lambda \in]0,1[$. If $f_i(y) = f_i(z)$ then the strict quasiconvexity of f_i implies $f_i(x) \leq f_i(z)$. If $f_i(y) \neq f_i(z)$ the strict quasiconvexity of f_i implies $f_i(x) < max(f_i(y), f_i(z))$ and with $f_i(x) \geq f_i(y)$ one has $f_i(x) < f_i(z)$. Therefore, in all cases, $f_i(x) \leq f_i(z), i = 1,\cdots,k$. As $z \in \mathit{Eff}\ P_I$ for some I such that $|I| \leq n$, we get also $x \in \mathit{Eff}\ P_I$, contradicting $x \notin U(P,n)$.

□

Theorem 3.7. *Let f_i, $i = 1,\cdots,k$ be strictly quasiconvex, continuous functions and suppose that $\mathit{WEff}\ P$ is bounded, then $\mathit{WEff}\ P = U(P,n) \cup S(P,n)$.*

Proof. The inclusion $\supset$ is clear from Lemma 3.6 and Theorem 3.2.

For $\subset$ observe that $\mathit{WEff}\ P = \text{bd}(\mathit{WEff}\ P) \cup \text{int}(\mathit{WEff}\ P)$. From Theorem 3.5 follows $\text{bd}(\mathit{WEff}\ P) \subset U(P,n)$.

Now we prove that $\text{int}(\mathit{WEff}\ P) \backslash U(P,n) \subset S(P,n)$.

Consider $x \in \text{int}(\mathit{WEff}\ P) \backslash U(P,n)$ and an halfline $r[x,v]$. Since $\mathit{WEff}\ P$ is bounded, there exists $z \in r[x,v] \cap \text{bd}(\mathit{WEff}(P))$, $z \neq x$. By Theorem 3.5, $z \in r[x,v] \cap U(P,n)$ and then $x \in S(P,n)$. □

4. Conclusion

The subset $S(P,n)$ is a kind of convex hull of $U(P,n)$ and the knowledge of $U(P,n)$ completely determines $S(P,n)$. Thus it is sufficient to solve subproblems P_I with at most n criteria, to obtain $WEff\, P$. In $\mathbb{R}^2$ a graphical representation of $WEff\, P$ can be obtained, even with a great number of criteria, as soon as bicriteria subproblems can be solved.

Acknowledgment. The authors wish to thank anonymous referees for their careful checking and their helpful suggestions that improved the quality of this paper.

References

[1] J.P. Crouzeix: Contribution à l'étude des fonctions quasi-convexes, Thèse de doctorat, U.E.R. des Sciences, Université de Clermont-Ferrand II, 1977, France.

[2] N. Popovici: Sur une notion de quasi-convexité pour des fonctions vectorielles, (préprint)

[3] T.J. Lowe, J.-F. Thisse, J.E. Ward and R.E. Wendell: On efficient solutions to multiple objective mathematical programs. Management Sci. 30, 1984, 1346–1349.

[4] R.T. Rockafellar: Convex Analysis. Princeton University Press, 1970, Princeton N.J.

[5] S. Schaible: Bibliography in fractional programming. Zeitschrift für Operations Research 26, 1982, 211–241.

[6] S. Schaible: A survey of fractional programming in : S. Schaible and W.T. Ziemba eds., Generalized Concavity in Optimization and Economics, (Acad. Press, New York 1981), 417–440.

[7] J. Ward: Structure of efficient sets for convex objectives. Math. of Oper. Res. 14, 1989, 249–257.