

GENERALIZED CO-COMPLEMENTARITY PROBLEMS IN p -UNIFORMLY SMOOTH BANACH SPACES

M. FIRDOSH KHAN AND SALAHUDDIN

Department of Mathematics
Aligarh Muslim University
Aligarh-202002, India

EMail: khan_mfk@yahoo.com



volume 7, issue 2, article 66,
2006.

*Received 14 October, 2005;
accepted 30 December, 2005.*

Communicated by: R.U. Verma

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


The objective of this paper is to study the iterative solutions of a class of generalized co-complementarity problems in p -uniformly smooth Banach spaces, with the devotion of sunny retraction mapping, p -strongly accretive, p -relaxed accretive and Lipschitzian (or more generally uniformly continuous) mappings. Our results are new and represents a significant improvement of previously known results. Some special cases are also discussed.

2000 Mathematics Subject Classification: 49J40, 90C33, 47H10.

Key words: Generalized co-complementarity problems, Iterative algorithm, Sunny retraction, Sunny nonexpansive mapping, p -strongly accretive, p -relaxed accretive mapping, Lipschitzian mapping, Hausdorff metric, p -uniformly smooth Banach spaces.

Contents

1	Introduction	3
2	Background of Problem Formulations	4
	2.1 Special Cases	7
3	The Characterization of Problem and Solutions	8
4	Iterative Algorithms and Pertinent Concepts	12
5	Main Results	15
	References	



Generalized Co-Complementarity Problems in p -Uniformly Smooth Banach Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 2 of 24


1. Introduction

The theory of complementarity problems initiated by Lemke [19] and Cottle and Dantzing [10] in the early sixties and later developed by other mathematicians see for example [6, 9, 11, 14, 17, 22] plays an important role and is fundamental in the study of a wide class of problems arising in optimization, game theory, economics and engineering sciences [3, 6, 8, 11, 15].

On the other hand, the accretive operators are of interest because several physically resolvent problems can be modeled by nonlinear evolution systems involving operators of the accretive type. Very closely related to the accretive operators is the class of dissipative operators, where an operator T is said to be dissipative if and only if $(-T)$ is accretive. The concepts of strictly strongly and m -(or sometimes hyper-) dissipativity are similarly defined.

These classes of operators have attracted a lot of interest because of their involvement in evolution systems modeling several real life problems. Consequently several authors have studied the existence, uniqueness and iterative approximations of solutions of nonlinear equations involving such operators, see [5, 12, 18] and the references cited therein.

It is our purpose in this article to establish the strong convergence of the iterative algorithm to a solution of the generalized co-complimentarity problems in p -uniformly smooth Banach spaces when the operators are accretive, strictly accretive, strongly accretive, relaxed accretive and Lipschitzian. Our iteration processes are simple and independent of the geometry of E and iteration parameters can be chosen at the start of the iteration process. Consequently, most important results known in this connection will be special cases of our problem.



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 3 of 24

2. Background of Problem Formulations

Throughout this article, we assume that E is a real Banach space whose norm is denoted by $\|\cdot\|$, E^* its topological dual space. $CB(E)$ denotes the family of all nonempty closed and bounded subsets of E . $D(\cdot, \cdot)$ is the Hausdorff metric on $CB(E)$ defined by

$$\max \left\{ \sup_{x \in A} d(x, B), \sup_{y \in B} d(A, y) \right\} = D(A, B),$$

where

$$d(x, B) = \inf_{y \in B} d(x, y) \quad \text{and} \quad d(A, y) = \inf_{x \in A} d(x, y),$$

d is the metric on E induced by the norm $\|\cdot\|$. As usual, $\langle \cdot, \cdot \rangle$ is the generalized duality pairing between E and E^* . For $1 < p < \infty$, the mapping $J_p : E \rightarrow 2^{E^*}$ defined by

$$J_p(x) = \{f^* \in E^* : \langle x, f^* \rangle = \|f\| \cdot \|x\| \text{ and } \|f\| = \|x\|^{p-1}\} \quad \text{for all } x \in E,$$

is called the duality mapping with gauge function $\phi(t) = t^{p-1}$. In particular for $p = 2$, the duality mapping J_2 with gauge function $\phi(t) = t$ is called the normalized duality mapping. It is known that $J_p(x) = \|x\|^{p-2} J_2(x)$ for all $x \neq 0$ and J_p is single valued if E^* is strictly convex. If $E = H$ is a Hilbert space, then J_2 becomes the identity mapping on H .

Proposition 2.1 ([7]). *Let E be a real Banach space. For $1 < p < \infty$, the duality mapping $J_p : E \rightarrow 2^{E^*}$ has the following basic properties:*

1. $J_p(x) \neq \emptyset$ for all $x \in E$ and $D(J_p)$ (the domain of J_p) = E ,



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 4 of 24

2. $J_p(x) = \|x\|^{p-2} J_2(x)$ for all $x \in E$, ($x \neq 0$),
3. $J_p(\alpha x) = \alpha^{p-1} J_p(x)$ for all $\alpha \in [0, \infty)$,
4. $J_p(-x) = -J_p(x)$,
5. J_p is bounded i.e., for any bounded subset $A \subset E$, $J_p(A)$ is a bounded subset in E^* ,
6. J_p can be equivalently defined as the subdifferential of the functional $\varphi(x) = p^{-1}\|x\|^p$ (Asplund [2]), i.e.,

$$J_p(x) = \partial\varphi(x) = \{f \in E^* : \varphi(y) - \varphi(x) \geq (f, y - x), \text{ for all } y \in E\}$$


7. E is a uniformly smooth Banach space (equivalently, E^* is a uniformly convex Banach space) if and only if J_p is single valued and uniformly continuous on any bounded subset of E (Xu and Roach [24]).

Definition 2.1. Let E be a real Banach space and K a nonempty subset of E . Let $T : K \rightarrow 2^E$ be a multivalued mapping

1. T is said to be accretive if for any $x, y \in K$, $u \in T(x)$ and $v \in T(y)$ there exists $j_2 \in J_2(x - y)$ such that

$$\langle u - v, j_2 \rangle \geq 0,$$

or equivalently, there exists $j_p \in J_p(x - y)$, $1 < p < \infty$, such that



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 5 of 24

2. T is said to be strongly accretive if for any $x, y \in K$, $u \in T(x)$ and $v \in T(y)$ there exists $j_2 \in J_2(x - y)$ such that

$$\langle u - v, j_2 \rangle \geq k \|x - y\|^2,$$

or equivalently, there exists $j_p \in J_p(x - y)$, $1 < p < \infty$ such that

$$\langle u - v, j_p \rangle \geq k \|x - y\|^p,$$

for some constant $k > 0$.

The concept of a single-valued accretive mapping was introduced independently by Browder [5] and Kato [18] in 1967. An early fundamental result in the theory of accretive mappings which is due to Browder states that the following initial value problem,

$$\frac{du(t)}{dt} + Tu(t) = 0, \quad u(0) = u_0,$$

is solvable if T is locally Lipschitzian and accretive on E .

More precisely, let $N : E \times E \rightarrow E$ and $m, g : E \rightarrow E$ be the single-valued mappings and $F, G, T : E \rightarrow CB(E)$ the multivalued mappings. Let X be a fixed closed convex cone of E . Define $K : E \rightarrow 2^E$ by


$$K(z) = m(z) + X \quad \text{for all } x \in E, z \in T(x).$$

We shall study the following generalized co-complementarity problem (**GCCP**):

Find $x \in E$, $u \in F(x)$, $v \in G(x)$, $z \in T(x)$ such that $g(x) \in K(z)$ and

$$(2.1) \quad N(u, v) \in (J(K(z) - g(x)))^*,$$

where $(J(K(z) - g(x)))^*$ is the dual cone of the set $J(K(z) - g(x))$.



Title Page

Contents



Go Back

Close

Quit

Page 6 of 24

2.1. Special Cases

- (i) If E is a Hilbert space, F, T are identity mappings and $N(u, v) = Bx + Av$, where B, A are single-valued mappings, then Problem (2.1) reduces to a problem of finding $x \in E, v \in G(x)$ such that $g(x) \in K(z)$ and

$$(2.2) \quad B(x) + A(v) \in (K(x) - g(x))^*$$

considered by Jou and Yao [16].

- (ii) If G and g are identity mappings, then (2.2) reduces to finding $x \in K(x)$ such that

$$(2.3) \quad B(x) + A(y) \in (K(x) - x)^*$$

which is called a strongly nonlinear quasi complementarity problem, studied by Noor [22].

- (iii) If m is a zero mapping, then (2.3) is equivalent to finding $x \in E$ such that


$$(2.4) \quad Bx + Ax \in E^* \quad \text{and} \quad \langle Bx + Ax, x \rangle = 0,$$

which is known as the mildly nonlinear complementarity problem, studied by Noor [21].

- (iv) If A is zero mapping, then (2.4) is equivalent to a problem of finding $x \in E$ such that $Bx \in E^*$ and

$$(2.5) \quad \langle Bx, x \rangle = 0,$$

considered by Habetler [14] and Karamardian [17].



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 7 of 24

3. The Characterization of Problem and Solutions

In this section, we briefly consider some basic concepts and results, which will be used throughout the paper. The real Banach space E is said to be uniformly smooth if its modulus of smoothness $\rho_E(\tau)$ defined by


$$\rho_E(\tau) = \sup \left\{ \frac{\|x + y\| + \|x - y\|}{2} - 1 : \|x\| = 1, \|y\| \leq \tau \right\}$$

satisfies $\frac{\rho_E(\tau)}{\tau} \rightarrow 0$ as $\tau \rightarrow 0$. It follows that E is uniformly smooth if and only if J_p is single-valued and uniformly continuous on any bounded subset of E and there exists a complete duality between uniform convexity and uniform smoothness. E is uniformly convex (smooth) if and only if E^* is uniformly smooth (convex). Recall that E is said to have the modulus of smoothness of power type $p > 1$ (and E is said to be p -uniformly smooth) if there exists a constant $c > 0$ such that

$$\rho_E(\tau) \leq c\tau^p \quad \text{for } 0 < \tau < \infty.$$

Remark 1. *It is known that all Hilbert spaces and Banach spaces, L_p, l_p and W_m^p ($1 < p < \infty$) are uniformly smooth and*

$$\rho_E(\tau) < \begin{cases} \frac{1}{p} \tau^p, & 1 < p < 2 \\ \frac{p-1}{2} \tau^2, & p \leq 2 \end{cases} \quad (E = L_p, l_p \text{ or } W_m^p)$$



Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 8 of 24

therefore E is a p -uniformly smooth Banach space with modulus of smoothness of power type $p < 1$ and J_p will always represent the single-valued duality mapping.

Definition 3.1 ([4, 13]). Let E be a p -uniformly smooth Banach space and let Ω a nonempty closed convex subset of E . A mapping $Q_\Omega : E \rightarrow \Omega$ is said to be

- (i) retraction on Ω if $Q_\Omega^p = Q_\Omega$;
- (ii) nonexpansive retraction if it satisfies the inequality

$$\|Q_\Omega(x) - Q_\Omega(y)\| \leq \|x - y\|, \quad \text{for all } x, y \in E;$$

- (iii) sunny retraction if for all $x \in E$ and for all $-\infty < t < \infty$

$$Q_\Omega(Q_\Omega(x) + t(x - Q_\Omega(x))) = Q_\Omega(x).$$


The following characterization of a sunny nonexpansive retraction mapping can be found in [4, 13].

Lemma 3.1 ([4, 13]). Q_Ω is sunny nonexpansive retraction if and only if for all $x, y \in E$,

$$\langle x - Q_\Omega^*, J(Q_\Omega x - y) \rangle \geq 0.$$

Lemma 3.2. Let E be a real Banach space and $J_p : E \rightarrow 2^{E^*}$, $1 < p < \infty$ a duality mapping. Then, for any given $x, y \in E$, we have

$$\|x + y\|^p \leq \|x\|^p + p \langle y, j_p \rangle, \quad \text{for all } j_p \in J_p(x + y).$$



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 9 of 24

Proof. From Proposition 2.1, it follows that $J_p(x) = \partial\varphi(x)$ (subdifferential of ψ), where $\psi(x) = p^{-1}\|x\|^p$. Also, it follows from the definition of the subdifferential of ψ that

$$\psi(x) - \psi(x + y) \geq \langle x - (x + y), j_p \rangle,$$

for all $j_p \in J_p(x + y)$. Substituting $\psi(x)$ by $p^{-1}\|x\|^p$, we have

$$\|x + y\|^p \leq \|x\|^p + p \langle y, j_p \rangle, \quad \text{for all } j_p \in J_p(x + y).$$

This completes the proof. \square

Theorem 3.3 ([9]). *Let E be a Banach space, Ω a nonempty closed convex subset of E , and $m : E \rightarrow E$. Then for all $x, y \in E$, we have*


$$Q_{\Omega+m(z)}(x) = m(z) + Q_{\Omega}(x - m(z)).$$

We mention the following characterization theorem for the solution of a generalized co-complementarity problem which can be easily proved by using Lemma 3.1 and the argument of [1, Theorem 8.1].

Theorem 3.4. *Let E be a real p -uniformly smooth Banach space and X a closed convex cone in E . Let $F, G, T : E \rightarrow CB(E)$ be the multivalued mappings, $m, g : E \rightarrow E$ the two single-valued mappings and $N : E \times E \rightarrow E$ the nonlinear mapping. Let $K : E \rightarrow 2^E$ and $K(z) = m(z) + X$ for $x \in E$. Then the following statements are equivalent:*

- (i) $x \in E$, $u \in F(x)$, $v \in G(x)$ and $z \in T(x)$ are solutions of the Problem (2.1), i.e., $g(x) \in K(z)$ and

$$N(u, v) \in (J(K(z) - g(x)))^*.$$



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 10 of 24

(ii) $x \in E, u \in F(x), v \in G(x)$ and $z \in T(x)$ and $\tau > 0$

$$g(x) = Q_{K(z)}[g(x) - \tau N(u, v)].$$

Combining Theorem 3.3 and 3.4, we have the following result.

Theorem 3.5. *Let E be a p -uniformly smooth Banach space and X a closed convex cone in B . Let $m, g : E \rightarrow E$ be the two single-valued mappings, $F, G, T : E \rightarrow CB(E)$ the multivalued mappings and $N : E \times E \rightarrow E$ a nonlinear mapping. Then the following statements are equivalent:*

(i) $x \in E, u \in F(x), v \in G(x)$ and $z \in T(x)$ are solutions of the Problem (2.1),

(ii) $x = x - g(x) + m(z) + Q_X[g(x) - \tau N(u, v) - m(z)]$, for some $\tau > 0$.

The following inequality will be used in our main results.


Lemma 3.6. *Let E be a real Banach space and $j_p : E \rightarrow 2^{E^*}, 1 < p < \infty$ a duality mapping. Then, for any given $x, y \in E$, we have*

$$\langle x - y, j_p(x) - j_p(y) \rangle \leq 2d^p \rho_E \left(\frac{4\|x - y\|}{d} \right),$$

where

$$d^p = \left(\frac{\|x\|^2 + \|y\|^2}{2} \right).$$

Proof. The proof of the above inequalities are the generalized form of the proof of Theorem 3.3, and hence will be omitted. \square



Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 11 of 24

4. Iterative Algorithms and Pertinent Concepts

We now propose the following iterative algorithm for computing the approximate solution of (GCCP).

Algorithm 1. Let $g, m : E \rightarrow E$ be the two single-valued mappings, $F, G, T : E \rightarrow CB(E)$ the multivalued mappings and $N : E \times E \rightarrow E$ a nonlinear mapping.

For any given $x_0 \in E$, $u_0 \in F(x_0)$, $v_0 \in G(x_0)$ and $z_0 \in T(x_0)$, let

$$x_1 = x_0 - g(x_0) + m(z_0) + Q_X[g(x_0) - \tau N(u_0, v_0) - m(z_0)]$$

where $\tau > 0$ is a constant.

Since $u_0 \in F(x_0) \in CB(E)$, $v_0 \in G(x_0) \in CB(E)$ and $z_0 \in T(x_0) \in CB(E)$, by Nadler's Theorem [20], there exists $u_1 \in F(x_1)$, $v_1 \in G(x_1)$ and $z_1 \in T(x_1)$ such that

$$\|u_0 - u_1\| \leq (1 + 1)D(F(x_0), F(x_1)),$$


$$\|v_0 - v_1\| \leq (1 + 1)D(G(x_0), G(x_1)),$$

$$\|z_0 - z_1\| \leq (1 + 1)D(T(x_0), T(x_1)),$$

where D is a Hausdorff metric on $CB(E)$.

Let

$$x_2 = x_1 - g(x_1) + m(z_1) + Q_X[g(x_1) - \tau N(u_1, v_1) - m(z_1)].$$



Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 12 of 24

Since $u_1 \in F(x_1) \in CB(E)$, $v_1 \in G(x_1) \in CB(E)$ and $z_1 \in T(x_1) \in CB(E)$, there exists $u_2 \in F(x_2)$, $v_2 \in G(x_2)$ and $z_2 \in T(x_2)$ such that

$$\begin{aligned}\|u_1 - u_2\| &\leq (1 + 2^{-1})D(F(x_1), F(x_2)), \\ \|v_1 - v_2\| &\leq (1 + 2^{-1})D(G(x_1), G(x_2)), \\ \|z_1 - z_2\| &\leq (1 + 2^{-1})D(T(x_1), T(x_2)).\end{aligned}$$

By induction, we can obtain $\{x_n\}$, $\{u_n\}$, $\{v_n\}$ and $\{z_n\}$ as

$$(4.1) \quad x_{n+1} = x_n - g(x_n) + m(z_n) + Q_X[g(x_n) - \tau N(u_n, v_n) - m(z_n)].$$

$$\begin{aligned}u_n \in F(x_n); \|u_n - u_{n+1}\| &\leq (1 + (1 + n)^{-1}) D(F(x_n), F(x_{n+1})), \\ v_n \in G(x_n); \|v_n - v_{n+1}\| &\leq (1 + (1 + n)^{-1}) D(G(x_n), G(x_{n+1})), \\ z_n \in T(x_n); \|z_n - z_{n+1}\| &\leq (1 + (1 + n)^{-1}) D(T(x_n), T(x_{n+1})),\end{aligned}$$

$n \geq 0$, where $\tau > 0$ is a constant.


These iteration processes have been extensively investigated by various authors for approximating either the fixed point of nonlinear mappings or solutions of nonlinear equations in Banach spaces or variational inequalities, variational inclusions, or complementarity problems in Hilbert spaces.

Definition 4.1. A single valued mapping $g : E \rightarrow E$ is said to be

(i) p -strongly accretive if for all $x, y \in E$ there exists $j_p \in J_p(x - y)$ such that

$$\langle g(x) - g(y), j_p(x - y) \rangle \geq \kappa \|x - y\|^p$$

for some real constant $\kappa \in (0, 1)$ and $1 < p < \infty$.



Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 13 of 24

(ii) Lipschitz continuous if for any $x, y \in E$, there exists constant $\beta > 0$, such that

$$\|g(x) - g(y)\| \leq \beta \|x - y\|.$$

Definition 4.2. A multivalued mapping $F : E \rightarrow CB(E)$ is said to be D -Lipschitz continuous if for any $x, y \in E$,

$$D(F(x), F(y)) \leq \mu \|x - y\|$$

for $\mu > 0$ and $D(\cdot, \cdot)$ is Hausdorff metric defined on $CB(E)$.

Definition 4.3. Let $F : E \rightarrow CB(E)$ be a multivalued mapping. A nonlinear mapping $N : E \times E \rightarrow E$ is said to be relaxed accretive with respect to the first argument of map F , if there exists a constant $\alpha > 0$ such that

$$\langle N(u_n, \cdot) - N(u_{n-1}, \cdot), j_p(x_n - x_{n-1}) \rangle \geq -\alpha \|x_n - x_{n-1}\|^p;$$

and N is Lipschitz continuous with respect to the first argument if

$$\|N(u, \cdot) - N(y, \cdot)\| \leq \sigma \|x - y\|, \quad \text{for } x, y \in E,$$

where $\sigma > 0$ is a constant.

Similarly, we define the Lipschitz continuity of N with respect to second argument.



Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 14 of 24

5. Main Results

In this section, we show that if E is a p -uniformly smooth Banach space, then the iterative process converges strongly to the given problem (2.1).

Theorem 5.1. *Let E be a p -uniformly smooth real Banach space with $\rho_E(\tau) \leq c\tau^p$ for some $c > 0$, $0 < \tau < \infty$ and $1 < p < \infty$. Let X be a closed convex cone of E . Let $m, g : E \rightarrow E$ be the two single-valued mappings, $F, G, T : E \rightarrow CB(E)$ the multivalued mappings. Let $K : E \rightarrow 2^E$ such that $K(z) = m(z) + X$ for all $x \in E$, $z \in T(x)$ and the following conditions hold.*


- (i) g and m are Lipschitz continuous;
- (ii) g is strongly accretive;
- (iii) F, G and T are D -Lipschitz continuous;
- (iv) N is Lipschitz continuous with respect to the first as well as the second argument;
- (v) N is p -relaxed accretive with respect to the first argument with mapping $F : E \rightarrow CB(E)$;

(vi)

$$q + (1 + p\alpha\tau + pc2^{2p+1}\tau^p\sigma^p\eta^p)^{1/p} + \tau\delta\xi < 1$$

and

$$(5.1) \quad q = 2(1 - p\kappa + c2^{2p+1}p\beta^p)^{1/p} + 2\mu\rho.$$



Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents




Go Back

Close

Quit

Page 15 of 24



Title Page

Contents



Go Back

Close

Quit

Page 16 of 24


Then for any $x_0 \in E$, $u_0 \in F(x_0)$, $v_0 \in G(x_0)$ and $z_0 \in T(x_0)$ the sequences x_n, u_n, v_n and z_n generated by Algorithm 1, converge strongly to some $x \in E$, $u \in F(x)$, $v \in G(x)$ and $z \in T(x)$, which solve the problem (2.1).

Proof. By the iterative schemes (4.1) and Definition 3.1, we have

$$\begin{aligned}
 & \|x_{n+1} - x_n\| \\
 &= \|x_n - g(x_n) + m(z_n) + Q_X[g(x_n) - \tau N(u_n, v_n) - m(z_n)] \\
 &\quad - x_{n-1} + g(x_{n-1}) - m(z_{n-1}) \\
 &\quad - Q_X[g(x_{n-1}) - \tau N(u_{n-1}, v_{n-1}) - m(z_{n-1})]\| \\
 &\leq \|x_n - x_{n-1} - (g(x_n) - g(x_{n-1}))\| + \|m(z_n) - m(z_{n-1})\| \\
 &\quad + \|Q_X[g(x_n) - \tau N(u_n, v_n) - m(z_n)] \\
 &\quad\quad - Q_X[g(x_{n-1}) - \tau N(u_{n-1}, v_{n-1}) - m(z_{n-1})]\| \\
 &\leq 2\|x_n - x_{n-1} - (g(x_n) - g(x_{n-1}))\| + 2\|m(z_n) - m(z_{n-1})\| \\
 &\quad + \|x_n - x_{n-1} - \tau(N(u_n, v_n) - N(u_{n-1}, v_{n-1}))\| \\
 &\leq 2\|x_n - x_{n-1} - (g(x_n) - g(x_{n-1}))\| + 2\|m(z_n) - m(z_{n-1})\| \\
 (5.2) \quad & + \|x_n - x_{n-1} - \tau(N(u_n, v_n) - N(u_{n-1}, v_{n-1}))\| \\
 & + \tau\|N(u_{n-1}, v_n) - N(u_{n-1}, v_{n-1})\|.
 \end{aligned}$$

By Lemmas 3.2, 3.6, p -strongly accretive, Lipschitz continuity of g and $j_p \in J_p(x + y)$, we have

$$\begin{aligned}
 & \|x_n - x_{n-1} - (g(x_n) - g(x_{n-1}))\|^p \\
 & \leq \|x_n - x_{n-1}\|^p + p\langle -(g(x_n) - g(x_{n-1})), j_p \rangle
 \end{aligned}$$



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 17 of 24

$$\begin{aligned}
 &\leq \|x_n - x_{n-1}\|^p - p\langle g(x_n) - g(x_{n-1}), \\
 &\quad j_p(x_n - x_{n-1} - (g(x_n) - g(x_{n-1}))) \rangle \\
 &\leq \|x_n - x_{n-1}\|^p - p\langle g(x_n) - g(x_{n-1}), j_p(x_n - x_{n-1}) \rangle \\
 &\quad - p\langle g(x_n) - g(x_{n-1}), j_p(x_n - x_{n-1} \\
 &\quad - (g(x_n) - g(x_{n-1}))) - j_p(x_n - x_{n-1}) \rangle \\
 &\leq \|x_n - x_{n-1}\|^p - p^k \|x_n - x_{n-1}\|^p \\
 &\quad + 2pd^p \rho_E \left(\frac{4\|g(x_n) - g(x_{n-1})\|}{d} \right) \\
 &\leq \|x_n - x_{n-1}\|^p - p^k \|x_n - x_{n-1}\|^p \\
 &\quad + \frac{2pd^p c4^p \|g(x_n) - g(x_{n-1})\|^p}{d^p} \\
 &\leq \|x_n - x_{n-1}\|^p - pk \|x_n - x_{n-1}\|^p + 2^{2p+1} cp\beta^p \|x_n - x_{n-1}\|^p \\
 (5.3) \quad &\leq (1 - pk + c2^{2p+1} p\beta^p) \|x_n - x_{n-1}\|^p.
 \end{aligned}$$

By the Lipschitz continuity of m and D -Lipschitz continuity of T , we have

$$\begin{aligned}
 (5.4) \quad \|m(z_n) - m(z_{n-1})\| &\leq \mu \|z_n - z_{n-1}\| \\
 &\leq \mu(1 + n^{-1})D(T(x_n), T(x_{n-1})) \\
 &\leq \mu(1 + n^{-1})\rho \|x_n - x_{n-1}\|.
 \end{aligned}$$

Since F is η -Lipschitz continuous, G is ξ -Lipschitz continuous and N is Lipschitz continuous with respect to the first and second arguments with positive constants σ and δ respectively. Using a similar argument to that of Xiaolin He [23], we have for every $u_n, u'_n \in F(x_n)$, $N(u_n, v) = N(u'_n, v)$. On the

other hand $u_{n-1} \in F(x_{n-1})$, and from the definition of Hausdorff metric and compactness of $F(x_n)$, there is a $u'_n \in F(x_n)$ such that

$$\|u'_n - u_{n-1}\| \leq D(F(x_n), F(x_{n-1})).$$

Hence


$$\begin{aligned} \|N(u_n, v_n) - N(u_{n-1}, v_n)\| &= \|N(u'_n, v_n) - N(u_{n-1}, v_n)\| \\ &\leq \sigma \|u'_n - u_{n-1}\| \\ &\leq \sigma(1 + n^{-1})D(F(x_n), F(x_{n-1})) \\ (5.5) \quad &\leq \sigma(1 + n^{-1})\eta \|x_n - x_{n-1}\|. \end{aligned}$$

Similarly, we get

$$(5.6) \quad \|N(u_{n-1}, v_n) - N(u_{n-1}, v_{n-1})\| \leq \delta \xi (1 + n^{-1}) \|x_n - x_{n-1}\|.$$

By using Lemma 3.6, the p -relaxed accretive mapping and (5.5), we have

$$\begin{aligned} &\|x_n - x_{n-1} - \tau(N(u_n, v_n) - N(u_{n-1}, v_n))\|^p \\ &\leq \|x_n - x_{n-1}\|^p \\ &\quad - p\tau \langle N(u_n, v_n) - N(u_{n-1}, v_n), \\ &\quad j_p(x_n - x_{n-1} - \tau(N(u_n, v_n) - N(u_{n-1}, v_n))) \rangle \\ &\leq \|x_n - x_{n-1}\|^p - p\tau \langle N(u_n, v_n) - N(u_{n-1}, v_n), j_p(x_n - x_{n-1}) \rangle \\ &\quad - p \langle \tau(N(u_n, v_n) - N(u_{n-1}, v_n)), \\ &\quad j_p(x_n - x_{n-1} - \tau(N(u_n, v_n) - N(u_{n-1}, v_n))) - j_p(x_n - x_{n-1}) \rangle \end{aligned}$$



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents




Go Back

Close

Quit

Page 18 of 24



$$\begin{aligned}
 &\leq \|x_n - x_{n-1}\|^p + p\tau\alpha\|x_n - x_{n-1}\|^p \\
 &\quad + 2pd^p\rho_E\left(\frac{\tau 4\|N(u_n, v_n) - N(u_{n-1}, v_n)\|}{d}\right) \\
 &\leq \|x_n - x_{n-1}\|^p + p\alpha\tau\|x_n - x_{n-1}\|^p \\
 &\quad + 2p\tau^p c 4^p\|N(u_n, v_n) - N(u_{n-1}, v_n)\|^p \\
 &\leq \|x_n - x_{n-1}\|^p + p\alpha\tau\|x_n - x_{n-1}\|^p \\
 &\quad + p\tau^p c 2^{2p+1}(1+n^{-1})^p\sigma^p\eta^p\|x_n - x_{n-1}\|^p \\
 (5.7) \quad &\leq (1 + p\alpha\tau + pc2^{2p+1}(1+n^{-1})^p\tau^p\sigma^p\eta^p)\|x_n - x_{n-1}\|^p.
 \end{aligned}$$

Now from (5.2) – (5.7), we get

$$\begin{aligned}
 &\|x_n - x_{n-1}\| \\
 &\leq 2(1 - pk + c2^{2p+1}p\beta^p)^{1/p}\|x_n - x_{n-1}\| \\
 &\quad + 2\mu\rho(1 + n^{-1})\|x_n - x_{n-1}\| \\
 &\quad + (1 + p\alpha\tau + pc2^{2p+1}(1 + n^{-1})^p\tau^p\sigma^p\eta^p)^{1/p}\|x_n - x_{n-1}\| \\
 &\quad + \tau\delta\xi(1 + n^{-1})\|x_n - x_{n-1}\| \\
 &\leq [2(1 - pk + c2^{2p+1}p\beta^p)^{1/p} + 2\mu\rho(1 + n^{-1}) \\
 &\quad + (1 + p\alpha\tau + pc2^{2p+1}(1 + n^{-1})^p\tau^p\sigma^p\eta^p)^{1/p} + \tau\delta\xi(1 + n^{-1})] \\
 &\leq [q_n + (1 + p\alpha\tau + pc2^{2p+1}(1 + n^{-1})^p\tau^p\sigma^p\eta^p)^{1/p} \\
 &\quad + \tau\delta\xi(1 + n^{-1})]\|x_n - x_{n-1}\| \\
 (5.8) \quad &\leq \theta_n\|x_n - x_{n-1}\|,
 \end{aligned}$$

Title Page

Contents



Go Back

Close

Quit

Page 19 of 24

where

$$\theta_n = q_n + (1 + p\alpha\tau + pc2^{2p+1}(1 + n^{-1})^p\tau^p\sigma^p\eta^p)^{1/p} + \tau\delta\xi(1 + n^{-1})$$

and

$$(5.9) \quad q_n = 2(1 - pk + c2^{2p+1}p\beta^p)^{1/p} + 2\mu\rho(1 + n^{-1}).$$

Letting

$$(5.10) \quad \theta = q + (1 + p\alpha\tau + pc2^{2p+1}\tau^p\sigma^p\eta^p)^{1/p} + \tau\delta\xi$$

and


$$q = 2(1 - pk + c2^{2p+1}p\beta^p)^{1/p} + 2\mu\rho.$$

We know that $\theta_n \rightarrow \theta$ as $n \rightarrow \infty$. From condition (5.1), it follows that $\theta < 1$. Hence $\theta_n < 1$, for n sufficiently large. Consequently $\{x_n\}$ is a Cauchy sequence and this converges to some $x \in E$. By Algorithm 1 and the D -Lipschitz continuity of F, G and T , it follows that

$$\begin{aligned} \|u_n - u_{n-1}\| &\leq (1 + n^{-1})D(F(x_n), F(x_{n-1})) \\ &\leq (1 + n^{-1})\eta\|x_n - x_{n-1}\|, \end{aligned}$$

$$\begin{aligned} \|v_n - v_{n-1}\| &\leq (1 + n^{-1})D(G(x_n), G(x_{n-1})) \\ &\leq (1 + n^{-1})\xi\|x_n - x_{n-1}\|, \end{aligned}$$

$$\begin{aligned} \|z_n - z_{n-1}\| &\leq (1 + n^{-1})D(T(x_n), T(x_{n-1})) \\ &\leq (1 + n^{-1})\rho\|x_n - x_{n-1}\|, \end{aligned}$$



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 20 of 24

which means that $\{u_n\}$, $\{v_n\}$ and $\{z_n\}$ are all Cauchy sequences in E . Therefore there exist $u \in E$, $v \in E$ and $z \in E$ such that $u_n \rightarrow u$, $v_n \rightarrow v$ and $z_n \rightarrow z$ as $n \rightarrow \infty$. Since g, m, F, G, T, N and Q_X are all continuous, we have

$$x = x - g(x) + Q_X[g(x) - \tau N(u, v) - m(z)].$$


Finally, we prove that $u \in F(x)$. In fact, since $u_n \in F(x_n)$ and

$$\begin{aligned} d(u_n, F(x)) &\leq \max \left\{ d(u_n, F(x)), \sup_{u \in F(x)} d(T(x_n), u) \right\} \\ &\leq \max \left\{ \sup_{y \in F(x_n)} d(y, F(x)), \sup_{u \in F(x)} d(F(x_n), u) \right\} \\ &= D(F(x_n), F(x)), \end{aligned}$$

We have

$$\begin{aligned} d(u, F(x)) &\leq \|u - u_n\| + d(u_n, F(x)) \\ &\leq \|u - u_n\| + D(F(x_n), F(x)) \\ &\leq \|u - u_n\| + \eta \|x_n - x\| \rightarrow 0 \quad \text{as } n \rightarrow \infty, \end{aligned}$$

which implies that $d(u, F(x)) = 0$. Since $F(x) \in CB(E)$, it follows that $u \in F(x)$. Similarly, we can prove that $G(x) \in CB(E)$ i.e., $v \in G(x)$ and $T(x) \in CB(E)$ i.e., $z \in T(x)$. Hence by Theorem 3.5, we get the conclusion. \square



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back


Close

Quit

Page 21 of 24

References

- [1] YA ALBER, Metric and generalized projection operators in Banach spaces, properties and applications, in *Theory and Applications of Non-linear Operators of Monotone and Accretive type*, Kartsatos (Ed.), Marcel Dekker, New York (1996), 15–50.
- [2] E. ASPLUND, Positivity of duality mappings, *Bull. Amer. Math. Soc.*, **73** (1967), 200–203.
- [3] C. BAIOCCHI AND A. CAPELO, *Variational and Quasivariational Inequalities, Application to Free Boundary Problems*, J. Wiley, New York, 1984.
- [4] Y. BANYAMINI AND J. LINDENSTRAUSS, Geometric nonlinear functional analysis, I, *American Mathematical Society Colloquium Publication*, **48** (2000), American Mathematical Society, Providence, R.I.
- [5] F.E. BROWDER, Nonlinear mappings of nonexpansive and accretive type in Banach spaces, *Bull. Amer. Math. Soc.*, **73** (1967), 875–885.
- [6] A. CARBONE AND G. ISAC, The generalized order complementarity problem: Application to economics and an existence result, *Nonlinear Studies*, **5**(2) (1998), 139–151.
- [7] S.S. CHANG, Y.J. CHO, B.S. LEE, J.S. JUNG AND S.M. KANG, Iterative approximation of fixed points and solutions for strongly accretive and strongly pseudo-contractive mappings in Banach spaces, *J. Math. Anal. Appl.*, **224** (1998), 149–165.



Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces

M. Firdosh Khan and Salahuddin

Title Page

Contents




Go Back

Close

Quit

Page 22 of 24

- [8] S.S. CHANG, J.K. KIM AND K.H. KIM, On the existence and iterative approximation problems of solutions for set-valued variational inclusions in Banach spaces, *J. Math. Anal. Appl.*, **268** (2002), 89–108.
- [9] J.Y. CHEN, N.C. WONG AND J.C. YAO, Algorithm for generalized co-complementarity problems in Banach spaces, *Comput. Math. Appl.*, **43** (2002), 49–54.
- [10] R.W. COTTLE AND G.B. DANTZING, Complementarity pivot theory of mathematical programming, *Linear Algebra Appl.*, **1** (1968), 163–185.
- [11] R.W. COTTLE J.S. PANG AND R.E. STONE, *The Linear Complementarity Problems*, Academic Press, New York, 1992.
- [12] L. DENG AND X.P. DING, Iterative approximation of Lipschitz strictly pseudo contractive mappings in uniformly smooth Banach spaces, *Nonlinear Analysis, TMA*, **24**(7) (1995), 981–987.
- [13] K. GOEBEL AND S. REICH, *Uniform Convexity, Hyperbolic Geometry and Nonexpansive Mappings*, Marcel Dekker, New York, 1984.
- [14] S.J. HABELTLER AND A.L. PRICE, Existence theory for generalized non-linear complementarity problems, *J Optim. Theory Appl.*, **7** (1971), 223–239.
- [15] D.H. HYERS, G. ISAC AND TH. M. RASSIAS, *Topics in Nonlinear Analysis and Applications*, World Scientific, Singapore, New Jersey, London, Hong Kong, 1997.



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents




Go Back

Close

Quit

Page 23 of 24

- [16] C.R. JOU AND J.C. YAO, Algorithm for generalized multivalued variational inequalities in Hilbert spaces, *Computers Math. Appl.*, **25**(9) (1993), 7–13.
- [17] S. KARAMARDIAN, Generalized complementarity problems, *J. Optim. Theo. Appl.*, **8** (1971), 161–168.
- [18] T. KATO, Nonlinear semigroups and evolution equations, *J. Math. Soc. Japan*, **18/19** (1967), 508–520.
- [19] C.E. LEMKE, Bimatrix equilibrium point and mathematical programming, *Management Science*, **11** (1965), 681–689.
- [20] S.B. NADLER Jr., Multivalued contraction mappings, *Pacific J. Math.*, **30** (1969), 475–488.
- [21] M.A. NOOR, On the nonlinear complementarity problem, *J. Math. Anal. Appl.*, **13** (1987), 455–460.
- [22] M.A. NOOR, Nonlinear quasi-complementarity problems, *Appl. Math. Lett.*, **2**(3)(1980), 251–254.
- [23] XIAOLIN HE, On Φ -strongly accretive mappings and some set-valued variational problems, *J. Math. Anal. Appl.*, **277**(2003), 504–511.
- [24] Z.B. XU AND G.F. ROACH, Characteristic inequalities in uniformly convex and uniformly smooth Banach spaces, *J. Math. Anal. Appl.*, **157** (1991), 189–210.



**Generalized
Co-Complementarity Problems
in p -Uniformly Smooth Banach
Spaces**

M. Firdosh Khan and Salahuddin

Title Page

Contents



Go Back

Close

Quit

Page 24 of 24