

RATE OF CONVERGENCE FOR A GENERAL SEQUENCE OF DURRMEYER TYPE OPERATORS

NIRAJ KUMAR

School of Applied Sciences
Netaji Subhas Institute of Technology
Sector 3, Dwarka
New Delhi 110045, India.

*E*Mail: neeraj@nsit.ac.in



volume 5, issue 3, article 79,
2004.

*Received 21 January, 2004;
accepted 14 June, 2004.*

Communicated by: C.P. Niculescu

[Abstract](#)

[Contents](#)




[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

In the present paper we give the rate of convergence for the linear combinations of the generalized Durrmeyer type operators which includes the well known Szasz-Durrmeyer operators and Baskakov-Durrmeyer operators as special cases.

2000 Mathematics Subject Classification: 41A16, 41A25.

Key words: Szasz-Durrmeyer operators, Baskakov-Durrmeyer operators, Rate of convergence, Order of approximation.

The author is extremely grateful to the referee for making valuable suggestions leading to the better presentation of the paper.

Contents

1	Introduction	3
2	Auxiliary Results	7
3	Rate Of Convergence	20
	References	

Rate Of Convergence For A General Sequence Of Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 2 of 23

1. Introduction

Durrmeyer [1] introduced the integral modification of Bernstein polynomials so as to approximate Lebesgue integrable functions on the interval $[0, 1]$. We now consider the general family of Durrmeyer type operators, which is defined by


$$(1.1) \quad S_n(f(t), x) = (n - c) \sum_{v=0}^{\infty} p_{n,v}(x) \int_0^{\infty} p_{n,v}(t) f(t) dt, \quad x \in I$$

where $n \in \mathbb{N}$, $n > \max\{0, -c\}$ and $p_{n,v}(x) = (-1)^v \frac{x^v}{v!} \varphi_n^{(v)}(x)$. Also $\{\phi_n\}$ is a sequence of real functions having the following properties on $[0, a]$, where $a > 0$ and for all $n \in \mathbb{N}$, $v \in \mathbb{N} \cup \{0\}$, we have

- (I) $\phi_n \in C^\infty[0, a]$, $\phi_n(0) = 1$.
- (II) ϕ_n is complete monotonic.
- (III) There exist $c \in \mathbb{N} : \phi_n^{(v+1)} = -n\phi_{n+c}^{(v)}$, $n > \max\{0, -c\}$.

Some special cases of the operators (1.1) are as follows:

1. If $c = 0$, $\phi_n(x) = e^{-nx}$, we get the Szasz-Durrmeyer operator.
2. If $c = 1$, $\phi_n(x) = (1 + x)^{-n}$, we obtain the Baskakov-Durrmeyer operator.
3. If $c > 1$, $\phi_n(x) = (1 + cx)^{\frac{-n}{c}}$, we obtain a general Baskakov-Durrmeyer operator.



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit


Page 3 of 23

4. If $c = -1$, $\phi_n(x) = (1-x)^n$, we obtain the Bernstein-Durrmeyer operator.

Very recently Srivastava and Gupta [9] studied a similar type of operators and obtained the rate of convergence for functions of bounded variation. It is easily verified that the operators (1.1) are linear positive operators and these operators reproduce the constant ones, while the operators studied in [9] reproduce every linear functional for the case $c = 0$. Several researchers studied different approximation properties on the special cases of the operators (1.1), the pioneer work on Durrmeyer type operators is due to S. Guo [3], Vijay Gupta (see e. g. [4], [5]), R. P. Sinha et al. [8] and Wang and Guo [11], etc. It turns out that the order of approximation for such type of Durrmeyer operators is at best $O(n^{-1})$, how so ever smooth the function may be. In order to improve the order of approximation, we have to slacken the positivity condition of the operators, for this we consider the linear combinations of the operators (1.1). The technique of linear combinations is described as follows:

$$S_{n,k}(f, x) = \begin{vmatrix} 1 & d_0^{-1} & d_0^{-2} & \dots & d_0^{-k} \\ 1 & d_1^{-1} & d_1^{-2} & \dots & d_1^{-k} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ 1 & d_k^{-1} & d_k^{-2} & \dots & d_k^{-k} \end{vmatrix}^{-1} \begin{vmatrix} S_{d_0n}(f, x) & d_0^{-1} & d_0^{-2} & \dots & d_0^{-k} \\ S_{d_1n}(f, x) & d_1^{-1} & d_1^{-2} & \dots & d_1^{-k} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ S_{d_kn}(f, x) & d_k^{-1} & d_k^{-2} & \dots & d_k^{-k} \end{vmatrix}.$$

Such types of linear combinations were first considered by May [7] to improve the order of approximation of exponential type operators. In the alternative form



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 4 of 23

the above linear combinations can be defined as

$$S_{n,k}(f, x) = \sum_{j=0}^k c(j, k) S_{d_j n}(f, x),$$

where


$$C(j, k) = \prod_{\substack{i=0 \\ i \neq j}}^k \frac{d_j}{d_j - d_i}, k \neq 0 \quad \text{and} \quad C(0, 0) = 1.$$

If $f \in L_p [0, \infty)$, $1 \leq p < \infty$ and $0 < a_1 < a_3 < a_2 < b_2 < b_3 < b_1 < \infty$ and $I_i = [a_i, b_i]$, $i = 1, 2, 3$, the Steklov mean $f_{\eta, m}$ of m^{th} order corresponding to f is defined as

$$f_{\eta, m}(t) = \eta^{-m} \int_{-\frac{\eta}{2}}^{\frac{\eta}{2}} \cdots \int_{-\frac{\eta}{2}}^{\frac{\eta}{2}} \left(f(t) + (-1)^{m-1} \Delta_{\sum_{i=1}^m t_i} f(t) \right) \prod_{i=1}^m dt_i; \quad t \in I_1.$$

It can be verified [6, 10] that

- (i) $f_{\eta, m}$ has derivative up to order m , $f_{\eta, m}^{(m-1)} \in AC(I_1)$ and $f_{\eta, m}^{(m)}$ exist almost every where and belongs to $L_p(I_1)$;
- (ii) $\|f_{\eta, m}^{(r)}\|_{L_p(I_1)} \leq K_1 \eta^{-r} \omega_r(f, \eta, p, I_1)$, $r = 1, 2, 3, \dots, m$;
- (iii) $\|f - f_{\eta, m}\|_{L_p(I_2)} \leq K_2 \omega_m(f, \eta, p, I_1)$;
- (iv) $\|f_{\eta, m}^m\|_{L_p(I_2)} \leq K_3 \eta^m \|f\|_{L_p(I_1)}$



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close


Quit

Page 5 of 23

$$(v) \left\| f_{\eta, m}^{(m)} \right\|_{L_p(I_1)} \leq K_4 \eta^{-r} \omega_r(f, \eta, p, I_1)$$

where K'_i 's, $i = 1, 2, 3, 4$ are certain constants independent of f and η .

In the present paper we establish the rate of convergence for the combinations of the generalized Durrmeyer type operators in L_p -norm.



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 6 of 23

2. Auxiliary Results

To prove the rate of convergence we need the following lemmas:

Lemma 2.1. Let $m \in \mathbb{N} \cup \{0\}$, also $\mu_{n,m}(x)$ is the m^{th} order central moment defined by

$$\mu_{n,m} \equiv S_n((t-x)^m, x) = (n-c) \sum_{v=0}^{\infty} p_{n,v}(x) \int_0^{\infty} p_{n,v}(t)(t-x)^m dt,$$

then

(i) $\mu_{n,m}(x)$ is a polynomial in x of degree m .

(ii) $\mu_{n,m}(x)$ is a rational function in n and for each $0 \leq x < \infty$ $\mu_{n,m}(x) = O(n^{-[(m+1)/2]})$.

Remark 2.1. Using Hölder's inequality, it can be easily verified that $S_n(|t-x|^r, x) = O(n^{-r/2})$ for each $r > 0$ and for every fixed $0 \leq x < \infty$.


Lemma 2.2. For sufficiently large n and $q \in \mathbb{N}$, there holds

$$S_{n,k}((t-x)^q, x) = n^{-(k+1)} \{F(q, k, x) + o(1)\},$$

where $F(q, k, x)$ are certain polynomials in x of degree q and $0 \leq x < \infty$ is arbitrary but fixed.

Proof. For sufficiently large values of n we can write, from Lemma 2.1

$$S_n((t-x)^q, x) = \frac{Q_0(x)}{n^{[(q+1)/2]}} + \frac{Q_1(x)}{n^{[(q+1)/2]+1}} + \dots + \frac{Q_{[q/2]}(x)}{n^q},$$



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 7 of 23

where $Q_i(x), i = 0, 1, 2, \dots$ are certain polynomials in x of at most degree q .

Therefore $S_{n,k}((t-x)^q, x)$ is given by


$$\begin{vmatrix} 1 & d_0^{-1} & d_0^{-2} & \dots & d_0^{-k} \\ 1 & d_1^{-1} & d_1^{-2} & \dots & d_1^{-k} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ 1 & d_k^{-1} & d_k^{-2} & \dots & d_k^{-k} \end{vmatrix}^{-1} \\ \times \begin{vmatrix} \frac{Q_0(x)}{(d_0n)^{\lfloor (q+1)/2 \rfloor}} + \frac{Q_1(x)}{(d_0n)^{\lfloor (q+1)/2 \rfloor + 1}} + \dots & d_0^{-1} & d_0^{-2} & \dots & d_0^{-k} \\ \frac{Q_0(x)}{(d_1n)^{\lfloor (q+1)/2 \rfloor}} + \frac{Q_1(x)}{(d_1n)^{\lfloor (q+1)/2 \rfloor + 1}} + \dots & d_1^{-1} & d_1^{-2} & \dots & d_1^{-k} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \frac{Q_0(x)}{(d_kn)^{\lfloor (q+1)/2 \rfloor}} + \frac{Q_1(x)}{(d_kn)^{\lfloor (q+1)/2 \rfloor + 1}} + \dots & d_k^{-1} & d_k^{-2} & \dots & d_k^{-k} \end{vmatrix} \\ = n^{-(k+1)} \{F(q, k, x) + o(1)\}, \quad 0 \leq x < \infty.$$

□

Lemma 2.3 ([4]). Let $1 \leq p < \infty$, $f \in L_p[a, b]$, $f^{(m)} \in AC[a, b]$ and $f^{(m+1)} \in L_p[a, b]$, then

$$\|f^{(j)}\|_{L_p[a,b]} \leq C \left\{ \|f^{(m+1)}\|_{L_p[a,b]} + \|f\|_{L_p[a,b]} \right\},$$

$j = 1, 2, 3, \dots, m$ and C is a constant depending on j, p, m, a, b .



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 8 of 23

Lemma 2.4. Let $f \in L_p[0, \infty)$, $p > 1$. If $f^{(2k+1)} \in AC(I_1)$ and $f^{(2k+2)} \in L_p(I_1)$, then for all n sufficiently large

$$(2.1) \quad \|S_{n,k}(f, \cdot) - f\|_{L_p(I_2)} \leq C_1 n^{-(k+1)} \left\{ \|f^{(2k+2)}\|_{L_p(I_2)} + \|f\|_{L_p[0, \infty)} \right\}.$$

Also if $f \in L_1[0, \infty)$, $f^{(2k+1)} \in L_1(I_1)$ with $f^{(2k)} \in AC(I_1)$ and $f^{(2k+1)} \in BV(I_1)$, then for all n sufficiently large


$$(2.2) \quad \|S_{n,k}(f, \cdot) - f\|_{L_p(I_2)} \leq C_2 n^{-(k+1)} \left\{ \|f^{(2k+1)}\|_{BV(I_1)} + \|f^{(2k+1)}\|_{L_1(I_2)} + \|f\|_{L_1[0, \infty)} \right\}.$$

Proof. First let $p > 1$. By the hypothesis, for all $t \in [0, \infty)$ and $x \in I_2$, we have

$$\begin{aligned} S_{n,k}(f, x) - f(x) &= \sum_{i=1}^{2k+1} \frac{f^{(i)}(x)}{i!} S_{n,k}((t-x)^i, x) \\ &\quad + \frac{1}{(2k+1)!} S_{n,k}(\phi(t) \int_x^t (t-w)^{2k+1} f^{(2k+2)}(w) dw, x) \\ &\quad + S_{n,k}(F(t, x)(1-\phi(t)), x) \\ &= E_1 + E_2 + E_3, \quad \text{say,} \end{aligned}$$

where $\phi(t)$ denotes the characteristic function of I_1 and

$$F(t, x) = f(t) - \sum_{i=0}^{2k+1} \frac{(t-x)^i}{i!} f^{(i)}(x).$$



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 9 of 23

Applying Lemma 2.2 and Lemma 2.3, we have


$$\begin{aligned} \|E_1\|_{L_p(I_2)} &\leq C_3 n^{-(k+1)} \sum_{i=1}^{2k+1} \|f^{(i)}\|_{L_p(I_2)} \\ &\leq C_4 n^{-(k+1)} \left\{ \|f\|_{L_p(I_2)} + \|f^{(2k+2)}\|_{L_p(I_2)} \right\}. \end{aligned}$$

Next we estimate E_2 . Let H_f be the Hardy Littlewood maximal function [10] of $f^{(2k+2)}$ on I_1 , using Hölder's inequality and Lemma 2.1, we have

$$\begin{aligned} |E_2| &\leq \frac{1}{(2k+2)!} S_{n,k} \left(\phi(t) \left| \int_x^t |t-w|^{2k+1} |f^{(2k+2)}(w)| dw \right|, x \right) \\ &\leq \frac{1}{(2k+2)!} S_{n,k} \left(\phi(t) |t-x|^{2k+1} \left| \int_x^t |f^{(2k+2)}(w)| dw \right|, x \right) \\ &\leq \frac{1}{(2k+2)!} S_{n,k} \left(\phi(t) |t-x|^{2k+2} |H_f(t)|, x \right) \\ &\leq \frac{1}{(2k+2)!} \left\{ S_{n,k} \left(\phi(t) |t-x|^{q(2k+2)}, x \right) \right\}^{\frac{1}{q}} \left\{ S_{n,k} \left(\phi(t) |H_f(t)|^p, x \right) \right\}^{\frac{1}{p}} \\ &\leq C_5 n^{-(k+1)} \left(\sum_{j=0}^k C(j,k) \int_{a_1}^{b_1} (d_j n - c) \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |H_f(t)|^p dt \right)^{\frac{1}{p}}. \end{aligned}$$

Next applying Fubini's theorem, we have

$$\begin{aligned} \|E_2\|_{L_p(I_2)}^p &\leq C_6 n^{-p(k+1)} \sum_{j=0}^k C(j,k) \int_{a_2}^{b_2} \int_{a_1}^{b_1} (d_j n - c) \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |H_f(t)|^p dt dx \end{aligned}$$



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 10 of 23

$$\begin{aligned} &\leq C_6 n^{-p(k+1)} \sum_{j=0}^k C(j, k) \int_{a_1}^{b_1} \left(\int_{a_2}^{b_2} (d_j n - c) \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) dx \right) |H_f(t)|^p dt \\ &\leq C_7 n^{-p(k+1)} \|H_f\|_{L_p(I_1)}^p \leq C_8 n^{-p(k+1)} \|f^{(2k+2)}\|_{L_p(I_1)}^p. \end{aligned}$$

Therefore


$$\|E_2\|_{L_p(I_2)} \leq C_8 n^{-(k+1)} \|f^{(2k+2)}\|_{L_p(I_1)}.$$

For $t \in [0, \infty) \setminus [a_1, b_1]$, $x \in I_2$ there exists a $\delta > 0$ such that $|t - x| \geq \delta$. Thus

$$\begin{aligned} E_3 &\equiv S_{n,k}(F(t, x)(1 - \phi(t)), x) \\ &\leq \delta^{-(2k+2)} \sum_{j=0}^k C(j, k) S_{d_j n}(|F(t, x)|(t - x)^{2k+2}, x) \\ &\leq \delta^{-(2k+2)} \sum_{j=0}^k C(j, k) \left[S_{d_j n}(|f(t)|(t - x)^{2k+2}, x) \right. \\ &\quad \left. + \sum_{i=0}^{2k+1} \frac{|f^{(i)}(x)|}{i!} S_{d_j n}(|t - x|^{2k+2+i}, x) \right] \\ &= E_{31} + E_{32}, \quad \text{say.} \end{aligned}$$

Applying Hölder's inequality and Lemma 2.1, we have

$$|E_{31}| \leq \delta^{-(2k+2)} \sum_{j=0}^k C(j, k) \{S_{d_j n}(|f(t)|^p, x)\}^{\frac{1}{p}} \{S_{d_j n}(|t - x|^{q(2k+2)}, x)\}^{\frac{1}{q}}$$



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 11 of 23

$$\leq C_9 \sum_{j=0}^k C(j, k) \{S_{d_j n}(|f(t)|^p, x)\}^{\frac{1}{p}} \frac{1}{(d_j n)^{k+1}}.$$

Finally by Fubini's theorem, we obtain

$$\begin{aligned} \|E_{31}\|_{L_p(I_2)}^p &= \int_{a_2}^{b_2} |E_{31}|^p dx \\ &\leq C_9 \sum_{j=0}^k C(j, k) (d_j n)^{-p(k+1)} \\ &\quad \times \int_{a_2}^{b_2} \int_0^\infty (d_j n - c) p_{d_j n, v}(x) p_{d_j n, v}(t) |f(t)|^p dt dx \\ &\leq C_{10} n^{-p(k+1)} \|f\|_{L_p[0, \infty)}^p. \end{aligned}$$

Again by Lemma 2.3, we have


$$\|E_{32}\|_{L_p(I_2)} \leq C_{11} n^{-(k+1)} \left\{ \|f\|_{L_p(I_2)} + \|f^{(2k+2)}\|_{L_p(I_2)} \right\}.$$

Thus

$$\|E_3\|_{L_p(I_2)} \leq C_{12} n^{-(k+1)} \left\{ \|f\|_{L_p[0, \infty)} + \|f^{(2k+2)}\|_{L_p(I_2)} \right\}.$$

Combining the estimates of E_1, E_2, E_3 , we get (2.1).

Next suppose $p = 1$. By the hypothesis, for almost all $x \in I_2$ and for all



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 12 of 23

$t \in [0, \infty)$, we have

$$\begin{aligned} S_{n,k}(f, x) - f(x) &= \sum_{i=1}^{2k+1} \frac{f^{(i)}(x)}{i!} S_{n,k}((t-x)^i, x) \\ &\quad + \frac{1}{(2k+1)!} S_{n,k}(\phi(t) \int_x^t (t-w)^{2k+1} f^{(2k+1)}(w) dw, x) \\ &\quad + S_{n,k}(F(t, x)(1-\phi(t)), x) \\ &= M_1 + M_2 + M_3, \quad \text{say,} \end{aligned}$$

where $\phi(t)$ denotes the characteristic function of I_1 and

$$F(t, x) = f(t) - \sum_{i=0}^{2k+1} \frac{(t-x)^i}{i!} f^{(i)}(x),$$


for almost all $x \in I_2$ and $t \in [0, \infty)$.

Applying Lemma 2.2 and Lemma 2.3, we have

$$\|M_1\|_{L_1(I_2)} \leq C_{13} n^{-(k+1)} \left\{ \|f\|_{L_1(I_2)} + \|f^{(2k+1)}\|_{L_1(I_2)} \right\}.$$

Next, we have

$$\begin{aligned} &\|M_2\|_{L_1(I_2)} \\ &\leq \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j, k)| \int_{a_2}^{b_2} \int_{a_1}^{b_1} (d_j n - c) \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |t-x|^{2k+1} \\ &\quad \times \left| \int_x^t |df^{(2k+1)}(w)| \right| dt dx. \end{aligned}$$



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 13 of 23

For each $d_j n$ there exists a non negative integer $r = r(d_j n)$ satisfying


$$r(d_j n)^{-1/2} < \max(b_1 - a_2, b_2 - a_1) \leq (r + 1)(d_j n)^{-1/2}.$$

Thus

$$\begin{aligned} \|M_2\|_{L_1(I_2)} &\leq \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j, k)| \sum_{l=0}^r \int_{a_2}^{b_2} \left\{ \int_{x+(l)(d_j n)^{-1/2}}^{x+(l+1)(d_j n)^{-1/2}} \phi(t)(d_j n - c) \right. \\ &\cdot \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |t - x|^{2k+1} \left. \right\} \left(\int_x^{x+(l+1)(d_j n)^{-1/2}} \phi(w) |df^{(2k+1)}(w)| \right) dt dx \\ &+ \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j, k)| \sum_{l=0}^r \int_{a_2}^{b_2} \left\{ \int_{x+(l+1)(d_j n)^{-1/2}}^{x-(l)(d_j n)^{-1/2}} \phi(t)(d_j n - c) \right. \\ &\times \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |t - x|^{2k+1} \left. \right\} \cdot \left(\int_{x-(l+1)(d_j n)^{-1/2}}^x \phi(w) |df^{(2k+1)}(w)| \right) dt dx. \end{aligned}$$

Suppose $\phi_{x,c,s}(w)$ denotes the characteristic function of the interval $[x - c(d_j n)^{-1/2}, x + s(d_j n)^{-1/2}]$, where c, s are nonnegative integers. Then we have

$$\begin{aligned} \|M_2\|_{L_1(I_2)} &\leq \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j, k)| \sum_{l=1}^r \int_{a_2}^{b_2} \left\{ \int_{x+(l)(d_j n)^{-1/2}}^{x+(l+1)(d_j n)^{-1/2}} \phi(t)(d_j n)^2 (d_j n - c) \right. \\ &\times \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) l^{-4} |t - x|^{2k+5} \left. \right\} \end{aligned}$$



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents




Go Back

Close

Quit

Page 14 of 23



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents




Go Back

Close

Quit

Page 15 of 23

$$\begin{aligned}
 & \times \left(\int_x^{x+(l+1)(d_j n)^{-1/2}} \phi(w) \phi_{x,0,l+1}(w) |df^{(2k+1)}(w)| \right) dt dx \\
 & + \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j,k)| \sum_{l=1}^r \int_{a_2}^{b_2} \left\{ \int_{x+(l+1)(d_j n)^{-1/2}}^{x-(l)(d_j n)^{-1/2}} \phi(t) (d_j n)^2 (d_j n - c) \right. \\
 & \times \left. \sum_{v=0}^{\infty} b_{d_j n, v}(x) p_{d_j n, v}(t) l^{-4} |t-x|^{2k+5} \right\} \\
 & \times \left(\int_{x-(l+1)(d_j n)^{-1/2}}^x \phi(w) \phi_{x,l+1,0}(w) |df^{(2k+1)}(w)| \right) dt dx \\
 & \times \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j,k)| \int_{a_2}^{b_2} \left\{ \int_{(d_j n)^{-1/2}}^{a_1+(d_j n)^{-1/2}} \phi(t) (d_j n - c) \right. \\
 & \times \left. \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |t-x|^{2k+1} \right\} \\
 & \times \left(\int_{x-(d_j n)^{-1/2}}^{x+(d_j n)^{-1/2}} \phi(w) \phi_{x,1,1}(w) |df^{(2k+1)}(w)| \right) dt dx \\
 & \leq \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j,k)| \sum_{l=1}^r l^{-4} \int_{a_2}^{b_2} \left\{ \int_{x+(l)(d_j n)^{-1/2}}^{x+(l+1)(d_j n)^{-1/2}} \phi(t) (d_j n - c) \right. \\
 & \times \left. \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |t-x|^{2k+5} \right\}
 \end{aligned}$$



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close


Quit

Page 16 of 23

$$\begin{aligned}
 & \times \left(\int_{a_1}^{b_1} \phi_{x,0,l+1}(w) \left| df^{(2k+1)}(w) \right| \right) dt dx \\
 & + \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j,k)| \sum_{l=1}^r l^{-4} \int_{a_2}^{b_2} \left\{ \int_{x+(l+1)(d_j n)^{-1/2}}^{x-l(d_j n)^{-1/2}} \phi(t)(d_j n - c) \right. \\
 & \times \left. \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |t-x|^{2k+5} \right\} \\
 & \times \left(\int_{a_1}^{b_1} \phi_{x,l+1,0}(w) \left| df^{(2k+1)}(w) \right| \right) dt dx \\
 & + \frac{1}{(2k+1)!} \sum_{j=0}^k |C(j,k)| \int_{a_2}^{b_2} \left\{ \int_{(d_j n)^{-1/2}}^{a_1+(d_j n)^{-1/2}} \phi(t)(d_j n - c) \right. \\
 & \times \left. \sum_{v=0}^{\infty} p_{d_j n, v}(x) p_{d_j n, v}(t) |t-x|^{2k+1} \right\} \\
 & \times \left(\int_{a_1}^{b_1} \phi(w) \phi_{x,1,1}(w) \left| df^{(2k+1)}(w) \right| \right) dt dx.
 \end{aligned}$$

Applying Lemma 2.1 and using Fubini's theorem we get

$$\begin{aligned}
 & \|M_2\|_{L_1(I_2)} \\
 & \leq C_{14} \sum_{j=0}^k |C(j,k)| (d_j n)^{-\frac{2k+1}{2}} \sum_{l=1}^r l^{-4} \left\{ \int_{a_2}^{b_2} \int_{a_1}^{b_1} \phi_{x,0,l+1}(w) \left| df^{(2k+1)}(w) \right| dx \right\}
 \end{aligned}$$



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 17 of 23

$$\begin{aligned}
 & + C_{14} \sum_{j=0}^k |C(j, k)|(d_j n)^{-\frac{2k+1}{2}} \sum_{l=1}^r l^{-4} \int_{a_2}^{b_2} \left\{ \int_{a_1}^{b_1} \phi_{x, l+1, 0}(w) |df^{(2k+1)}(w)| dx \right. \\
 & \left. + \int_{a_1}^{b_1} \phi_{x, 1, 1}(w) |df^{(2k+1)}(w)| dx \right\} \\
 \leq & C_{14} \sum_{j=0}^k |C(j, k)|(d_j n)^{-\frac{2k+1}{2}} \sum_{l=1}^r l^{-4} \left\{ \int_{a_1}^{b_1} \left(\int_{a_2}^{b_2} \phi_{x, 0, l+1}(w) dx \right) |df^{(2k+1)}(w)| \right\} \\
 & + C_{14} \sum_{j=0}^k |C(j, k)|(d_j n)^{-\frac{2k+1}{2}} \sum_{l=1}^r l^{-4} \int_{a_1}^{b_1} \left\{ \int_{a_2}^{b_2} \phi_{x, l+1, 0}(w) |df^{(2k+1)}(w)| dx \right. \\
 & \left. + \int_{a_2}^{b_2} \phi_{x, 1, 1}(w) |df^{(2k+1)}(w)| dx \right\} \\
 \leq & C_{15} \sum_{j=0}^k |C(j, k)|(d_j n)^{-\frac{2k+1}{2}} \sum_{l=1}^r l^{-4} \left\{ \int_{a_1}^{b_1} \left(\int_{w-(l+1)(d_j n)^{-\frac{1}{2}}}^w dx \right) |df^{(2k+1)}(w)| \right\} \\
 & + C_{15} \sum_{j=0}^k |C(j, k)|(d_j n)^{-\frac{2k+1}{2}} \sum_{l=1}^r l^{-4} \int_{a_1}^{b_1} \left\{ \left(\int_w^{w+(l+1)(d_j n)^{-\frac{1}{2}}} dx \right) |df^{(2k+1)}(w)| \right. \\
 & \left. + \int_{a_1}^{b_1} \left(\int_{w-(d_j n)^{-\frac{1}{2}}}^{w+(d_j n)^{-\frac{1}{2}}} dx \right) |df^{(2k+1)}(w)| \right\} \\
 \leq & C_{16} n^{-(k+1)} \|f^{(2k+1)}\|_{B.V.(I_1)}.
 \end{aligned}$$


Finally we estimate M_3 . It is sufficient to choose the expression without the

linear combinations. For all $t \in [0, \infty) \setminus [a_1, b_1]$ and all $x \in I_2$, we choose a $\delta > 0$ such that $|t - x| \geq \delta$. Thus

$$\begin{aligned} & \|S_n(F(t, x)(1 - \phi(t)), x)\|_{L_1(I_2)} \\ & \leq \int_{a_2}^{b_2} \int_0^\infty (n - c) \sum_{v=0}^\infty p_{n,v}(x)p_{n,v}(t) |f(t)| (1 - \phi(t)) dt dx \\ & \leq \sum_{i=0}^{2k+1} \frac{1}{i!} \int_{a_2}^{b_2} \int_0^\infty (n - c) \sum_{v=0}^\infty p_{n,v}(x)p_{n,v}(t) |f^{(i)}(t)| \cdot |t - x|^i (1 - \phi(t)) dt dx \\ & = M_4 + M_5, \quad \text{say.} \end{aligned}$$

For sufficiently large t there exist positive constants C_{17}, C_{18} such that $\frac{(t-x)^{2k+2}}{t^{2k+2}+1} > C_{17}$ for all $t \geq C_{18}, t \in I_2$. Applying Fubini's theorem and Lemma 2.1, we obtain

$$\begin{aligned} M_4 & = \left(\int_0^{C_{18}} \int_{a_2}^{b_2} + \int_{C_{18}}^\infty \int_{a_2}^{b_2} \right) \sum_{v=0}^\infty (n - c) p_{n,v}(x) p_{n,v}(t) |f(t)| (1 - \phi(t)) dt dx \\ & \leq C_{19} n^{-(k+1)} \left(\int_0^{C_{20}} |f(t)| dt \right) \\ & \quad + \frac{1}{C_{17}} \int_{C_{20}}^\infty \int_{a_2}^{b_2} \sum_{v=0}^\infty p_{n,v}(x) p_{n,v}(t) \frac{(t-x)^{2k+2}}{(t^{2k+2}+1)} |f(t)| dx dt \\ & \leq C_{20} n^{-(k+1)} \left\{ \left(\int_0^{C_{20}} |f(t)| dt \right) + \left(\int_{C_{20}}^\infty |f(t)| dt \right) \right\} \\ & \leq C_{21} n^{-(k+1)} \|f\|_{L_1[0, \infty)}. \end{aligned}$$



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 18 of 23


Finally by the Lemma 2.3, we have

$$M_5 \leq C_{22} n^{-(k+1)} \left\{ \|f\|_{L_1(I_2)} + \|f^{(2k+1)}\|_{L_1(I_2)} \right\}.$$

Combining M_4 and M_5 , we obtain

$$M_3 \leq C_{23} n^{-(k+1)} \left\{ \|f\|_{L_1[0,\infty)} + \|f^{(2k+1)}\|_{L_1(I_2)} \right\}.$$

This completes the proof of (2.2) of the lemma. □



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 19 of 23

3. Rate Of Convergence

Theorem 3.1. Let $f \in L_p[0, \infty)$, $p \geq 1$. Then for n sufficiently large

$$\|S_{n,k}(f, \cdot) - f\|_{L_p(I_2)} \leq C_{24} \left\{ \omega_{2k+2}(f, n^{-1/2}, p, I_1) + n^{-(k+1)} \|f\|_{L_p[0, \infty)} \right\},$$

where C_{24} is a constant independent of f and n .

Proof. We can write

$$\begin{aligned} & \|S_{n,k}(f, *) - f\|_{L_p(I_2)} \\ & \leq \|S_{n,k}(f - f_{\eta, 2k+2}, *)\|_{L_p(I_2)} \\ & \quad + \|S_{n,k}(f_{\eta, 2k+2}, *) - f_{\eta, 2k+2}\|_{L_p(I_2)} + \|(f_{\eta, 2k+2} - f)\|_{L_p(I_2)} \\ & = E_1 + E_2 + E_3, \quad \text{say.} \end{aligned}$$

It is well known that

$$\left\| f_{\eta, 2k+2}^{(2k+1)} \right\|_{B.V.(I_3)} = \left\| f_{\eta, 2k+2}^{(2k+2)} \right\|_{L_1(I_3)}.$$

Therefore from Lemma 2.4 ($p > 1$) and ($p = 1$) we have

$$\begin{aligned} E_2 & \leq C_{25} n^{-(k+1)} \left(\left\| f_{\eta, 2k+2}^{(2k+2)} \right\|_{L_p(I_3)} + \|f_{\eta, 2k+2}\|_{L_p[0, \infty)} \right) \\ & \leq C_{26} n^{-(k+1)} \left(n^{-(k+2)} \omega_{2k+2}(f, \eta, p, I_1) + \|f\|_{L_p[0, \infty)} \right), \end{aligned}$$

which follows from the properties of Steklov means.



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 20 of 23

Let $\phi(t)$ be the characteristic function of I_3 , we have

$$\begin{aligned} S_n((f - f_{\eta,2k+2})(t), x) &= S_n(\phi(t)(f - f_{\eta,2k+2})(t), x) + S_n((1 - \phi(t))(f - f_{\eta,2k+2})(t), x) \\ &= E_4 + E_5, \quad \text{say.} \end{aligned}$$

By Hölder's inequality

$$\int_{a_2}^{b_2} |E_4|^p dx \leq \int_{a_2}^{b_2} \int_{a_1}^{b_1} (n - c) \sum_{v=0}^{\infty} p_{n,v}(x)p_{n,v}(t) |(f - f_{\eta,2k+2})(t)|^p dt dx.$$

Applying Fubini's theorem, we have

$$\int_{a_2}^{b_2} |E_4|^p dx \leq \|f - f_{\eta,2k+2}\|_{L_p(I_2)}.$$

Similarly, for all $p \geq 1$

$$\|E_5\|_{L_p(I_2)} \leq C_{27} \eta^{-(k+1)} \|f - f_{\eta,2k+2}\|_{L_p(0,\infty)}.$$

Consequently, via the property of Steklov means, we find that


$$\|S_n(f - f_{\eta,2k+2}, \cdot)\|_{L_p(I_2)} \leq C_{28} \left\{ \omega_{2k+2}(f, \eta, p, I_1) + \eta^{-(k+1)} \|f\|_{L_p[0,\infty)} \right\}.$$

Hence

$$E_1 \leq C_{29} \left\{ \omega_{2k+2}(f, \eta, p, I_1) + \eta^{-(k+1)} \|f\|_{L_p[0,\infty)} \right\}.$$

Thus, with $\eta = n^{-1/2}$, the result follows.

This completes the proof of the theorem. □



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 21 of 23

References

- [1] J.L. DURMEYER, Une Formula d' inversion de la Transformee de Laplace: Application a la Theorie des Moments, *These de 3c cycle, Faculte des Science de l' University de Paris*, 1967.
- [2] S. GOLDBERG AND V. MEIR, Minimum moduli of ordinary differential operators, *Proc. London Math. Soc.*, **23** (3) (1971), 1–15.
- [3] S. GUO, On the rate of convergence of Durrmeyer operator for functions of bounded variation, *J. Approx. Theory*, **51**(1987), 183–192.
- [4] V. GUPTA, Simultaneous approximation by Szasz-Durrmeyers operators, *The Math. Student*, **64** (1-4) (1995), 27–36.
- [5] V. GUPTA, A note on modified Baskakov operators, *Approx.Theory and its Appl.*, **10**(3) (1994), 74–78.
- [6] E. HEWITT AND K. STROMBERG, *Real and Abstract Analysis*, McGraw Hill, New York (1956).
- [7] C.P. MAY, Saturation and inverse theorem for combination of a class of exponential type operators, *Canad. J. Math.*, **28** (1976), 1224–1250.
- [8] R.P. SINHA, P.N. AGRAWAL AND V. GUPTA, On simultaneous approximation by modified Baskakov operators, *Bull Soc. Math Belg. Ser. B.*, **44** (1991), 177–188.
- [9] H.M. SRIVASTAVA AND V. GUPTA, A certain family of summation integral type operators, *Math and Comput. Modelling*, **37** (2003), 1307–1315.



Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators

Niraj Kumar

Title Page

Contents




Go Back

Close

Quit

Page 22 of 23

- [10] A.F. TIMAN, *Theory of Approximation of Functions of a Real Variable*, Hindustan Publ. Corp., Delhi 1966.
- [11] Y. WANG AND S. GUO, Rate of approximation of function of bounded variation by modified Lupas operators, *Bull. Austral. Math. Soc.*, **44** (1991), 177–188.



**Rate Of Convergence For A
General Sequence Of
Durrmeyer Type Operators**

Niraj Kumar

Title Page

Contents



Go Back

Close

Quit

Page 23 of 23