

DERIVATIVES OF CATALAN RELATED SUMS

ANTHONY SOFO

School of Engineering and Science
Victoria University, PO Box 14428 Melbourne
VIC 8001, Australia.

EMail: anthony.sofa@vu.edu.au

Received: 19 March, 2009

Accepted: 29 July, 2009

Communicated by: **J. Sándor**

2000 AMS Sub. Class.: Primary 05A10. Secondary 26A09, 26D20, 33C20.

Key words: Catalan numbers, Binomial coefficients, Estimates, Integral identities.

Abstract: We investigate the representation of sums of the derivative of the reciprocal of Catalan type numbers in integral form. We show that for various parameter values the sums maybe expressed in closed form. Finally we give bounds for the sums under investigation, in terms of the parameters.





Catalan Related Sums

Anthony Sofo

vol. 10, iss. 3, art. 69, 2009

[Title Page](#)

[Contents](#)



Page 1 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	Catalan Related Sums	7



Catalan Related Sums

Anthony Sofo

vol. 10, iss. 3, art. 69, 2009

Title Page

Contents



Page 2 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



1. Introduction

We define $C_n(j)$, the Catalan related numbers, for $j \in \mathbb{N} \cup \{0\}$, $\mathbb{N} = \{1, 2, 3, \dots\}$ as

$$(1.1) \quad C_n(j) = \begin{cases} \frac{1}{n+1} \binom{2n+j}{n} = \frac{1}{2n+j+1} \binom{2n+j+1}{n+1}, & \text{for } n = 1, 2, 3, \dots \\ 1, & \text{for } n = 0, \end{cases}$$

and in particular, from (1.1), the Catalan numbers $C_n = C_n(0)$ are defined by

$$C_n = \begin{cases} \frac{1}{n+1} \binom{2n}{n} = \frac{1}{2n+1} \binom{2n+1}{n}, & \text{for } n = 1, 2, 3, \dots \\ 1, & \text{for } n = 0. \end{cases}$$

Catalan numbers have many representations, see Adamchik [1], in particular Pen-son and Sixdeniers [3], by employing some ideas of the Mellin transform gave the integral representation,

$$(1.2) \quad C_n = \frac{1}{2\pi} \int_0^4 t^{n-1/2} \sqrt{4-t} dt.$$

By the change of variable $tw^2 = 4 - t$, in (1.2) we find that

$$C_n = \frac{2^{2n+2}}{\pi} \int_0^\infty \frac{w^2}{(1+w^2)^{n+2}} dw,$$

which, see Bailey et.al. [2], is closely related to cb_n , the central binomial coefficient

$$cb_n = \binom{2n}{n} = \frac{2^{2n+1}}{\pi} \int_0^\infty \frac{1}{(1+w^2)^{n+1}} dw.$$

Moreover, it can be seen that $cb_n = (n+1)C_n$.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 4 of 16

Go Back

Full Screen

Close

Lemma 1.1. For $j \geq 0$ and $n \in \mathbb{N}$, let the Catalan related numbers

$$C_n(j) = \begin{cases} \frac{1}{n+1} \binom{2n+j}{n} = \frac{1}{2n+j+1} \binom{2n+j+1}{n+1}, & \text{for } n = 1, 2, 3, \dots \\ 1, & \text{for } n = 0 \end{cases}$$

be an analytic function in j then

$$\begin{aligned} (1.3) \quad \frac{d}{dj} \left(\frac{1}{C_n(j)} \right) &= \left(\frac{1}{C_n(j)} \right)' \\ &= -\frac{n+1}{\binom{2n+j}{n}} \sum_{r=1}^n \frac{1}{r+j+n} \\ &= -\frac{n+1}{\binom{2n+j}{n}} [\Psi(1+j+2n) - \Psi(1+j+n)], \end{aligned}$$

where the Psi(or digamma function)

$$\Psi(z) = \frac{d}{dz} \ln(\Gamma(z)) = \frac{(\Gamma(z))'}{\Gamma(z)}$$

and the Gamma function

$$\Gamma(w) = \int_0^{\infty} t^{w-1} e^{-t} dt,$$

for $\Re(w) > 0$.



[Title Page](#)

[Contents](#)



Page 5 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

Proof. For the first derivative of $\frac{1}{C_n(j)}$ with respect to j , let for integer n ,

$$\begin{aligned}\frac{1}{C_n(j)} &= \frac{n+1}{\binom{2n+j}{n}} \\ &= \frac{(n+1)\Gamma(n+1)\Gamma(n+j+1)}{\Gamma(2n+j+1)} \\ &= \frac{(n+1)\Gamma(n+1)}{\prod_{r=1}^n (n+r+j)}.\end{aligned}$$

Taking the logs on both sides we have

$$\ln \left[\frac{1}{C_n(j)} \right] = \ln(n+1) + \ln[\Gamma(n+1)] - \ln \left[\sum_{r=1}^n (n+r+j) \right]$$

and differentiating with respect to j we obtain the result (1.3).

It may be seen that for $j=0$, we obtain

$$\left(\frac{1}{C_n(j)} \right)' \Big|_{j=0} = -\frac{n+1}{\binom{2n}{n}} \sum_{r=1}^n \frac{1}{r+n} = -\frac{n+1}{\binom{2n}{n}} [H_{2n+1}^{(1)} - H_{n+1}^{(1)}],$$

where the n^{th} Harmonic number

$$H_n^{(1)} = H_n = \int_{t=0}^1 \frac{1-t^n}{1-t} dt = \sum_{r=1}^n \frac{1}{r} = \gamma + \Psi(n+1),$$

and γ denotes the Euler-Mascheroni constant, defined by

$$\gamma = \lim_{n \rightarrow \infty} \left(\sum_{r=1}^n \frac{1}{r} - \log(n) \right) = -\Psi(1) \approx 0.577215664901532860606512.....$$


An extension of the n^{th} harmonic numbers is introduced and studied by Sandor [4]. As an aside it is possible to consider higher derivatives of (1.3). \square

Some numbers of (1.3), without the negative sign are

$$\left[\begin{array}{c} n \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ \dots \end{array} \quad \begin{array}{c} \left(\frac{1}{C_n(j)} \right)' \\ \frac{2}{(j+2)^2} \\ \frac{3!(2j+7)}{(j+3)^2(j+4)^2} \\ \frac{4!(3j^2+30j+74)}{(j+4)^2(j+5)^2(j+6)^2} \\ \frac{5!(2j^3+39j^2+251j+533)}{(j+5)^2(j+6)^2(j+7)^2(j+8)^2} \\ \frac{6!(5j^4+160j^3+1905j^2+10000j+19524)}{(j+6)^2(j+7)^2(j+8)^2(j+9)^2(j+10)^2} \\ \frac{7!(6j^5+285j^4+5380j^3+50445j^2+234908j+434568)}{(j+7)^2(j+8)^2(j+9)^2(j+10)^2(j+11)^2(j+12)^2} \\ \dots \end{array} \right]$$

In the next theorems we give integral representations for various summation expressions of Catalan type numbers with parameters. In some particular cases we give closed form values of the sums and then determine upper and lower bounds in terms of the given parameters.

The following theorem is proved.



Catalan Related Sums

Anthony Sofo

vol. 10, iss. 3, art. 69, 2009

Title Page

Contents



Page 6 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



2. Catalan Related Sums

Theorem 2.1. Let the Catalan related numbers, with parameter $j = 0, 1, 2, 3, 4, \dots$, $C_n(j) = \frac{1}{n+1} \binom{2n+j}{n}$ and $|t| < 4$, then

$$\begin{aligned}
 S_j(t) &= \sum_{n=1}^{\infty} \frac{t^n}{C_n(j)} \sum_{r=1}^n \frac{1}{r+j+n} \\
 &= \sum_{n=1}^{\infty} \frac{t^n (n+1)}{\binom{2n+j}{n}} \sum_{r=1}^n \frac{1}{r+j+n} \\
 &= \sum_{n=1}^{\infty} \frac{t^n (n+1)}{\binom{2n+j}{n}} [\Psi(1+j+2n) - \Psi(1+j+n)] \\
 (2.1) \quad &= -2t \int_0^1 \frac{(1-x)^{j+1} \log(1-x)}{(1-tx(1-x))^3} dx.
 \end{aligned}$$

Proof. Consider

$$\begin{aligned}
 \sum_{n=1}^{\infty} \frac{t^n}{C_n(j)} &= \sum_{n=1}^{\infty} \frac{t^n (n+1)}{\binom{2n+j}{n}} \\
 &= \sum_{n=1}^{\infty} \frac{t^n (n+1) \Gamma(n+1) \Gamma(n+j+1)}{\Gamma(2n+j+1)}.
 \end{aligned}$$

Now by the use of the Gamma property $\Gamma(n+1) = n\Gamma(n)$ we have

$$\sum_{n=1}^{\infty} \frac{t^n n (n+1) \Gamma(n) \Gamma(n+j+1)}{\Gamma(2n+j+1)} = \sum_{n=1}^{\infty} t^n n (n+1) B(n, n+j+1),$$

where

$$\begin{aligned} B(\alpha, \beta) &= \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta + 1)} \\ &= \int_0^1 (1-y)^{\alpha-1} y^{\beta-1} dy \\ &= \int_0^1 (1-y)^{\beta-1} y^{\alpha-1} dy \end{aligned}$$

for $\alpha > 0$ and $\beta > 0$ is the classical Beta function, therefore


$$\begin{aligned} \sum_{n=1}^{\infty} t^n n(n+1) B(n, n+j+1) &= \sum_{n=1}^{\infty} t^n n(n+1) \int_0^1 (1-x)^{n+j} x^{n-1} dx \\ &= \int_0^1 \frac{(1-x)^j}{x} \sum_{n=1}^{\infty} n(n+1) (tx(1-x))^n dx \end{aligned}$$

by interchanging sum and integral. Now applying Lemma 1.1 we obtain

$$S_j(t) = -2t \int_0^1 \frac{(1-x)^{j+1} \log(1-x)}{(1-tx(1-x))^3} dx,$$

which is the result (2.1). □

Other integral representations involving binomial coefficients and Harmonic numbers can be seen in [5], [6], [7] and [8].



Catalan Related Sums

Anthony Sofo

vol. 10, iss. 3, art. 69, 2009

Title Page

Contents



Page 8 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 9 of 16

Go Back

Full Screen

Close

Remark 1. For $j \in \mathbb{N}$ we obtain for $t = 2$

$$\begin{aligned} S_j(2) &= \sum_{n=1}^{\infty} \frac{2^n (n+1)}{\binom{2n+j}{n}} \sum_{r=1}^n \frac{1}{r+j+n} \\ &= -4 \int_0^1 \frac{(1-x)^{j+1} \log(1-x)}{(1-2x(1-x))^3} dx \\ &= \alpha_1 G + \alpha_2 \zeta(2) + \alpha_3 \pi \ln(2) + \alpha_4 \pi + \alpha_5, \end{aligned}$$

where the Catalan constant

$$G = \int_0^{\frac{\pi}{4}} \ln(\cot(x)) dx = .0.91596559\dots,$$

and $\zeta(z)$ is the Zeta function. Some specific cases of $S_j(2)$ are

j	α_1	α_2	α_3	α_4	α_5
0	3	0	$\frac{3}{4}$	$\frac{1}{4}$	0
6	$\frac{3}{8}$	$-\frac{27}{32}$	$\frac{3}{32}$	$-\frac{7}{32}$	$\frac{7}{4}$
13	$\frac{35}{32}$	$-\frac{27}{256}$	$\frac{35}{128}$	$-\frac{3}{32}$	$-\frac{6920563}{6350400}$
.....

Remark 2. For $j \in \mathbb{N}$ we obtain for $t = -\frac{1}{2}$

$$S_j\left(-\frac{1}{2}\right) = \sum_{n=1}^{\infty} \frac{\left(-\frac{1}{2}\right)^n (n+1)}{\binom{2n+j}{n}} \sum_{r=1}^n \frac{1}{r+j+n}$$

$$\begin{aligned}
&= \int_0^1 \frac{(1-x)^{j+1} \log(1-x)}{\left(1 + \frac{1}{2}x(1-x)\right)^3} dx \\
&= \beta_1 \zeta(2) + \beta_2 (\ln(2))^2 + \beta_3 \ln(2) + \beta_4.
\end{aligned}$$

Some specific cases of $S_j\left(-\frac{1}{2}\right)$ are

$$\begin{bmatrix}
j & \beta_1 & \beta_2 & \beta_3 & \beta_4 \\
0 & -\frac{8}{81} & \frac{4}{81} & -\frac{2}{27} & 0 \\
2 & \frac{8}{81} & -\frac{8}{81} & -\frac{4}{27} & -\frac{2}{9} \\
8 & -\frac{31412}{81} & \frac{31744}{81} & \frac{6596}{27} & \frac{5045}{18} \\
\dots & \dots & \dots & \dots & \dots
\end{bmatrix}.$$

Next we shall give upper and lower bounds for the series $S_j(t)$ given by (2.1).

Theorem 2.2. For $j = 0, 1, 2, 3, \dots$, and $0 < t < 4$

$$\begin{aligned}
(2.2) \quad &\frac{2t}{(j+2)^2} < S_j(t) \\
&\leq \begin{cases} \frac{2t}{(j+2)^2} \left(\frac{4}{4-t}\right)^3, \\ 2t \sqrt{\frac{2}{(2j+3)^3}} \left[\frac{2t^4 - 41t^3 + 342t^2 - 1490t + 3860}{5(4-t)^5} + \frac{1008 \arcsin\left(\frac{\sqrt{t}}{2}\right)}{\sqrt{t(4-t)^{11}}} \right]^{\frac{1}{2}}. \end{cases}
\end{aligned}$$

Proof. Consider the integral inequality

$$\int_{x_0}^{x_1} |f(x)g(x)| dx \leq \sup_{x \in [x_0, x_1]} |f(x)| \int_{x_0}^{x_1} |g(x)| dx,$$



Catalan Related Sums

Anthony Sofo

vol. 10, iss. 3, art. 69, 2009

Title Page

Contents



Page 10 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

[Title Page](#)[Contents](#)

Page 11 of 16

[Go Back](#)[Full Screen](#)[Close](#)

and from the integral (2.1) we can identify

$$(2.3) \quad \int_{x_0}^{x_1} |g(x)| dx = \int_0^1 (1-x)^{j+1} \ln(1-x) dx = \frac{1}{(j+2)^2}.$$

Similarly

$$(2.4) \quad f(x) = \frac{2t}{(1-tx(1-x))^3}$$

is monotonic on $x \in [0, 1]$ with $\lim_{x \rightarrow 0} f(x) = 2t$, $\lim_{x \rightarrow 1} f(x) = 0$, hence

$$\sup_{x \in [0,1]} f(x) = 2t \left(\frac{4}{4-t} \right)^3.$$

The series (2.1) is one of positive terms and its lower bound is given by the first term, hence combining these results we obtain the first part of the inequality (2.2). For the second part of the inequality (2.2), consider the Euclidean norm where for $\alpha = \beta = 2$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, and for $f(x)$ and $g(x)$ defined by (2.4) and (2.3) respectively we have that $|f(x)|^2$ and $|g(x)|^2$ are integrable functions defined on $x \in [0, 1]$. From (2.1) and by Hölder's integral inequality, which is a special case of the Cauchy-Buniakowsky-Schwarz inequality,

$$S_j(t) \leq \left(\int_0^1 |f(x)|^2 dx \right)^{\frac{1}{2}} \left(\int_0^1 |g(x)|^2 dx \right)^{\frac{1}{2}},$$

where

$$\left(\int_0^1 |g(x)|^2 dx \right)^{\frac{1}{2}} = \left(\int_0^1 \left| (1-x)^{j+1} \ln(1-x) \right|^2 dx \right)^{\frac{1}{2}} = \sqrt{\frac{2}{(2j+3)^3}}$$



[Title Page](#)

[Contents](#)



Page 12 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

and

$$\left(\int_0^1 |f(x)|^2 dx \right)^{\frac{1}{2}} = 2t \left[\frac{2t^4 - 41t^3 + 342t^2 - 1490t + 3860}{5(4-t)^5} + \frac{1008 \arcsin\left(\frac{\sqrt{t}}{2}\right)}{\sqrt{t(4-t)^{11}}} \right]^{\frac{1}{2}},$$

so that

$$S_j(t) \leq 2t \sqrt{\frac{2}{(2j+3)^3}} \left[\frac{2t^4 - 41t^3 + 342t^2 - 1490t + 3860}{5(4-t)^5} + \frac{1008 \arcsin\left(\frac{\sqrt{t}}{2}\right)}{\sqrt{t(4-t)^{11}}} \right]^{\frac{1}{2}}$$

and the second part of (2.2) follows. \square

Remark 3. For a particular value of t and $j \geq 0$ we can plot the exact value of $S_j(t)$, (2.1) against the upper bounds from (2.2)

$$(2.5) \quad A_j(t) = \frac{2t}{(j+2)^2} \left(\frac{4}{4-t} \right)^3$$

and

$$(2.6) \quad B_j(t) = 2t \sqrt{\frac{2}{(2j+3)^3}} \times \left[\frac{2t^4 - 41t^3 + 342t^2 - 1490t + 3860}{5(4-t)^5} + \frac{1008 \arcsin\left(\frac{\sqrt{t}}{2}\right)}{\sqrt{t(4-t)^{11}}} \right]^{\frac{1}{2}}.$$




Figure 1: A plot of $S_j(1.5)$, $A_j(1.5)$ and $B_j(1.5)$ showing the crossover point at about $j = 4.5$.

Remark 4. For $t = 1.5$ we have the graph, Figure 1, showing that $B_j(1.5)$ is a better estimator of $S_j(1.5)$, (the lower curve in Figure 1), than $A_j(1.5)$ up to about $j = 4.5$. The exact value of j for a given value of t can be exactly calculated from (2.5) and (2.6).

The graph in Figure 1 suggests that the sum in (2.1) may be convex. We prove the convexity of (2.1) in the following theorem.

Theorem 2.3. For $j \geq 0$ and $0 < t < 4$ the function $j \mapsto S_j(t)$, as given in Theorem 2.1 is strictly decreasing and convex with respect to the parameter $j \in [0, \infty)$ for every $x \in [0, 1]$.



Title Page

Contents



Page 13 of 16

Go Back

Full Screen

Close



Title Page

Contents



Page 14 of 16

Go Back

Full Screen

Close

Proof. Let

$$g_j(x, t) = \frac{(1-x)^{j+1} \ln(1-x)}{(1-tx(1-x))^3}$$

be an integrable function for $x \in [0, 1]$ and put

$$S_j(t) = -2t \int_0^1 g_j(x, t) dx,$$

so that

$$S_0(t) = -2t \int_0^1 g_0(x, t) dx = -2t \int_0^1 \frac{(1-x) \ln(1-x)}{(1-tx(1-x))^3} dx.$$

Applying the Leibniz rule for differentiation under the integral sign, we have that

$$\begin{aligned} S'_j(t) &= \int_0^1 \frac{\partial}{\partial j} g_j(x, t) dx \\ &= -2t \int_0^1 \frac{(1-x)^{j+1} (\ln(1-x))^2}{(1-tx(1-x))^3} dx. \end{aligned}$$

Since $j \geq 0$ and $0 < t < 4$

$$\frac{(1-x)^{j+1} (\ln(1-x))^2}{(1-tx(1-x))^3} > 0 \quad \text{for } x \in (0, 1),$$

then $S'_j(t) < 0$, so that the sum in (2.1), is a strictly decreasing sum with respect to the parameter j for $x \in [0, 1]$.



Title Page

Contents



Page 15 of 16

Go Back

Full Screen

Close

Now

$$\begin{aligned} S_j''(t) &= \int_0^1 \frac{\partial^2}{\partial t^2} g_j(x, t) dx \\ &= -2t \int_0^1 \frac{(1-x)^{j+1} (\ln(1-x))^3}{(1-tx(1-x))^3} dx, \end{aligned}$$


and since

$$\frac{(1-x)^{j+1} (\ln(1-x))^3}{(1-tx(1-x))^3} < 0,$$

then $S_j''(t) > 0$ so that (2.1) is a convex function for $x \in [0, 1]$. □

References

- [1] V. ADAMCHIK, Certain series associated with Catalan's constant, *J. Anal. and Appl.*, **21**(3) (2002), 818–826.
- [2] D.H. BAILEY, J.M. BORWEIN, N.J. CALKIN, R. GIRGENSOHN, D.R. LUKE AND V.H. MOLL, *Experimental Mathematics in Action*, Wellesley, MA, AK Peters, 2007.
- [3] K.A. PENSON AND J.M. SIXDENIERS, Integral representations of Catalan and related numbers, *J. Int. Sequences*, **4** (2001), Article 01.2.5.
- [4] J. SÁNDOR, Remark on a function which generalizes the harmonic series, *C. R. Bulg. Acad. Sci.*, **41**(5) (1988), 19–21.
- [5] A. SOFO, *Computational Techniques for the Summation of Series*, Kluwer Academic/Plenum Publishers, 2003.
- [6] A. SOFO, Integral identities for sums, *Math. Comm.*, **13** (2008), 303–309.
- [7] A. SOFO, Integral forms of sums associated with harmonic numbers, *Appl. Math. Comput.*, **207** (2009), 365–372.
- [8] A. SOFO, Some more identities involving rational sums, *Appl. Anal. Disc. Maths.*, **2** (2008), 56–66.
- [9] O. MARICHEV, J. SONDOW AND E.W. WEISSTEIN, *Catalan's Constant*, from Mathworld-A Wolfram Web Resource. <http://mathworld.wolfram.com/CatalansConstant.html>



Catalan Related Sums

Anthony Sofo

vol. 10, iss. 3, art. 69, 2009

Title Page

Contents



Page 16 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756