


# Journal of Inequalities in Pure and Applied Mathematics

## ON A CERTAIN RETARDED INTEGRAL INEQUALITY AND ITS APPLICATIONS

B.G. PACHPATTE

57 Shri Niketan Colony  
Near Abhinay Talkies  
Aurangabad 431 001  
(Maharashtra) India.  
*EMail:* [bgpachpatte@hotmail.com](mailto:bgpachpatte@hotmail.com)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
120-03



---

volume 5, issue 1, article 19,  
2004.

*Received 16 September, 2003;  
accepted 15 December, 2003.*

*Communicated by: D. Bainov*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

In the present paper explicit bound on a new retarded integral inequality in two independent variables is established. Applications are given to illustrate the usefulness of the inequality.

*2000 Mathematics Subject Classification:* 26D10, 26D15.

*Key words:* Retarded integral inequality, explicit bound, two independent variables, Volterra-Fredholm integral equation, uniqueness of solutions, continuous dependence of solution.

## Contents

1	Introduction .....	3
2	Main Result .....	5
3	Applications .....	10
	References	



---

### On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 22

# 1. Introduction

In the study of differential, integral and finite difference equations, one has often to deal with certain integral and finite difference inequalities, which provide explicit bounds on the unknown functions. A detailed account on such inequalities and some of their applications can be found in [2, 3, 6, 7, 9]. In [8] the present author has established the following useful integral inequality.

**Lemma 1.1.** *Let  $u(t) \in C(I, \mathbb{R}_+)$ ,  $a(t, s), b(t, s) \in C(D, \mathbb{R}_+)$  and  $a(t, s), b(t, s)$  are nondecreasing in  $t$  for each  $s \in I$ , where  $I = [\alpha, \beta]$ ,  $\mathbb{R}_+ = [0, \infty)$ ,  $D = \{(t, s) \in I^2 : \alpha \leq s \leq t \leq \beta\}$  and suppose that*

$$u(t) \leq k + \int_{\alpha}^t a(t, s) u(s) ds + \int_{\alpha}^{\beta} b(t, s) u(s) ds,$$

for  $t \in I$ , where  $k \geq 0$  is a constant. If


$$p(t) = \int_{\alpha}^{\beta} b(t, s) \exp\left(\int_{\alpha}^s a(s, \sigma) d\sigma\right) ds < 1,$$

for  $t \in I$ , then

$$u(t) \leq \frac{k}{1 - p(t)} \exp\left(\int_{\alpha}^t a(t, s) ds\right),$$

for  $t \in I$ .

A version of the above inequality when  $a(t, s) = a(s), b(t, s) = b(s)$  is first given in [2, p. 11]. In a recent paper [10] a useful general retarded version of the above inequality is given. The aim of the present paper is to establish a



---

On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

---

Title Page

Contents




Go Back

Close

Quit

Page 3 of 22

general two independent variable retarded version of the above inequality which can be used as a tool to study the behavior of solutions of a general retarded Volterra-Fredholm integral equation in two independent variables. Applications are given to convey the importance of our result to the literature.



---

## On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 22

## 2. Main Result

In what follows,  $\mathbb{R}$  denotes the set of real numbers,  $\mathbb{R}_+ = [0, \infty)$ ,  $I_1 = [x_0, M]$  and  $I_2 = [y_0, N]$  are the given subsets of  $\mathbb{R}$ . Let  $\Delta = I_1 \times I_2$  and

$$E = \{(x, y, s, t) \in \Delta^2 : x_0 \leq s \leq x \leq M, y_0 \leq t \leq y \leq N\}.$$


Our main result is established in the following theorem.

**Theorem 2.1.** *Let  $u(x, y) \in C(\Delta, \mathbb{R}_+)$ ,  $a(x, y, s, t), b(x, y, s, t) \in C(E, \mathbb{R}_+)$  and  $a(x, y, s, t), b(x, y, s, t)$  be nondecreasing in  $x$  and  $y$  for each  $s \in I_1, t \in I_2, \alpha \in C^1(I_1, I_1), \beta \in C^1(I_2, I_2)$  be nondecreasing with  $\alpha(x) \leq x$  on  $I_1, \beta(y) \leq y$  on  $I_2$  and suppose that*

$$(2.1) \quad u(x, y) \leq c + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(x, y, s, t) u(s, t) dt ds \\ + \int_{\alpha(x_0)}^{\alpha(M)} \int_{\beta(y_0)}^{\beta(N)} b(x, y, s, t) u(s, t) dt ds,$$

for  $x \in I_1, y \in I_2$ , where  $c \geq 0$  is a constant. If

$$(2.2) \quad p(x, y) = \int_{\alpha(x_0)}^{\alpha(M)} \int_{\beta(y_0)}^{\beta(N)} b(x, y, s, t) \\ \times \exp \left( \int_{\alpha(x_0)}^{\alpha(s)} \int_{\beta(y_0)}^{\beta(t)} a(s, t, \sigma, \tau) d\tau d\sigma \right) dt ds < 1,$$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 5 of 22



Title Page

Contents



Go Back

Close

Quit

Page 6 of 22

for  $x \in I_1, y \in I_2$ , then

$$(2.3) \quad u(x, y) \leq \frac{c}{1-p(x, y)} \exp \left( \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(x, y, s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ .

*Proof.* Fix any arbitrary  $(X, Y) \in \Delta$ . Then for  $x_0 \leq x \leq X, y_0 \leq y \leq Y$  we have

$$(2.4) \quad u(x, y) \leq c + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(X, Y, s, t) u(s, t) dt ds \\ + \int_{\alpha(x_0)}^{\alpha(M)} \int_{\beta(y_0)}^{\beta(N)} b(X, Y, s, t) u(s, t) dt ds.$$

Let

$$(2.5) \quad k = c + \int_{\alpha(x_0)}^{\alpha(M)} \int_{\beta(y_0)}^{\beta(N)} b(X, Y, s, t) u(s, t) dt ds,$$

then (2.4) can be restated as

$$(2.6) \quad u(x, y) \leq k + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(X, Y, s, t) u(s, t) dt ds,$$

for  $x_0 \leq x \leq X, y_0 \leq y \leq Y$ . Now a suitable application of the inequality ( $c_1$ ) given in Theorem 3 in [9, p. 51] to (2.6) yields

$$(2.7) \quad u(x, y) \leq k \exp \left( \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(X, Y, s, t) dt ds \right),$$

for  $x_0 \leq x \leq X, y_0 \leq y \leq Y$ . Since  $(X, Y) \in \Delta$  is arbitrary, from (2.7) and (2.5) with  $X$  and  $Y$  replaced by  $x$  and  $y$  we have

$$(2.8) \quad u(x, y) \leq k \exp \left( \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(x, y, s, t) dt ds \right),$$

where

$$(2.9) \quad k = c + \int_{\alpha(x_0)}^{\alpha(M)} \int_{\beta(y_0)}^{\beta(N)} b(x, y, s, t) u(s, t) dt ds,$$

for all  $x \in I_1, y \in I_2$ . Using (2.8) on the right side of (2.9) and in view of (2.2) we have


$$(2.10) \quad k \leq \frac{c}{1 - p(x, y)}.$$

Using (2.10) in (2.8) we get the desired inequality in (2.3). The proof is complete.  $\square$

By taking  $b(x, y, s, t) = 0$  in Theorem 2.1, we get the following useful inequality.

**Corollary 2.2.** *Let  $u(x, y), a(x, y, s, t), \alpha(x), \beta(y)$  and  $c$  be as in Theorem 2.1. If*

$$(2.11) \quad u(x, y) \leq c + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(x, y, s, t) u(s, t) dt ds,$$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 7 of 22



for  $x \in I_1, y \in I_2$ , then

$$(2.12) \quad u(x, y) \leq c \exp \left( \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} a(x, y, s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ .

The following corollaries of Theorem 2.1 and Corollary 2.2 are also useful in certain applications.

**Corollary 2.3.** Let  $u(x, y)$ ,  $a(x, y, s, t)$ ,  $b(x, y, s, t)$  and  $c$  be as in Theorem 2.1 and suppose that

$$(2.13) \quad u(x, y) \leq c + \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) u(s, t) dt ds \\ + \int_{x_0}^M \int_{y_0}^N b(x, y, s, t) u(s, t) dt ds,$$

for  $x \in I_1, y \in I_2$ . If

$$(2.14) \quad q(x, y) \\ = \int_{x_0}^M \int_{y_0}^N b(x, y, s, t) \exp \left( \int_{x_0}^s \int_{y_0}^t a(s, t, \sigma, \tau) d\tau d\sigma \right) dt ds < 1,$$

for  $x \in I_1, y \in I_2$ , then

$$(2.15) \quad u(x, y) \leq \frac{c}{1 - q(x, y)} \exp \left( \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ .

---

On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 22



**Corollary 2.4.** Let  $u(x, y)$ ,  $a(x, y, s, t)$  and  $c$  be as in Corollary 2.2. If


$$(2.16) \quad u(x, y) \leq c + \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) u(s, t) dt ds,$$

for  $x \in I_1, y \in I_2$ , then

$$(2.17) \quad u(x, y) \leq c \exp \left( \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ .

The proofs of Corollaries 2.3 and 2.4 follow by taking  $\alpha(x) = x, \beta(y) = y$  in Theorem 2.1 and Corollary 2.2.



---

### On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 9 of 22

### 3. Applications

In this section, we present applications of Theorem 2.1 to study certain properties of solutions of the retarded Volterra-Fredholm integral equation in two independent variables of the form

$$(3.1) \quad z(x, y) = f(x, y) + \int_{x_0}^x \int_{y_0}^y A(x, y, s, t, z(s - h_1(s), t - h_2(t))) dt ds + \int_{x_0}^M \int_{y_0}^N B(x, y, s, t, z(s - h_1(s), t - h_2(t))) dt ds,$$

where  $z, f \in C(\Delta, \mathbb{R})$ ,  $A, B \in C(E \times \mathbb{R}, \mathbb{R})$  and  $h_1 \in C(I_1, \mathbb{R}_+)$ ,  $h_2 \in C(I_2, \mathbb{R}_+)$ , are nonincreasing,  $x - h_1(x) \geq 0$ ,  $y - h_2(y) \geq 0$ ,  $x - h_1(x) \in C^1(I_1, I_1)$ ,  $y - h_2(y) \in C^1(I_2, I_2)$ ,  $h_1'(x) < 1$ ,  $h_2'(x) < 1$ ,  $h_1(x_0) = h_2(y_0) = 0$ .


The following theorem gives the bound on the solution of equation (3.1).

**Theorem 3.1.** *Suppose that the functions  $f, A, B$  in equation (3.1) satisfy the conditions*

$$(3.2) \quad |f(x, y)| \leq c,$$

$$(3.3) \quad |A(x, y, s, t, z)| \leq a(x, y, s, t) |z|,$$

$$(3.4) \quad |B(x, y, s, t, z)| \leq b(x, y, s, t) |z|,$$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 10 of 22

where  $c, a(x, y, s, t), b(x, y, s, t)$  are as in Theorem 2.1. Let

$$(3.5) \quad M_1 = \max_{x \in I_1} \frac{1}{1 - h'_1(x)}, \quad M_2 = \max_{y \in I_2} \frac{1}{1 - h'_2(y)},$$

and

$$(3.6) \quad \bar{p}(x, y) = \int_{\phi(x_0)}^{\phi(M)} \int_{\psi(y_0)}^{\psi(N)} \bar{b}(x, y, s, t) \\ \times \exp \left( \int_{\phi(x_0)}^{\phi(s)} \int_{\psi(y_0)}^{\psi(t)} \bar{a}(s, t, \sigma, \tau) d\tau d\sigma \right) dt ds < 1,$$

where  $\phi(x) = x - h_1(x), x \in I_1, \psi(y) = y - h_2(y), y \in I_2$  and

$$\bar{a}(x, y, \sigma, \tau) = M_1 M_2 a(x, y, \sigma + h_1(s), \tau + h_2(t)),$$

$$\bar{b}(x, y, \sigma, \tau) = M_1 M_2 b(x, y, \sigma + h_1(s), \tau + h_2(t)).$$

If  $z(x, y)$  is a solution of equation (3.1) on  $\Delta$ , then

$$(3.7) \quad |z(x, y)| \leq \frac{c}{1 - \bar{p}(x, y)} \exp \left( \int_{\phi(x_0)}^{\phi(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{a}(x, y, \sigma, \tau) d\tau d\sigma \right),$$

for  $x \in I_1, y \in I_2$ .

*Proof.* Since  $z(x, y)$  is a solution of equation (3.1), from (3.1) – (3.4) we have

$$(3.8) \quad |z(x, y)| \leq c + \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) |z(s - h_1(s), t - h_2(t))| dt ds \\ + \int_{x_0}^M \int_{y_0}^N b(x, y, s, t) |z(s - h_1(s), t - h_2(t))| dt ds.$$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 11 of 22

Now by making the change of variables on the right side of (3.8) and using (3.5) we have

$$(3.9) \quad |z(x, y)| \leq c + \int_{\phi(x_0)}^{\phi(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{a}(x, y, \sigma, \tau) |z(\sigma, \tau)| d\tau d\sigma \\ + \int_{\phi(x_0)}^{\phi(M)} \int_{\psi(y_0)}^{\psi(N)} \bar{b}(x, y, \sigma, \tau) |z(\sigma, \tau)| d\tau d\sigma.$$

A suitable application of Theorem 2.1 to (3.9) yields (3.7). □

The next result deals with the uniqueness of solutions of (3.1).


**Theorem 3.2.** *Suppose that the functions  $A, B$  in equation (3.1) satisfy the conditions*

$$(3.10) \quad |A(x, y, s, t, z) - A(x, y, s, t, \bar{z})| \leq a(x, y, s, t) |z - \bar{z}|,$$

$$(3.11) \quad |B(x, y, s, t, z) - B(x, y, s, t, \bar{z})| \leq b(x, y, s, t) |z - \bar{z}|,$$

where  $a(x, y, s, t), b(x, y, s, t)$  are as in Theorem 2.1. Let  $M_1, M_2, \phi, \psi, \bar{a}, \bar{b}, \bar{p}$  be as in Theorem 3.1. Then the equation (3.1) has at most one solution on  $\Delta$ .

*Proof.* Let  $z(x, y)$  and  $\bar{z}(x, y)$  be two solutions of equation (3.1) on  $\Delta$ . From




---

On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 12 of 22

(3.1), (3.10), (3.11) we have

$$\begin{aligned}
 (3.12) \quad & |z(x, y) - \bar{z}(x, y)| \\
 & \leq \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) |z(s - h_1(s), t - h_2(t)) \\
 & \quad - \bar{z}(s - h_1(s), t - h_2(t))| dt ds \\
 & + \int_{x_0}^x \int_{y_0}^y b(x, y, s, t) |z(s - h_1(s), t - h_2(t)) \\
 & \quad - \bar{z}(s - h_1(s), t - h_2(t))| dt ds.
 \end{aligned}$$


By making the change of variables on the right side of (3.12) and using (3.5) we have

$$\begin{aligned}
 (3.13) \quad & |z(x, y) - \bar{z}(x, y)| \\
 & \leq \int_{\phi(x_0)}^{\phi(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{a}(x, y, \sigma, \tau) |z(\sigma, \tau) - \bar{z}(\sigma, \tau)| d\tau d\sigma \\
 & \quad + \int_{\phi(x_0)}^{\phi(M)} \int_{\psi(y_0)}^{\psi(N)} \bar{b}(x, y, \sigma, \tau) |z(\sigma, \tau) - \bar{z}(\sigma, \tau)| d\tau d\sigma.
 \end{aligned}$$

Now a suitable application of Theorem 2.1 to (3.13) yields

$$|z(x, y) - \bar{z}(x, y)| \leq 0.$$

Therefore  $z(x, y) = \bar{z}(x, y)$ , i.e. there is at most one solution to the equation (3.1).  $\square$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 13 of 22

The following theorem deals with the continuous dependence of solution of equation (3.1) on the right side.

Consider the equation (3.1) and the following equation

$$(3.14) \quad w(x, y) = g(x, y) + \int_{x_0}^x \int_{y_0}^y F(x, y, s, t, w(s - h_1(s), t - h_2(t))) dt ds + \int_{x_0}^M \int_{y_0}^N G(x, y, s, t, w(s - h_1(s), t - h_2(t))) dt ds,$$

where  $w, g \in C(\Delta, \mathbb{R})$ ,  $F, G \in C(E \times \mathbb{R}, \mathbb{R})$  and  $h_1, h_2$  are as in equation (3.1).

**Theorem 3.3.** *Suppose that the functions  $A, B$  in equation (3.1) satisfy the conditions (3.10), (3.11) in Theorem 3.2 and further assume that*

$$(3.15) \quad |f(x, y) - g(x, y)| \leq \varepsilon,$$

$$(3.16) \quad \int_{x_0}^x \int_{y_0}^y |A(x, y, s, t, w(s - h_1(s), t - h_2(t))) - F(x, y, s, t, w(s - h_1(s), t - h_2(t)))| dt ds \leq \varepsilon,$$

$$(3.17) \quad \int_{x_0}^M \int_{y_0}^N |B(x, y, s, t, w(s - h_1(s), t - h_2(t))) - G(x, y, s, t, w(s - h_1(s), t - h_2(t)))| dt ds \leq \varepsilon,$$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close


Quit

Page 14 of 22

where  $\varepsilon > 0$  is an arbitrary small constant, and let  $M_1, M_2, \phi, \psi, \bar{a}, \bar{b}, \bar{p}$  be as in Theorem 3.1. Then the solution of equation (3.1) depends continuously on the functions involved on the right side of equation (3.1).

*Proof.* Let  $z(x, y)$  and  $w(x, y)$  be the solutions of (3.1) and (3.14) respectively. Then we have

$$\begin{aligned}
 (3.18) \quad & z(x, y) - w(x, y) \\
 &= f(x, y) - g(x, y) \\
 &+ \int_{x_0}^x \int_{y_0}^y \{A(x, y, s, t, z(s - h_1(s), t - h_2(t))) \\
 &\quad - A(x, y, s, t, w(s - h_1(s), t - h_2(t)))\} dt ds \\
 &+ \int_{x_0}^x \int_{y_0}^y \{A(x, y, s, t, w(s - h_1(s), t - h_2(t))) \\
 &\quad - F(x, y, s, t, w(s - h_1(s), t - h_2(t)))\} dt ds \\
 &+ \int_{x_0}^M \int_{y_0}^N \{B(x, y, s, t, z(s - h_1(s), t - h_2(t))) \\
 &\quad - B(x, y, s, t, w(s - h_1(s), t - h_2(t)))\} dt ds \\
 &+ \int_{x_0}^M \int_{y_0}^N \{B(x, y, s, t, w(s - h_1(s), t - h_2(t))) \\
 &\quad - G(x, y, s, t, w(s - h_1(s), t - h_2(t)))\} dt ds.
 \end{aligned}$$




---

**On A Certain Retarded Integral Inequality And Its Applications**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 15 of 22

Using (3.10), (3.11), (3.15) – (3.17) in (3.18) we get


$$\begin{aligned}
 (3.19) \quad & |z(x, y) - w(x, y)| \\
 & \leq 3\varepsilon + \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) |z(s - h_1(s), t - h_2(t)) \\
 & \quad - w(s - h_1(s), t - h_2(t))| dt ds \\
 & + \int_{x_0}^M \int_{y_0}^N b(x, y, s, t) |z(s - h_1(s), t - h_2(t)) \\
 & \quad - w(s - h_1(s), t - h_2(t))| dt ds.
 \end{aligned}$$

By making the change of variables on the right side of (3.19) and using (3.5) we get

$$\begin{aligned}
 (3.20) \quad & |z(x, y) - w(x, y)| \\
 & \leq 3\varepsilon + \int_{\phi(x_0)}^{\phi(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{a}(x, y, s, t) |z(\sigma, \tau) - w(\sigma, \tau)| d\tau d\sigma \\
 & \quad + \int_{\phi(x_0)}^{\phi(M)} \int_{\psi(y_0)}^{\psi(N)} \bar{b}(x, y, s, t) |z(\sigma, \tau) - w(\sigma, \tau)| d\tau d\sigma.
 \end{aligned}$$

Now a suitable application of Theorem 2.1 to (3.20) yields

$$\begin{aligned}
 (3.21) \quad & |z(x, y) - w(x, y)| \\
 & \leq 3\varepsilon \left[ \frac{1}{1 - \bar{p}(x, y)} \exp \left( \int_{\phi(x_0)}^{\phi(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{a}(x, y, \sigma, \tau) d\tau d\sigma \right) \right],
 \end{aligned}$$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 16 of 22



for  $x \in I_1, y \in I_2$ . On the compact set, the quantity in square brackets in (3.21) is bounded by some positive constant  $\overline{M}$ . Therefore  $|z(x, y) - w(x, y)| \leq 3\overline{M}\varepsilon$  on the set, so the solution to equation (3.1) depends continuously on the functions involved on the right side of equation (3.1). If  $\varepsilon \rightarrow 0$ , then  $|z(x, y) - w(x, y)| \rightarrow 0$  on the set.  $\square$


We next consider the following retarded Volterra-Fredholm integral equations

$$(3.22) \quad z(x, y) = f(x, y) + \int_{x_0}^x \int_{y_0}^y A(x, y, s, t, z(s - h_1(s)), t - h_2(t)), \mu) dt ds + \int_{x_0}^M \int_{y_0}^N B(x, y, s, t, z(s - h_1(s)), t - h_2(t)), \mu) dt ds,$$

$$(3.23) \quad z(x, y) = f(x, y) + \int_{x_0}^x \int_{y_0}^y A(x, y, s, t, z(s - h_1(s)), t - h_2(t)), \mu_0) dt ds + \int_{x_0}^M \int_{y_0}^N B(x, y, s, t, z(s - h_1(s)), t - h_2(t)), \mu_0) dt ds,$$

where  $z, f \in C(\Delta, \mathbb{R})$ ,  $A, B \in C(E \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$  and  $\mu, \mu_0$  are real parameters.

The following theorem shows the dependency of solutions of equations (3.22) and (3.23) on parameters.



**On A Certain Retarded Integral Inequality And Its Applications**

B.G. Pachpatte

Title Page	
Contents	
◀◀	▶▶
◀	▶
Go Back	
Close	
Quit	
Page 17 of 22	

**Theorem 3.4.** *Suppose that*

$$(3.24) \quad |A(x, y, s, t, z, \mu) - A(x, y, s, t, \bar{z}, \mu)| \leq a(x, y, s, t) |z - \bar{z}|,$$

$$(3.25) \quad |A(x, y, s, t, \bar{z}, \mu) - A(x, y, s, t, \bar{z}, \mu_0)| \leq r(x, y, s, t) |\mu - \mu_0|,$$

$$(3.26) \quad |B(x, y, s, t, z, \mu) - B(x, y, s, t, \bar{z}, \mu)| \leq b(x, y, s, t) |z - \bar{z}|,$$


$$(3.27) \quad |B(x, y, s, t, \bar{z}, \mu) - B(x, y, s, t, \bar{z}, \mu_0)| \leq e(x, y, s, t) |\mu - \mu_0|,$$

where  $a(x, y, s, t)$ ,  $b(x, y, s, t)$  are as in Theorem 2.1 and  $r, e \in C(E, \mathbb{R}_+)$  are such that

$$(3.28) \quad \int_{x_0}^x \int_{y_0}^y r(x, y, s, t) dt ds \leq k_1,$$

$$(3.29) \quad \int_{x_0}^M \int_{y_0}^N e(x, y, s, t) dt ds \leq k_2,$$

where  $k_1, k_2$  are positive constants. Let  $M_1, M_2, \phi, \psi, \bar{a}, \bar{b}, \bar{p}$  be as in Theorem 3.1. Let  $z_1(x, y)$  and  $z_2(x, y)$  be the solutions of (3.22) and (3.23) respectively.




---

**On A Certain Retarded Integral Inequality And Its Applications**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 22


Then

$$(3.30) \quad |z_1(x, y) - z_2(x, y)| \leq \frac{(k_1 + k_2) |\mu - \mu_0|}{1 - \bar{p}(x, y)} \exp \left( \int_{\phi(x_0)}^{\phi(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{a}(x, y, s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ .

*Proof.* Let  $z(x, y) = z_1(x, y) - z_2(x, y)$ ,  $(x, y) \in \Delta$ . Then

$$(3.31) \quad z(x, y) = \int_{x_0}^x \int_{y_0}^y \{A(x, y, s, t, z_1(s - h_1(s)), t - h_2(t)), \mu) - A(x, y, s, t, z_2(s - h_1(s)), t - h_2(t)), \mu)\} dt ds + \int_{x_0}^x \int_{y_0}^y \{A(x, y, s, t, z_2(s - h_1(s)), t - h_2(t)), \mu) - A(x, y, s, t, z_2(s - h_1(s)), t - h_2(t)), \mu_0)\} dt ds + \int_{x_0}^M \int_{y_0}^N \{B(x, y, s, t, z_1(s - h_1(s)), t - h_2(t)), \mu) - B(x, y, s, t, z_2(s - h_1(s)), t - h_2(t)), \mu)\} dt ds + \int_{x_0}^M \int_{y_0}^N \{B(x, y, s, t, z_2(s - h_1(s)), t - h_2(t)), \mu) - B(x, y, s, t, z_2(s - h_1(s)), t - h_2(t)), \mu_0)\} dt ds.$$



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 19 of 22

Using (3.24) – (3.29) in (3.31) we get


$$(3.32) \quad |z(x, y)| \leq |\mu - \mu_0| k_1 + |\mu - \mu_0| k_2 \\ + \int_{x_0}^x \int_{y_0}^y a(x, y, s, t) |z(s - h_1(s), t - h_2(t))| dt ds \\ + \int_{x_0}^M \int_{y_0}^N b(x, y, s, t) |z(s - h_1(s), t - h_2(t))| dt ds.$$

By using the change of variables on the right side of (3.32) and (3.5) we get

$$(3.33) \quad |z(x, y)| \leq (k_1 + k_2) |\mu - \mu_0| \\ + \int_{\phi(x_0)}^{\phi(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{a}(x, y, \sigma, \tau) |z(\sigma, \tau)| d\tau d\sigma \\ + \int_{\phi(x_0)}^{\phi(M)} \int_{\psi(y_0)}^{\psi(N)} \bar{b}(x, y, \sigma, \tau) |z(\sigma, \tau)| d\tau d\sigma.$$

Now a suitable application of Theorem 2.1 to (3.33) yields (3.30), which shows the dependency of solutions of (3.22) and (3.23) on parameters.  $\square$

In conclusion, we note that the results given in this paper can be extended very easily to functions involving many independent variables. Since the formulations of such results are quite straightforward in view of the results given above (see also [6]) and hence we omit the details. For the study of behavior of solutions of Volterra-Fredholm integral equations involving functions of one independent variable, see [1, 4, 5].



On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 20 of 22

## References

- [1] S. ASIROV AND Ja.D. MAMEDOV, Investigation of solutions of nonlinear Volterra-Fredholm operator equations, *Dokl. Akad. Nauk SSSR*, **229** (1976), 982–986.
- [2] D. BAINOV AND P. SIMEONOV, *Integral Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, 1992.
- [3] O. LIPOVAN, A retarded Gronwall-like inequality and its applications, *J. Math. Anal. Appl.*, **252** (2000), 389–401.
- [4] R.K. MILLER, J.A. NOHEL AND J.S.W. WONG, A stability theorem for nonlinear mixed integral equations, *J. Math. Anal. Appl.*, **25** (1969), 446–449.
- [5] B.G. PACHPATTE, On the existence and uniqueness of solutions of Volterra-Fredholm integral equations, *Math. Seminar Notes*, **10** (1982), 733–742.
- [6] B.G. PACHPATTE, *Inequalities for Differential and Integral Equations*, Academic Press, New York, 1998.
- [7] B.G. PACHPATTE, *Inequalities for Finite Difference Equations*, Marcel Dekker, Inc. New York, 2002.
- [8] B.G. PACHPATTE, A note on certain integral inequality, *Tamkang J. Math.*, **33** (2002), 353–358.
- [9] B.G. PACHPATTE, Explicit bounds on certain integral inequalities, *J. Math. Anal. Appl.*, **267** (2002), 48–61.



---

On A Certain Retarded Integral Inequality And Its Applications

B.G. Pachpatte

---

Title Page

Contents




Go Back

Close

Quit

Page 21 of 22

[10] B.G. PACHPATTE, Explicit bound on a retarded integral inequality, *Math. Inequal. Appl.*, **7** (2004), 7–11.



---

**On A Certain Retarded Integral Inequality And Its Applications**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 22 of 22