



INEQUALITIES FOR THE POLAR DERIVATIVE OF A POLYNOMIAL

Polar Derivative of
a Polynomial

K.K. Dewan and C.M. Upadhye
vol. 9, iss. 4, art. 119, 2008

[Title Page](#)

[Contents](#)



Page 1 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

K.K. DEWAN

Department of Mathematics
Faculty of Natural Science
Jamia Milia Islamia (Central University)
New Delhi-110025 (INDIA)
EMail:

C.M. UPADHYE

Gargi College
University of Delhi
Siri Fort Road,
New Delhi-110049 (INDIA)
EMail: c_upadhye@rediffmail.com

Received: 17 April, 2007

Accepted: 15 May, 2008

Communicated by: **D. Stefanescu**

2000 AMS Sub. Class.: 30A10, 30C15.

Key words: Polynomials, Inequality, Polar Derivative.

Abstract: In this paper we obtain new results concerning maximum modules of the polar derivative of a polynomial with restricted zeros. Our results generalize and refine upon the results of Aziz and Rather [3] and Jagjeet Kaur [9].

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction and Statement of Results	3
2	Lemmas	8
3	Proofs of the Theorems	11



**Polar Derivative of
a Polynomial**

K.K. Dewan and C.M. Upadhye

vol. 9, iss. 4, art. 119, 2008

Title Page

Contents



Page 2 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 3 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

1. Introduction and Statement of Results

Let $p(z)$ be a polynomial of degree n and $p'(z)$ its derivative. It was proved by Turán [11] that if $p(z)$ has all its zeros in $|z| \leq 1$, then

$$(1.1) \quad \max_{|z|=1} |p'(z)| \geq \frac{n}{2} \max_{|z|=1} |p(z)|.$$

The result is best possible and equality holds in (1.1) if all the zeros of $p(z)$ lie on $|z| = 1$.

For the class of polynomials having all its zeros in $|z| \leq k$, $k \geq 1$, Govil [7] proved:

Theorem A. *If $p(z) = \sum_{v=0}^n a_v z^v$ is a polynomial of degree n having all the zeros in $|z| \leq k$, $k \geq 1$, then*

$$(1.2) \quad \max_{|z|=1} |p'(z)| \geq \frac{n}{1+k^n} \max_{|z|=1} |p(z)|.$$

Inequality (1.2) is sharp. Equality holds for $p(z) = z^n + k^n$.

Let $D_\alpha\{p(z)\}$ denote the polar derivative of the polynomial $p(z)$ of degree n with respect to α , then

$$D_\alpha\{p(z)\} = np(z) + (\alpha - z)p'(z).$$

The polynomial $D_\alpha\{p(z)\}$ is of degree at most $n - 1$ and it generalizes the ordinary derivative in the sense that

$$\lim_{\alpha \rightarrow \infty} \frac{D_\alpha p(z)}{\alpha} = p'(z).$$

Aziz and Rather [3] extended (1.2) to the polar derivative of a polynomial and proved the following:



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 4 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

Theorem B. *If the polynomial $p(z) = \sum_{v=0}^n a_v z^v$ has all its zeros in $|z| \leq k$, $k \geq 1$, then for every real or complex number α with $|\alpha| \geq k$,*

$$(1.3) \quad \max_{|z|=1} |D_\alpha p(z)| \geq n \left(\frac{|\alpha| - k}{1 + k^n} \right) \max_{|z|=1} |p(z)|.$$

The bound in Theorem B depends only on the zero of largest modulus and not on the other zeros even if some of them are close to the origin. Therefore, it would be interesting to obtain a bound, which depends on the location of all the zeros of a polynomial. In this connection we prove the following:

Theorem 1.1. *Let*

$$p(z) = \sum_{v=0}^n a_v z^v = a_n \prod_{v=1}^n (z - z_v), \quad a_n \neq 0,$$

be a polynomial of degree n , $|z_v| \leq k_v$, $1 \leq v \leq n$, let $k = \max(k_1, k_2, \dots, k_n) \geq 1$. Then for every real or complex number $|\alpha| \geq k$,

$$(1.4) \quad \max_{|z|=1} |D_\alpha p(z)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{k}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{k^n + 1} \right) \min_{|z|=k} |p(z)| \right].$$

Dividing both sides of (1.4) by $|\alpha|$ and letting $|\alpha| \rightarrow \infty$, we get the following refinement of a result due to Aziz [1].

Corollary 1.2. *Let $p(z) = \sum_{v=0}^n a_v z^v = a_n \prod_{v=1}^n (z - z_v)$, $a_n \neq 0$, be a polynomial*



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 5 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

of degree n , $|z_v| \leq k_v$, $1 \leq v \leq n$, let $k = \max(k_1, k_2, \dots, k_n) \geq 1$. Then,

$$(1.5) \quad \max_{|z|=1} |p'(z)| \geq \sum_{v=1}^n \frac{k}{k+k_v} \left[\frac{2}{1+k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n-1}{k^n+1} \right) \min_{|z|=k} |p(z)| \right].$$

Since $\frac{k}{k+k_v} \geq \frac{1}{2}$ for $1 \leq v \leq n$, Theorem 1.1 gives the following result, which is an improvement of Theorem B.

Corollary 1.3. *If $p(z) = a_n \prod_{v=1}^n (z - z_v)$, $a_n \neq 0$, is a polynomial of degree n , having all its zeros in $|z| \leq k$, $k \geq 1$, then for every real or complex number $|\alpha| \geq k$,*


$$(1.6) \quad \max_{|z|=1} |D_\alpha p(z)| \geq n(|\alpha| - k) \left[\frac{1}{1+k^n} \max_{|z|=1} |p(z)| + \frac{1}{2k^n} \left(\frac{k^n-1}{k^n+1} \right) \min_{|z|=k} |p(z)| \right].$$

Dividing both sides of (1.6) by $|\alpha|$ and letting $|\alpha| \rightarrow \infty$, we obtain the following refinement of Theorem A due to Govil [7].

Corollary 1.4. *If $p(z) = a_n \prod_{v=1}^n (z - z_v)$, $a_n \neq 0$, is a polynomial of degree n , having all its zeros in $|z| \leq k$, $k \geq 1$, then*

$$(1.7) \quad \max_{|z|=1} |p'(z)| \geq n \left[\frac{1}{1+k^n} \max_{|z|=1} |p(z)| + \frac{1}{2k^n} \left(\frac{k^n-1}{k^n+1} \right) \min_{|z|=k} |p(z)| \right].$$

The bound in Theorem 1.1 can be further improved for polynomials of degree $n \geq 2$. More precisely, we prove the following:



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 6 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

Theorem 1.5. Let $p(z) = \sum_{v=0}^n a_v z^v = a_n \prod_{v=1}^n (z - z_v)$, $a_n \neq 0$, be a polynomial of degree $n \geq 2$, $|z_v| \leq k_v$, $1 \leq v \leq n$, and let $k = \max(k_1, k_2, \dots, k_n) \geq 1$. Then for every real or complex number $|\alpha| \geq k$,

$$(1.8) \quad \max_{|z|=1} |D_\alpha p(z)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{k}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{k^n + 1} \right) \right. \\ \times \min_{|z|=k} |p(z)| + \frac{2|a_{n-1}|}{k(1 + k^n)} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n - 2} \right) \Big] \\ \left. + \left(1 - \frac{1}{k^2} \right) |na_0 + \alpha a_1| \quad \text{for } n > 2 \right.$$

and

$$(1.9) \quad \max_{|z|=1} |D_\alpha p(z)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{k}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{k^n + 1} \right) \right. \\ \left. \times \min_{|z|=k} |p(z)| + |a_1| \frac{(k - 1)^n}{k(1 + k^n)} \right] + \left(1 - \frac{1}{k} \right) |na_0 + \alpha a_1| \quad \text{for } n = 2.$$

Since $\frac{k}{k+k_v} \geq \frac{1}{2}$ for $1 \leq v \leq n$, the above theorem gives in particular:

Corollary 1.6. If $p(z) = a_n \prod_{v=1}^n (z - z_v)$, $a_n \neq 0$, is a polynomial of degree n having all its zeros in $|z| \leq k$, $k \geq 1$, then for every real or complex number $|\alpha| \geq k$,

$$(1.10) \quad \max_{|z|=1} |D_\alpha p(z)| \geq n(|\alpha| - k) \left[\frac{1}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{2k^n} \left(\frac{k^n - 1}{k^n + 1} \right) \right. \\ \left. \times \min_{|z|=k} |p(z)| + \frac{|a_{n-1}|}{k(1 + k^n)} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n - 2} \right) \right] \\ \left. + \left(1 - \frac{1}{k^2} \right) |na_0 + \alpha a_1| \quad \text{for } n > 2,$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 7 of 18

Go Back

Full Screen

Close

and

$$(1.11) \quad \max_{|z|=1} |D_{\alpha} p(z)| \geq (|\alpha| - k) \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{k^n + 1} \right) \right. \\ \left. \times \min_{|z|=k} |p(z)| + \frac{|a_1|(k-1)^n}{k(1+k^n)} \right] + \left(1 - \frac{1}{k} \right) |na_0 + \alpha a_1|, \text{ for } n = 2.$$

Now it is easy to verify that if $k > 1$ and $n > 2$, then $\left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n-2} \right) > 0$. Hence for polynomials of degree $n \geq 2$, the above corollary is a refinement of Theorem 1.1. In fact, except the case when $p(z)$ has all the zeros on $|z| = k$, $a_0 = 0$, $a_1 = 0$, and $a_{n-1} = 0$, the bound obtained by Theorem 1.5 is always sharper than the bound obtained by Theorem 1.1.

Remark 1. Dividing both sides of inequalities (1.8), (1.9), (1.10) and (1.11) by $|\alpha|$ and letting $|\alpha| \rightarrow \infty$, we get the results due to Jagjeet Kaur [9]. In addition to this, if $\min_{|z|=k} |p(z)| = 0$ i.e. if a zero of a polynomial lies on $|z| = k$, then we obtain the results due to Govil [8].

Finally, as an application of Theorem 1.1 we prove the following:

Theorem 1.7. *If $p(z) = \sum_{v=0}^n a_v z^v = a_n \prod_{v=1}^n (z - z_v)$, $a_n \neq 0$, is a polynomial of degree n , $|z_v| \geq k_v$, $1 \leq v \leq n$, and $k = \min(k_1, k_2, \dots, k_n) \leq 1$, then for every real or complex number δ with $|\delta| \leq k$,*

$$(1.12) \quad \max_{|z|=1} |D_{\delta} p(z)| \\ \geq (k - |\delta|) k^{n-1} \sum_{v=1}^n \frac{k_v}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1 - k^n}{k^n(1 + k^n)} m \right],$$

where $m = \min_{|z|=k} |p(z)|$.



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 8 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

2. Lemmas

For the proofs of the theorems, we need the following lemmas.

Lemma 2.1. *If $p(z)$ is a polynomial of degree n , then for $R \geq 1$*

$$(2.1) \quad \max_{|z|=R} |p(z)| \leq R^n \max_{|z|=1} |p(z)|.$$

The above lemma is a simple consequence of the Maximum Modulus Principle [10].

Lemma 2.2. *If $p(z) = \sum_{v=0}^n a_v z^v$ is a polynomial of degree n , then for all $R > 1$,*

$$(2.2) \quad \max_{|z|=R} |p(z)| \leq R^n \max_{|z|=1} |p(z)| - (R^n - R^{n-2})|p(0)| \quad \text{for } n \geq 2$$

and

$$(2.3) \quad \max_{|z|=1} |p(z)| \leq R \max_{|z|=1} |p(z)| - (R - 1)|p(0)| \quad \text{for } n = 1.$$

This result is due to Frappier, Rahman and Ruscheweyh [5].

Lemma 2.3. *If $p(z) = a_n \prod_{v=1}^n (z - z_v)$, is a polynomial of degree $n \geq 2$, $|z_v| \geq 1$ for $1 \leq v \leq n$, then*

$$(2.4) \quad \max_{|z|=R \geq 1} |p(z)| \leq \frac{R^n + 1}{2} \max_{|z|=1} |p(z)| - |a_1| \left(\frac{R^n - 1}{n} - \frac{R^{n-2}}{n-2} \right) - \frac{R^n - 1}{2} \min_{|z|=1} |p(z)|, \quad \text{if } n > 2$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 9 of 18

Go Back

Full Screen

Close

and

$$(2.5) \quad \max_{|z|=R \geq 1} |p(z)| \leq \frac{R^2 + 1}{2} \max_{|z|=1} |p(z)| \\ - |a_1| \frac{(R-1)^2}{2} - \frac{R^2 - 1}{2} \min_{|z|=1} |p(z)|, \quad \text{if } n = 2.$$

The above lemma is due to Jagjeet Kaur [9].

Lemma 2.4. *If $p(z) = a_n \prod_{v=1}^n (z - z_v)$, $a_n \neq 0$, is a polynomial of degree n , such that $|z_v| \leq 1$, $1 \leq v \leq n$, then*

$$(2.6) \quad \max_{|z|=1} |p'(z)| \geq \sum_{v=1}^n \frac{1}{1 + |z_v|} \max_{|z|=1} |p(z)|.$$

This lemma is due to Giroux, Rahman and Schmeisser [6].

Lemma 2.5. *If $p(z)$ is a polynomial of degree n , which has all its zeros in the disk $|z| \leq k$, $k \geq 1$, then*

$$(2.7) \quad \max_{|z|=k} |p(z)| \geq \frac{2k^n}{1 + k^n} \max_{|z|=1} |p(z)|.$$

Inequality (2.7) is best possible and equality holds for $p(z) = z^n + k^n$.

The above result is due to Aziz [1].

Lemma 2.6. *If $p(z)$ is a polynomial of degree n , having all its zeros in the disk $|z| \leq k$, $k \geq 1$, then*

$$(2.8) \quad \max_{|z|=k} |p(z)| \geq \frac{2k^n}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{k^n - 1}{1 + k^n} \min_{|z|=k} |p(z)|.$$

The result is best possible and equality holds for $p(z) = z^n + k^n$.



Title Page

Contents



Page 10 of 18

Go Back

Full Screen

Close

Proof of Lemma 2.6. For $k = 1$, there is nothing to prove. Therefore it is sufficient to consider the case $k > 1$.

Let $m = \min_{|z|=k} |p(z)|$. Then $m \leq |p(z)|$ for $|z| = k$.

Since all the zeros of $p(z)$ lie in $|z| \leq k$, $k > 1$, by Rouché's theorem, for every λ with $|\lambda| < 1$, the polynomial $p(z) + \lambda m$ has all its zeros in $|z| \leq k$, $k > 1$. Applying Lemma 2.5 to the polynomial $p(z) + \lambda m$, we get

$$\max_{|z|=k} |p(z) + \lambda m| \geq \frac{2k^n}{1 + k^n} \max_{|z|=1} |p(z) + \lambda m|.$$

Choosing the argument of λ such that $|p(z) + \lambda m| = |p(z)| + |\lambda|m$ and letting $|\lambda| \rightarrow 1$, we get


$$\max_{|z|=k} |p(z)| \geq \frac{2k^n}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{k^n - 1}{1 + k^n} \min_{|z|=k} |p(z)|.$$

This completes the proof of Lemma 2.6. \square

Lemma 2.7. *If $p(z)$ is a polynomial of degree n and α is any real or complex number with $|\alpha| \neq 0$, then*

$$(2.9) \quad |D_{\alpha q}(z)| = |n\bar{\alpha}p(z) + (1 - \bar{\alpha}z)p'(z)| \quad \text{for } |z| = 1.$$

Lemma 2.7 is due to Aziz [2].



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 11 of 18

Go Back

Full Screen

Close

3. Proofs of the Theorems

Proof of Theorem 1.1. Let $G(z) = p(kz)$. Since the zeros of $p(z)$ are z_v , $1 \leq v \leq n$, the zeros of the polynomial $G(z)$ are z_v/k , $1 \leq v \leq n$, and because all the zeros of $p(z)$ lie in $|z| \leq k$, all the zeros of $G(z)$ lie in $|z| \leq 1$, therefore applying Lemma 2.4 to the polynomial $G(z)$, we get

$$(3.1) \quad \max_{|z|=1} |G'(z)| \geq \sum_{v=1}^n \frac{1}{1 + \frac{|z_v|}{k}} \max_{|z|=1} |G(z)|.$$

Let $H(z) = z^n \overline{G(1/\bar{z})}$. Then it can be easily verified that

$$(3.2) \quad |H'(z)| = |nG(z) - zG'(z)|, \quad \text{for } |z| = 1.$$

The polynomial $H(z)$ has all its zeros in $|z| \geq 1$ and $|H(z)| = |G(z)|$ for $|z| = 1$, therefore, by the result of de Bruijn [4]

$$(3.3) \quad |H'(z)| \leq |G'(z)| \quad \text{for } |z| = 1.$$

Now for every real or complex number α with $|\alpha| \geq k$, we have


$$\begin{aligned} |D_{\alpha/k}G(z)| &= \left| nG(z) - zG'(z) + \frac{\alpha}{k}G'(z) \right| \\ &\geq |\alpha/k| |G'(z)| - |nG(z) - zG'(z)|. \end{aligned}$$

This gives with the help of (3.2) and (3.3) that

$$(3.4) \quad \max_{|z|=1} |D_{\alpha/k}G(z)| \geq \frac{|\alpha| - k}{k} \max_{|z|=1} |G'(z)|.$$



Using (3.1) in (3.4), we get

$$\max_{|z|=1} |D_{\alpha/k}G(z)| \geq \frac{|\alpha| - k}{k} \sum_{v=1}^n \frac{k}{k + |z_v|} \max_{|z|=1} |G(z)|.$$



Title Page

Contents



Page 12 of 18

Go Back

Full Screen

Close

Replacing $G(z)$ by $p(kz)$, we get

$$\max_{|z|=1} |D_{\alpha/k} p(kz)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{1}{k + |z_v|} \max_{|z|=1} |p(kz)|$$

which implies

$$\max_{|z|=1} \left| np(kz) + \left(\frac{\alpha}{k} - z \right) kp'(kz) \right| \geq (|\alpha| - k) \sum_{v=1}^n \frac{1}{k + |z_v|} \max_{|z|=1} |p(kz)|,$$

which gives


$$\max_{|z|=k} |D_{\alpha} p(z)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{1}{k + |z_v|} \max_{|z|=k} |p(z)|.$$

Using Lemma 2.6 in the above inequality, we get

$$(3.5) \quad \max_{|z|=k} |D_{\alpha} p(z)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{1}{k + |z_v|} \left[\frac{2k^n}{1 + k^n} \max_{|z|=1} |p(z)| \right. \\ \left. + \left(\frac{k^n - 1}{1 + k^n} \right) \min_{|z|=k} |p(z)| \right].$$



Since $D_{\alpha} p(z)$ is a polynomial of degree at most $n - 1$ and $k \geq 1$, applying Lemma 2.1 to the polynomial $D_{\alpha} p(z)$, we get

$$(3.6) \quad \max_{|z|=k} |D_{\alpha} p(z)| \leq k^{n-1} \max_{|z|=1} |D_{\alpha} p(z)|.$$



Title Page

Contents



Page 13 of 18

Go Back

Full Screen

Close

Combining (3.5) and (3.6), we get

$$\begin{aligned} & \max_{|z|=1} |D_\alpha p(z)| \\ & \geq (|\alpha| - k) \sum_{v=1}^n \frac{k}{k + |z_v|} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{1 + k^n} \right) \min_{|z|=k} |p(z)| \right] \\ & \geq (|\alpha| - k) \sum_{v=1}^n \frac{k}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{1 + k^n} \right) \min_{|z|=k} |p(z)| \right] \end{aligned}$$


which is the required result. Hence the proof Theorem 1.1 is complete. \square

Proof of Theorem 1.5. Let $G(z) = p(kz)$. Since the zeros of $p(z)$ are z_v , $1 \leq v \leq n$, the zeros of the polynomial $G(z)$ are z_v/k , $1 \leq v \leq n$, and because all the zeros of $p(z)$ lie in $|z| \leq k$, all the zeros of $G(z)$ lie in $|z| \leq 1$, therefore applying Lemma 2.4 to the polynomial $G(z)$ and proceeding in the same way as in Theorem 1.1, we obtain

$$(3.7) \quad \max_{|z|=k} |D_\alpha p(z)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{1}{k + |z_v|} \max_{|z|=k} |p(z)|.$$

Now let $q(z) = z^n p(\frac{1}{z})$ be the reciprocal polynomial of $p(z)$. Since the polynomial $p(z)$ has all its zeros in $|z| \leq k$, $k \geq 1$ the polynomial $q(\frac{z}{k})$ has all its zeros in $|z| \geq 1$. Hence applying (2.4) of Lemma 2.3 to the polynomial $q(\frac{z}{k})$, $k \geq 1$, we get

$$\begin{aligned} \max_{|z|=k} \left| q\left(\frac{z}{k}\right) \right| & \leq \frac{k^n + 1}{2} \max_{|z|=1} \left| q\left(\frac{z}{k}\right) \right| - \left(\frac{k^n - 1}{2} \right) \min_{|z|=1} \left| q\left(\frac{z}{k}\right) \right| \\ & \quad - \frac{|a_{n-1}|}{k} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n - 2} \right), \end{aligned}$$



Title Page

Contents



Page 14 of 18

Go Back

Full Screen

Close

which gives

$$\max_{|z|=1} |p(z)| \leq \frac{k^n + 1}{2k^n} \max_{|z|=k} |p(z)| - \left(\frac{k^n - 1}{2k^n} \right) \min_{|z|=k} |p(z)| - \frac{|a_{n-1}|}{k} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n-2} \right),$$

which is equivalent to


$$(3.8) \quad \max_{|z|=k} |p(z)| \geq \frac{2k^n}{1+k^n} \max_{|z|=1} |p(z)| + \left(\frac{k^n - 1}{1+k^n} \right) \min_{|z|=k} |p(z)| + \frac{2|a_{n-1}|k^{n-1}}{1+k^n} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n-2} \right).$$

Using (3.8) in (3.7) we get

$$(3.9) \quad \max_{|z|=k} |D_\alpha p(z)| \geq (|\alpha| - k) \sum_{v=1}^n \frac{1}{k + |z_v|} \left[\frac{2k^n}{1+k^n} \max_{|z|=1} |p(z)| + \left(\frac{k^n - 1}{1+k^n} \right) \times \min_{|z|=k} |p(z)| + \frac{2|a_{n-1}|k^{n-1}}{1+k^n} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n-2} \right) \right] \quad \text{if } n > 2.$$

Since $D_\alpha p(z)$ is a polynomial of degree $n - 1$ and $k \geq 1$, from (2.2) of Lemma 2.2, we get

$$(3.10) \quad \max_{|z|=k} |D_\alpha p(z)| \leq k^{n-1} \max_{|z|=1} |D_\alpha p(z)| - (k^{n-1} - k^{n-3}) |D_\alpha p(0)|, \quad \text{if } n > 2.$$



Title Page

Contents

◀ ▶

◀ ▶

Page 15 of 18

Go Back

Full Screen

Close

Combining (3.9) and (3.10) we have

$$\begin{aligned} \max_{|z|=1} |D_{\alpha} p(z)| &\geq (|\alpha| - k) \sum_{v=1}^n \frac{k}{k + |z_v|} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{1 + k^n} \right) \min_{|z|=k} |p(z)| \right. \\ &\quad \left. + \frac{2|a_{n-1}|}{k(1 + k^n)} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n - 2} \right) \right] \\ &\quad + \left(1 - \frac{1}{k^2} \right) |na_0 + \alpha a_1|, \quad \text{if } n > 2 \\ &\geq (|\alpha| - k) \sum_{v=1}^n \frac{k}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1}{k^n} \left(\frac{k^n - 1}{1 + k^n} \right) \min_{|z|=k} |p(z)| \right. \\ &\quad \left. + \frac{2|a_{n-1}|}{k(1 + k^n)} \left(\frac{k^n - 1}{n} - \frac{k^{n-2} - 1}{n - 2} \right) \right] \\ &\quad + \left(1 - \frac{1}{k^2} \right) |na_0 + \alpha a_1|, \quad \text{if } n > 2 \end{aligned}$$

which completes the proof of (1.8).



The proof of (1.9) follows on the same lines as the proof of (1.8) but instead of (2.2) and (2.4) we use inequalities (2.3) and (2.5) respectively. We omit the details. \square

Proof of Theorem 1.7. By hypothesis, the zeros of $p(z)$ satisfy $|z_v| \geq k_v$ for $1 \leq v \leq n$ such that $k = \min(k_1, k_2, \dots, k_n) \leq 1$. It follows that the zeros of the polynomial $q(z) = z^n p(1/\bar{z})$ satisfy $1/|z_v| \leq 1/k_v$, $1 \leq v \leq n$ such that $1/k = \max(1/k_1, 1/k_2, \dots, 1/k_n) \geq 1$. On applying Theorem 1.1 to the polynomial $q(z)$,



Title Page

Contents



Page 16 of 18

Go Back

Full Screen

Close

we get

$$(3.11) \quad \max_{|z|=1} |D_\alpha q(z)| \geq k^{n-1} (|\alpha| - 1/k) \sum_{v=1}^n \frac{1}{1/k + 1/k_v} \left[\frac{2/k^n}{1 + 1/k^n} \max_{|z|=1} |q(z)| + \frac{1/k^n - 1}{1 + 1/k^n} \min_{|z|=1/k} |q(z)| \right], \quad |\alpha| \geq \frac{1}{k}.$$

Now from Lemma 2.7 it follows that

$$|D_\alpha q(z)| = |\alpha| |D_{1/\bar{\alpha}} p(z)| \quad \text{for } |z| = 1.$$

Using the above equality in (3.11), we get for $|\alpha| \geq 1/k$,

$$(3.12) \quad |\alpha| \max_{|z|=1} |D_{1/\bar{\alpha}} p(z)| \geq k^{n-1} (|\alpha| - 1/k) \sum_{v=1}^n \frac{k k_v}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1 - k^n}{(1 + k^n) k^n} \min_{|z|=k} |p(z)| \right].$$


Replacing $\frac{1}{\alpha}$ by δ , so that $|\delta| \leq k$, we get from (3.12)

$$|1/\delta| \max_{|z|=1} |D_\delta p(z)| \geq k^{n-1} (|1/\delta| - 1/k) \sum_{v=1}^n \frac{k k_v}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1 - k^n}{(1 + k^n) k^n} \min_{|z|=k} |p(z)| \right],$$

or

$$\begin{aligned} & \max_{|z|=1} |D_\delta p(z)| \\ & \geq k^{n-1}(k - |\delta|) \sum_{v=1}^n \frac{k_v}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1 - k^n}{(1 + k^n)k^n} \min_{|z|=k} |p(z)| \right], \\ & \geq k^{n-1}(k - |\delta|) \sum_{v=1}^n \frac{k_v}{k + k_v} \left[\frac{2}{1 + k^n} \max_{|z|=1} |p(z)| + \frac{1 - k^n}{(1 + k^n)k^n} \min_{|z|=k} |p(z)| \right], \end{aligned}$$

which is (1.12). Hence the proof of Theorem 1.7 is complete. \square





**Polar Derivative of
a Polynomial**

K.K. Dewan and C.M. Upadhye

vol. 9, iss. 4, art. 119, 2008

Title Page

Contents



Page 17 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

References

- [1] A. AZIZ, Inequalities for the derivative of a polynomial, *Proc. Amer. Math. Soc.*, **89** (1983), 259–266.
- [2] A. AZIZ, Inequalities for the polar derivative of a polynomial, *J. Approx. Theory*, **55** (1988), 183–193.
- [3] A. AZIZ AND N.A. RATHER, A refinement of a theorem of Paul Turán concerning polynomials, *J. Math. Ineq. Appl.*, **1** (1998), 231–238.
- [4] N.G. de BRUIJN, Inequalities concerning polynomials in the complex domain, *Nederl. Akad. Wetensch. Proc. Ser. A*, **50** (1947), 1265–1272; *Indag. Math.*, **9** (1947), 591–598.
- [5] C. FRAPPIER, Q.I. RAHMAN AND St. RUSCHEWEYH, New inequalities for polynomials, *Trans. Amer. Math. Soc.*, **288** (1985), 69–99.
- [6] A. GIROUX, Q.I. RAHMAN AND G. SCHMEISSER, On Bernstein's inequality, *Canad. J. Math.*, **31** (1979), 347–353.
- [7] N.K. GOVIL, On the derivative of a polynomial, *Proc. Amer. Math. Soc.*, **41** (1973), 543–546.
- [8] N.K. GOVIL, Inequalities for the derivative of a polynomial, *J. Approx. Theory*, **63** (1990), 65–71.
- [9] J. KAUR, Inequalities for the derivative of a polynomial, *Pure and Applied Mathematical Sciences*, **XII** (1–2) (1995), 45–51.
- [10] PÓLYA AND G. SZEGÖ, *Ausgaben und Lehratze ous der Analysis*, Springer-Verlag, Berlin, 1925.
- [11] P. TURÁN, Über die ableitung von polynomen, *Compositio Math.*, **7** (1939), 89–95.





Polar Derivative of
a Polynomial

K.K. Dewan and C.M. Upadhye

vol. 9, iss. 4, art. 119, 2008

Title Page

Contents



Page 18 of 18

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756