

A NOTE ON GLOBAL IMPLICIT FUNCTION THEOREM

MIHAI CRISTEA

University of Bucharest
Faculty of Mathematics
Str. Academiei 14, R-010014,
Bucharest, Romania
EMail: mcristea@fmi.unibuc.ro

Received: 21 April, 2006

Accepted: 01 May, 2006

Communicated by: **G. Kohr**

2000 AMS Sub. Class.: 26B10, 46C05.

Key words: Global implicit function, Boundary behaviour of a maximal implicit function.

Abstract: We study the boundary behaviour of some certain maximal implicit function. We give estimates of the maximal balls on which some implicit functions are defined and we consider some cases when the implicit function is globally defined. We extend in this way an earlier result from [3] concerning an inequality satisfied by the partial derivatives $\frac{\partial h}{\partial x}$ and $\frac{\partial h}{\partial y}$ of the map h which verifies the global implicit function problem

$$h(t, x) = h(a, b), \quad x(a) = b.$$



Global Implicit Function Theorem

Mihai Cristea

vol. 8, iss. 4, art. 100, 2007

[Title Page](#)

[Contents](#)



Page 1 of 28

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



The implicit function theorem is a classical result in mathematical analysis. Local versions can be found in [1],[8], [10], [13], [15], [17], [18] and some papers deal with some global versions (see [2], [3], [9], [16]).

We first give some local versions of the implicit function theorem, using our local homeomorphism theorem from [4].

Theorem 1. *Let E, F be Banach spaces, $\dim F < \infty$, $U \subset E$ open, $V \subset F$ open, $h : U \times V \rightarrow F$ continuous such that there exists $K \subset U \times V$ countable such that h is differentiable on $(U \times V) \setminus K$ and $\frac{\partial h}{\partial y}(x, y) \in \text{Isom}(F, F)$ for every $(x, y) \in (U \times V) \setminus K$ and let $A = \text{Pr}_1 K \subset U$. Then, for every $(a, b) \in U \times V$ there exists $r, \delta > 0$ and a unique continuous map $\varphi : B(a, r) \rightarrow B(b, \delta)$ such that*

$$\varphi(a) = b \text{ and } h(x, \varphi(x)) = h(a, b) \text{ for every } x \in B(a, r)$$

and φ is differentiable on $B(a, r) \setminus A$.

Proof. Let $(a, b) \in U \times V$ be fixed and $f : U \times V \rightarrow E \times F$ be defined by

$$f(x, y) = (x, h(x, y) + b - h(a, b)) \text{ for } (x, y) \in U \times V.$$

Also, let $T : U \times V \rightarrow E \times F$,

$$T(x, y) = (0, y - h(x, y) + h(a, b) - b) \text{ for } (x, y) \in U \times V.$$

Then $\text{Im}T \subset F$, hence T is compact and we see that $f = I - T$, f is differentiable on $(U \times V) \setminus K$ and $f'(x, y) \in \text{Isom}(E \times F, E \times F)$ for every $(x, y) \in (U \times V) \setminus K$. Using the local inversion theorem from [4], we see that f is a local homeomorphism on $U \times V$. Let $W \in \mathcal{V}((a, b))$ and $\delta > 0$ be such that

$$f|_{B(a, \delta) \times B(b, \delta)} : B(a, \delta) \times B(b, \delta) \rightarrow W$$

Title Page

Contents



Page 2 of 28

Go Back

Full Screen

Close

is a homeomorphism and let

$$g = (g_1, g_2) : W \rightarrow B(a, \delta) \times B(b, \delta)$$

be its inverse. We take $\ell > 0$ such that $Q = B(a, b) \times (b, \ell) \subset W$ and let $r = \min \{\ell, \delta\}$. We have

$$\begin{aligned} (x, z) &= f(g(x, z)) \\ &= f(g_1(x, z), g_2(x, z)) \\ &= (g_1(x, z), h(g_1(x, z), g_2(x, z)) + b - h(a, b)) \end{aligned}$$


for every $(x, z) \in Q$, hence

$$x = g_1(x, z), \quad h(x, g_2(x, z)) = z + h(a, b) - b$$

for $x \in B(a, r)$, $z \in B(b, \ell)$.

We define now $\varphi : B(a, r) \rightarrow B(b, \delta)$ by $\varphi(x) = g_2(x, b)$ for every $x \in B(a, r)$ and we see that $h(x, \varphi(x)) = h(a, b)$ for every $x \in B(a, r)$. We have $f(a, b) = (a, b) = f(a, \varphi(a))$ and using the injectivity of f on $B(a, \delta) \times B(b, \delta)$, we see that $\varphi(a) = b$. Also, if $\psi : B(a, r) \rightarrow B(b, \delta)$ is continuous and $\psi(a) = b, h(x, \psi(x)) = h(a, b)$ for every $x \in B(a, r)$, then $f(x, \varphi(x)) = (x, b) = f(x, \psi(x))$ for every $x \in B(a, r)$ and using again the injectivity of the map f on $B(a, \delta) \times B(b, \delta)$, we find that $\varphi = \psi$ on $B(a, r)$.

Let now $x_0 \in B(a, r) \setminus A$. Then $(x_0, b) = f(x_0, \beta)$, with $(x_0, \beta) \in (B(a, r) \times B(b, \delta) \setminus K)$, hence f is differentiable in (x_0, β) , $f'(x_0, \beta) \in \text{Isom}(E \times F, E \times F)$ and since f is a homeomorphism on $B(a, r) \times B(b, \delta)$, it results that g is also differentiable in $(x_0, b) = f(x_0, \beta)$ and $g'(x_0, b) = [f'(x_0, \beta)^{-1}]$, and we see that φ is differentiable in x_0 . \square



[Title Page](#)

[Contents](#)




Page 3 of 28

[Go Back](#)

[Full Screen](#)

[Close](#)



Theorem 2. Let E be an infinite dimensional Banach space, $\dim F < \infty$, $U \subset E$, $V \subset F$ be open sets, $h : U \times V \rightarrow F$ be continuous such that there exists $K \subset U \times V$,

$$K = \bigcup_{p=1}^{\infty} K_p$$

with K_p compact sets for $p \in \mathbb{N}$ such that h is differentiable on $(U \times V) \setminus K$, there exists $\frac{\partial h}{\partial y}$ on $U \times V$ and $\frac{\partial h}{\partial y}(x, y) \in \text{Isom}(F, F)$ for every $(x, y) \in U \times V$ and let $A = \bigcup_{r_1} K$. Then, for every $(a, b) \in U \times V$ there exists $r, \delta > 0$ and a unique continuous implicit function $\varphi : B(a, r) \rightarrow B(b, \delta)$ differentiable on $B(a, r) \setminus A$ such that $\varphi(a) = b$ and $h(x, \varphi(x)) = h(a, b)$ for every $x \in B(a, r)$.

Proof. We apply Theorem 11 of [8]. We see that in an infinite dimensional Banach space E , a set K which is a countable union of compact sets is a "thin" set, i.e., $\text{int } K = \emptyset$ and $B \setminus K$ is connected and simply connected for every ball B from E . Also, since A is a countable union of compact sets, we see that $\text{int } A = \emptyset$.

If E, F are Banach spaces and $A \in \mathcal{L}(E, F)$, we let

$$\|A\| = \sup_{\|x\|=1} \|A(x)\|$$

and

$$\ell(A) = \inf_{\|x\|=1} \|A(x)\|$$

and if $D \subset E$, $\lambda > 0$, we let

$$\lambda D = \{x \in E \mid \text{there exists } y \in D \text{ such that } x = \lambda y\}.$$

If X, Y are Banach spaces, $D \subset X$ is open, $x \in D$ and $f : D \rightarrow Y$ is a map, we let

$$D^+ f(x) = \limsup_{y \rightarrow x} \frac{\|f(y) - f(x)\|}{\|y - x\|}$$

Title Page

Contents



Page 4 of 28

Go Back

Full Screen

Close



and we say that f is a light map if for every $x \in D$ and every $U \in \mathcal{V}(x)$, there exists $Q \in \mathcal{V}(x)$ such that $\bar{Q} \subset U$ and $f(x) \notin f(\partial Q)$. \square

Remark 1. We can replace in Theorem 1 and Theorem 2 the condition " $\dim F < \infty$ " by "There exists $\frac{\partial h}{\partial y}$ on $U \times V$ and it is continuous on $U \times V$ and $\frac{\partial h}{\partial y}(x, y) \in \text{Isom}(F, F)$ for every $(x, y) \in U \times V$ " to obtain the same conclusion, and this is the classical implicit function theorem. Also, keeping the notations from Theorem 1 and Theorem 2, we see that if $(\alpha, \beta) \in B(a, r) \times B(b, \delta)$ is such that $h(\alpha, \beta) = h(a, b)$, then $\beta = \varphi(\alpha)$.

We shall use the following lemma from [7].

Lemma 3. Let $a > 0$, $f : [0, a] \rightarrow [0, \infty)$ be continuous and let $\omega : [0, \infty) \rightarrow [0, \infty)$ be continuous such that $\omega > 0$ on $(0, \infty)$ and

$$|f(b) - f(c)| \leq \int_b^c \omega(f(t)) dt \text{ for every } 0 < b < c \leq a.$$

Then, if

$$m = \inf_{t \in [0, a]} f(t), \quad M = \sup_{t \in [0, a]} f(t),$$

it results that

$$\int_m^M \frac{ds}{\omega(s)} \leq a.$$

We obtain now the following characterization of the boundary behaviour of the solutions of some differential inequalities.

Theorem 4. Let E, F be Banach spaces, $U \subset E$ a domain, $K \subset U$ at most countable, $\varphi : U \rightarrow F$ continuous on U and differentiable on $U \setminus K$ such that there exists

$\omega : [0, \infty) \rightarrow [0, \infty)$ continuous with $\|\varphi'(x)\| \leq \omega(\|\varphi(x)\|)$ for every $x \in U \setminus K$.
Then, if $\alpha \in \partial U$ and $C \subset U$ is convex such that $\alpha \in \overline{C}$, either there exists

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \varphi(x) = \ell \in F \text{ or } \lim_{\substack{x \rightarrow \alpha \\ x \in C}} \|\varphi(x)\| = \infty$$

or, if $\omega > 0$ on $(1, \infty)$ and

$$\int_1^\infty \frac{ds}{\omega(s)} = \infty,$$

there exists

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \varphi(x) = \ell \in F.$$

If $\omega > 0$ on $(0, 1)$ and

$$\int_0^1 \frac{ds}{\omega(s)} = \infty$$

and there exists $\alpha \in U$ such that $\varphi(\alpha) = 0$, it results that $\varphi(x) = 0$, for every $x \in U$.

Proof. Replacing, if necessary, ω by $\omega + \lambda$ for some $\lambda > 0$, we can suppose that $\omega > 0$ on $[0, \infty)$. Let $\alpha \in \partial U$, $C \subset U$ convex such that $\alpha \in \overline{C}$ and let $q : [0, 1) \rightarrow C$ be a path such that

$$\lim_{t \rightarrow 1} q(t) = \alpha$$

and there exists $L > 0$ such that $D_q^+(t) \leq L$ for every $t \in [0, 1)$. Then

$$\|q(s) - q(t)\| \leq L \cdot (s - t)$$

for every $0 \leq t < s < 1$ and let $0 \leq c < d < 1$ be fixed,

$$A = \text{co}(q([c, d]))$$



Title Page

Contents



Page 6 of 28

Go Back

Full Screen

Close

and $\varepsilon > 0$. Let $g : A \rightarrow \mathbb{R}$ be defined by

$$g(z) = \omega(\|\varphi(z)\|) \text{ for every } z \in A.$$

Then A is compact and convex and g is uniformly continuous on A , hence we can find $\delta'_\varepsilon > 0$ such that


$$|g(z_1) - g(z_2)| \leq \varepsilon \text{ for } z_1, z_2 \in A$$

with $\|z_1 - z_2\| \leq \delta'_\varepsilon$. Since $q : [c, d] \rightarrow C$ is uniformly continuous, we can find $\delta_\varepsilon > 0$ such that $\|q(t) - q(s)\| \leq \delta'_\varepsilon$ if $s, t \in [c, d]$ are such that $|s - t| \leq \delta_\varepsilon$. Let now

$$\Delta = (c = t_0 < t_1 < \dots < t_m = d) \in \mathcal{D}([c, d])$$

be such that $\|\Delta\| \leq \delta_\varepsilon$. Using Denjoi-Bourbaki's theorem we have

$$\begin{aligned} \left| \|\varphi(q(d))\| - \|\varphi(q(c))\| \right| &\leq \|\varphi(q(d)) - \varphi(q(c))\| \\ &\leq \left\| \sum_{k=0}^{m-1} \varphi(q(t_{k+1})) - \varphi(q(t_k)) \right\| \\ &\leq \sum_{k=0}^{m-1} \|q(t_{k+1}) - q(t_k)\| \cdot \sup_{z \in [q(t_k), q(t_{k+1})] \setminus K} \|\varphi'(z)\| \\ &\leq L \cdot \sum_{k=0}^{m-1} (t_{k+1} - t_k) \cdot \sup_{z \in [q(t_k), q(t_{k+1})]} \omega(\|\varphi(z)\|) \\ &\leq L \cdot \sum_{k=0}^{m-1} (t_{k+1} - t_k) \cdot (\omega(\|\varphi(q(t_k))\|) + \varepsilon). \end{aligned}$$



Title Page

Contents



Page 7 of 28

Go Back

Full Screen

Close



Letting $\|\Delta\| \rightarrow 0$ and then $\varepsilon \rightarrow 0$, we obtain

$$(1) \quad \begin{aligned} | \|\varphi(q(d))\| - \|\varphi(q(c))\| | &\leq \|\varphi(q(d)) - \varphi(q(c))\| \\ &\leq L \cdot \int_c^d \omega(\|\varphi(q(t))\|) dt \text{ for } 0 \leq c < d < 1. \end{aligned}$$

If

$$m = \inf_{t \in [0,1]} \|\varphi(q(t))\|, \quad M = \sup_{t \in [0,1]} \|\varphi(q(t))\|,$$

we obtain from Lemma 3 and (1)

$$(2) \quad \int_m^M \frac{ds}{\omega(s)} \leq L.$$

Let now $z_p \rightarrow \alpha$ be such that

$$\|z_p - \alpha\| \leq \frac{1}{2^p}, \quad z_p \in C \text{ for } p \in \mathbb{N}$$

and suppose that there exists $\rho > 0$ such that $\|\varphi(z_p)\| \leq \rho$ for every $p \in \mathbb{N}$. We take

$$0 = t_0 < t_1 < \dots < t_k < t_{k+1} < \dots < 1$$

such that $t_k \nearrow 1$ and we define $q : [0, 1) \rightarrow C$ by

$$q(t) = \frac{z_k(t_{k+1} - t) + z_{k+1}(t - t_k)}{t_{k+1} - t_k} \text{ for } t \in [t_k, t_{k+1}], \quad k \in \mathbb{N}.$$

Then

$$D^+q(t) = c_k = \frac{\|z_{k+1} - z_k\|}{t_{k+1} - t_k} \text{ for } t \in [t_k, t_{k+1}]$$

Global Implicit Function Theorem

Mihai Cristea

vol. 8, iss. 4, art. 100, 2007

Title Page

Contents



Page 8 of 28

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-575b

© 2007 Victoria University. All rights reserved.

and taking $t_k = \frac{k}{k+1}$ for $k \in \mathbb{N}$, we see that $c_k \rightarrow 0$. Then

$$\alpha_p = \sup_{t \in [t_p, 1)} D^+ q(t) = \sup_{k \geq p} c_k \rightarrow 0$$

and let

$$a_p = \inf_{t \in [t_p, 1)} \|\varphi(q(t))\|, \quad b_p = \sup_{t \in [t_p, 1)} \|\varphi(q(t))\| \text{ for } p \in \mathbb{N}.$$

Using (2) we obtain that

$$\int_{a_p}^{b_p} \frac{ds}{\omega(s)} \leq \alpha_p \text{ for } p \in \mathbb{N}$$

and let $p_0 \in \mathbb{N}$ be such that

$$\alpha_p < \int_{\rho}^{\infty} \frac{ds}{\omega(s)} \text{ for } p \geq p_0.$$

Suppose that there exists $p \geq p_0$ such that $b_p = \infty$. Then, since $q(t_k) = z_k$, we see that

$$a_k \leq \|\varphi(q(t_k))\| = \|\varphi(z_k)\| \leq \rho \text{ for } k \in \mathbb{N},$$

hence


$$0 < \int_{\rho}^{\infty} \frac{ds}{\omega(s)} \leq \int_{a_p}^{b_p} \frac{ds}{\omega(s)} \leq \alpha_p < \int_{\rho}^{\infty} \frac{ds}{\omega(s)}$$

and we have reached a contradiction.

It results that $b_p < \infty$ for $p \geq p_0$ and let

$$K_p = \sup_{t \in [0, b_p]} \omega(t)$$





for $p \geq p_0$. Then $K_p < \infty$ and we see from (1) that

$$\|\varphi(q(d)) - \varphi(q(c))\| \leq \alpha_p \cdot K_p \cdot |d - c|$$

for $t_p \leq c < d < 1$ and $p \geq p_0$, and this implies that

$$\lim_{t \rightarrow 1} \varphi(q(t)) = \ell \in F.$$

It results that

$$\lim_{p \rightarrow \infty} \varphi(z_p) = \lim_{p \rightarrow \infty} \varphi(q(t_p)) = \ell \in F.$$

Now, if the case

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \|\varphi(x)\| = \infty$$

does not hold, there exists $\rho > 0$ and $x_p \rightarrow \alpha$, $x_p \in C$ with

$$\|\varphi(x_p)\| \leq \rho \text{ and } \|x_p - \alpha\| \leq \frac{1}{2^p}$$

for every $p \in \mathbb{N}$, and from what we have proved before, it results that

$$\lim_{p \rightarrow \infty} \varphi(x_p) = \ell \in F.$$

If $a_p \in C$, $\|a_p - \alpha\| \leq \frac{1}{2^p}$ for every $p \in \mathbb{N}$, then

$$\lim_{p \rightarrow \infty} \varphi(a_p) = \ell_1 \in F.$$

Let $z_{2p} = x_p$, $z_{2p+1} = a_p$ for $p \in \mathbb{N}$. We see that

$$\lim_{p \rightarrow \infty} \varphi(z_p) = \ell_2 \in F,$$



hence

$$\ell = \lim_{p \rightarrow \infty} \varphi(x_p) = \lim_{p \rightarrow \infty} \varphi(z_{2p}) = \ell_2$$

and

$$\ell_1 = \lim_{p \rightarrow \infty} \varphi(a_p) = \lim_{p \rightarrow \infty} \varphi(z_{2p+1}) = \ell_2,$$

hence $\ell = \ell_1 = \ell_2$. We have proved that if $a_p \in C$, $\|a_p - \alpha\| \leq \frac{1}{2^p}$ for every $p \in \mathbb{N}$, then

$$\lim_{p \rightarrow \infty} \varphi(a_p) = \ell.$$

We show now that if $a_p \in C$, $a_p \rightarrow \alpha$, then

$$\lim_{p \rightarrow \infty} \varphi(a_p) = \ell.$$

Indeed, if this is false, there exists $\varepsilon > 0$ and $(a_{p_k})_{k \in \mathbb{N}}$ such that

$$\|\varphi(a_{p_k}) - \ell\| > \varepsilon$$

for every $k \in \mathbb{N}$. Let $(a_{p_{k_q}})_{q \in \mathbb{N}}$ be a subsequence such that

$$\|a_{p_{k_q}} - \alpha\| < \frac{1}{2^q}$$

for every $q \in \mathbb{N}$. From what we have proved before it results that

$$\lim_{q \rightarrow \infty} \varphi(a_{p_{k_q}}) = \ell$$

and we have reached a contradiction.

We have therefore proved that either

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \varphi(x) = \ell \in F,$$

or

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \|\varphi(x)\| = \infty.$$

Suppose now

$$\int_1^\infty \frac{ds}{\omega(s)} = \infty$$

and let $\alpha \in \partial U$. We take $x \in C$ and let $q : [0, 1) \rightarrow C$ be defined by

$$q(t) = (1 - t)x + t\alpha$$

for $t \in [0, 1)$ and

$$m = \inf_{t \in [0, 1)} \|\varphi(q(t))\|, \quad M = \sup_{t \in [0, 1)} \|\varphi(q(t))\|.$$

Since $D^+q(t) = \|x - \alpha\|$ for every $t \in (0, 1)$, we see from (2)

$$\int_{\|x\|}^M \frac{ds}{\omega(s)} \leq \int_m^M \frac{ds}{\omega(s)} \leq \|x - \alpha\|$$

and this implies that $M < \infty$. Let

$$b = \sup_{t \in [0, M]} \omega(t).$$

Then $b < \infty$ and using (1), we see that

$$\|\varphi(q(d)) - \varphi(q(c))\| \leq b \cdot \|x - \alpha\| \cdot |d - c|$$

for $0 \leq c < d < 1$ and this implies that

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \varphi(x) = \ell \in F.$$



Global Implicit Function Theorem

Mihai Cristea

vol. 8, iss. 4, art. 100, 2007

Title Page

Contents



Page 12 of 28

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.



It results that the case

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \|\varphi(x)\| = \infty$$

cannot hold, hence

$$\lim_{x \rightarrow \alpha} \varphi(x) = \ell \in F.$$

Suppose now that

$$\int_0^1 \frac{ds}{\omega(s)} = \infty$$

and that there exists $\alpha \in U$ such that $\varphi(\alpha) = 0$. Let $r = d(\alpha, \partial U)$, $y \in B(\alpha, r)$ and $q : [0, 1] \rightarrow B(\alpha, r)$, $q(t) = (1-t)\alpha + ty$ for $t \in [0, 1]$. Then $D^+q(t) = \|y - \alpha\|$ for $t \in [0, 1]$ and let

$$m = \inf_{t \in [0, 1]} \|\varphi(q(t))\|$$

and

$$M = \sup_{t \in [0, 1]} \|\varphi(q(t))\|.$$

Then $m = 0$ and we see from (2) that

$$\int_0^M \frac{ds}{\omega(s)} \leq \|y - \alpha\|.$$

This implies that $M = 0$ and hence $\varphi(y) = 0$. We proved that $\varphi \equiv 0$ on $B(\alpha, r)$ and since U is a domain, we see that $\varphi \equiv 0$ on U . \square

Remark 2. We proved that if φ is as in Theorem 2 and

$$\int_1^\infty \frac{ds}{\omega(s)} = \infty,$$

then φ has angular limits in every point $\alpha \in \partial U$.

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 13 of 28

Go Back

Full Screen

Close



We now obtain the following characterization of the boundary behaviour of some implicit function.

Theorem 5. *Let E, F be Banach spaces, $U \subset E$ a domain, $K \subset U \times F$ such that $A = \text{Pr}_1 K$ is at most countable and let $h : U \times F \rightarrow F$ be continuous on $U \times F$, differentiable on $(U \times F) \setminus K$ such that*

$$\ell \left(\frac{\partial h}{\partial y} (x, y) \right) > 0 \text{ on } (U \times F) \setminus K$$

and there exists $\omega : [0, \infty) \rightarrow [0, \infty)$ continuous such that

$$\left\| \frac{\partial h}{\partial x} (x, y) \right\| / \ell \left(\frac{\partial h}{\partial y} (x, y) \right) \leq \omega (\|y\|)$$

for every $(x, y) \in (U \times F) \setminus K$. Suppose that $\varphi : U \rightarrow F$ is continuous on U , differentiable on $U \setminus A$, $\varphi(a) = b$ and

$$h(x, \varphi(x)) = h(a, b)$$

for every $x \in U$.

Then, if $\alpha \in \partial U$ and $C \subset U$ is convex such that $\alpha \in \overline{C}$, either

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \|\varphi(x)\| = \ell \in F,$$

or

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \|\varphi(x)\| = \infty.$$

Also, if $\omega > 0$ on $(1, \infty)$ and

$$\int_1^\infty \frac{ds}{\omega(s)} = \infty,$$

[Title Page](#)

[Contents](#)



Page 14 of 28

[Go Back](#)

[Full Screen](#)

[Close](#)

then

$$\lim_{\substack{x \rightarrow \alpha \\ x \in C}} \varphi(x) = \ell \in F.$$

Proof. We see that if $x \in U \setminus A$, then $(x, \varphi(x)) \in (U \times F) \setminus K$, hence h is differentiable in $(x, \varphi(x))$ and we have

$$\frac{\partial h}{\partial x}(x, \varphi(x)) + \frac{\partial h}{\partial y}(x, \varphi(x)) \cdot \varphi'(x) = 0$$

and we see that

$$\|\varphi'(x)\| \cdot \ell \left(\frac{\partial h}{\partial y}(x, \varphi(x)) \right) \leq \left\| \frac{\partial h}{\partial y}(x, \varphi(x)) (\varphi'(x)) \right\| = \left\| \frac{\partial h}{\partial x}(x, \varphi(x)) \right\|.$$

It results that

$$\|\varphi'(x)\| \leq \left\| \frac{\partial h}{\partial x}(x, \varphi(x)) \right\| / \ell \left(\frac{\partial h}{\partial y}(x, \varphi(x)) \right) \leq \omega(\|\varphi(x)\|)$$

for every $x \in U \setminus A$ and we now apply Theorem 4. □

Remark 3. If E is an infinite dimensional Banach space and

$$K = \bigcup_{p=1}^{\infty} K_p \text{ with } K_p \subset E$$

are compact sets for every $p \in \mathbb{N}$ and $y \in E$, then the set

$$M(K, y) = \{w \in E \mid \text{there exists } t > 0 \text{ and } x \in K \text{ such that } w = tx\}$$

is also a countable union of compact sets and hence a "thin" set. Keeping the notations from Theorem 4, we see that the basic inequality

$$(3) \quad \|\varphi(z_1) - \varphi(z_2)\| \leq \sup_{z \in [z_1, z_2]} \omega(\|\varphi(z)\|) \quad \text{if } [z_1, z_2] \subset U$$



Title Page

Contents




Page 15 of 28

Go Back

Full Screen

Close



is also valid for K a countable union of compact sets and φ as in Theorem 4.

If $\dim E = n$ and $K \subset E$ has a σ -finite $(n - 1)$ -dimensional measure (i.e. $K = \bigcup_{p=1}^{\infty} K_p$, with $m_{n-1}(K_p) < \infty$ for every $p \in \mathbb{N}$, where m_q is the q -Hausdorff measure from \mathbb{R}^n), a theorem of Gross shows that if $H \subset E$ is a hyperplane and $P : E \rightarrow H$ is the projection on H , then $P^{-1}(z) \cap K$ is at most countable with the possible exception of a set $B \subset H$, with $m_{n-1}(B) = 0$. Applying as in Theorem 4 the theorem of Denjoi and Bourbaki on each interval from $P^{-1}(z) \cap K$ for every $z \in H \setminus B$ and using a natural limiting process, we see that if $\dim E = n$ and $K \subset E$ has a σ -finite $(n - 1)$ -dimensional measure, then the inequality (3) also holds. It is easy to see now that Theorem 4 and Theorem 5 hold if the set K , respectively the set $A = \text{Pr}_1 K$ are chosen to be a countable union of compact sets if $\dim E = \infty$ and having σ -finite $(n - 1)$ -dimensional measure if $\dim E = n$.

The following theorem is the main theorem of the paper and it gives some cases when the implicit function is globally defined or some estimates of the maximal balls on which some implicit function is defined.

We say that a domain D from a Banach space is starlike with respect to the point $a \in D$ if $[a, x] \subset D$ for every $x \in D$, and if D is a domain in the Banach space E and $a \in D$. We set

$$D_a = \{x \in D \mid [a, x] \subset D\}.$$

Theorem 6. *Let E, F be Banach spaces, $\dim F < \infty$, $D \subset E$ a domain, $K \subset D \times F$ at most countable, and $A = \text{Pr}_1 K$. Also, let $h : D \times F \rightarrow F$ be continuous on $D \times F$ and differentiable on $(D \times F) \setminus K$ such that*

$$\ell \left(\frac{\partial h}{\partial y} (x, y) \right) > 0 \text{ on } (D \times F) \setminus K.$$

In addition, there exists $\omega : [0, \infty) \rightarrow [0, \infty)$ continuous such that $\omega > 0$ on $(0, \infty)$

[Title Page](#)

[Contents](#)



Page 16 of 28

[Go Back](#)

[Full Screen](#)

[Close](#)



and

$$\left\| \frac{\partial h}{\partial x}(x, y) \right\| / \ell \left(\frac{\partial h}{\partial y}(x, y) \right) \leq \omega(\|y\|)$$

for every $(x, y) \in (D \times F) \setminus K$. Then, if $(a, b) \in D \times F$ and

$$Q_{a,b} = D_a \cap B \left(a, \int_{\|b\|}^{\infty} \frac{ds}{\omega(s)} \right),$$

there exists a unique continuous map $\varphi : Q_{a,b} \rightarrow F$, differentiable on $Q_{a,b} \setminus A$ such that $h(x, \varphi(x)) = h(a, b)$ for every $x \in Q_{a,b}$. If D is starlike with respect to a and

$$\int_1^{\infty} \frac{ds}{\omega(s)} = \infty,$$


then $Q_{a,b} = D$ and $\varphi : D \rightarrow F$ is globally defined on D .

Proof. Let $z \in Q_{a,b}$ and let $B = \{x \in [a, z] \mid \text{there exists an open, convex domain } D_x \subset Q_{a,b} \text{ such that } [a, x] \subset D_x\}$ and a continuous implicit function $\varphi_x : D_x \rightarrow F$, differentiable on $D_x \setminus A$ such that $\varphi_x(a) = b$ and $h(u, \varphi_x(u)) = h(a, b)$ for every $u \in D_x$. We see that B is open, and from Theorem 1, $B \neq \emptyset$. We show that B is a closed set. Let $x_p \in B$, $x_p \rightarrow x$, and we can suppose that $x_p \in [a, x]$ for every $p \in \mathbb{N}$, and let

$$Q = \bigcup_{p=1}^{\infty} D_{x_p}.$$

We define $\Psi : Q \rightarrow F$ by $\Psi(u) = \varphi_{x_p}(u)$ for $u \in D_{x_p}$ and $p \in \mathbb{N}$ and the definition is correct. Indeed, if $p, q \in \mathbb{N}$, $p \neq q$ let $U_{pq} = D_{x_p} \cap D_{x_q}$. Then U_{pq} is a nonempty, open and convex set, hence it is a nonempty domain. If

$$V_{pq} = \{u \in U_{pq} \mid \varphi_{x_q}(u) = \varphi_{x_p}(u)\},$$



we see that $a \in V_{pq}$, hence $V_{pq} \neq \emptyset$ and we see that V_{pq} is a closed set in U_{pq} . Using the property of the local unicity of the implicit function from Theorem 1, we obtain that V_{pq} is also an open set. Since U_{pq} is a domain, it results that $U_{pq} = V_{pq}$ and hence that Ψ is correctly defined. We also see immediately that $\Psi(a) = b$, and Ψ is continuous on Q and differentiable on $Q \setminus A$.

Let $q : [0, 1] \rightarrow Q$ be defined by

$$q(t) = (1-t)a + tx \text{ for } t \in [0, 1].$$

Then

$$D^+q(t) = \|x - a\| < \int_{\|b\|}^{\infty} \frac{ds}{\omega(s)}.$$

Let

$$m = \inf_{t \in [0,1]} \|\Psi(q(t))\|, \quad M = \sup_{t \in [0,1]} \|\Psi(q(t))\|.$$

As in Theorem 5, we see that

$$\|\Psi'(u)\| \leq \omega(\|\Psi(u)\|) \text{ for every } u \in Q \setminus A,$$

hence

$$\|\Psi(z_1) - \Psi(z_2)\| \leq \sup_{z \in [z_1, z_2]} \omega(\|\Psi(z)\|)$$

if $[z_1, z_2] \subset Q$. This implies that relations (1) and (2) from Theorem 4 also hold and we see that

$$\int_m^M \frac{ds}{\omega(s)} \leq \|x - a\|.$$

Then

$$\int_{\|b\|}^M \frac{ds}{\omega(s)} \leq \int_m^M \frac{ds}{\omega(s)} \leq \|x - a\|$$

Title Page

Contents



Page 18 of 28

Go Back

Full Screen

Close



and this implies that $M < \infty$, hence

$$\ell = \sup_{t \in [0, M]} \omega(t) < \infty.$$

Using (1) and Theorem 4, we see that

$$\|\Psi(q(d)) - \Psi(q(c))\| \leq \ell \cdot \|x - a\| \cdot |d - c| \text{ for every } 0 \leq c < d < 1$$

and this implies that

$$\lim_{\substack{u \rightarrow x \\ u \in \text{Im } q}} \Psi(u) = w \in F.$$

Using Theorem 1, we can find $r, \delta > 0$ and a unique continuous implicit function $\Psi_x : B(x, r) \rightarrow B(w, \delta)$, differentiable on $B(x, r) \setminus A$ such that

$$\Psi_x(x) = w \text{ and } h(u, \Psi_x(u)) = h(a, b)$$

for every $u \in B(x, r)$. Let $0 < \varepsilon < r$ and $p_\varepsilon \in \mathbb{N}$ be such that

$$\|x_p - x\| < \varepsilon \text{ and } \|\Psi(x_p) - w\| < \delta$$

for $p \geq p_\varepsilon$ and let $p \geq p_\varepsilon$ be fixed. Since

$$\varphi_{x_p}(x_p) = \Psi(x_p) \in B(w, \delta),$$

we see from Remark 1 that $\varphi_{x_p}(x_p) = \Psi_x(x_p)$ and hence the set

$$U = \{u \in D_{x_p} \cap B(x, \varepsilon) \mid \varphi_{x_p}(u) = \Psi_x(u)\}$$

is nonempty. We also see that $D_{x_p} \cap B(x, \varepsilon)$ is an open, nonempty, convex set, hence it is a domain and U is an open, closed and nonempty subset of $D_{x_p} \cap B(x, \varepsilon)$, and this implies that

$$U = D_{x_p} \cap B(x, \varepsilon).$$



Title Page

Contents



Page 20 of 28

Go Back

Full Screen

Close

Let $U_0 = D_{x_p} \cup B(x, \varepsilon)$. We can now correctly define $\Phi : U_0 \rightarrow F$ by

$$\Phi(u) = \varphi_{x_p}(u) \text{ if } u \in D_{x_p}$$

and

$$\Phi(u) = \Psi_x(u) \text{ if } u \in B(x, \varepsilon)$$

and we see that Φ is continuous on U_0 , differentiable on $U_0 \setminus A$, $\Phi(a) = b$ and $h(u, \Phi(u)) = h(a, b)$ for every $u \in U_0$. It results that $x \in B$, hence B is also a closed set and since $[a, z]$ is a connected set, we see that $B = [a, z]$.

We have therefore proved that for every $z \in Q_{a,b}$ there exists a convex domain D_z such that $[a, z] \subset D_z$ and a unique continuous implicit function $\varphi_z : D_z \rightarrow F$, differentiable on $D_z \setminus A$ such that

$$\varphi_z(a) = b, h(u, \varphi_z(u)) = h(a, b) \text{ for every } u \in D_z.$$

We now define $\varphi : Q_{a,b} \rightarrow F$ by $\varphi(x) = \varphi_z(x)$ for $x \in D_z$ and we see, as before, that the definition is correct, that $\varphi(a) = b$, φ is continuous on $Q_{a,b}$, differentiable on $Q_{a,b} \setminus A$ and $h(x, \varphi(x)) = h(a, b)$ for every $x \in Q_{a,b}$. \square

Remark 4. The result from Theorem 6 extends a global implicit function theorem from [3]. The result from [3] also involves an inequality containing

$$\left\| \frac{\partial h}{\partial x}(x, y) \right\| \quad \text{and} \quad \ell \left(\frac{\partial h}{\partial y}(x, y) \right)$$

and it says that if E, F are Banach spaces, $h : E \times F \rightarrow F$ is a C^1 map such that $\frac{\partial h}{\partial y}(x, y) \in \text{Isom}(F, F)$ for every $(x, y) \in E \times F$ and there exists $\omega : [0, \infty) \rightarrow (0, \infty)$ continuous such that

$$\left(1 + \left\| \frac{\partial h}{\partial x}(x, y) \right\| \right) / \ell \left(\frac{\partial h}{\partial y}(x, y) \right) \leq \omega(\max(\|x\|, \|y\|))$$



for every $(x, y) \in E \times F$, then, for $(x_0, y_0) \in E \times F$, $z_0 = h(x_0, y_0)$ and

$$r = \int_{\max(\|x_0\|, \|y_0\|)}^{\infty} \frac{ds}{1 + \omega(s)},$$

there exists a C^1 map $\varphi : B(x_0, r) \times B(z_0, r) \rightarrow F$ such that $h(x, \varphi(x, z)) = z$ for every $(x, z) \in B(x_0, r) \times B(z_0, r)$. The main advantage of our new global implicit function theorems is that these theorems hold even if the map h is defined on a proper subset of $E \times F$, namely, on a set $D \times F \subset E \times F$, where $D \subset E$ is an open starlike domain.

Example 1. A known global inversion theorem of Hadamard, Lévy and John says that if E, F are Banach spaces, $f : E \rightarrow F$ is a C^1 map such that $f'(x) \in \text{Isom}(E, F)$ for every $x \in E$ and there exists $\omega : [0, \infty) \rightarrow (0, \infty)$ continuous such that

$$\int_1^{\infty} \frac{ds}{\omega(s)} = \infty \quad \text{and} \quad \|f'(x)^{-1}\| \leq \omega(\|x\|)$$

for every $x \in E$, then it results that $f : E \rightarrow F$ is a C^1 diffeomorphism (see [11], [14], [12], [3], [7]). If $E = F = \mathbb{R}^n$ or if $\dim E = \infty$ and $f = I - T$ with T compact, we can drop the continuity of the derivative on E and we can impose the essential condition " $f'(x) \in \text{Isom}(F, F)$ " with the possible exception of a "thin" set (see [4], [5],[6]) and we will still obtain that $f : E \rightarrow F$ is a homeomorphism.

Now, let E, F be Banach spaces, $D \subset E$ a domain, $a \in D$, $b \in F$, $g : D \rightarrow F$ be differentiable on D , $f : F \rightarrow F$ be differentiable on F such that $f'(y) \in \text{Isom}(F, F)$ for every $y \in F$ and there exists $\omega : [0, \infty) \rightarrow (0, \infty)$ continuous such that

$$\|f'(y)^{-1}\| \leq \omega(\|y\|) \quad \text{for every } y \in F.$$

Title Page

Contents

◀▶


◀▶

Page 21 of 28

Go Back

Full Screen

Close



Let $h : D \times F \rightarrow F$ be defined by $h(x, y) = f(y) - g(x)$ for $x \in D, y \in F$, $r_0 = d(a, \partial D)$ and suppose that

$$M_r = \sup_{x \in B(a, r)} \|g'(x)\| < \infty \text{ for every } 0 < r \leq r_0.$$

Then

$$\left\| \frac{\partial h}{\partial x}(x, y) \right\| \left/ \ell \left(\frac{\partial h}{\partial y}(x, y) \right) \right. \leq \|g'(x)\| \cdot \|f'(y)^{-1}\| \leq M_r \cdot \omega(\|y\|)$$

if $(x, y) \in B(a, r) \times F$ and $0 < r \leq r_0$ and let

$$\delta_r = \min \left\{ r, \frac{1}{M_r} \cdot \int_{\|b\|}^{\infty} \frac{ds}{\omega(s)} \right\} \text{ for } 0 < r \leq r_0.$$

Using Theorem 6, we see that there exists a unique differentiable map $\varphi : B(a, \delta_r) \rightarrow F$ such that $\varphi(a) = b$ and $h(x, \varphi(x)) = h(a, b)$ for every $x \in B(a, \delta_r)$, i.e. $f(\varphi(x)) = g(x) + h(a, b)$ for every $x \in B(a, r)$ and every $0 < r \leq r_0$.

If

$$r_0 \cdot M_{r_0} < \int_{\|b\|}^{\infty} \frac{ds}{\omega(s)},$$

then φ is defined on $B(a, r_0)$. Additionally, if $D = B(a, r_0)$, then φ is globally defined on D and $f \circ \varphi = g + h(a, b)$ on D . In the special case $D = E, g(x) = x$ for every $x \in E$,

$$\int_1^{\infty} \frac{ds}{\omega(s)} = \infty \quad \text{and} \quad b = f(a),$$

then $f(\varphi(x)) = x$ for every $x \in E$, and φ is defined on E and is the inverse of f and it results that $f : F \rightarrow F$ is a homeomorphism. In this way we obtain an alternative proof of the Hadamard-Lévy-John theorem.

Title Page

Contents




Page 22 of 28

Go Back

Full Screen

Close



Remark 5. The global implicit function problem

$$h(t, x) = h(a, b), \quad x(a) = b$$

considered before has two basic properties:

1. It satisfies the differential inequality $\|\varphi'(x)\| \leq \omega(\|\varphi(x)\|)$.
2. It has the property of the local existence and local unicity of the solutions around each point (t_0, x_0) .

This shows that by considering some other conditions of local existence and local unicity of the implicit function instead of the conditions from Theorem 1, we can produce corresponding global implicit function results. Using the conditions of local existence and local unicity from Theorem 11 of [8], we obtain the following corresponding version of Theorem 6.

Theorem 7. Let E, F be Banach spaces, $\dim E = \infty$, $\dim F < \infty$, $D \subset E$ a domain, $K \subset D \times F$,


$$K = \bigcup_{p=1}^{\infty} K_p$$

with K_p compact sets for every $p \in \mathbb{N}$, $A = \text{Pr}_1 K$, $h : D \times F \rightarrow F$ continuous on $D \times F$ and differentiable on $(D \times F) \setminus K$ such that

$$\ell \left(\frac{\partial h}{\partial y}(x, y) \right) > 0 \text{ on } (D \times F) \setminus K,$$

and there exists $\omega : [0, \infty) \rightarrow [0, \infty)$ continuous such that $\omega > 0$ on $(0, \infty)$ and

$$\left\| \frac{\partial h}{\partial x}(x, y) \right\| / \ell \left(\frac{\partial h}{\partial y}(x, y) \right) \leq \omega(\|y\|)$$



[Title Page](#)

[Contents](#)



Page 24 of 28

[Go Back](#)

[Full Screen](#)

[Close](#)

for every $(x, y) \in (D \times F) \setminus K$. Suppose that the map $y \rightarrow h(x, y)$ is a light map on F for every $x \in D$. Then, if $a, b \in D \times F$ and

$$Q_{a,b} = D_a \cap B \left(a, \int_{\|b\|}^{\infty} \frac{ds}{\omega(s)} \right),$$

there exists a unique continuous implicit function $\varphi : Q_{a,b} \rightarrow F$, differentiable on $Q_{a,b} \setminus A$ such that $\varphi(a) = b$ and $h(x, \varphi(x)) = h(a, b)$ for every $x \in Q_{a,b}$ and if D is starlike with respect to a and

$$\int_1^{\infty} \frac{ds}{\omega(s)} = \infty,$$

then $Q_{a,b} = D$ and $\varphi : D \rightarrow F$ is globally defined on D .

Remark 6. The condition "the map $y \rightarrow h(x, y)$ is a light map on F for every $x \in D$ " is satisfied if $\frac{\partial h}{\partial y}$ exists on $D \times F$ and

$$\ell \left(\frac{\partial h}{\partial y}(x, y) \right) > 0 \text{ for every } (x, y) \in D \times F.$$

Using the conditions of local existence and local unicity from Theorem 7 of [8], we obtain the following global implicit function theorem.

Theorem 8. Let $n \geq 2$, $D \subset \mathbb{R}^n$ be a domain, $h : D \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be differentiable and let $K \subset D \times \mathbb{R}^m$,

$$K = \bigcup_{p=1}^{\infty} K_p$$

with K_p closed sets such that $m_{n-2}(\text{Pr}_1 K_p) = 0$ for every $p \in \mathbb{N}$, $A = \text{Pr}_1 K$, such that $\frac{\partial h}{\partial y}(x, y) \in \text{Isom}(\mathbb{R}^m, \mathbb{R}^m)$ for every $(x, y) \in (D \times \mathbb{R}^m) \setminus K$ and the map



$y \rightarrow h(x, y)$ is a light map on \mathbb{R}^m for every $x \in D$. Suppose that there exists $\omega : [0, \infty) \rightarrow [0, \infty)$ continuous such that $\omega > 0$ on $(0, \infty)$ and

$$\left\| \frac{\partial h}{\partial x}(x, y) \right\| / \ell \left(\frac{\partial h}{\partial y}(x, y) \right) \leq \omega(\|y\|) \text{ for every } (x, y) \in (D \times F) \setminus K.$$

Then, if $(a, b) \in D \times F$ and

$$Q_{a,b} = D_a \cap B \left(a, \int_{\|b\|}^{\infty} \frac{ds}{\omega(s)} \right),$$

there exists a unique continuous implicit function $\varphi : Q_{a,b} \rightarrow F$, differentiable on $Q_{a,b} \setminus A$ such that $\varphi(a) = b$ and $h(x, \varphi(x)) = h(a, b)$ for every $x \in Q_{a,b}$, and if D is starlike with respect to a and

$$\int_1^{\infty} \frac{ds}{\omega(s)} = \infty,$$

then $Q_{a,b} = D$ and $\varphi : D \rightarrow F$ is globally defined on D .

Proof. We see that $m_{m+n-2}(K_p) = 0$ for every $p \in \mathbb{N}$ and A has σ -finite $(n-1)$ -dimensional measure. We now apply the local implicit function theorem from Theorem 7 of [8], Remark 3 and the preceding arguments. \square

Using the classical implicit function theorem, we obtain the following global implicit function theorem

Theorem 9. Let E, F be Banach spaces, $D \subset E$ a domain, $h : D \times F \rightarrow F$ be continuous such that $\frac{\partial h}{\partial y}$ exists on $D \times F$, it is continuous on $D \times F$ and

$$\ell \left(\frac{\partial h}{\partial y}(x, y) \right) > 0 \text{ for every } (x, y) \in D \times F.$$

Title Page

Contents



Page 25 of 28

Go Back

Full Screen

Close



Also, let $K \subset D \times F$ be such that $A = \text{Pr}_1 K$ is a countable union of compact sets if $\dim E = \infty$ and has σ -finite $(n - 1)$ -dimensional measure if $\dim E = n$. Additionally, suppose that h is differentiable on $(D \times F) \setminus K$ and there exists $\omega : [0, \infty) \rightarrow [0, \infty)$ continuous such that $\omega > 0$ on $(0, \infty)$ and

$$\left\| \frac{\partial h}{\partial x}(x, y) \right\| / \ell \left(\frac{\partial h}{\partial y}(x, y) \right) \leq \omega(\|y\|)$$

for every $(x, y) \in (D \times F) \setminus K$. Then, if $(a, b) \in D \times F$ and

$$Q_{a,b} = D_a \cap B \left(a, \int_{\|b\|}^{\infty} \frac{ds}{\omega(s)} \right),$$

there exists a unique continuous implicit function $\varphi : Q_{a,b} \rightarrow F$, differentiable on $Q_{a,b} \setminus A$ such that $\varphi(a) = b$ and $h(x, \varphi(x)) = h(a, b)$ for every $x \in Q_{a,b}$. If D is starlike with respect to a and

$$\int_1^{\infty} \frac{ds}{\omega(s)} = \infty,$$

then $Q_{a,b} = D$ and $\varphi : D \rightarrow F$ is globally defined.

Global Implicit Function Theorem

Mihai Cristea

vol. 8, iss. 4, art. 100, 2007

Title Page

Contents



Page 26 of 28

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

References

- [1] A.V. ARUTYNOV, The implicit function theorem and abnormal points, *Doklady. Math.*, **60**(2) (1999), 231–234.
- [2] I. BLOT, On global implicit function, *Nonlinear Analysis TMA*, **17** (1992), 947–959.
- [3] M. CRISTEA, A note on global inversion theorems and applications to differential equations, *Nonlinear Analysis TMA*, **5**(11) (1981), 1155–1161.
- [4] M. CRISTEA, Local inversion theorems without assuming continuous differentiability, *J. Math. Analysis and Appl.*, **143**(1) (1989), 259–263.
- [5] M. CRISTEA, Some conditions for the local or global injectivity of a mapping between two n -manifolds, *Rev. Roumaine Math. Pures Appl.*, **33**(10) (1988), 861–869.
- [6] M. CRISTEA, Local and global inversion theorems without assuming continuous differentiability, *Bull. Math. Soc. Sci. Roumaine*, **33**(3) (1989), 233–238.
- [7] M. CRISTEA, Some conditions for the openness, local injectivity and global injectivity of the mappings between two Banach spaces, *Rev. Roumaine Math. Pures Appl.*, **36**(1-2) (1991), 67–79.
- [8] M. CRISTEA, Local inversion theorems and implicit function theorems without assuming continuous differentiability, *Bull. Math. Soc. Sci. Roumaine*, **36**(24) (1992), 227–236.
- [9] G. DARBO, Teoremi di esplicitazione in grande, *Rend. Acc. Naz. dei Lincei*, **IX**(4) (1985), 43–52.



Global Implicit Function Theorem

Mihai Cristea

vol. 8, iss. 4, art. 100, 2007

Title Page

Contents



Page 27 of 28


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [10] V.F. DEMYANOV, Theorem of an implicit function in quasidifferential calculus, *J. Math. Sciences*, **78**(5) (1996), 556–562.
- [11] J. HADAMARD, Sur les transformations ponctuelles, *Bull. Math. Sci. France*, **34** (1906), 71–84.
- [12] F. JOHN, On quasiisometric mappings I, *Comm. Pures Appl. Math.*, **21** (1968), 77–110.
- [13] M.D. LEBORGNE, Sur les théorème des fonctions implicites, *C.R. Acad. Sci. Paris*, **280**(3) (1985), 641–643.
- [14] P. LÉVY, Sur les fonctions des lignes implicites, *Bull. Soc. Math. France*, **48** (1920), 13–27.
- [15] K.J. LORENZ, An implicit function theorem in locally convex spaces, *Math. Nach.*, **129** (1986), 91–101.
- [16] M. RĂDULESCU AND S. RĂDULESCU, Global inversion theorems and applications to differential equations, *Nonlinear Analysis, TMA*, **4**(5) (1980), 951–965.
- [17] J. WARGA, Fat homeomorphisms and unbounded derivative containers, *J. Math. Analysis and Appl.*, **81** (1981), 545–560.
- [18] I.V. ZHURAVLEV, A.Yu. IGUMNOV AND V.M. MIKLYUKOV, On an implicit function theorem, Preprint 346, 2003, Univ. of Helsinki.

[Title Page](#)

[Contents](#)



Page 28 of 28

[Go Back](#)

[Full Screen](#)

[Close](#)