

AN EXAMPLE OF A STABLE FUNCTIONAL EQUATION WHEN THE HYERS METHOD DOES NOT WORK

ZOLTÁN KAISER AND ZSOLT PÁLES

Institute of Mathematics
University of Debrecen
H-4010 Debrecen, Pf. 12, Hungary

EMail: kaiserz@math.klte.hu

EMail: pales@math.klte.hu



volume 6, issue 1, article 14,
2005.

*Received 15 December, 2004;
accepted 08 February, 2005.*

Communicated by: J.M. Rassias

[Abstract](#)

[Contents](#)




[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

We show that the functional equation

$$g\left(\frac{x+y}{2}\right) = \sqrt[4]{g(x)g(y)}$$

is stable in the classical sense on arbitrary \mathbb{Q} -algebraically open convex sets, but the Hyers method does not work.

For the convenience of the reader, we have included an extensive list of references where stability theorems for functional equations were obtained using the direct method of Hyers.

2000 Mathematics Subject Classification: 39B82

Key words: Cauchy's functional equation, Stability, Hyers iteration

This research has been supported by the Hungarian Scientific Research Fund (OTKA) Grant Nos. T-043080, T-038072.

Contents

1	Introduction	3
2	Results	6
	References	

An Example of a Stable Functional Equation When the Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 2 of 25

1. Introduction


The basic problem of the stability of functional equations asks whether an 'approximate solution' of the Cauchy functional equation $g(x + y) = g(x) + g(y)$ 'can be approximated' by a solution of this equation. This problem was formulated (and also solved) in Gy. Pólya and G. Szegő's book [60] (Teil I, Aufgabe 99) for functions defined on the set of positive integers, it was reformulated in a more general form by S. Ulam in 1940 (see [86], [87]). In 1941, D. H. Hyers [40] gave the following solution to this problem: *If X and Y are Banach spaces, ε is a nonnegative real number and a function $f : X \rightarrow Y$ fulfills the inequality $\|f(x + y) - f(x) - f(y)\| \leq \varepsilon$ ($x, y \in X$), then there exists a unique solution $g : X \rightarrow Y$ of the Cauchy equation for which $\|f(x) - g(x)\| \leq \varepsilon$ ($x \in X$).* Stability problems of this type were investigated by several authors during the last decades, most of them used the idea of Hyers, which will be described below. For surveys on these developments see, e.g., the papers by Forti [16], Ger [30], Székelyhidi [85] and the book [43].

Let X and Y be non-empty sets and let $*$, \diamond be binary operations on X and Y , respectively. The Cauchy equation concerning these general structures is the functional equation

$$(1.1) \quad g(x * y) = g(x) \diamond g(y) \quad (x, y \in X),$$

where $g : X \rightarrow Y$ is considered as an unknown function.

Assuming, in addition, that Y is a metric space with metric d , we can speak about approximate solutions of (1.1): A function $f : X \rightarrow Y$ is called an ε -approximate solution of (1.1) if it satisfies the following so-called stability



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 3 of 25

inequality

$$(1.2) \quad d(f(x * y), f(x) \diamond f(y)) \leq \varepsilon \quad (x, y \in X)$$

for some $\varepsilon \geq 0$.

The Cauchy equation (1.1) is said to be *stable in the sense of Hyers and Ulam* if, for all positive δ , there exists $\varepsilon > 0$ such that, for an arbitrary solution f of (1.2), there exists a solution g of (1.1) satisfying $d(f(x), g(x)) \leq \delta$ for all $x \in X$.

The most general results concerning this stability problem were obtained in the context of square-symmetric groupoids, i.e., when the operations $*$ and \diamond satisfy the algebraic identities

$$(x * y) * (x * y) = (x * x) * (y * y) \quad \text{and} \quad (u \diamond v) \diamond (u \diamond v) = (u \diamond u) \diamond (v \diamond v)$$


for all x, y in X and u, v in Y . (Cf. [78], [15], [59], [58], [2].)

Let us denote $x*x$ by $\sigma_*(x)$ ($x \in X$), and $u \diamond u$ by $\sigma_\diamond(u)$ ($u \in Y$) (i.e., σ_* and σ_\diamond stand for the squaring in the corresponding structures). The square-symmetry of the operations $*$ and \diamond simply means that σ_* and σ_\diamond are endomorphisms. Substituting $x = y$ into (1.1) and (1.2), we get the following single-variable functional equation and functional inequality:

$$(1.3) \quad g \circ \sigma_*(x) = \sigma_\diamond \circ g(x) \quad (x \in X),$$

and

$$(1.4) \quad d(f \circ \sigma_*(x), \sigma_\diamond \circ f(x)) \leq \varepsilon \quad (x \in X).$$



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 4 of 25

Assuming that a function $f : X \rightarrow Y$ satisfies (1.2), we see that it also satisfies (1.4). In order to construct the solution g of (1.1) which is close to f , the idea of the Hyers method is to consider one of the following two iterations:


$$(1.5) \quad g_1 := f, \quad g_{n+1} = \sigma_{\diamond} \circ g_n \circ \sigma_*^{-1} \quad (n \in \mathbb{N}),$$

$$(1.6) \quad g_1 := f, \quad g_{n+1} = \sigma_{\diamond}^{-1} \circ g_n \circ \sigma_* \quad (n \in \mathbb{N}).$$

(assuming that σ_* and σ_{\diamond} is invertible, respectively) and then to show that for all solutions f of (1.4), one of these sequences of functions converges to a limit function g , which is a solution of (1.3) and of (1.1), moreover, $d(f(x), g(x)) \leq c\varepsilon$ for some $c \in \mathbb{R}$.

Results concerning stability of various functional equations in several variables using these kinds of iterations can be found in a huge number of recent works (see the extensive list of references at the end of the paper).

In this note we present an example of a stable Cauchy-type functional equation with square-symmetric operations, where for all solutions f of (1.4), the limit function of the corresponding Hyers-sequences either does not exist or is a solution of the single variable functional equation (1.3) but it does not solve (1.1) and it is not close to the original function f .



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 5 of 25

2. Results

Let X denote a vector space over the field of rational numbers throughout this paper. In what follows, we deal with the stability of the two-variable functional equation

$$(2.1) \quad g\left(\frac{x+y}{2}\right) = \sqrt[4]{g(x)g(y)} \quad (x, y \in H),$$

where H is a midpoint-convex set of X , i.e., $\frac{x+y}{2} \in H$ for all $x, y \in H$. A function $f : H \rightarrow [0, \infty[$ is called an ε -approximate solution of (2.1) if it satisfies the functional inequality

$$(2.2) \quad \left| f\left(\frac{x+y}{2}\right) - \sqrt[4]{f(x)f(y)} \right| \leq \varepsilon \quad (x, y \in H).$$

Observe that, with the notations

$$x * y := \frac{x+y}{2} \quad \text{and} \quad x \diamond y := \sqrt[4]{xy},$$


the operations $*$ and \diamond are square-symmetric (over H and $[0, \infty[$), furthermore, (2.1) and (2.2) are particular cases of (1.1) and (1.2), respectively.

With the substitution $y = x$ one obtains the following single variable functional equation and functional inequality from (2.1) and (2.2):

$$(2.3) \quad g(x) = \sqrt{g(x)} \quad (x \in H),$$

and

$$(2.4) \quad |f(x) - \sqrt{f(x)}| \leq \varepsilon \quad (x \in H),$$



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 6 of 25

respectively.

In this setting, for the iteration (1.6), we get

$$g_n(x) = (f(x))^{2^{n-1}} \quad (x \in X, n \in \mathbb{N}),$$

which is not convergent for those elements $x \in X$ where $f(x) > 1$, otherwise

$$g(x) = \lim_{n \rightarrow \infty} g_n(x) = \begin{cases} 0 & \text{if } f(x) < 1, \\ 1 & \text{if } f(x) = 1. \end{cases}$$

Clearly, g is a solution of (2.3). Assume that H has at least two elements and $0 < \varepsilon \leq 1$. Let $x_0 \in H$ be fixed. Define $f_1 : H \rightarrow [0, \infty[$ by


$$f_1(x) = \begin{cases} 1 & \text{if } x = x_0, \\ 1 + \varepsilon & \text{if } x \neq x_0, \end{cases}$$

and $f_2 : H \rightarrow [0, \infty[$ by

$$(2.5) \quad f_2(x) = \begin{cases} 1 & \text{if } x = x_0, \\ 1 - \varepsilon & \text{if } x \neq x_0. \end{cases}$$

It is not so difficult to prove, that f_1 and f_2 satisfy inequality (2.2). It is clear, that the corresponding iteration (1.6) referring to f_1 is not convergent when $x \neq x_0$. The iteration (1.6) referring to f_2 converges to

$$g(x) = \begin{cases} 1 & \text{if } x = x_0, \\ 0 & \text{if } x \neq x_0, \end{cases}$$



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 7 of 25

which is a solution of (2.3), but as we see later, does not necessarily solve (2.1). It is obvious that there does not exist a $c \in \mathbb{R}$, for which $d(f_2(x), g(x)) \leq c\varepsilon$ for an arbitrary ε . Moreover, if $\varepsilon \approx 0$, then $d(f_2(x), g(x)) \approx 1$, when $x \neq x_0$.

Similarly, for the iteration (1.5), we get

$$g_n(x) = (f(x))^{\frac{1}{2^n-1}} \quad (x \in X, n \in \mathbb{N}),$$

so we have

$$g(x) := \lim_{n \rightarrow \infty} g_n(x) = \begin{cases} 0 & \text{if } f(x) = 0, \\ 1 & \text{if } f(x) \neq 0, \end{cases}$$


which is a solution of (2.3). Assume that H has at least two elements, $0 < \varepsilon \leq 1$ and define $f_3 : H \rightarrow [0, \infty[$ by

$$(2.6) \quad f_3(x) = \begin{cases} 0 & \text{if } x = x_0, \\ \varepsilon^2 & \text{if } x \neq x_0, \end{cases}$$

where $x_0 \in H$ is fixed. It is obvious, that (2.2) holds for the function f_3 , but the Hyers iteration now converges to

$$g(x) = \begin{cases} 0 & \text{if } x = x_0, \\ 1 & \text{if } x \neq x_0, \end{cases}$$

which solves (2.3) but does not necessarily solve (2.1). Again as before, there does not exist a $c \in \mathbb{R}$, for which $d(f_3(x), g(x)) \leq c\varepsilon$ for an arbitrary ε , because if ε is approximately zero, then $d(f_3(x), g(x))$ is approximately 1 when $x \neq x_0$.



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 8 of 25

In what follows, we prove the stability of the functional equations (2.3) and (2.1). It can be immediately seen that the solutions of (2.3) are functions with values 0 and 1, that is, characteristic functions of a certain subset of H . The next result shows that if f is a solution of (2.4) then it is close to a certain characteristic function as well. Thus, the functional equation (2.3) is stable in the Hyers-Ulam sense.

Theorem 2.1. *Let H be a nonempty set and let $f : H \rightarrow [0, \infty[$ be a solution of the functional inequality (2.4) with $0 \leq \varepsilon \leq \varepsilon_0 := 4/25$. Then, for all $x \in H$,*

$$\text{either } f(x) \leq \frac{25\varepsilon^2}{16} \quad \text{or} \quad -\frac{9\varepsilon}{4} \leq f(x) - 1 \leq 2\varepsilon.$$


Proof. Define the subsets A and B of H by

$$A := \left\{ x \in H : -\frac{9\varepsilon}{4} \leq f(x) - 1 \leq 2\varepsilon \right\}, \quad B := \left\{ x \in H : f(x) \leq \frac{25\varepsilon^2}{16} \right\}.$$

The proof of the theorem is equivalent to showing that A and B form a partition of H .

Let $x \in H$ be arbitrary. Inequality (2.4) is equivalent to the quadratic inequalities

$$(2.7) \quad -\varepsilon \leq \left(\sqrt{f(x)} \right)^2 - \sqrt{f(x)} \leq \varepsilon.$$



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 9 of 25

Since $\varepsilon \leq \varepsilon_0 = 4/25$, we have the estimate

$$\begin{aligned}
 (2.8) \quad \frac{1 - \sqrt{1 - 4\varepsilon}}{2} &= \frac{4\varepsilon}{2(1 + \sqrt{1 - 4\varepsilon})} \\
 &\leq \frac{2\varepsilon}{1 + \sqrt{1 - 4\varepsilon_0}} \\
 &= \frac{2\varepsilon}{1 + \sqrt{9/25}} \leq \frac{5}{4}\varepsilon.
 \end{aligned}$$

From (2.7), using (2.8), we obtain that either


$$0 \leq f(x) = (\sqrt{f(x)})^2 \leq \left(\frac{1 - \sqrt{1 - 4\varepsilon}}{2}\right)^2 \leq \frac{25\varepsilon^2}{16},$$

i.e., $x \in B$, or

$$\frac{1 + \sqrt{1 - 4\varepsilon}}{2} \leq \sqrt{f(x)} \leq \frac{1 + \sqrt{1 + 4\varepsilon}}{2},$$

consequently, in view of the estimate (2.8) and

$$(2.9) \quad \frac{\sqrt{1 + 4\varepsilon} - 1}{2} = \frac{4\varepsilon}{2(\sqrt{1 + 4\varepsilon} + 1)} \leq \varepsilon,$$



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 10 of 25

we have

$$\begin{aligned}
 1 - \frac{9\varepsilon}{4} &\leq 1 - \frac{1 - \sqrt{1 - 4\varepsilon}}{2} - \varepsilon = \left(\frac{1 + \sqrt{1 - 4\varepsilon}}{2} \right)^2 \\
 &\leq f(x) \\
 &\leq \left(\frac{1 + \sqrt{1 + 4\varepsilon}}{2} \right)^2 = 1 + \frac{\sqrt{1 + 4\varepsilon} - 1}{2} + \varepsilon \leq 1 + 2\varepsilon,
 \end{aligned}$$

which means that $x \in A$.


Thus we have showed that $A \cup B = H$. On the other hand, since $\varepsilon \leq \varepsilon_0 = 4/25$, it easily follows that $A \cap B = \emptyset$. \square

In order to investigate the stability of the two-variable functional equation (2.1), we need the notion of an ideal of midpoint-convex sets. We say that a set $I \subset H$ is an *ideal in the midpoint-convex set H with respect to the midpoint operation* if

$$(2.10) \quad x \in H \text{ and } y \in I \implies \frac{x+y}{2} \in I.$$

Trivially, \emptyset and H are always ideals in H . However, in general, there could exist further ideals in H . For instance, if H is the closed unit interval $H = [0, 1] \subset \mathbb{R}$ then the sets $]0, 1[$, $[0, 1[$, and $]0, 1]$ are also ideals for H . As we shall see below, if H enjoys a certain openness property then it can have only trivial ideals.

We say that a set H is \mathbb{Q} -*algebraically open* if, for each point $p \in H$ and vector $v \in X$, there exists a positive number τ such that $p + tv \in H$ for all $t \in [0, \tau] \cap \mathbb{Q}$. It is obvious, that every open set (of a topological linear space) is \mathbb{Q} -algebraically open, but the reversed statement is not true in general.



An Example of a Stable Functional Equation When the Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 11 of 25

Lemma 2.2. Let $H \subset X$ be a \mathbb{Q} -algebraically open midpoint-convex set. Then H has only trivial ideals, i.e., the only ideals in H are the sets \emptyset and H .

Proof. Assume that $I \subset H$ is a nonempty ideal with respect to the midpoint operation, and let $y \in I$ be fixed. It easily follows by induction that $\frac{2^n-1}{2^n}x + \frac{1}{2^n}y \in I$ for all $x \in H$.

Now let $x \in H$ be arbitrary. Since H is \mathbb{Q} -algebraically open, for large $n \in \mathbb{N}$, we have that

$$x_n = x + \frac{1}{2^n-1}(x-y) \in H.$$

Then

$$\frac{2^n-1}{2^n}x_n + \frac{1}{2^n}y = x,$$

which, in view of the ideal property of I , yields that $x \in I$. Therefore, $H \subset I$ follows. \square


Our next result concerns the stability of the functional equation (2.1).

Theorem 2.3. Let $H \subset X$ be a \mathbb{Q} -algebraically open midpoint-convex set and let $f : H \rightarrow [0, \infty[$ be a solution of the functional inequality (2.2) with $0 \leq \varepsilon \leq \varepsilon_0 := 4/25$. Then, either

$$f(x) \leq \frac{25\varepsilon^2}{16} \quad (x \in H)$$

or

$$-\frac{9\varepsilon}{4} \leq f(x) - 1 \leq 2\varepsilon \quad (x \in H).$$



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 12 of 25

Proof. Define the sets A and B as in the proof of Theorem 2.1. Then, by Theorem 2.1, these sets form a partition of H . In order to complete the proof, we have to show that one of the sets A or B is empty. To do that, we prove that B is an ideal in H with respect to the midpoint operation.


Let $x \in H$ and $y \in B$. It suffices to prove that $\frac{x+y}{2} \notin A$, because then we have $\frac{x+y}{2} \in H \setminus A = B$. From inequality (2.2) it follows that

$$\begin{aligned} f\left(\frac{x+y}{2}\right) &\leq \varepsilon + \sqrt{\sqrt{f(x)}\sqrt{|f(y)|}} \\ &\leq \varepsilon + \sqrt{\sqrt{1+2\varepsilon}\sqrt{\frac{25\varepsilon^2}{16}}} \\ &\leq \varepsilon_0 + \sqrt{\sqrt{1+2\varepsilon_0}\frac{5\varepsilon_0}{4}} \\ &= \frac{4}{25} + \frac{\sqrt[4]{33}}{5} < \frac{16}{25} = 1 - \frac{9\varepsilon_0}{4} \leq 1 - \frac{9\varepsilon}{4}. \end{aligned}$$

In view of Lemma 2.2, we have that B is a trivial ideal, i.e., either $B = H$ or $B = \emptyset$ which means that $A = H$, and the statement of the theorem follows from this. \square

The functions $g \equiv 0$ and $g \equiv 1$ are trivially the solutions of the functional equation (2.1). Choosing $\varepsilon = 0$ in Theorem 2.3, we immediately get that the reversed statement is also true, i.e., we have the following result:

Corollary 2.4. *Let X be a real linear space, $H \subset X$ be a \mathbb{Q} -algebraically open midpoint-convex set. Then a function $g : H \rightarrow [0, \infty[$ is a solution of the functional equation (2.1) if and only if either $g \equiv 0$ or $g \equiv 1$.*



An Example of a Stable Functional Equation When the Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back


Close

Quit

Page 13 of 25

Now Theorem 2.3 can be interpreted as the stability theorem of (2.1) since it states that if f solves the stability inequality (2.2), then it is close to one of the solutions of the functional equation (2.1). Thus, (2.1) is stable in the Hyers-Ulam sense.

On the other hand, Corollary 2.4 shows that if we consider equation (2.1) over a \mathbb{Q} -algebraically open midpoint-convex set, then the limits of the corresponding Hyers-sequences referring to the functions f_2 and f_3 defined in (2.5) and (2.6) are not solutions of (2.1), so the stability of this functional equation cannot be proved via the Hyers-method.



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back


Close

Quit

Page 14 of 25

References

- [1] J.-H. BAE AND H.-M. KIM, On the generalized Hyers-Ulam stability of a quadratic mapping, *Far East J. Math. Sci.*, **3**(4) (2001), 599–608. MR 2002d:39043
- [2] K. BARON AND P. VOLKMANN, On functions close to homomorphisms between square symmetric structures, *Seminar LV* (<http://www.mathematik.uni-karlsruhe.de/~semlv/>) (2002), no. 14, 12 pp. (electronic).
- [3] C. BORELLI FORTI AND G.-L. FORTI, On a general Hyers-Ulam stability result, *Internat. J. Math. Math. Sci.*, **18**(2) (1995), 229–236. MR 96c:39019
- [4] D.G. BOURGIN, Approximate isometries, *Bull. Amer. Math. Soc.*, **52** (1946), 704–714. MR 8,157f
- [5] D.G. BOURGIN, Approximately isometric and multiplicative transformations on continuous function rings, *Duke Math. J.*, **16** (1949), 385–397. MR 11,115e
- [6] D.G. BOURGIN, Classes of transformations and bordering transformations, *Bull. Amer. Math. Soc.*, **57** (1951), 223–237. MR 13,138d
- [7] I.-S. CHANG AND H.-M. KIM, On the Hyers-Ulam stability of quadratic functional equations, *J. Inequal. Pure Appl. Math.*, **3**(3) (2002), Art. 33, 12 pp. (electronic). [ONLINE: <http://jipam.vu.edu.au/article.php?sid=185>] MR 2003f:39092



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 15 of 25

- [8] I.-S. CHANG, E. H. LEE AND H.-M. KIM, On Hyers-Ulam-Rassias stability of a quadratic functional equation, *Math. Inequal. Appl.*, **6**(1) (2003), 87–95. MR 2003m:39039
- [9] P.W. CHOLEWA, Remarks on the stability of functional equations, *Aequationes Math.*, **27**(1-2) (1984), 76–86. MR 86d:39016
- [10] P.W. CHOLEWA, The stability problem for a generalized Cauchy type functional equation, *Rev. Roumaine Math. Pures Appl.*, **29**(6) (1984), 457–460. MR 86c:39013
- [11] S. CZERWIK, On the stability of the quadratic mapping in normed spaces, *Abh. Math. Sem. Univ. Hamburg*, **62** (1992), 59–64. MR 94e:39026
- [12] S. CZERWIK, The stability of the quadratic functional equation, Stability of mappings of Hyers-Ulam type, *Hadronic Press Collect. Orig. Artic.*, Hadronic Press, Palm Harbor, FL, 1994, pp. 81–91. MR 95i:39025
- [13] S. CZERWIK, On the stability of the quadratic functional equation and related topics, *Recent Progress in Inequalities* (Niš, 1996), *Math. Appl.*, Vol. 430, Kluwer Acad. Publ., Dordrecht, 1998, pp. 449–455. MR 99a:39064
- [14] S. CZERWIK, *Functional Equations and Inequalities in Several Variables*, World Scientific Publishing Co. Inc., River Edge, NJ, 2002. MR 2003b:39027
- [15] G.-L. FORTI, An existence and stability theorem for a class of functional equations, *Stochastica*, **4**(1) (1980), 23–30. MR 81i:39006



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 16 of 25

- [16] G.-L. FORTI, Hyers-Ulam stability of functional equations in several variables, *Aequationes Math.*, **50**(1-2) (1995), 143–190. MR 96i:39033
- [17] G.-L. FORTI, Comments on the core of the direct method for proving Hyers-Ulam stability of functional equations, *J. Math. Anal. Appl.*, **295**(1) (2004), 127–133.
- [18] G.-L. FORTI AND J. SCHWAIGER, Stability of homomorphisms and completeness, *C. R. Math. Rep. Acad. Sci. Canada*, **11**(6) (1989), 215–220. MR 90k:20095
- [19] Z. GAJDA, On stability of additive mappings, *Internat. J. Math. Math. Sci.*, **14**(3) (1991), 431–434. MR 92e:39029
- [20] P. GĂVRUȚĂ, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, *J. Math. Anal. Appl.*, **184**(3) (1994), 431–436. MR 95e:47089
- [21] P. GĂVRUȚĂ, On the stability of some functional equations, Stability of mappings of Hyers-Ulam type, *Hadronic Press Collect. Orig. Artic.*, Hadronic Press, Palm Harbor, FL, 1994, pp. 93–98. MR 95k:39027
- [22] P. GĂVRUȚĂ, On the Hyers-Ulam-Rassias stability of mappings, *Recent Progress in Inequalities* (Niš, 1996), *Math. Appl.*, vol. 430, Kluwer Acad. Publ., Dordrecht, 1998, pp. 465–469.
- [23] P. GĂVRUȚĂ, Hyers-Ulam stability of Hosszú's equation, *Functional Equations and Inequalities*, *Math. Appl.*, vol. 518, Kluwer Acad. Publ., Dordrecht, 2000, pp. 105–110.



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 17 of 25

- [24] P. GĂVRUȚĂ, On a problem of G. Isac and Th. M. Rassias concerning the stability of mappings, *J. Math. Anal. Appl.*, **261**(2) (2001), 543–553. MR 2002k:39032
- [25] P. GĂVRUȚĂ, On the Hyers-Ulam-Rassias stability of the quadratic mappings, *Nonlinear Funct. Anal. Appl.*, **9**(3) (2004), 415–428.
- [26] P. GĂVRUȚĂ AND L. CĂDARIU, Inequalities for approximately additive mappings, *Bul. Științ. Univ. Politeh. Timiș. Ser. Mat. Fiz.*, **43**(57)(1) (1998), 24–30. MR 99k:39074
- [27] P. GĂVRUȚĂ AND L. CĂDARIU, General stability of the cubic functional equation, *Bul. Științ. Univ. Politeh. Timiș. Ser. Mat. Fiz.*, **47**(61)(1) (2002), 59–70. MR 2004a:39057
- [28] P. GĂVRUȚĂ, M. HOSSU, D. POPESCU AND C. CĂPRĂU, On the stability of mappings and an answer to a problem of Th. M. Rassias, *Ann. Math. Blaise Pascal*, **2**(2) (1995), 55–60. MR 97f:39035
- [29] R. GER, On functional inequalities stemming from stability questions, *General Inequalities*, **6** (Oberwolfach, 1990) (W. Walter, ed.), *International Series of Numerical Mathematics*, Birkhäuser, Basel–Boston, 1992, pp. 227–240. MR 94b:39042
- [30] R. GER, A survey of recent results on stability of functional equations, *Proc. of the 4th International Conference on Functional Equations and Inequalities* (Cracow), Pedagogical University of Cracow, 1994, pp. 5–36.



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 18 of 25

- [31] A. GILÁNYI, Charakterisierung von monomialen Funktionen und Lösung von Funktionalgleichungen mit Computern, Diss., Universität Karlsruhe, Karlsruhe, Germany, 1995.
- [32] A. GILÁNYI, A characterization of monomial functions, *Aequationes Math.*, **54**(3) (1997), 289–307. MR 99g:39027
- [33] A. GILÁNYI, On Hyers-Ulam stability of monomial functional equations, *Abh. Math. Sem. Univ. Hamburg*, **68** (1998), 321–328. MR 99k:39063
- [34] A. GILÁNYI, On the stability of the square-norm equation, *Publ. Math. Debrecen*, **52**(3-4) (1998), 419–427. MR 99e:39082
- [35] A. GILÁNYI, Hyers-Ulam stability of monomial functional equations on a general domain, *Proc. Natl. Acad. Sci. USA*, **96**(19) (1999), 10588–10590.
- [36] A. GILÁNYI, On the stability of monomial functional equations, *Publ. Math. Debrecen*, **56**(1-2) (2000), 201–212. MR 2001b:39025
- [37] A. GILÁNYI, Über die Stabilität monomialer Funktionalgleichungen, Kummulative Habilitationsschrift, Universität Karlsruhe, Karlsruhe, Germany, 2000.
- [38] A. GILÁNYI, On approximately monomial functions, *Functional Equations – Results and Advances* (Z. Daróczy and Zs. Páles, eds.), *Advances in Mathematics*, vol. 3, Kluwer Acad. Publ., Dordrecht, 2002, pp. 99–111. MR 2003e:39062
- [39] A. GILÁNYI, Z. KAISER AND Zs. PÁLES, Estimates to the stability of the Cauchy equation, manuscript.



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 19 of 25

- [40] D.H. HYERS, On the stability of the linear functional equation, *Proc. Natl. Acad. Sci. U.S.A.*, **27** (1941), 222–224. MR 2,315a
- [41] D.H. HYERS, The stability of homomorphisms and related topics, *Global Analysis – Analysis on Manifolds*, Teubner-Texte Math., vol. 57, Teubner, Leipzig, 1983, pp. 140–153. MR 86a:39004
- [42] D.H. HYERS, G. ISAC AND Th.M. RASSIAS, On the asymptoticity aspect of Hyers-Ulam stability of mappings, *Proc. Amer. Math. Soc.*, **126**(2) (1998), 425–430. MR 98d:39004
- [43] D.H. HYERS, G. ISAC AND Th.M. RASSIAS, Stability of Functional Equations in Several Variables, *Progress in Nonlinear Differential Equations and their Applications*, 34, Birkhäuser Boston Inc., Boston, MA, 1998. MR 99i:39035
- [44] K.-W. JUN AND H.-M. KIM, Remarks on the stability of additive functional equation, *Bull. Korean Math. Soc.*, **38**(4) (2001), 679–687. MR 2003a:39034
- [45] K.-W. JUN AND H.-M. KIM, The generalized Hyers-Ulam-Rassias stability of a cubic functional equation, *J. Math. Anal. Appl.*, **274**(2) (2002), 267–278. MR 2003j:39077
- [46] K.-W. JUN AND H.-M. KIM, On the Hyers-Ulam-Rassias stability of a general cubic functional equation, *Math. Inequal. Appl.*, **6**(2) (2003), 289–302. MR 2004c:39061



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 20 of 25

- [47] S.-M. JUNG AND P.K. SAHOO, On the Hyers-Ulam stability of a functional equation of Davison, *Kyungpook Math. J.*, **40**(1) (2000), 87–92. MR 2001d:39027
- [48] S.-M. JUNG AND P.K. SAHOO, A functional equation characterizing cubic polynomials and its stability, *Int. J. Math. Math. Sci.*, **27**(5) (2001), 301–307. MR 2002m:39023
- [49] S.-M. JUNG AND P.K. SAHOO, Hyers-Ulam stability of the quadratic equation of Pexider type, *J. Korean Math. Soc.*, **38**(3) (2001), 645–656. MR 2003d:39047
- [50] S.-M. JUNG AND P.K. SAHOO, Hyers-Ulam stability of a generalized Hosszú functional equation, *Glas. Mat. Ser. III*, **37**(57)(2) (2002), 283–292. MR 2003k:39038
- [51] Z. KAISER, On stability of the Cauchy equation in normed spaces over fields with valuation, *Publ. Math. Debrecen*, **64**(1-2) (2004), 189–200.
- [52] G.H. KIM, A generalization of the Hyers-Ulam-Rassias stability of the beta functional equation, *Publ. Math. Debrecen*, **59**(1-2) (2001), 111–119. MR 2002i:39016
- [53] G.H. KIM, On the stability of functional equations with square-symmetric operation, *Math. Inequal. Appl.*, **4**(2) (2001), 257–266. MR 2003d:39044
- [54] G.H. KIM, On the stability of the quadratic mapping in normed spaces, *Int. J. Math. Math. Sci.*, **25**(4) (2001), 217–229. MR 2002b:39020



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 21 of 25

- [55] G.H. KIM, Y.W. LEE AND K.S. JI, Modified Hyers-Ulam-Rassias stability of functional equations with square-symmetric operation, *Commun. Korean Math. Soc.*, **16**(2) (2001), 211–223. MR 2002h:39036
- [56] M. KUCZMA, *An Introduction to the Theory of Functional Equations and Inequalities*, Prace Naukowe Uniwersytetu Śląskiego w Katowicach, vol. 489, Państwowe Wydawnictwo Naukowe — Uniwersytet Śląski, Warszawa–Kraków–Katowice, 1985. MR 86i:39008
- [57] Zs. PÁLES, Generalized stability of the Cauchy functional equation, *Aequationes Math.*, **56**(3) (1998), 222–232. MR 99k:39076
- [58] Zs. PÁLES, Hyers-Ulam stability of the Cauchy functional equation on square-symmetric grupoids, *Publ. Math. Debrecen*, **58**(4) (2001), 651–666. MR 2002f:39058
- [59] Zs. PÁLES, P. VOLKMANN AND R.D. LUCE, Hyers-Ulam stability of functional equations with a square-symmetric operation, *Proc. Natl. Acad. Sci. USA*, **95**(22) (1998), 12772–12775 (electronic). MR 2000a:39030
- [60] Gy. PÓLYA AND G. SZEGŐ, *Aufgaben und Lehrsätze aus der Analysis*, Vol. I, Springer, Berlin, 1954. (2nd edition) MR 15,512a
- [61] D. POPA, Functional inclusions on square-symmetric groupoids and Hyers-Ulam stability, *Math. Inequal. Appl.*, **7**(3) (2004), 419–428.
- [62] J.M. RASSIAS, On approximation of approximately linear mappings by linear mappings, *J. Funct. Anal.*, **46**(1) (1982), 126–130. MR 84e:47027



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 22 of 25

- [63] J.M. RASSIAS, Solution of a problem of Ulam, *J. Approx. Theory*, **57**(3) (1989), 268–273. MR 90d:46034
- [64] J.M. RASSIAS, Solution of a stability problem of Ulam, *Discuss. Math.*, **12** (1992), 95–103 (1993). MR 94d:39010
- [65] J.M. RASSIAS, Complete solution of the multi-dimensional problem of Ulam, *Discuss. Math.*, **14** (1994), 101–107. MR 96g:39019
- [66] J.M. RASSIAS, On the stability of a multi-dimensional Cauchy type functional equation, *Geometry, Analysis and Mechanics*, World Sci. Publishing, River Edge, NJ, 1994, pp. 365–376. MR 96g:39018
- [67] J.M. RASSIAS, Solution of a stability problem of Ulam, *Functional Analysis, Approximation Theory and Numerical Analysis*, World Sci. Publishing, River Edge, NJ, 1994, pp. 241–249. MR 96b:47066
- [68] J.M. RASSIAS, Solution of a Cauchy-Jensen stability Ulam type problem, *Arch. Math.* (Brno), **37**(3) (2001), 161–177. MR 2002h:39034
- [69] J.M. RASSIAS, Solution of a Cauchy-Jensen stability Ulam type problem, *Cienc. Mat.* (Havana), **19**(1) (2001), 45–60. MR 2002f:39061
- [70] J.M. RASSIAS AND M.J. RASSIAS, On the Ulam stability of Jensen and Jensen type mappings on restricted domains, *J. Math. Anal. Appl.*, **281**(2) (2003), 516–524. MR 2004g:39052
- [71] Th.M. RASSIAS, On the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.*, **72**(2) (1978), 297–300. MR 80d:47094



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 23 of 25

- [72] Th.M. RASSIAS, On a modified Hyers-Ulam sequence, *J. Math. Anal. Appl.*, **158**(1) (1991), 106–113. MR 92d:46046
- [73] Th.M. RASSIAS, On a problem of S. M. Ulam and the asymptotic stability of the Cauchy functional equation with applications, *General Inequalities*, 7 (Oberwolfach, 1995), *Internat. Ser. Numer. Math.*, vol. 123, Birkhäuser, Basel, 1997, pp. 297–309. MR 99e:39092
- [74] Th.M. RASSIAS, On the stability of functional equations and a problem of Ulam, *Acta Appl. Math.*, **62**(1) (2000), 23–130. MR 2001j:39042
- [75] Th.M. RASSIAS AND P. ŠEMRL, On the Hyers-Ulam stability of linear mappings, *J. Math. Anal. Appl.*, **173**(2) (1993), 325–338. MR 94d:39011
- [76] Th.M. RASSIAS AND J. TABOR, What is left of Hyers-Ulam stability?, *J. Natur. Geom.*, **1**(2) (1992), 65–69. MR 93b:47131
- [77] Th.M. RASSIAS AND J. TABOR (eds.), Stability of mappings of Hyers-Ulam type, *Hadronic Press Collection of Original Articles*, Hadronic Press Inc., Palm Harbor, FL, 1994. MR 95f:39001
- [78] J. RÄTZ, On approximately additive mappings, *General Inequalities*, 2 (Oberwolfach, 1978) (Basel–Boston) (E. F. Beckenbach, ed.), *International Series in Numerical Mathematics*, vol. 47, Birkhäuser, 1980, pp. 233–251. MR 83e:90131
- [79] J. SCHWAIGER, Functional equations for homogeneous polynomials arising from multilinear mappings and their stability, *Ann. Math. Sil.*, **8** (1994), 157–171, Polish-Austrian Seminar on Functional Equations and Iteration Theory (Cieszyn, 1994). MR 96e:39020



**An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work**

Zoltán Kaiser and Zsolt Páles

Title Page

Contents




Go Back

Close

Quit

Page 24 of 25

- [80] J. SCHWAIGER, The functional equation of homogeneity and its stability properties, *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II*, **205** (1996), 3–12 (1997). MR 99j:39024
- [81] F. SKOF, Local properties and approximation of operators, *Rend. Sem. Mat. Fis. Milano*, **53** (1983), 113–129 (1986), *Geometry of Banach spaces and related topics* (Italian) (Milan, 1983). MR 87m:39009
- [82] F. SKOF, On the approximation of locally δ -additive mappings, *Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur.*, **117**(4-6) (1983), 377–389 (1986). MR 89a:39015
- [83] F. SKOF, Approximation of δ -quadratic functions on a restricted domain, *Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur.*, **118**(1-2) (1984), 58–70. MR 86k:39005
- [84] F. SKOF, On δ -isometries in normed spaces, *Rend. Mat. Appl.*, (7) **10**(4) (1990), 853–866 (1991). MR 92j:46020
- [85] L. SZÉKELYHIDI, Ulam's problem, Hyers's solution — and to where they led, *Functional Equations and Inequalities* (Th. M. Rassias, ed.), *Math. Appl.*, vol. 518, Kluwer Acad. Publ., Dordrecht, 2000, pp. 259–285. MR 2003a:39035
- [86] S.M. ULAM, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience Publishers, New York-London, 1960. MR 22 #10884
- [87] S.M. ULAM, *Problems in Modern Mathematics*, John Wiley & Sons, New York, 1964. MR 43 #6031



An Example of a Stable
Functional Equation When the
Hyers Method Does Not Work

Zoltán Kaiser and Zsolt Páles

Title Page

Contents



Go Back

Close

Quit

Page 25 of 25