



Symbolic Dynamics of $(-\beta)$ -Expansions

K. Dajani and S. D. Ramawadh

Department of Mathematics

Utrecht University

Postbus 80.010

3508 TA Utrecht

The Netherlands

k.dajani1@uu.nl

S.D.Ramawadh@uu.nl

Abstract

We study shift spaces associated with a family of transformations generating expansions in base $-\beta$, with $\beta > 1$. We give a complete characterization when these shift spaces are sofic or even shifts of finite type.

1 Introduction

In 1957, Rényi studied the expansions of numbers in base $\beta > 1$. He only considered the numbers in $[0, 1)$, and only considered one specific rule, the β -transformation, with which we could find these expansions [12]. Many other mathematicians have studied these expansions since then, of which the most famous result is a paper by Parry [10], in which he gave quite an extensive description of both ergodic and symbolic properties of the β -transformation posed by Rényi. Later on several others looked at other types of β -expansions. Erdős [6], for example, studied lazy β -expansions (the opposite, the greedy β -expansions were studied by Rényi). Dajani and Kraaikamp studied a whole class of β -expansions [2] and also random β -expansions [3, 4].

In 2009, Ito and Sadahiro [8] studied $(-\beta)$ -expansions in base $-\beta$, $\beta > 1$. They introduced a transformation, the $(-\beta)$ -transformation, which allowed them to find an expansion of the form

$$x = \sum_{n=1}^{\infty} \frac{x_n}{(-\beta)^n}, \quad x_n \in \{0, 1, \dots, \lfloor \beta \rfloor\} \quad (1)$$

of any number $x \in I := \left[-\frac{\beta}{\beta+1}, \frac{1}{\beta+1}\right)$. After having introduced this dynamical system, they continued their study of these expansions by introducing an order on the set of expansions and studying the resulting $(-\beta)$ -shift. This approach was later continued by Frougny and Lai [7], who gave a detailed description of the symbolic properties of the $(-\beta)$ -shift. Recently, others have considered different transformations generating $(-\beta)$ -expansions. Dombek, Masáková and Pelantová [5] considered a generalization of the aforementioned approach. Dajani and Kalle [1], took an even more general approach by studying a whole class of transformations generating $(-\beta)$ -expansions for $1 < \beta < 2$.

The aim of this paper is to study shift spaces associated with various transformations generating $(-\beta)$ -expansions and to give a complete characterization when these shift spaces are sofic or shifts of finite type.

2 The dynamical system

Let $\beta > 1$ be given. The set of numbers $x \in \mathbb{R}$ that satisfy (1) for certain coefficients $x_n \in \{0, 1, \dots, \lfloor \beta \rfloor\}$ is contained in the interval

$$I_{-\beta} := \left[-\frac{\beta \cdot \lfloor \beta \rfloor}{\beta^2 - 1}, \frac{\lfloor \beta \rfloor}{\beta^2 - 1}\right].$$

We call the set $\mathcal{A} := \{0, 1, \dots, \lfloor \beta \rfloor\}$ the *canonical alphabet*. If x has an expansion of the form (1), then we may refer to the right-hand side of (1) as a $(-\beta)$ -*expansion of x* . Moreover, we can refer to the digits of the expansion explicitly as follows:

$$d_{-\beta}(x) := (x_1, x_2, \dots) \quad \text{and} \quad d_{-\beta}(x, n) := x_n.$$

We may also write $x = (.x_1x_2\cdots)_{-\beta}$. Moreover, if a part is periodic we can show this by overlining it. For example, $(1, \overline{0, 2}) = (1, 0, 2, 0, 2, 0, 2, \dots)$. If a sequence consists of only a periodic part, we will call it *purely periodic*. If it consists of a pre-periodic and a periodic part, we will call it *eventually periodic*. Now, suppose that $x \in I_{-\beta}$ with expansion $d_{-\beta}(x) = (x_1, x_2, \dots)$. It follows that

$$-\beta x - x_1 = \sum_{n=1}^{\infty} \frac{x_{n+1}}{(-\beta)^n}.$$

In particular, we see that $-\beta x - x_1 \in I_{-\beta}$ as the right-hand side is another $(-\beta)$ -expansion. Therefore, one should look at the family of maps $-\beta x - a$, $a \in \mathcal{A}$, on the interval $I_{-\beta}$. Figure 1 shows what happens for a specific choice for β . Notice that there is a small subinterval for which there is more than one possibility for a such that $-\beta x - a$ remains in $I_{-\beta}$ for all points in this subinterval. Such a subinterval will be called a *choice region*. Other subintervals, where there is only one choice for $a \in \mathcal{A}$, are called *uniqueness regions*. Note that this behaviour is not restricted to this specific base $-\beta$; in fact, it happens for all non-integer base $-\beta$. This is a consequence of our alphabet \mathcal{A} .

In any case, we need to specify a function $d : I_{-\beta} \rightarrow \mathcal{A}$, the *digit function*, such that for any $x \in I_{-\beta}$ it follows that also $-\beta x - d(x) \in I_{-\beta}$. Henceforth, we will work with the following assumptions:




Figure 1: The maps $-\beta x - a$ with $a \in \mathcal{A}$ on $I_{-\beta}$ for $\beta = \frac{1+\sqrt{5}}{2}$.

Assumption 1. The digit function $d : I_{-\beta} \rightarrow \mathcal{A}$ satisfies the following properties:

- i. $d(x)$ is such that $x \in I_{-\beta} \Rightarrow -\beta x - d(x) \in I_{-\beta}$,
- ii. d is either left- or right-continuous everywhere,
- iii. d is nonincreasing on $I_{-\beta}$.

The idea behind the assumptions on the digit function is that it divides the interval $I_{-\beta}$ into at most $\lfloor \beta \rfloor + 1$ smaller intervals, where the intervals are all either left- or right-closed and each of the intervals has a digit from the alphabet \mathcal{A} assigned to them. These digits are assigned in such a way that, from left to right, these digits are decreasing. This implies that all digits of \mathcal{A} are actually used.

Property i of Assumption 2.1 is worth considering more carefully. As a consequence of our choice of alphabet \mathcal{A} , we have that for any $\beta > 1$ and any $x \in I_{-\beta}$ there is at least one $a \in \mathcal{A}$ such that $-\beta x - a \in I_{-\beta}$. To see why, fix $x \in I_{-\beta}$ and consider the set $S(x) = \{-\beta x - a : a \in \mathcal{A}\}$. The smallest element of $S(x)$ cannot be greater than $\frac{\lfloor \beta \rfloor}{\beta^2 - 1}$ and the largest element of $S(x)$ cannot be smaller than $-\frac{\beta \cdot \lfloor \beta \rfloor}{\beta^2 - 1}$. Moreover, $I_{-\beta}$ has length strictly greater than 1, whereas the difference between two consecutive digits in \mathcal{A} is 1. These observations imply that $S(x) \cap I_{-\beta} \neq \emptyset$.

The points where d is discontinuous are called *cut points*. The cut points always lie in choice regions, and the cut points divide these regions into smaller regions on which the digit function dictates which map $-\beta x - a$ to choose.

Using this digit function, we can finally pose our dynamical system. We define, for a given digit function d , the map

$$T_{-\beta}(x) := -\beta x - d(x)$$

on $I_{-\beta}$. If we want to emphasize whether the underlying digit function is left- or right-continuous, we may write $T_{L,-\beta}(x)$ and $T_{R,-\beta}(x)$ respectively.

Proposition 2. *Let $x \in I_{-\beta}$ be arbitrary and let $x_n := d(T_{-\beta}^{n-1}(x))$. Then we have $x = (.x_1 x_2 \dots)_{-\beta}$.*

Proof. We show, by induction on n , that for all $n \geq 1$ we have

$$x = \frac{T_{-\beta}^n(x)}{(-\beta)^n} + \sum_{k=1}^n \frac{x_k}{(-\beta)^k}.$$

By definition, we have $T_{-\beta}(x) = -\beta x - x_1$, which in turn is equivalent to $x = \frac{T_{-\beta}(x)}{-\beta} + \frac{x_1}{-\beta}$. Hence, the statement holds for $n = 1$.

Now assume that the statement is true for some positive integer $m - 1$. Since we know that $T_{-\beta}^m(x) = -\beta T_{-\beta}^{m-1}(x) - x_m$, we also have

$$T_{-\beta}^{m-1}(x) = \frac{x_m}{-\beta} + \frac{T^m(x)}{-\beta}$$

and hence

$$\begin{aligned} x &= \frac{T_{-\beta}^{m-1}(x)}{(-\beta)^{m-1}} + \sum_{k=1}^{m-1} \frac{x_k}{(-\beta)^k} \\ &= \frac{\frac{x_m}{-\beta} + \frac{T^m(x)}{-\beta}}{(-\beta)^{m-1}} + \sum_{k=1}^{m-1} \frac{x_k}{(-\beta)^k} \\ &= \frac{T_{-\beta}^m(x)}{(-\beta)^m} + \sum_{k=1}^m \frac{x_k}{(-\beta)^k}. \end{aligned}$$

We see that the statement is true for m as well. The proof is finished by noting that $\frac{T_{-\beta}^m(x)}{(-\beta)^m} \rightarrow 0$ as $m \rightarrow \infty$. \square

The following example shows that the original $(-\beta)$ -transformation posed by Ito and Sadahiro [8] can be viewed as restrictions of our maps.

Example 3. Let the base be given by $\beta = \frac{1+\sqrt{5}}{2}$ and consider the map $T_{L,-\beta}$ with cut point $2 - \sqrt{5}$. This map is shown in Figure 2. The smaller box in Figure 2 is the $(-\beta)$ -




Figure 2: The map $T_{L,-\beta}$ from Example 3.

transformation as defined by Ito and Sadahiro. Hence, for this particular β and choice of cut point, our map $T_{L,-\beta}$ is an extension of their $(-\beta)$ -transformation.

Finding $(-\beta)$ -expansions using this map $T_{L,-\beta}$ can be done using Proposition 2. For example, we find $-\frac{1}{2} = (.100)_{-\beta}$, which agrees with the result of Ito and Sadahiro. However, we can also find $-\frac{3}{4} = (.1011110)_{-\beta}$; a result that cannot be found using the original $(-\beta)$ -transformation.

3 Admissibility and shift spaces

Henceforth, whenever we refer to the map $T_{-\beta}$, we assume that it has been fixed beforehand. A sequence $(x_1, x_2, \dots) \in \mathcal{A}^{\mathbb{N}}$ is called $T_{-\beta}$ -admissible if we have $d_{-\beta}(x) = (x_1, x_2, \dots)$ for some $x \in I_{-\beta}$ and, moreover, the $(-\beta)$ -expansion of x generated by the map $T_{-\beta}$ (see Proposition 2) has coefficients x_1, x_2, \dots . A finite block is called $T_{-\beta}$ -admissible if it appears in a $T_{-\beta}$ -admissible sequence.

In order to nicely characterize the $T_{-\beta}$ -admissible sequences, we introduce an order on the set of $T_{-\beta}$ -admissible sequences: the *alternate (lexicographical) order*. We write $(x_1, x_2, \dots) \prec (y_1, y_2, \dots)$ if and only if there exists some integer $k \geq 1$ such that $x_i = y_i$ for all $i < k$ and $(-1)^k(x_k - y_k) < 0$. Also, $(x_1, x_2, \dots) \preceq (y_1, y_2, \dots)$ means that either $(x_1, x_2, \dots) \prec (y_1, y_2, \dots)$ or $(x_1, x_2, \dots) = (y_1, y_2, \dots)$ holds.

Proposition 4. *Let $T_{-\beta}$ be given. Let $x, y \in I_{-\beta}$ with $x \neq y$ be arbitrary and let $d_{-\beta}(x)$ and $d_{-\beta}(y)$ be the $(-\beta)$ -expansions of respectively x and y generated by $T_{-\beta}$. Then $x < y$ if and only if $d_{-\beta}(x) \prec d_{-\beta}(y)$.*

Proof. This has been shown in [1, Lemma 3.1] for $1 < \beta < 2$. We give the proof here for completeness. Write $d_{-\beta}(x) = (x_1, x_2, \dots)$ and $d_{-\beta}(y) = (y_1, y_2, \dots)$. Let $k \geq 1$ be the smallest integer for which $x_k \neq y_k$. Then we have

$$x = \frac{T_{-\beta}^{k-1}(x)}{(-\beta)^{k-1}} + \sum_{i=1}^{k-1} \frac{x_i}{(-\beta)^i} < \frac{T_{-\beta}^{k-1}(y)}{(-\beta)^{k-1}} + \sum_{i=1}^{k-1} \frac{y_i}{(-\beta)^i} = y$$

if and only if

$$\frac{T_{-\beta}^{k-1}(x)}{(-\beta)^{k-1}} < \frac{T_{-\beta}^{k-1}(y)}{(-\beta)^{k-1}}.$$

This latter inequality is, regardless of the parity of k , equivalent to the inequality

$$(-1)^k(x_k - y_k) < 0,$$

since $x_k = d(T_{-\beta}^{k-1}(x))$ (similar for y_k). The statement now follows. \square

We see that the alternate order respects the natural order on \mathbb{R} (as long as we compare $(-\beta)$ -expansions generated by the same map $T_{-\beta}$). The behaviour of the map $T_{-\beta}$, and hence the generated $(-\beta)$ -expansions, are determined by our choice for the cut points; two

maps with the same base β but different cut points will produce different $(-\beta)$ -expansions for at least one point $x \in I_{-\beta}$. Therefore, we should pay extra attention to our cut points. Let c be any cut point of a given map $T_{-\beta}$, then we define $d_{L,-\beta}(c)$ to be $(-\beta)$ -expansion generated by the left-continuous version of $T_{-\beta}$, regardless of the continuity of $T_{-\beta}$ itself (we define $d_{R,-\beta}(c)$ similarly).

Example 5. We continue Example 3. Our cut point is $2 - \sqrt{5}$, and the $(-\beta)$ -expansion of this point generated by the map is $2 - \sqrt{5} = (.11\bar{0})_{-\beta}$. Since the map in Example 3 is left-continuous, we know that $d_{L,-\beta}(2 - \sqrt{5}) = (1, 1, \bar{0})$. To determine $d_{R,-\beta}(2 - \sqrt{5})$ we modify the map in Example 3 such that it is right-continuous at $2 - \sqrt{5}$ (all other points remain unchanged). The $(-\beta)$ -expansion of $2 - \sqrt{5}$ generated by this new map is $d_{R,-\beta}(2 - \sqrt{5})$. The reader is invited to verify that $d_{R,-\beta}(2 - \sqrt{5}) = (0, 0, 1, \bar{0})$.

In order to state the following characterization of $T_{-\beta}$ -admissible sequences neatly, we introduce a little more notation. Recall that the digit function divides $I_{-\beta}$ into smaller subintervals. A subinterval on which the digit function is constant with value a will be denoted by I_a . Its left and right endpoint will be denoted by respectively ℓ_a and r_a . We can now characterize $T_{-\beta}$ -admissible sequences as follows:

Proposition 6. *The following hold:*

- (a) *If (x_1, x_2, \dots) is $T_{L,-\beta}$ -admissible, then for all $n \geq 1$:*
 - *if $x_n = \lfloor \beta \rfloor$, then $d_{-\beta}(\ell_{\lfloor \beta \rfloor}) \preceq (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}(r_{\lfloor \beta \rfloor})$;*
 - *if $x_n = a \neq \lfloor \beta \rfloor$, then $d_{R,-\beta}(\ell_a) \prec (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}(r_a)$.*
- (b) *If (x_1, x_2, \dots) is $T_{R,-\beta}$ -admissible, then for all $n \geq 1$:*
 - *if $x_n = 0$, then $d_{R,-\beta}(\ell_0) \preceq (x_n, x_{n+1}, \dots) \preceq d_{-\beta}(r_0)$;*
 - *if $x_n = a \neq 0$, then $d_{R,-\beta}(\ell_a) \preceq (x_n, x_{n+1}, \dots) \prec d_{L,-\beta}(r_a)$.*

Proof. We will only prove (a), since the proof of (b) is similar. If $x_n = \lfloor \beta \rfloor$, the result follows from Proposition 4.

Now, suppose that $x_n = a \neq \lfloor \beta \rfloor$. Then, by Proposition 4 we only need to show that $d_{R,-\beta}(\ell_a) \prec (x_n, x_{n+1}, \dots)$. Since the map is left-continuous, we know that $d_{L,-\beta}(\ell_a, 1) \neq a$ and hence $\ell_a < T_{L,-\beta}^{n-1}(x)$. Hence, there is a smallest integer $k \geq 1$ such that $d_{R,-\beta}(\ell_a, k) \neq d_{R,-\beta}(T_{L,-\beta}^{n-1}(x), k)$. Regardless of the parity of k , it follows that

$$(-1)^k (d_{R,-\beta}(\ell_a, k) - d_{R,-\beta}(T_{L,-\beta}^{n-1}(x), k)) < 0,$$

which proves $d_{R,-\beta}(\ell_a) \prec (x_n, x_{n+1}, \dots)$. □

The converse of Proposition 6 is not generally true. It is true in the case of Example 3 (we will see this later). We present the following Example in which the converse is not true.

Example 7. Consider the map $T_{L,-\beta}$ with $\beta = \frac{1+\sqrt{5}}{2}$ and cut point 0, that is, the map $T_{L,-\beta}$ on $I_{-\beta} = [-1, \beta - 1]$ given by:

$$T_{L,-\beta}(x) = \begin{cases} -\beta x - 1, & \text{if } -1 \leq x \leq 0; \\ -\beta x, & \text{if } 0 < x < \beta - 1. \end{cases}$$

Then, according to Proposition 6, a sequence (x_1, x_2, \dots) is $T_{L,-\beta}$ -admissible if

$$(\overline{1,0}) \preceq (x_n, x_{n+1}, \dots) \preceq (1, \overline{1,0}) \quad \text{or} \quad (\overline{0}) \prec (x_n, x_{n+1}, \dots) \preceq (\overline{0,1})$$

holds for all $n \geq 1$. The sequence $(0, 1, \overline{1,0})$ also satisfies this condition, but it is not $T_{L,-\beta}$ -admissible. To see this, note that $(.01\overline{10})_{-\beta} = 0$, whereas the $(-\beta)$ -expansion of 0 generated by the map $T_{L,-\beta}$ is $d_{-\beta}(0) = (1, \overline{1,0})$.

Hence, the set of sequences that satisfy the series of inequalities in Proposition 6 might be a bit larger than the set of $T_{-\beta}$ -admissible sequences. The problem arises whenever a cut point gets mapped onto some (not necessarily different) cut point after an odd amount of applications of the map $T_{-\beta}$.

Let $T_{-\beta}$ be given and let c be any cut point of $T_{-\beta}$. We define

$$d_{L,-\beta}^*(c) := \lim_{x \uparrow c} d_{-\beta}(x) \quad \text{and} \quad d_{R,-\beta}^*(c) := \lim_{x \downarrow c} d_{-\beta}(x).$$

Notice that the definition of $d_{L,-\beta}^*(c)$ does not require that the underlying map is left-continuous (similar for $d_{R,-\beta}^*(c)$). Instead, these sequences should be thought of as ‘‘improvements’’ of respectively $d_{L,-\beta}(c)$ and $d_{R,-\beta}(c)$. The improvement follows from the following lemma:

Lemma 8. *Let $T_{-\beta}$ be given and let c be any cut point of $T_{-\beta}$. The following hold:*

- (a) *There exists no $T_{L,-\beta}$ -admissible sequence (x_1, x_2, \dots) such that we have $d_{R,-\beta}(c) \prec (x_1, x_2, \dots) \prec d_{R,-\beta}^*(c)$.*
- (b) *There exists no $T_{R,-\beta}$ -admissible sequence (x_1, x_2, \dots) such that we have $d_{L,-\beta}^*(c) \prec (x_1, x_2, \dots) \prec d_{L,-\beta}(c)$.*

Proof. We only prove (a), since the proof of (b) is similar. Suppose that the statement is false and let (x_1, x_2, \dots) be a $T_{L,-\beta}$ -admissible sequence satisfying $d_{R,-\beta}(c) \prec (x_1, x_2, \dots) \prec d_{R,-\beta}^*(c)$ and $x = (.x_1 x_2 \dots)_{-\beta}$. By assumption, we have $x_1 = d_{R,-\beta}(c, 1)$, and hence $x > c$.

Let $(x^n)_{n=1}^\infty$ be a decreasing sequence for which $d_{-\beta}(x^n, 1) = d_{R,-\beta}^*(c, 1)$ for all $n \geq 1$, where $d_{-\beta}(x^n)$ is the $(-\beta)$ -expansion of x^n generated by $T_{-\beta}$. For all $m, n \geq 1$ with $m < n$ we have $d_{-\beta}(x^n) \prec d_{-\beta}(x^m)$ by Proposition 4. It follows that $d_{R,-\beta}^*(c) \preceq d_{-\beta}(x^n)$ for all $n \geq 1$. Moreover, the sequence converges to c and hence for some $k \geq 1$ we have $x^k < c$ and $d_{R,-\beta}^*(c) \preceq d_{-\beta}(x^k) \prec d_{-\beta}(x)$. This latter is, however, a contradiction and hence the sequence (x_1, x_2, \dots) cannot exist. \square

For any given map $T_{-\beta}$, computing the sequences $d_{L,-\beta}^*(c)$ and $d_{R,-\beta}^*(c)$, where c is a cut point, is not hard. Fix c and suppose we want to find $d_{L,-\beta}^*(c)$. We present the following algorithm:

1. Start with $k := 1$, $\hat{T}_1 := T_{L,-\beta}$ and $c_0 := c$.
2. Compute $c_k := \hat{T}_k(c_{k-1})$ and set

$$d_{L,-\beta}^*(c, k) := \begin{cases} d^L(c_{k-1}), & \text{if } \hat{T}_k = T_{L,-\beta}; \\ d^R(c_{k-1}), & \text{if } \hat{T}_k = T_{R,-\beta}; \end{cases}$$

where d^L and d^R are the digit functions of $T_{L,-\beta}$ and $T_{R,-\beta}$ respectively.

3. Let $m := \max\{0 \leq i < k : c_i \text{ is a cut point}\}$. If c_k is a cut point and if $k - m$ is odd, then:

- if $\hat{T}_k = T_{L,-\beta}$, then $\hat{T}_{k+1} := T_{R,-\beta}$;
- if $\hat{T}_k = T_{R,-\beta}$, then $\hat{T}_{k+1} := T_{L,-\beta}$;

and else $\hat{T}_{k+1} := \hat{T}_k$.

4. Increase k by one and go back to step 2.

If we wish to find $d_{R,-\beta}^*(c)$ instead, all that changes is that we should start with $\hat{T}_1 := T_{R,-\beta}$.

Proposition 9. *The method described above for finding $d_{L,-\beta}^*(c)$ and $d_{R,-\beta}^*(c)$ is correct.*

Proof. We only prove correctness for $d_{L,-\beta}^*(c)$. Call the sequence obtained by the algorithm above $s_{L,-\beta}(c)$, then we need to show that $s_{L,-\beta}(c) = \lim_{x \uparrow c} d_{-\beta}(x)$. For any integer $n \geq 1$, define (recall that $\ell_{s_{L,-\beta}(c,k)}$ denotes the left endpoint of the subinterval on which the digit function equals the k th digit of $s_{L,-\beta}(c)$):

$$\begin{aligned} \epsilon_n &:= \frac{1}{\beta^{n-1}} \min \left(\left\{ |c_k - \ell_{s_{L,-\beta}(c,k+1)}| : 0 \leq k \leq n, k \text{ even} \right\} \right. \\ &\quad \left. \cup \left\{ |c_k - r_{s_{L,-\beta}(c,k+1)}| : 0 \leq k \leq n, k \text{ odd} \right\} \right). \end{aligned}$$

It follows that any $x \in (c - \epsilon_n, c)$ satisfies $d_{L,-\beta}(x, i) = s_{L,-\beta}(c, i)$ for all $i \leq n$. It suffices to show that $\epsilon_n \neq 0$ for all $n \geq 1$. Suppose that $k \geq 1$ is the smallest integer such that $\epsilon_k = 0$ and assume that k is odd (the proof is similar for k even). It follows that $c_k = r_{s_{L,-\beta}(c,k+1)}$, and, since $\hat{T}_{k+1} = T_{R,-\beta}$, we must hit $r_{-\beta}$ after an odd number of iterations (it cannot be any other cut point since that would imply $c_k = \ell_{s_{L,-\beta}(c,k+1)} \neq r_{s_{L,-\beta}(c,k+1)}$). It follows that c_{k-1} is either a cut point or $-\frac{\beta \cdot \lfloor \beta \rfloor}{\beta^2 - 1}$. By iterating this argument it follows that c must be an endpoint of $I_{-\beta}$. We have reached a contradiction since the endpoints are no cut points. It follows that $\epsilon_n > 0$ for all $n \geq 1$. \square

Example 10. We continue Example 7. Since the orbit of 0 never hits 0 again under $T_{L,-\beta}$, we conclude that $d_{L,-\beta}^*(0) = d_{L,-\beta}(0) = (1, \overline{1}, 0)$. Under $T_{R,-\beta}$, however, 0 hits itself after one iteration. Hence, we have $d_{R,-\beta}^*(0, 1) = 0$ and continue iterating $T_{L,-\beta}$. At this point, we know the orbit of 0 under $T_{L,-\beta}$ and therefore $d_{R,-\beta}^*(0)$ is the concatenation of the single-digit block (0) with $(1, \overline{1}, 0)$: $d_{R,-\beta}^*(0) = (0, 1, \overline{1}, 0)$.

Example 11. We continue Example 5. Since the orbit of $2 - \sqrt{5}$ never hits itself under both $T_{L,-\beta}$ and $T_{R,-\beta}$, we have $d_{L,-\beta}^*(2 - \sqrt{5}) = d_{L,-\beta}(2 - \sqrt{5}) = (1, 1, \bar{0})$ and $d_{R,-\beta}^*(2 - \sqrt{5}) = d_{R,-\beta}(2 - \sqrt{5}) = (0, 0, 1, \bar{0})$.

Remark 12. The sequences $d_{L,-\beta}^*(c)$ and $d_{R,-\beta}^*(c)$ are concatenations of blocks of sequences of the form $d_{L,-\beta}(\ell_a)$, $d_{R,-\beta}(\ell_a)$, $d_{L,-\beta}(r_a)$ and $d_{R,-\beta}(r_a)$ where $a \in \mathcal{A}$.

The following lemma is very important:

Lemma 13. *Let $(x_1, x_2, \dots) \in \mathcal{A}^{\mathbb{N}}$. The following hold:*

(a) *If for all $n \geq 1$ we have*

$$d_{R,-\beta}^*(\ell_{x_n}) \preceq (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}(r_{x_n}),$$

then every finite subblock of (x_1, x_2, \dots) is $T_{L,-\beta}$ -admissible.

(b) *If for all $n \geq 1$ we have*

$$d_{R,-\beta}(\ell_{x_n}) \preceq (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}^*(r_{x_n}),$$

then every finite subblock of (x_1, x_2, \dots) is $T_{R,-\beta}$ -admissible.

Proof. We only prove (a), as the proof of (b) is similar. Suppose that (x_1, x_2, \dots) satisfies the given condition and define $x^{(k)} := \sum_{n=k}^{\infty} \frac{x_n}{(-\beta)^{n-k+1}}$. Assume that there exist integers $s(k), t(k) > k$ such that:

$$\begin{aligned} x_k \cdots x_{s(k)-1} &= d_{R,-\beta}^*(\ell_{x_k}, 1) \cdots d_{R,-\beta}^*(\ell_{x_k}, s(k) - k), \\ (-1)^{s(k)-k+1} (d_{R,-\beta}^*(\ell_{x_k}, s(k) - k + 1) - x_{s(k)}) &< 0, \end{aligned}$$

and

$$\begin{aligned} x_k \cdots x_{t(k)-1} &= d_{L,-\beta}(r_{x_k}, 1) \cdots d_{L,-\beta}(r_{x_k}, t(k) - k), \\ (-1)^{t(k)-k+1} (x_{t(k)} - d_{L,-\beta}(r_{x_k}, t(k) - k + 1)) &< 0. \end{aligned}$$

If either $s(k)$ or $t(k)$ does not exist, then there is nothing to prove. Let y be a point such that $d_{L,-\beta}(y, i) = d_{R,-\beta}^*(\ell_{x_k}, i)$ for $1 \leq i \leq s(k) - k + 1$, then

$$\begin{aligned} x^{(k)} - \ell_{x_k} &> x^{(k)} - y \\ &= \sum_{n=s(k)}^{\infty} \frac{x_n}{(-\beta)^{n-k+1}} - \frac{T_{L,-\beta}^{s(k)-k}(y)}{(-\beta)^{s(k)-k}} \\ &> \frac{1}{(-\beta)^{s(k)-k}} \left(x^{s(k)} - T_{L,-\beta}^{s(k)-k}(y) \right), \end{aligned}$$

from which it follows that

$$x^{(k)} - \ell_{x_k} > \begin{cases} \frac{x^{s(k)} - \ell_{x_{s(k)}}}{(-\beta)^{s(k)-k+1}}, & \text{if } s(k) - k + 1 \text{ is odd;} \\ \frac{x^{s(k)} - r_{x_{s(k)}}}{(-\beta)^{s(k)-k+1}}, & \text{if } s(k) - k + 1 \text{ is even.} \end{cases} \quad (2)$$

Similarly, one can show that

$$r_{x_k} - x^{(k)} > \begin{cases} \frac{r_{x_{t(k)}} - x^{t(k)}}{(-\beta)^{t(k)-k+1}}, & \text{if } t(k) - k + 1 \text{ is odd;} \\ \frac{\ell_{x_{t(k)}} - x^{t(k)}}{(-\beta)^{t(k)-k+1}}, & \text{if } t(k) - k + 1 \text{ is even.} \end{cases} \quad (3)$$

By iteratively applying the bounds in (2) and (3) we get a sequence of bounds that approach zero (as the numerator is bounded and the denominator goes to infinity). It follows that $l_{x_k} < x^{(k)} < r_{x_k}$ for all $k \geq 1$, which implies $d_{L,-\beta}(x^{(1)}) = (x_1, x_2, \dots)$. \square

Using these new objects, we can make the inequalities in Proposition 6 more strict and obtain an equivalence.

Theorem 14. *The following hold:*

(a) *The sequence (x_1, x_2, \dots) is $T_{L,-\beta}$ -admissible if and only if for all $n \geq 1$:*

- *if $x_n = \lfloor \beta \rfloor$, then $d_{-\beta}(\ell_{\lfloor \beta \rfloor}) \preceq (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}(r_{\lfloor \beta \rfloor})$;*
- *if $x_n = a \neq \lfloor \beta \rfloor$, then $d_{R,-\beta}^*(\ell_a) \prec (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}(r_a)$.*

(b) *The sequence (x_1, x_2, \dots) is $T_{R,-\beta}$ -admissible if and only if for all $n \geq 1$:*

- *if $x_n = 0$, then $d_{R,-\beta}(\ell_0) \preceq (x_n, x_{n+1}, \dots) \preceq d_{-\beta}(r_0)$;*
- *if $x_n = a \neq 0$, then $d_{R,-\beta}(\ell_a) \preceq (x_n, x_{n+1}, \dots) \prec d_{L,-\beta}^*(r_a)$.*

Proof. This has been proven for $1 < \beta < 2$ in [1, Theorem 3.1]. We give the proof for completeness. We only prove (a), since the proof of (b) is similar. The "only if" part follows from Proposition 6 and Lemma 8.

To prove the "if" part, suppose that (x_1, x_2, \dots) satisfies the given inequalities. From Lemma 13 it follows that every finite subblock (x_n, \dots, x_{n+k}) is $T_{L,-\beta}$ -admissible. It is sufficient to check that $(x_n, x_{n+1}, \dots) \neq d_{R,-\beta}^*(\ell_{x_n})$ for all $n \geq 1$ such that $x_n \neq \lfloor \beta \rfloor$. However, this follows directly from the given inequalities. \square

Let $T_{-\beta}$ be a $(-\beta)$ -transformation such that $T_{-\beta}$ restricted to the interval $\left[-\frac{\beta}{\beta+1}, \frac{1}{\beta+1}\right)$ is the Ito-Sadahiro transformation. While our condition in Theorem 14 is different from the condition found by Ito and Sadahiro [8, Theorem 10], the two actually agree. This is because their transformation implicitly specifies the cut points. Moreover, these cut points are all mapped to $-\frac{\beta}{\beta+1}$. The following example illustrates this:

Example 15. Consider the map of Example 3 restricted to the interval $\left[-\frac{\beta}{\beta+1}, \frac{1}{\beta+1}\right)$. Our condition in Theorem 14(a), when restricted to this interval, becomes:

$$(1, \bar{0}) \preceq (x_n, x_{n+1}, \dots) \preceq (1, 1, \bar{0}) \quad \vee \quad (0, 0, 1, \bar{0}) \prec (x_n, x_{n+1}, \dots) \prec (0, 1, \bar{0}) \quad (4)$$

for all $n \geq 1$. On the other hand, by Ito and Sadahiro [8, Theorem 10] we have:

$$(1, \bar{0}) \preceq (x_n, x_{n+1}, \dots) \prec (0, 1, \bar{0})$$

for all $n \geq 1$, which suggests that the sequences corresponding to the cut point are redundant. To see why this is the case, suppose that (x_1, x_2, \dots) satisfies (4) for all $n \geq 1$. Fix $k \geq 1$ and assume for now that $x_k = 1$, then we have $(x_k, x_{k+1}, \dots) \preceq (1, 1, \bar{0})$ from which it follows that $(1, \bar{0}) \preceq (x_{k+1}, x_{k+2}, \dots)$. This condition, however, is already satisfied which means that we can reduce the first part of (4) to $(1, \bar{0}) \preceq (x_k, x_{k+1}, \dots)$. Similarly, we can reduce the second part of (4) whenever $x_k = 0$ and get

$$\begin{cases} (1, \bar{0}) \preceq (x_n, x_{n+1}, \dots), & \text{whenever } x_n = 1; \\ (x_n, x_{n+1}, \dots) \prec (0, 1, \bar{0}), & \text{whenever } x_n = 0, \end{cases}$$

which in turn can be neatly written as

$$(1, \bar{0}) \preceq (x_n, x_{n+1}, \dots) \prec (0, 1, \bar{0}),$$

which is the result by Ito and Sadahiro.

4 Shifts and graphs

In Theorem 14 we gave a characterization of the shift-invariant set of $T_{-\beta}$ -admissible sequences. This set will be denoted with S_T . In order to turn S_T into a shift space, we need to consider its closure. This closure will be called the $(-\beta)$ -shift (for the underlying map $T_{-\beta}$) and will be denoted by $S_{-\beta}$. The context should make it clear what the underlying map $T_{-\beta}$ is. If we wish to emphasize whether $T_{-\beta}$ is left- or right-continuous, we may write $S_{L,-\beta}$ and $S_{R,-\beta}$ respectively. The most straightforward characterization of the $(-\beta)$ -shift is the following.

Lemma 16. *Let $T_{-\beta}$ be given and let $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$. Then $\mathbf{x} \in S_{-\beta}$ if and only if $(\mathbf{x}_p, \dots, \mathbf{x}_q)$ is $T_{-\beta}$ -admissible for all $1 \leq p \leq q$.*

We use this lemma to prove the following much more useful characterization.

Theorem 17. *Let $T_{-\beta}$ be given and let $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$. The following hold:*

- (a) *We have $\mathbf{x} \in S_{L,-\beta}$ if and only if $d_{R,-\beta}^*(\ell_{x_n}) \preceq (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}(r_{x_n})$ for all $n \geq 1$.*
- (b) *We have $\mathbf{x} \in S_{R,-\beta}$ if and only if $d_{R,-\beta}(\ell_{x_n}) \preceq (x_n, x_{n+1}, \dots) \preceq d_{L,-\beta}^*(r_{x_n})$ for all $n \geq 1$.*

Proof. We only prove (a), as the proof of (b) is similar. The "if" part follows from Lemma 13 and Lemma 16. Let $N \geq 1$ be arbitrary and fixed. If (x_N, x_{N+1}, \dots) is $T_{L,-\beta}$ -admissible, Theorem 14 applies. Thus, assume it is not $T_{L,-\beta}$ -admissible. By Lemma 16, every finite subblock (x_N, \dots, x_{N+k}) , where $k \geq 1$, is $T_{L,-\beta}$ -admissible. Hence, we have

$$(x_N, \dots, x_{N+k}) \preceq (d_{L,-\beta}(r_{x_N}, 1), \dots, d_{L,-\beta}(r_{x_N}, k+1))$$

which follows immediately and, by Lemma 8, we also have

$$(d_{R,-\beta}^*(\ell_{x_N}, 1), \dots, d_{R,-\beta}^*(\ell_{x_N}, k+1)) \preceq (x_N, \dots, x_{N+k}).$$

This proves the theorem. □

Based on this theorem, we can classify the shift space $S_{-\beta}$. The behaviour of these shift spaces depend on the sequences $d_{R,-\beta}^*(\ell_a)$ and $d_{L,-\beta}(r_a)$ (where $a \in \mathcal{A}$) if the underlying map is left-continuous and $d_{R,-\beta}(\ell_a)$ and $d_{L,-\beta}^*(r_a)$ if the underlying map is right-continuous. The sequences will be called *boundary sequences*.

The next theorem states that eventually periodic boundary sequences are a necessary and sufficient condition for the shift space to be sofic. In order to prove this, we will give an algorithm, based on the algorithm in [8], that construct a labeled graph recognizing the shift. For now, we will assume that all boundary sequences are eventually periodic.

Assume that our shift space has a left-continuous underlying map. The shift space has at most $2(\lfloor \beta \rfloor + 1)$ boundary sequences, two for each $a \in \mathcal{A}$ used by the underlying digit function: one corresponding to ℓ_a and the other to r_a . In what follows we will assume that all digits in \mathcal{A} are used. Hence, we will need two sets for each digit, m_a^ℓ and m_a^r , which are subsets of $\mathbb{Z}_{\geq 0}$. These sets keep track of the (finite) memory of a machine that reads the sequence (x_1, x_2, \dots) one digit at a time. The meaning of $k \in m_a^\ell$ is that the current memory contains a finite subblock starting with a that equals the prefix of the boundary sequence corresponding to ℓ_a and where the next digit needs to be matched against the $(k+1)$ -th digit of the boundary sequence corresponding to ℓ_a (similarly for m_a^r). In fact, we can give an upper bound for each of these memory sets by using the fact that all boundary sequences are eventually periodic. Let the boundary sequence be eventually periodic with pre-periodic part of length $q \geq 0$ and periodic part of length $p \geq 1$, then the upper bounds, denoted by U_a^ℓ and U_a^r respectively, are equal to

$$\begin{cases} 2p, & \text{if } q = 0; \\ 2p + q - 1, & \text{if } q \neq 0 \text{ and } p \text{ is odd;} \\ p + q - 1, & \text{if } q \neq 0 \text{ and } p \text{ is even.} \end{cases}$$

These upper bounds tell us how many different prefixes we may expect.

Next, assume that we read a new digit of the sequence (x_1, x_2, \dots) , say d . Our memory has now changed and hence we need to update our memory sets. If $k \in m_a^\ell$ and we read the digit d , we change the index k into another index $m_a^\ell(k/d)$, which is defined as:

$$m_a^\ell(k/d) := \begin{cases} k + 1, & \text{if } k < U_a^\ell \text{ and } d = d_{R,-\beta}^*(\ell_a, k + 1); \\ 1, & \text{if } k = U_a^\ell, d = d_{R,-\beta}^*(\ell_a, k + 1) \text{ and } q = 0; \\ q, & \text{if } k = U_a^\ell, d = d_{R,-\beta}^*(\ell_a, k + 1) \text{ and } q \neq 0; \\ 0, & \text{if } k \neq 0 \text{ and } (-1)^{k+1} (d_{R,-\beta}^*(\ell_a, k + 1) - d) < 0; \\ 0, & \text{if } k = 0 \text{ and } d \neq a; \\ 1, & \text{if } k = 0 \text{ and } d = a; \\ F, & \text{otherwise.} \end{cases} \quad (5)$$

Let us shortly explain what each of these seven cases mean. If our memory contains a subblock matching the prefix of the boundary sequence of ℓ_a but does not contain one complete period, then we check digit d against the next digit of the prefix (case 1). If the boundary sequence is purely periodic and if our memory contains two complete periods (we need two complete periods in the case p is odd), then move back to the beginning (case

2). If the boundary sequence is eventually periodic and our memory contains the maximum number of complete periods (one if p is even, otherwise two), then return to the beginning of the periodic part (case 3). If our memory contains a subblock that, together with the digit d , can be accepted because of Theorem 17, then accept it and "forget" about this block (case 4). If our memory does not contain the digit a and if the new digit also is not a , do absolutely nothing (case 5). On the other hand, if the new digit equals a , we have a subblock of length 1 matching the prefix of equal length of the boundary sequence ℓ_a (case 6). If none of the above happens, then we are neither undecided (cases 1,2,3,5 and 6) nor can we accept any subblock (case 4), and hence we must reject it (case 7).

Similarly, we define

$$m_a^r(k/d) := \begin{cases} k+1, & \text{if } k < U_a^r \text{ and } d = d_{L,-\beta}(r_a, k+1); \\ 1, & \text{if } k = U_a^r, d = d_{L,-\beta}(r_a, k+1) \text{ and } q = 0; \\ q, & \text{if } k = U_a^r, d = d_{L,-\beta}(r_a, k+1) \text{ and } q \neq 0; \\ 0, & \text{if } k \neq 0 \text{ and } (-1)^{k+1} (d - d_{L,-\beta}(r_a, k+1)) < 0; \\ 0, & \text{if } k = 0 \text{ and } d \neq a; \\ 1, & \text{if } k = 0 \text{ and } d = a; \\ F, & \text{otherwise.} \end{cases} \quad (6)$$

We state the following result, which follows directly from equations (5) and (6):

Corollary 18. *Let (z_1, \dots, z_n) be our current memory containing no forbidden blocks. Then for some $a \in \mathcal{A}$ and $i \leq U_a^\ell$ (or U_a^r) either $m_a^\ell(i/d)$ or $m_a^r(i/d)$ equals F if and only if (z_1, \dots, z_n, d) contains a forbidden block.*

To construct a labeled graph recognizing the shift $S_{L,-\beta}$ whose boundary sequences are all eventually periodic, we perform the following steps:

1. Let \mathcal{V} be the set of vertices where each vertex $v \in \mathcal{V}$ is labeled $(m_0^\ell, m_0^r, \dots, m_{[\beta]}^\ell, m_{[\beta]}^r)$ such that m_a^ℓ is a nonempty subset of $\{0, 1, \dots, U_a^\ell\}$ and m_a^r is a nonempty subset of $\{0, 1, \dots, U_a^r\}$.
2. For each vertex $v \in \mathcal{V}$ and $d \in \mathcal{A}$ draw an edge with label d from vertex v to vertex $w = (w_1, w_2, \dots, w_{2[\beta]+2})$, where

$$w_k = \begin{cases} \varphi_{\ell, (k-1)/2}(m_{(k-1)/2}^\ell, d), & \text{if } k \text{ is odd;} \\ \varphi_{\ell, (k-2)/2}(m_{(k-2)/2}^r, d), & \text{if } k \text{ is even;} \end{cases}$$

and the maps $\varphi_{\ell, a}$ and $\varphi_{r, a}$ are defined as

$$\varphi_{\ell, a}(v, d) := \begin{cases} F, & \text{if } F \in \bigcup_{i \in v} \{m_a^\ell(i/d)\}; \\ \bigcup_{i \in v} \{m_a^\ell(i/d)\} \setminus \{0\}, & \text{if } \{0\} \subsetneq \bigcup_{i \in v} \{m_a^\ell(i/d)\} \\ & \text{and } F \notin \bigcup_{i \in v} \{m_a^\ell(i/d)\}; \\ \{1\} \cup \bigcup_{i \in v} \{m_a^\ell(i/d)\} \setminus \{0\}, & \text{if } 1, F \notin \bigcup_{i \in v} \{m_a^\ell(i/d)\} \\ & \text{and } d = a; \\ \bigcup_{i \in v} \{m_a^\ell(i/d)\}, & \text{otherwise.} \end{cases} \quad (7)$$

The definition of $\varphi_{r,a}$ is nearly identical; simply replaces all instances of ℓ in (7) with r .

3. The connected component (i.e., each vertex has at least one outgoing edge) containing the zero vertex is the labeled graph recognizing $S_{L,-\beta}$.

Since our memory sets may contain more than one digit, we need to keep track of the way each of these indices react to the new digit d and, should they arise, remove any inconsistencies. This is the job of $\varphi_{\ell,a}$ and $\varphi_{r,a}$. Let us explain the four cases in (7) in more detail. If, because of the new digit d , our memory contains a forbidden subblock, automatically reject it (case 1). If we cannot reject it, then suppose our memory contains at least two subblocks on which we are undecided, but can accept some (not all) of them after we read the new digit d . In that case adding a 0 to the memory set would be redundant (case 2). If the new digit equals a , then add 1 to the memory set, as we have a new subblock (of length 1) starting with a (case 3). If none of these three cases apply, there are no inconsistencies (case 4).

In practice, one starts with the zero vertex and adds vertices along the way. Since none of the vertices carry the label F , we do not draw an edge if the update rule in (7) implies that the edge should point towards such a vertex.

Theorem 19. *The shift space $S_{-\beta}$ is sofic if and only if its boundary sequences are all eventually periodic.*

Proof. Assume, without loss of generality, that the underlying map is left-continuous (in the remainder of the proof we will write $S_{L,-\beta}$). Let \mathcal{G} be a right-resolving labeled graph recognizing $S_{L,-\beta}$ (recall [9, Theorem 3.3.2]) and let \mathcal{L} be the labeling. Fix $a \in \mathcal{A}$ and consider $d_{R,-\beta}^*(\ell_a)$ (the argument is similar for $d_{L,-\beta}(r_a)$). There exists a path $\xi = e_1 e_2 \cdots$ in \mathcal{G} such that $(\mathcal{L}(e_1), \mathcal{L}(e_2), \dots) = d_{R,-\beta}^*(\ell_a)$. By Theorem 17, $d_{R,-\beta}^*(\ell_a)$ is the smallest sequence with respect to the order \prec starting with a and hence ξ is the path that first chooses the edge with label a and then the edge with the smallest possible label (with respect to \prec). Since \mathcal{G} is finite, the path ξ passes some vertex v infinitely often. Hence, ξ contains a loop of even length, which implies that for some $k, n \geq 1$ we have $\mathcal{L}(e_i) = \mathcal{L}(e_{i+2k})$. It follows that $d_{R,-\beta}^*(\ell_a)$ is eventually periodic.

Conversely, let $S_{L,-\beta}$ be given, \mathcal{G} the labeled graph that follows from our construction and let L_S and $L_{\mathcal{G}}$ be the language of the shift space $S_{L,-\beta}$ and the edge shift on \mathcal{G} respectively. By Corollary 18 and the construction of \mathcal{G} it immediately follows that $L_S \subset L_{\mathcal{G}}$. If $(a_1, \dots, a_n, d) \notin L_S$ and $(a_1, \dots, a_n) \in L_S$ (we may assume this without loss of generality), then from Corollary 18 it follows that either $m_a^\ell(i/d)$ or $m_a^r(i/d)$ equals F for some i and a . Hence, by construction, there is a finite path $\zeta = (e_1, \dots, e_n)$ such that $\mathcal{L}(e_i) = a_i$. Say the path ζ ends at vertex v , then there is no outgoing edge from v with label d . For if it did exist, it would point to a vertex with label F as one of its coordinates by (7). Such vertices, however, do not exist (recall step 1 of the construction). It follows that $L_S = L_{\mathcal{G}}$. \square

Remark 20. The construction of the labeled graph is also valid if at least one boundary sequence is not eventually periodic. In this case, the upper bound U_a^ℓ or U_a^r corresponding to this boundary sequence, as well as p for this specific boundary sequence, should be equal to ∞ (as there is no periodicity to exploit). Using our construction, the number of vertices in the labeled graphs of these shifts might be countably infinite.

Let us illustrate the results of this section with two examples.

Example 21. We continue the case of Example 3. In Examples 5 and 11 we determined the boundary sequences and hence, by Theorem 4.2, a sequence belongs to the shift space $S_{-\beta}$ if and only if

$$(\overline{1,0}) \preceq (x_n, x_{n+1}, \dots) \preceq (1, 1, \overline{0}) \quad \vee \quad (0, 0, 1, \overline{0}) \preceq (x_n, x_{n+1}, \dots) \preceq (\overline{0,1})$$

holds for all $n \geq 1$. We can simplify this as

$$(x_n, x_{n+1}, \dots) \preceq (1, 1, \overline{0}) \quad \vee \quad (0, 0, 1, \overline{0}) \preceq (x_n, x_{n+1}, \dots), \quad (8)$$

since the other two inequalities are always satisfied, as $(\overline{1,0})$ and $(\overline{0,1})$ are the smallest and the greatest sequences with respect to the alternate order respectively. It is useful to create tables stating all possible outcomes of $m_a^\ell(i/d)$ and $m_a^r(i/d)$ for all $a, d \in \mathcal{A}$ and all allowed indices i . In this specific case, we find:

$m_0^\ell(i/d)$	0	1		$m_1^r(i/d)$	0	1
0	1	0		0	0	1
1	2	0		1	0	2
2	0	3		2	3	0
3	4	0		3	2	F
4	3	F				

The entries labeled F correspond to forbidden blocks, though not all forbidden blocks can be found explicitly this way. We see that $m_0^\ell(4/1) = F$, which means we can find a forbidden block by taking the prefix of length 4 of $d_{R,-\beta}^*(2 - \sqrt{5})$ and adding a fifth digit to it, in this case 1. So, $(0, 0, 1, 0, 1)$ is a forbidden block, which is clear from (8). Similarly, $m_1^r(3/1) = F$ tells us that $(1, 1, 0, 1)$ is a forbidden block. Figure 3 shows the labeled graph recognizing the shift space.

Example 22. Now consider the map given in Example 7. By Example 10, $d_{R,-\beta}^*(0) = (0, 1, \overline{1,0})$, and thus a sequence belongs to the shift space generated by this map if and only if

$$(x_n, x_{n+1}, \dots) \preceq (1, \overline{1,0}) \quad \vee \quad (0, 1, \overline{1,0}) \preceq (x_n, x_{n+1}, \dots) \quad (9)$$

holds for all $n \geq 1$. Once again, we omitted two inequalities since they are redundant. We can proceed as in the previous example and create the following tables:

$m_0^\ell(i/d)$	0	1		$m_1^r(i/d)$	0	1
0	1	0		0	0	1
1	F	2		1	0	2
2	0	3		2	1	0
3	2	0				

The tables now only reveal one forbidden block: $(0,0)$. As we will see in a later section, this is the only forbidden block. For the moment, assume we are not aware of this. We construct a labeled graph that recognizes the shift based on our lengthy exposition earlier. This graph is shown in Figure 4.




Figure 3: A labeled graph recognizing the shift space given by (8).

As in the case of expansions in positive base, whenever β is Pisot we can improve Theorem 19 a little. Recall that a *Pisot number* is a positive algebraic integer whose other conjugates lie in the unit circle in the complex plane. The following theorem shows why Pisot bases are nice to work with.

Theorem 23. *Let β be Pisot, $x \in I_{-\beta}$ and $T_{-\beta}$ be given and let $d_{-\beta}(x)$ be the $(-\beta)$ -expansion of x generated by $T_{-\beta}$. If $x \in \mathbb{Q}(\beta)$, then $d_{-\beta}(x)$ is eventually periodic.*

Proof. We will not give the complete proof here, as it is nearly identical to the proof of [5, Theorem 24]. In our case, replace $[l, r)$ with $I_{-\beta}$ and note that we have the bound $|r_n| < \frac{\beta \cdot |\beta|}{\beta^2 - 1}$. \square

As an immediate corollary, we have:

Theorem 24. *Let β be Pisot and $S_{-\beta}$ be given. Then $S_{-\beta}$ is sofic if and only if the cut points of the underlying map $T_{-\beta}$ all lie in $\mathbb{Q}(\beta)$.*

5 Forbidden blocks

Example 22 showed that, while technically correct, our construction of the labeled graph can yield us quite large graphs. However, by carefully looking at Examples 21 and 22 we see why the first one is only sofic and the latter is actually a shift of finite type. In the first example, the forbidden blocks contain both a part of the pre-periodic and a part of the periodic part of the boundary sequence, where in the second example the forbidden block only contained part of the pre-periodic part. This observation quickly leads us to a conjecture: a shift space of the form given in Theorem 17 is a shift of finite type if and only if its forbidden blocks only contain part of either pre-periodic or periodic parts of boundary sequences, but not both. Indeed, if the latter does happen, we can abuse the periodicity of the boundary sequences




Figure 4: A labeled graph recognizing the shift space given by (9).

to generate infinitely many forbidden blocks that cannot be covered by a finite set (because of the pre-periodic part). Based on these observations, we state the following

Theorem 25. *Let $S_{-\beta}$ be a sofic shift. Then $S_{-\beta}$ is a shift of finite type if and only if the periodic part of every eventually periodic boundary sequence is a purely periodic boundary sequence.*

Proof. Once again, we assume that the underlying map is left-continuous and write $S_{L,-\beta}$ instead of $S_{-\beta}$. Let $S_{L,-\beta}$ be a shift of finite type. Every shift of finite type is sofic and hence by Theorem 19 we conclude that all boundary sequences are eventually periodic. Assume that there exists a left-boundary sequence (the proof is similar for right-boundary sequences) $(a_1, \dots, a_k, \overline{a_{k+1}, \dots, a_{k+p}})$, where $k, p \geq 1$, whose periodic part does not equal some other boundary sequence. Moreover, assume that k is odd (the proof is similar for k even). By assumption, there exists a smallest integer $1 \leq j \leq p$ such that there exists some $b \in \mathcal{A}$ for which we have $(-1)^{k+j}(b - a_{k+j}) < 0$. It follows that the block $(a_1, \dots, a_{k+j-1}, b)$ is forbidden. Moreover, the blocks $w^{(n)} := a_1 \cdots a_k (a_{k+1} \cdots a_{k+p})^{2n} a_{k+1} \cdots a_{k+j-1} b$ are forbidden for all $n \geq 0$.

Suppose that there exists some $1 \leq i \leq j - 1$ such that the subblock $a_{k+i} \cdots b$ of length at most j is forbidden. Then the blocks $(a_{k+1} \cdots a_{k+p})^{2n} a_{k+1} \cdots a_{k+j-1} b$ are forbidden for all $n \geq 0$ and thus we conclude that

$$(\overline{a_{k+1} \cdots a_{k+p}}) \prec (a_{k+1} \cdots a_{k+p})^{2n} a_{k+1} \cdots a_{k+j-1} b$$

holds for all $n \geq 1$. This, however, implies that $(\overline{a_{k+1}, \dots, a_{k+p}})$ is a right-boundary sequence, which contradicts our initial assumption. Hence, there exists some integer $1 \leq m \leq k$ such that the blocks $a_m \cdots a_k (a_{k+1} \cdots a_{k+p})^{2n} a_{k+1} \cdots a_{k+j-1} b$ are forbidden for all $n \geq 0$, but any of their subblocks are admissible. Hence, $S_{-\beta}$ is not a shift of finite type.

Next, assume that all boundary sequences satisfy the property given in Theorem 25 and let (a_1, a_2, \dots) be an arbitrary but fixed left-boundary sequence (the proof is similar for

right-boundary sequences). If it is purely periodic with period p , then the forbidden blocks are given by

$$\{(x_1, \dots, x_{2p}) \in \mathcal{A}^{2p} : (x_1, \dots, x_{2p}) \prec (a_1, \dots, a_{2p}) \wedge x_1 = a_1\}. \quad (10)$$

If it is eventually periodic with pre-periodic part of length q and period p , then the set

$$\{(x_1, \dots, x_{q+1}) \in \mathcal{A}^{q+1} : (x_1, \dots, x_{q+1}) \prec (a_1, \dots, a_{q+1}) \wedge x_1 = a_1\} \quad (11)$$

is sufficient. To see why, let (y_1, y_2, \dots) be a sequence such that $y_1 = a_1$ and $(y_1, y_2, \dots) \prec (a_1, a_2, \dots)$. Let m be the smallest integer such that $y_i = a_i$ for $i < m$ and $(-1)^m(y_m - a_m) < 0$ and assume that $m > q + 1$. Moreover, assume that q is odd (the argument is similar for q even). Then $y_1 \cdots y_{q+1} = a_1 \cdots a_{q+1}$ and

$$(\overline{a_{q+1}, \dots, a_{q+p}}) \prec (y_{k+1}, y_{k+2}, \dots).$$

Since the purely periodic sequence on the left-hand side is another boundary sequence, it follows that the forbidden blocks of (y_1, y_2, \dots) are contained within the subsequence $(y_{q+1}, y_{q+2}, \dots)$. But these forbidden blocks are already covered by a set of the form (10). Hence, it is sufficient to consider forbidden blocks of the form in (11). Since there are finitely many boundary sequences and since each of these boundary sequences only have a finite number of forbidden blocks, the shift space is a shift of finite type. \square

We can use the result of this theorem to reduce our graphs quite a bit as follows. Consider any boundary sequence. If its periodic part is a boundary sequence of the same shift space, then remove this periodic part. If our memory contains the whole pre-periodic part of this boundary sequence, then accept it and forget about it (i.e., the new memory index in this specific case would be a zero). The following two examples illustrate this.

Example 26. Reconsider Example 22. Since $(\overline{1, 0})$ is a boundary sequence (the smallest with respect to \prec), we can forget about this periodic part. Moreover, this turns $(1, \overline{1, 0})$ into the single digit boundary block (1), which itself is redundant, and hence we are left with the following condition:

$$(0, 1) \preceq (x_n, x_{n+1})$$

for all $n \geq 1$ such that $x_n = 0$. Although not necessary at this point, we still fill in the table:

$m_0^\ell(i/d)$	0	1
0	1	0
1	F	0

The resulting graph is shown in Figure 5.

The same approach can be applied to any shift space of the form in Theorem 17 as long as at least one eventually periodic boundary sequence has a periodic part equal to a periodic boundary sequence.




Figure 5: Another labeled graph recognizing the shift space of Example 22.

Example 27. Let $\beta = 2$ and consider the left-continuous map with cut points $-\frac{5}{6}$ and $-\frac{1}{6}$. Theorem 17 tells us that $(x_1, x_2, \dots) \in S_{-\beta}$ if and only if for all $n \geq 1$ we have:

$$\begin{aligned} (\overline{2, 0}) &\preceq (x_n, x_{n+1}, \dots) \preceq (2, \overline{1}), \\ (1, \overline{0, 2}) &\preceq (x_n, x_{n+1}, \dots) \preceq (1, \overline{1, 0}), \\ \text{or } (0, \overline{0, 1}) &\preceq (x_n, x_{n+1}, \dots) \preceq (\overline{0, 2}). \end{aligned}$$

Careful inspection reveals that the boundary sequences $(\overline{2, 0})$ and $(\overline{0, 2})$ are redundant (the smallest and largest possible sequence with respect to \preceq). Moreover, we may reduce $(1, \overline{0, 2})$ to the single digit block (1), which is once again redundant. We are left with:

$$\begin{cases} (x_n, x_{n+1}, \dots) \preceq (2, \overline{1}), & \text{whenever } x_n = 2; \\ (x_n, x_{n+1}, \dots) \preceq (1, \overline{1, 0}), & \text{whenever } x_n = 1; \\ (0, \overline{0, 1}) \preceq (x_n, x_{n+1}, \dots), & \text{whenever } x_n = 0. \end{cases} \quad (12)$$

We create tables one more time:

$m_0^l(i/d)$	0	1	2	$m_1^r(i/d)$	0	1	2	$m_2^r(i/d)$	0	1	2
0	1	0	0	0	0	1	0	0	0	0	1
1	2	0	0	1	0	2	F	1	0	2	F
2	0	1	F	2	1	0	0	2	F	1	0

and use these to produce the labeled graph in Figure 6.

6 Discussion

Our goal was to consider a large class of $(-\beta)$ -transformations and to prove statements analogous to what Ito and Sadahiro for their $(-\beta)$ -transformation. Our class of $(-\beta)$ -transformations include transformations that, when restricted to the interval $\left[-\frac{\beta}{\beta+1}, \frac{1}{\beta+1}\right)$, are the Ito-Sadahiro transformation. As to be expected, our results agree with results on their transformation (compare Theorem 14 and 19 with [8, Theorem 10 & 12] and Theorem 25 with [7, Theorem 4]). While we only considered the canonical alphabet, there are many other alphabets to be considered. In fact, there is a rather large class of alphabets satisfying property i of Assumption 1 (compare with [11, Proposition 2.1]) of which we assume the results in this paper will carry over.

Of the many open problems surrounding $(-\beta)$ -expansions we want to highlight two. The first problem is based on the examples in sections 4 and 5 illustrating the output of our algorithm. Can this algorithm be improved upon? The second problem follows from [8]: given two sequences $\mathbf{x}, \mathbf{y} \in \mathcal{A}^{\mathbb{N}}$ (or any other alphabet), is there a $(-\beta)$ -transformation such that \mathbf{x} and \mathbf{y} are boundary sequences of this transformation?

References

- [1] K. Dajani and C. Kalle, Transformations generating negative β -expansions, *Integers* **11B** (2011).
- [2] K. Dajani and C. Kraaikamp, From greedy to lazy expansions and their driving dynamics, *Expo. Math.* **20** (2002), 315–327.
- [3] K. Dajani and C. Kraaikamp, Random β -expansions, *Ergodic Theory Dynam. Systems* **23** (2003), 461–479.
- [4] K. Dajani and M. de Vries, Measures of maximal entropy for random β -expansions, *J. Eur. Math. Soc. (JEMS)* **7** (2005), 51–68.
- [5] D. Dombek, Z. Masáková, and E. Pelantová, Number representation using generalized $(-\beta)$ -transformation, *Theoret. Comput. Sci.* **412** (2011), 6653–6665.
- [6] P. Erdős and I. Joó, On the expansion $1 = \sum q^{-n_i}$, *Period. Math. Hungar.* **23** (1991), 27–30.
- [7] C. Frougny and A. C. Lai, On negative bases, in *Proc. DLT 09*, Lecture Notes in Comp. Sci., Vol. 5583, Springer, 2009, pp. 252–263.
- [8] S. Ito and T. Sadahiro, Beta-expansions with negative bases, *Integers* **9** (2009), 239–259.
- [9] D. Lind and B. Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge University Press, 1995.
- [10] W. Parry, On the β -expansions of real numbers, *Acta. Math. Acad. Sci. Hung.* **11** (1960), 401–416.
- [11] M. Pedicini, Greedy expansions and sets with deleted digits, *Theoret. Comput. Sci.* **332** (2005), 313–336.
- [12] A. Rényi, Representations for real numbers and their ergodic properties, *Acta. Math. Acad. Sci. Hung.* **8** (1957), 477–493.

2010 *Mathematics Subject Classification*: Primary 37B10; Secondary 11A63.

Keywords: $(-\beta)$ -expansions, sofic shift, shift of finite type.

Received October 10 2011; revised version received January 11 2012. Published in *Journal of Integer Sequences*, January 14 2012.

Return to [Journal of Integer Sequences home page](#).




Figure 6: A labeled graph recognizing the shift from Example 27.