



Dyck Paths With No Peaks At Height k

Paul Peart and Wen-Jin Woan

Department of Mathematics
Howard University
Washington, D.C. 20059, USA

Email addresses: pp@scs.howard.edu, wwoan@howard.edu

Abstract

A Dyck path of length $2n$ is a path in two-space from $(0, 0)$ to $(2n, 0)$ which uses only steps $(1, 1)$ (north-east) and $(1, -1)$ (south-east). Further, a Dyck path does not go below the x -axis. A peak on a Dyck path is a node that is immediately preceded by a north-east step and immediately followed by a south-east step. A peak is at height k if its y -coordinate is k . Let $G_k(x)$ be the generating function for the number of Dyck paths of length $2n$ with no peaks at height k with $k \geq 1$. It is known that $G_1(x)$ is the generating function for the Fine numbers (sequence [A000957](#) in [6]). In this paper, we derive the recurrence

$$G_k(x) = \frac{1}{1 - xG_{k-1}(x)}, \quad k \geq 2, \quad G_1(x) = \frac{2}{1 + 2x + \sqrt{1 - 4x}}.$$


It is interesting to see that in the case $k = 2$ we get $G_2(x) = 1 + xC(x)$, where $C(x)$ is the generating function for the ubiquitous Catalan numbers ([A000108](#)). This means that the number of Dyck paths of length $2n + 2$, $n \geq 0$, with no peaks at height 2 is the Catalan number $c_n = \frac{1}{n+1} \binom{2n}{n}$. We also provide a combinatorial proof for this last fact by introducing a bijection between the set of all Dyck paths of length $2n + 2$ with no peaks at height 2 and the set of all Dyck paths of length $2n$.

Keywords: Dyck paths, Catalan number, Fine number, generating function.

1 Introduction

In [1] it was shown that Fine numbers ([A000957](#)) count Dyck paths with no peaks at height 1. One of the results of this paper is that the Catalan numbers ([A000108](#)) count Dyck paths with no peaks at height 2. This provides yet another combinatorial setting for the Catalan numbers (cf. [4], [5], [6], [7]).

A Dyck path is a path in two-space which starts at the origin, stays above the x -axis, and allows only steps of $(1, 1)$ (i.e. north-east) and $(1, -1)$ (i.e. south-east). A Dyck path ends on the x -axis. A Dyck path therefore has even length with the number of north-east steps equal to the number of south-east steps. A lattice point on the path is called a peak if it is immediately preceded by a north-east step and immediately followed by a south-east step. A peak is at height k if its y -coordinate is k . Here are two Dyck paths each of length 10:



The first path has one peak at height 2 and two peaks at height 3. It has no peaks at height 1. The second path has one peak at height 1 and two at height 3. It has no peaks at height 2. Reference [1] contains much information about Dyck paths. It is known that the number of Dyck paths of length $2n$ is c_n , the n^{th} Catalan number, given by

$$c_n = \frac{1}{n+1} \binom{2n}{n}.$$

We will prove that the number of these paths with no peaks at height 2 is c_{n-1} . It is known [1] that the number of these paths with no peaks at height 1 is f_n , the n^{th} Fine number with generating function

$$F(x) = \frac{1}{1 - x^2 C^2(x)} = 1 + x^2 + 2x^3 + 6x^4 + 18x^5 + 57x^6 + 186x^7 + O(x^8)$$

where $C(x) = \frac{1 - \sqrt{1 - 4x}}{2x}$ is the generating function for the Catalan numbers. See [1], [2], and [3] for further information about the Fine numbers. Theorem 2 below contains a proof that the Fine numbers count Dyck paths with no peaks at height 1. In Theorem 1, we obtain the recurrence

$$G_k(x) = \frac{1}{1 - xG_{k-1}(x)}, \quad k \geq 2,$$

where $G_k(x)$ is the generating function for the number of Dyck paths of length $2n$ with no peaks at height k . In Section 3 we introduce a bijection between the set of all Dyck paths of length $2n$ and the set of all Dyck paths of length $2n + 2$ with no peaks at height 2. This bijection provides a combinatorial proof that $G_2(x) = 1 + xC(x)$.

2 Theorems

We will use the fact that

$$F(x) = \sum_{n \geq 0} f_n x^n = \frac{C(x)}{1 + xC(x)}.$$

Theorem 1: Let $G_m(x) = \sum_{n \geq 0} g(m, n)x^n$ be the generating function for Dyck paths of length $2n$ with no peaks at height m , $m \geq 1$. Then

$$G_m(x) = \frac{1}{1 - xG_{m-1}(x)} \quad ; \quad m \geq 2 .$$

PROOF. The set of all Dyck paths of length $2n$, $n \geq 0$, with no peaks at height m consists of the trivial path (the origin) and paths with general form shown in the diagram.

$$\begin{array}{c} A \\ \times \quad B \end{array}$$

It starts with a north-east step followed by a segment labeled A which represents any Dyck path of length $2k$, $0 \leq k \leq n - 1$, with no peaks at height $m - 1$. A is followed by a south-east step followed by a segment labeled B which represents any Dyck path of length $2n - 2 - 2k$ with no peaks at height m . Therefore

$$g(m, 0) = 1, \quad g(m, n) = \sum_{k=0}^{n-1} g(m-1, k)g(m, n-1-k) = [x^{n-1}] \{G_{m-1}(x)G_m(x)\}.$$

$$i.e. \quad g(m, 0) = 1, \quad g(m, n) = [x^n] \{xG_{m-1}(x)G_m(x)\}; \quad n \geq 1,$$

where $[x^k]$ denotes "coefficient of x^k in ". That is,

$$G_m(x) = 1 + x G_{m-1}(x)G_m(x) ,$$

or equivalently,

$$G_m(x) = \frac{1}{1 - xG_{m-1}(x)}$$

■

Theorem 2: The number of Dyck paths of length $2n$ with no peaks at height 1 is the Fine number f_n for $n \geq 0$.

PROOF. With the notation of Theorem 1, we will prove that

$$G_1 = \sum_{n=0}^{\infty} g(1, n)x^n = \frac{1}{1 - x^2C^2}$$

Obviously, $g(1, 0) = 1$ and $g(1, 1) = 0$. For $n \geq 2$, a Dyck path of length $2n$ with no peaks at height 1 has the form of the diagram in the proof of Theorem 1 with A any Dyck path of length $2k$, $1 \leq k \leq n - 1$, and B a Dyck path of length $2n - 2k - 2$ with no peaks at height 1. Therefore, for $n \geq 2$, we have

$$\begin{aligned} g(1, n) &= \sum_{k=1}^{n-1} c_k g(1, n-k-1) = [x^{n-1}] \{C(x)G_1(x)\} - g(1, n-1) \\ &= [x^n] \{xC(x)G_1(x)\} - g(1, n-1) \end{aligned}$$

Therefore

$$\begin{aligned} G_1(x) &= 1 + \sum_{n \geq 2} g(1, n)x^n = 1 + xC(x)G_1(x) - x - xG_1(x) + x \\ &= 1 + xG_1(x)(C(x) - 1) = 1 + xG_1(x)x^2C^2(x) \end{aligned}$$

That is,

$$G_1(x) = \frac{1}{1 - x^2 C^2(x)}$$

■

Theorem 3: The number of Dyck paths of length $2n$ with no peaks at height 2 is the Catalan number c_{n-1} , for $n \geq 1$.

PROOF. From Theorem 1,

$$G_2(x) = \frac{1}{1 - xG_1(x)} = \frac{1}{1 - x \frac{C(x)}{1+xC(x)}} = 1 + xC(x)$$

■

Remark: In [1] it was shown that

$$\frac{f_{n-1}}{c_n} \rightarrow \frac{1}{9} \quad \text{as } n \rightarrow \infty$$

Therefore

$$\frac{f_n}{c_n} \rightarrow \frac{4}{9} \quad \text{as } n \rightarrow \infty$$

Since

$$\frac{c_{n-1}}{c_n} \rightarrow \frac{1}{4} \quad \text{as } n \rightarrow \infty$$

we see that, for sufficiently large n , approximately $\frac{4}{9}$ of the Dyck paths of length $2n$ have no peaks at height 1, while approximately $\frac{1}{4}$ have no peaks at height 2.

Remark: $G_3(x) = \frac{2}{2-3x+x\sqrt{(1-4x)}} = 1 + x + 2x^2 + 4x^3 + 9x^4 + 22x^5 + 58x^6 + 163x^7 + 483x^8 + 1494x^9 + O(x^{10})$ (sequence [A059019](#) in [6]).

3 A bijection between two Catalan families


We end with a bijection between the two Catalan families mentioned in this paper. Let Φ be the set of all Dyck paths of length $2n$ and let Ψ be the set of all Dyck paths of length $2n + 2$ with no peaks at height 2. We define a bijection between Φ and Ψ as follows. First, starting with a Dyck path P from Φ , we obtain a Dyck path \hat{P} from Ψ using the following steps.

- (1) Attach a Dyck path of length 2 to the left of P to produce P^* .
- (2) Let S^* be a maximal sub-Dyck path of P^* with S^* having no peaks at height 1. To each such S^* add a north-east step at the beginning and a south-east step at the end to produce sub-Dyck path \tilde{S} . This step produces a Dyck path \tilde{P} .
- (3) From \tilde{P} eliminate each Dyck path of length 2 that is to the immediate left of each \tilde{S} . We now have a unique element \hat{P} of Ψ .

To obtain P from \hat{P} , we reverse the steps as follows:

- (1) Let \widehat{S} be a sub-Dyck path of \widehat{P} between two consecutive points on the x -axis with \widehat{S} having no peaks at height 1. To each \widehat{S} add a Dyck path of length 2 immediately to the left. This step produces a Dyck path \widetilde{P} .
- (2) Let \widetilde{S} be a maximal sub-Dyck path of \widetilde{P} . From each such \widetilde{S} remove the left-most north-east step and the right-most south-east step to produce a sub-Dyck path S^* . This step produces a Dyck path P^* of length $2n+2$.
- (3) From P^* , remove the left-most Dyck path of length 2 to produce P .

For example, we obtain a Dyck path of length 18 with no peaks at height 2 starting with a Dyck path of length 16 as follows:



It is now easy to show that the Catalan numbers count parallelogram polyominoes (or Fat Path Pairs) with no columns at height 2 (see [7], p. 257).

References

- [1] E. Deutsch. *Dyck Path Enumeration*. Discrete Math. 204 (1999), no. 1-3, 167-202.
- [2] E. Deutsch & L. W. Shapiro. *Fine Numbers*. Preprint.
- [3] T. Fine. *Extrapolation when very little is known about the source*. Information and Control 16 (1970) 331-359.

- [4] H. W. Gould. *Bell & Catalan Numbers: Research Bibliography of Two Special Number Sequences*, 6th ed. Morgantown, WV: Math Monongliae, 1985.
- [5] L. W. Shapiro. *A Catalan Triangle*. *Discrete Math.* 14 (1976) 83-90.
- [6] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences. Published electronically at <http://www.research.att.com/~njas/sequences/> .
- [7] R. P. Stanley. *Enumerative Combinatorics*. Vol. 2. Cambridge University Press, 1999.

(Concerned with sequences [A000108](#), [A000957](#), [A059019](#), [A059027](#).)

Received October 16, 2000; revised version received February 8, 2001; published in *Journal of Integer Sequences*, May 12, 2001.

Return to [Journal of Integer Sequences home page](#).
