

*Changhong Wu and Lanzhe Liu*

**BOUNDEDNESS FOR MULTILINEAR COMMUTATOR OF  
 LITTLEWOOD-PALEY OPERATOR ON HARDY AND  
 HERZ-HARDY SPACES**

(submitted by D. Kh. Mushtari)

**ABSTRACT.** In this paper, the  $(H_{\vec{b}}^p, L^p)$  and  $(H\dot{K}_{q,\vec{b}}^{\alpha,p}, \dot{K}_q^{\alpha,p})$  type boundedness for the multilinear commutator associated with the Littlewood-Paley operator are obtained.

**1. INTRODUCTION AND DEFINITION**

Let  $L_{loc}^q(R^n) = \{f^q \text{ is locally integrable on } R^n\}$ . Suppose  $f \in L_{loc}^1(R^n)$ ,  $B = B(x_0, r) = \{x \in R^n : |x - x_0| < r\}$  denotes a ball of  $R^n$  centered at  $x_0$  and having radius  $r$ , write  $f_B = |B|^{-1} \int_B f(x)dx$  and  $f^\#(x) = \sup_{x \in B} |B|^{-1} \int_B |f(x) - f_B|dx < \infty$ . The function  $f$  is said to belong to  $BMO(R^n)$  if  $f^\# \in L^\infty(R^n)$ , and define  $\|f\|_{BMO} = \|f^\#\|_{L^\infty}$ .

Let  $T$  be a linear operator and  $K$  be a function on  $R^n \times R^n$ ,

$$T(f)(x) = \int_{R^n} K(x, y) f(y) dy \quad \text{for } f \in C_0^\infty,$$

where  $K$  satisfies:

- (1)  $|K(x, y + h) - K(x, y)| \leq C \cdot |h|^\alpha \cdot |x - y|^{-n-\alpha}$  for  $2|h| < |x - y|$ ,  $0 < \alpha \leq 1$ ;
- (2)  $\|T(f)\|_{L^{p_0}} \leq C \|f\|_{L^{p_0}}$  for some  $1 < p_0 \leq \infty$ ;

*2000 Mathematical Subject Classification.* 42B20, 42B25.

*Key words and phrases.* Littlewood-Paley operator, Multilinear commutator, BMO, Hardy space, Herz-Hardy space.

Then we call  $T$  is the Calderón-Zygmund singular integral operator.

Let  $b \in BMO(R^n)$  and  $T$  be the Calderón-Zygmund singular integral operator. The commutator  $[b, T]$  generated by  $b$  and  $T$  is defined by

$$[b, T](f)(x) = b(x)T(f)(x) - T(bf)(x).$$

A classical result of Coifman, Rochberg and Weiss (see[2]) proved that the commutator  $[b, T]$  is bounded on  $L^p(R^n)$  ( $1 < p < \infty$ ). However, it was observed that the  $[b, T]$  is not bounded, in general, from  $H^p(R^n)$  to  $L^p(R^n)$ . But if  $H^p(R^n)$  is replaced by a suitable atomic space  $H_{\vec{b}}^p(R^n)$  and  $H\dot{K}_{q,\vec{b}}^{\alpha,p}(R^n)$ (see[1][7][12]), then  $[b, T]$  maps continuously  $H_{\vec{b}}^p(R^n)$  into  $L^p(R^n)$  and  $H\dot{K}_{q,\vec{b}}^{\alpha,p}(R^n)$  into  $\dot{K}_q^{\alpha,p}$ . In addition we have easily known that  $H_{\vec{b}}^p(R^n) \subset H^p(R^n)$ ,  $\dot{K}_{q,\vec{b}}^{\alpha,p}(R^n) \subset H\dot{K}_q^{\alpha,p}(R^n)$ . The main purpose of this paper is to consider the continuity of the multilinear commutators related to the Littlewood-Paley operators and  $BMO(R^n)$  functions on certain Hardy and Herz-Hardy spaces. Let us first introduce some definitions(see [1][3-10][12][13]).

Given a positive integer  $m$  and  $1 \leq j \leq m$ , we denote by  $C_j^m$  the family of all finite subsets  $\sigma = \{\sigma(1), \dots, \sigma(j)\}$  of  $\{1, \dots, m\}$  of  $j$  different elements. For  $\sigma \in C_j^m$ , set  $\sigma^c = \{1, \dots, m\} \setminus \sigma$ . For  $\vec{b} = (b_1, \dots, b_m)$  and  $\sigma = \{\sigma(1), \dots, \sigma(j)\} \in C_j^m$ , set  $\vec{b}_\sigma = (b_{\sigma(1)}, \dots, b_{\sigma(j)})$ ,  $b_\sigma = b_{\sigma(1)} \cdots b_{\sigma(j)}$  and  $\|\vec{b}_\sigma\|_{BMO} = \|b_{\sigma(1)}\|_{BMO} \cdots \|b_{\sigma(j)}\|_{BMO}$ .

**Definition 1.** Let  $b_i$  ( $i = 1, \dots, m$ ) be a locally integrable functions and  $0 < p \leq 1$ . A bounded measurable function  $a$  on  $R^n$  is called a  $(p, \vec{b})$  atom, if

- (1)  $\text{supp } a \subset B = B(x_0, r)$
- (2)  $\|a\|_{L^\infty} \leq |B(x_0, r)|^{-1/p}$
- (3)  $\int_B a(y) dy = \int_B a(y) \prod_{l \in \sigma} b_l(y) dy = 0$  for any  $\sigma \in C_j^m$ ,  $1 \leq j \leq m$ .

A temperate distribution(see[14][15])  $f$  is said to belong to  $H_{\vec{b}}^p(R^n)$ , if, in the Schwartz distribution sense, it can be written as

$$f(x) = \sum_{j=1}^{\infty} \lambda_j a_j(x).$$

where  $a'_j$ s are  $(p, \vec{b})$  atoms,  $\lambda_j \in C$  and  $\sum_{j=1}^{\infty} |\lambda_j|^p < \infty$ . Moreover,  $\|f\|_{H_{\vec{b}}^p} \approx (\sum_{j=1}^{\infty} |\lambda_j|^p)^{1/p}$ .

**Definition 2.** Let  $0 < p, q < \infty$ ,  $\alpha \in R$ . For  $k \in Z$ , set  $B_k = \{x \in R^n : |x| \leq 2^k\}$  and  $C_k = B_k \setminus B_{k-1}$ , and  $\chi_k = \chi_{C_k}$  for  $k \in Z$ , where  $\chi_{C_k}$  is

the characteristic function of set  $C_k$ . Denote the characteristic function of  $B_0$  by  $\chi_0$ .

(1) The homogeneous Herz space is defined by

$$\dot{K}_q^{\alpha,p}(R^n) = \left\{ f \in L_{loc}^q(R^n \setminus \{0\}) : \|f\|_{\dot{K}_q^{\alpha,p}} < \infty \right\}.$$

where

$$\|f\|_{\dot{K}_q^{\alpha,p}} = \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|f\chi_k\|_{L^q}^p \right]^{1/p}.$$

(2) The nonhomogeneous Herz space is defined by

$$K_q^{\alpha,p}(R^n) = \left\{ f \in L_{loc}^q(R^n) : \|f\|_{K_q^{\alpha,p}} < \infty \right\}.$$

where

$$\|f\|_{K_q^{\alpha,p}} = \left[ \sum_{k=1}^{\infty} 2^{k\alpha p} \|f\chi_k\|_{L^q}^p + \|f\chi_0\|_{L^q}^p \right]^{1/p}.$$

**Definition 3.** Let  $\alpha \in R^n$ ,  $1 < q < \infty$ ,  $\alpha \geq n(1 - 1/q)$ ,  $b_i \in BMO(R^n)$ ,  $1 \leq i \leq m$ . A function  $a(x)$  is called a central  $(\alpha, q, \vec{b})$ -atom (or a central  $(\alpha, q, \vec{b})$ -atom of restrict type), if

- (1)  $\text{supp } a \subset B = B(x_0, r)$  (or for some  $r \geq 1$ ),
- (2)  $\|a\|_{L^q} \leq |B(x_0, r)|^{-\alpha/n}$ ,
- (3)  $\int_B a(x)x^\beta dx = \int_B a(x)x^\beta \prod_{i \in \sigma} b_i(x) dx = 0$  for any  $\sigma \in C_j^m$ ,  $1 \leq j \leq m$ ,  $0 \leq |\beta| \leq \alpha$ , where  $\beta = (\beta_1, \dots, \beta_n)$  is the multi-indices with  $\beta_i \in N$  for  $1 \leq i \leq n$  and  $|\beta| = \sum_{i=1}^n \beta_i$ .

A temperate distribution  $f$  is said to belong to  $H\dot{K}_{q,\vec{b}}^{\alpha,p}(R^n)$  (or  $HK_{q,\vec{b}}^{\alpha,p}(R^n)$ ), if it can be written as  $f = \sum_{j=-\infty}^{\infty} \lambda_j a_j$  (or  $f = \sum_{j=0}^{\infty} \lambda_j a_j$ ), in the  $S'(R^n)$  sense, where  $a_j$  is a central  $(\alpha, q, \vec{b})$ -atom (or a central  $(\alpha, q, \vec{b})$ -atom of restrict type) supported on  $B(0, 2^j)$  and  $\sum_{-\infty}^{\infty} |\lambda_j|^p < \infty$  (or  $\sum_{j=0}^{\infty} |\lambda_j| < \infty$ ). Moreover,

$$\|f\|_{H\dot{K}_{q,\vec{b}}^{\alpha,p}} \quad (\text{or } \|f\|_{HK_{q,\vec{b}}^{\alpha,p}}) = \inf \left( \sum_j |\lambda_j|^p \right)^{1/p},$$

where the infimum are taken over all the decompositions of  $f$  as above.

**Definition 4.** Let  $\varepsilon > 0$  and  $\psi$  be a fixed function which satisfies the following properties:

- (1)  $\int_{R^n} \psi(x) dx = 0$ ,
- (2)  $|\psi(x)| \leq C(1 + |x|)^{-(n+1)}$ ,
- (3)  $|\psi(x+y) - \psi(x)| \leq C|y|^\varepsilon(1 + |x|)^{-(n+1+\varepsilon)}$  when  $2|y| < |x|$ ;

The Littlewood-Paley multilinear commutator is defined by

$$S_{\psi}^{\vec{b}}(f)(x) = \left[ \int \int_{\Gamma(x)} |F_t^{\vec{b}}(f)(x, y)|^2 \frac{dy dt}{t^{n+1}} \right]^{1/2},$$

where

$$F_t^{\vec{b}}(f)(x, y) = \int_{R^n} \left[ \prod_{j=1}^m (b_j(x) - b_j(z)) \right] \psi_t(y - z) f(z) dz,$$

$\Gamma(x) = \{(y, t) \in R_+^{n+1} : |x - y| < t\}$  and  $\psi_t(x) = t^{-n} \psi(x/t)$  for  $t > 0$ . Set  $F_t(f)(y) = \int_{R^n} \psi_t(y - x) f(x) dx$ . We also define that

$$S_{\psi}(f)(x) = \left( \int \int_{\Gamma(x)} |F_t(f)(x, y)|^2 \frac{dy dt}{t^{n+1}} \right)^{1/2},$$

which is the Littlewood-Paley operator (see [15]).

## 2. THEOREMS AND PROOFS

**Theorem 1.** Let  $\varepsilon > 0, b_i \in BMO, 1 \leq i \leq m, \vec{b} = (b_1, \dots, b_m), n/(n+\varepsilon) < p \leq 1$ . Then the multilinear commutator  $S_{\psi}^{\vec{b}}$  is bounded from  $H_{\vec{b}}^p(R^n)$  to  $L^p(R^n)$ .

**Proof.** It suffices to show that there exist a constant  $C > 0$ , such that for every  $(p, \vec{b})$  atom  $a$ ,

$$\|S_{\psi}^{\vec{b}}(a)\|_{L^p} \leq C.$$

Let  $a$  be a  $(p, \vec{b})$  atom supported on a ball  $B = B(x_0, r)$ . When  $m = 1$  see [7], and now we prove  $m > 1$ . Write

$$\begin{aligned} & \int_{R^n} |S_{\psi}^{\vec{b}}(a)(x)|^p dx \\ &= \int_{|x-x_0| \leq 2r} |S_{\psi}^{\vec{b}}(a)(x)|^p dx + \int_{|x-x_0| > 2r} |S_{\psi}^{\vec{b}}(a)(x)|^p dx = I + II. \end{aligned}$$

For  $I$ , taking  $q > 1$ , by Hölder's inequality and the  $L^q$ -boundedness of  $S_{\psi}^{\vec{b}}$ , we have

$$\begin{aligned} I &\leq \left( \int_{|x-x_0| \leq 2r} |S_{\psi}^{\vec{b}}(a)(x)|^q dx \right)^{p/q} \cdot |B(x_0, 2r)|^{1-p/q} \\ &\leq C \|S_{\psi}^{\vec{b}}(a)(x)\|_{L^q}^p \cdot |B(x_0, 2r)|^{1-p/q} \\ &\leq C \|\vec{b}\|_{BMO}^p \|a\|_{L^q}^p |B|^{1-p/q} \leq C \|\vec{b}\|_{BMO}^p. \end{aligned}$$

For  $II$ , denoting  $\lambda = (\lambda_1, \dots, \lambda_m)$  with  $\lambda_i = (b_i)_B$ ,  $1 \leq i \leq m$ , where  $(b_i)_B = |B(x_0, r)|^{-1} \int_{B(x_0, r)} b_i(x) dx$ , by Hölder's inequality and the vanishing moment of  $a$ , we get

$$\begin{aligned} II &= \sum_{k=1}^{\infty} \int_{2^{k+1}r \geq |x-x_0| > 2^k r} |S_{\psi}^{\vec{b}}(a)(x)|^p dx \\ &\leq C \sum_{k=1}^{\infty} |B(x_0, 2^{k+1}r)|^{1-p} \left( \int_{2^{k+1}r \geq |x-x_0| > 2^k r} |S_{\psi}^{\vec{b}}(a)(x)| dx \right)^p \\ &\leq C \sum_{k=1}^{\infty} |B(x_0, 2^{k+1}r)|^{1-p} \times \left[ \int_{2^{k+1}r \geq |x-x_0| > 2^k r} \left( \int_{\Gamma(x)} \left| \int_B \prod_{j=1}^m (b_j(x) - b_j(z)) \psi_t(y-z) a(z) dz \right|^2 \frac{dy dt}{t^{n+1}} \right)^{1/2} dx \right]^p \\ &\leq C \sum_{k=1}^{\infty} |B(x_0, 2^{k+1}r)|^{1-p} \times \left[ \int_{2^{k+1}r \geq |x-x_0| > 2^k r} \left( \int_{\Gamma(x)} \left( \int_B |\psi_t(y-z) - \psi_t(y-x_0)| \prod_{j=1}^m |(b_j(x) - b_j(z))| |a(z)| dz \right)^2 \frac{dy dt}{t^{n+1}} \right)^{1/2} dx \right]^p; \end{aligned}$$

noting that  $z \in B$ ,  $x \in B(x_0, 2^{k+1}r) \setminus B(x_0, 2^k r)$ , then

$$\begin{aligned} S_{\psi}^{\vec{b}}(a)(x) &= \left[ \int_{\Gamma(x)} \left( \int_B |\psi_t(y-z) - \psi_t(y-x_0)| \prod_{j=1}^m |b_j(x) - b_j(z)| |a(z)| dz \right)^2 \frac{dy dt}{t^{n+1}} \right]^{1/2} \\ &\leq C \left[ \int_{\Gamma(x)} \left( \int_B t^{-n} |a(z)| \prod_{j=1}^m |b_j(x) - b_j(z)| \frac{(|x_0 - z|/t)^{\varepsilon}}{(1 + |x_0 - y|/t)^{n+1+\varepsilon}} dz \right)^2 \frac{dy dt}{t^{n+1}} \right]^{1/2} \\ &\leq C \left( \int_{\Gamma(x)} \frac{t^{1-n}}{(t + |x_0 - y|)^{2(n+1+\varepsilon)}} dy dt \right)^{1/2} \\ &\quad \times \int_B \prod_{j=1}^m |b_j(x) - b_j(z)| |x_0 - z|^{\varepsilon} |a(z)| dz \\ &\leq C \left( \int_{\Gamma(x)} \frac{t^{1-n} 2^{2(n+1+\varepsilon)}}{(2t + 2|x_0 - y|)^{2(n+1+\varepsilon)}} dy dt \right)^{1/2} \\ &\quad \times \int_B \prod_{j=1}^m |b_j(x) - b_j(z)| |x_0 - z|^{\varepsilon} |a(z)| dz; \end{aligned}$$

Notice that  $2t + |x_0 - y| > 2t + |x_0 - x| - |x - y| > t + |x_0 - x|$  when  $|x - y| < t$ , and it is easy to calculate that

$$\int_0^\infty \frac{tdt}{(t + |x - x_0|)^{2(n+1+\varepsilon)}} = C|x - x_0|^{-2(n+\varepsilon)};$$

then, we deduce

$$\begin{aligned} S_\psi^{\vec{b}}(a)(x) &\leq C \left( \int \int_{\Gamma(x)} \frac{t^{1-n}}{(2t + |x_0 - y|)^{2(n+1+\varepsilon)}} dy dt \right)^{1/2} \\ &\quad \times \int_B \prod_{j=1}^m |b_j(x) - b_j(z)| |x_0 - z|^\varepsilon |a(z)| dz \\ &\leq C \left( \int \int_{\Gamma(x)} \frac{t^{1-n}}{(t + |x - x_0|)^{2(n+1+\varepsilon)}} dy dt \right)^{1/2} \\ &\quad \times \int_B \prod_{j=1}^m |b_j(x) - b_j(z)| |x_0 - z|^\varepsilon |a(z)| dz \\ &\leq C \left( \int_0^\infty \frac{tdt}{(t + |x - x_0|)^{2(n+1+\varepsilon)}} \right)^{1/2} \\ &\quad \times \int_B \prod_{j=1}^m |b_j(x) - b_j(z)| |x_0 - z|^\varepsilon |a(z)| dz \\ &\leq C|B|^{\varepsilon/n-1/p} |x - x_0|^{-(n+\varepsilon)} \int_B \prod_{j=1}^m |b_j(x) - b_j(z)| dz. \end{aligned}$$

So

$$\begin{aligned} II &\leq C|B|^{\varepsilon/n-1/p} \sum_{k=1}^\infty |B(x_0, 2^{k+1}r)|^{1-p} \\ &\quad \times \left[ \int_{2^{k+1}r \geq |x - x_0| > 2^k r} |x - x_0|^{-(n+\varepsilon)} \left( \int_B \prod_{j=1}^m |b_j(x) - b_j(z)| dz \right) dx \right]^p \\ &\leq C|B|^{\varepsilon/n-1/p} \sum_{k=1}^\infty |B(x_0, 2^{k+1}r)|^{1-p} \\ &\quad \times \left[ \sum_{j=0}^m \sum_{\sigma \in C_j^m} \int_{2^{k+1}r \geq |x - x_0| > 2^k r} |x - x_0|^{-(n+\varepsilon)} |(\vec{b}(x) - \lambda)_\sigma| dx \int_B |(\vec{b}(z) - \lambda)_{\sigma^c}| dz \right]^p \end{aligned}$$

$$\begin{aligned}
&\leq C|B|^{\varepsilon/n-1/p} \sum_{j=0}^m \sum_{\sigma \in C_j^m} \left( \int_B |(\vec{b}(z) - \lambda)_{\sigma^c}| dz \right)^p \sum_{k=1}^{\infty} |B(x_0, 2^{k+1}r)|^{1-p} \\
&\quad \times \left[ \int_{2^{k+1}r \geq |x-x_0| > 2^k r} |x-x_0|^{-(n+\varepsilon)} |(\vec{b}(x) - \lambda)_{\sigma}| dx \right]^p \\
&\leq C \sum_{j=0}^m \sum_{\sigma \in C_j^m} \|\vec{b}_{\sigma^c}\|_{BMO}^p \\
&\quad \cdot \|\vec{b}_{\sigma}\|_{BMO}^p \sum_{k=1}^{\infty} |B(x_0, 2^{k+1}r)|^{1-(n+\varepsilon)p/n} k^p |B|^{(1+\varepsilon/n-1/p)p} \\
&\leq C \|\vec{b}\|_{BMO}^p \sum_{k=1}^{\infty} k^p \cdot 2^{kn(1-p(n+\varepsilon)/n)} \leq C \|\vec{b}\|_{BMO}.
\end{aligned}$$

This finishes the proof of Theorem 1.

**Theorem 2.** Let  $0 < p < \infty$ ,  $1 < q < \infty$ ,  $n(1 - 1/q) \leq \alpha < n(1 - 1/q) + \varepsilon$ ,  $\varepsilon > 0$ , and  $b_i \in BMO(R^n)$ ,  $1 \leq i \leq m$ ,  $\vec{b} = (b_1, \dots, b_m)$ . Then  $S_{\psi}^{\vec{b}}$  is bounded from  $H\dot{K}_{q,\vec{b}}^{\alpha,p}(R^n)$  to  $\dot{K}_q^{\alpha,p}(R^n)$ .

**Proof.** Let  $f \in H\dot{K}_{q,\vec{b}}^{\alpha,p}(R^n)$  and  $f(x) = \sum_{j=-\infty}^{\infty} \lambda_j a_j(x)$  be the atomic decomposition for  $f$  as in Definition 3, we write

$$\begin{aligned}
\|S_{\psi}^{\vec{b}}(f)(x)\|_{\dot{K}_q^{\alpha,p}} &= \left( \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|S_{\psi}^{\vec{b}}(f)\chi_k\|_{L_q}^p \right)^{1/p} \\
&\leq C \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{\infty} |\lambda_j| \|S_{\psi}^{\vec{b}}(a_j)\chi_k\|_{L_q} \right)^p \right]^{1/p} \\
&\leq C \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{k-3} |\lambda_j| \|S_{\psi}^{\vec{b}}(a_j)\chi_k\|_{L_q} \right)^p \right]^{1/p} \\
&\quad + C \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=k-2}^{\infty} |\lambda_j| \|S_{\psi}^{\vec{b}}(a_j)\chi_k\|_{L_q} \right)^p \right]^{1/p} \\
&= I + II.
\end{aligned}$$

For  $II$ , by the boundedness of  $S_{\psi}^{\vec{b}}$  on  $L^q$  and the Hölder's inequality, we have

$$II \leq C \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=k-2}^{\infty} |\lambda_j| \|S_{\psi}^{\vec{b}}(a_j)\chi_j\|_{L_q} \right)^p \right]^{1/p}$$

$$\begin{aligned}
&\leq C \left[ \sum_{-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=k-2}^{\infty} |\lambda_j| \|a_j\|_{L^q} \right)^p \right]^{1/p} \\
&\leq C \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=k-2}^{\infty} |\lambda_j| \cdot 2^{-j\alpha} \right)^p \right]^{1/p} \\
&\leq C \begin{cases} \left[ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \sum_{k=-\infty}^{j+2} 2^{(k-j)\alpha p} \right]^{1/p}, & 0 < p \leq 1 \\ \left[ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \sum_{k=-\infty}^{j+2} 2^{(k-j)p/2} \right]^{1/p}, & 1 < p < \infty \end{cases} \\
&\leq C \left( \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right)^{1/p} \\
&\leq C \|f\|_{H\dot{K}_{q,\tilde{b}}^{\alpha,p}}.
\end{aligned}$$

For  $I$ , when  $m=1$ , let  $b_j^1 = |B_j|^{-1} \int_{B_j} b_1(x) dx$ . We have

$$\begin{aligned}
&S_\psi^{b_1}(a_j)(x) \\
&= \left[ \int \int_{\Gamma(x)} \left| \int_{B_j} (b_1(x) - b_1(z)) \psi_t(y-z) a_j(z) dz \right|^2 \frac{dy dt}{t^{n+1}} \right]^{1/2} \\
&\leq \left[ \int \int_{\Gamma(x)} \left( \int_{B_j} |\psi_t(y-z) - \psi_t(y-x_0)| |b_1(y) - b_1(z)| |a_j(z)| dz \right)^2 \frac{dy dt}{t^{n+1}} \right]^{1/2} \\
&\leq \left[ \int \int_{\Gamma(x)} \left( \int_{B_j} t^{-n} |a_j(z)| |b_1(x) - b_1(z)| \frac{(|x_0 - z|/t)^\varepsilon}{(1 + |x_0 - y|/t)^{n+1+\varepsilon}} dy \right)^2 \frac{dy dt}{t^{n+1}} \right]^{1/2} \\
&\leq C \left( \int \int_{\Gamma(x)} \frac{t^{1-n}}{(t + |x_0 - y|)^{2(n+1+\varepsilon)}} dy dt \right)^{1/2} \\
&\quad \int_{B_j} |x_0 - z|^\varepsilon |a_j(z)| |b_1(x) - b_1(z)| dz \\
&\leq C \left( \int \int_{\Gamma(x)} \frac{t^{1-n}}{(2t + |x_0 - y|)^{2(n+1+\varepsilon)}} dy dt \right)^{1/2} \\
&\quad \int_{B_j} |x_0 - z|^\varepsilon |a_j(z)| |b_1(x) - b_1(z)| dz \\
&\leq C \left( \int \int_{\Gamma(x)} \frac{t^{1-n}}{(t + |x - x_0|)^{2(n+1+\varepsilon)}} dy dt \right)^{1/2} \\
&\quad \int_{B_j} |x_0 - z|^\varepsilon |a_j(z)| |b_1(x) - b_1(z)| dz
\end{aligned}$$

$$\begin{aligned}
 &\leq C \left( \int_0^\infty \frac{tdt}{(t+|x-x_0|)^{2(n+1+\varepsilon)}} \right)^{1/2} \left( \int_{B_j} |x_0-z|^\varepsilon |a_j(z)| |b_1(x) - b_1(z)| dz \right) \\
 &\leq C|x-x_0|^{-(n+\varepsilon)} \int_{B_j} |x_0-z|^\varepsilon |a_j(z)| |b_1(x) - b_1(z)| dz \\
 &\leq C|x-x_0|^{-(n+\varepsilon)} \int_{B_j} |x_0-z|^\varepsilon |a_j(z)| |b_1(x) - b_j^1| dz \\
 &+ C|x-x_0|^{-(n+\varepsilon)} \int_{B_j} |x_0-z|^\varepsilon |a_j(z)| |b_1(z) - b_j^1| dz \\
 &\leq C|x-x_0|^{-(n+\varepsilon)} (|b_1(x) - b_j^1| 2^{j(\varepsilon+n(1-1/q)-\alpha)} + 2^{j(\varepsilon+n(1-1/q)-\alpha)} \|b_1\|_{BMO}).
 \end{aligned}$$

So

$$\begin{aligned}
 &\|S_\psi^{b_1}(a_j)\chi_k\|_{L_q} \\
 &\leq C 2^{j(\varepsilon+n(1-1/q)-\alpha)} \left[ \left( \int_{B_k} |b_1(x) - b_j^1| |x-x_0|^{-q(n+\varepsilon)} dx \right)^{1/q} \right. \\
 &\quad \left. + \left( \int_{B_k} |x-x_0|^{-q(n+\varepsilon)} dx \right)^{1/q} \|b_1\|_{BMO} \right] \\
 &\leq C 2^{j(\varepsilon+n(1-1/q)-\alpha)} [2^{-k(n+\varepsilon)} \cdot |B_k|^{1/q} \|b_1\|_{BMO} \\
 &\quad + 2^{-k(n+\varepsilon)} \cdot |B_k|^{1/q} \|b_1\|_{BMO}] \\
 &\leq C \|b_1\|_{BMO} 2^{[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]} ;
 \end{aligned}$$

thus

$$\begin{aligned}
 I &= \\
 C \left[ \sum_{j=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{k-3} |\lambda_j| \|S_\psi^{b_1}(a_j)\chi_k\|_{L^q} \right)^p \right]^{1/p} \\
 &\leq C \|b_1\|_{BMO} \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{k-3} |\lambda_j| 2^{[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]} \right)^p \right]^{1/p} \\
 &\leq C \|b_1\|_{BMO} \\
 &\times \begin{cases} \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \sum_{j=-\infty}^{k-3} |\lambda_j|^p 2^{[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]p} \right]^{1/p}, \\ 0 < p \leq 1 \\ \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{k-3} |\lambda_j|^p 2^{[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]/2} \right)^{p/p'} \right]^{1/p}, \\ 1 < p < \infty \end{cases}
 \end{aligned}$$

$$\begin{aligned}
&\leq C\|b_1\|_{BMO} \\
&\times \begin{cases} \left[ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \sum_{k=j+3}^{\infty} 2^{(j-k)(\varepsilon+n(1-1/q)-\alpha)p} \right]^{1/p}, & 0 < p \leq 1 \\ \left[ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \sum_{k=j+3}^{\infty} 2^{(j-k)(\varepsilon+n(1-1/q)-\alpha)p/2} \right]^{1/p}, & 1 < p < \infty \end{cases} \\
&\leq C\|b_1\|_{BMO} \left( \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right)^{1/p} \leq C\|f\|_{H\vec{K}_{q,\vec{b}}^{\alpha,p}}.
\end{aligned}$$

When  $m > 1$ , Let  $b_j^i = |B_j|^{-1} \int_{B_j} b_i(x) dx$ ,  $1 \leq i \leq m$ ,  $\vec{b} = (b_j^1, \dots, b_j^m)$ .

We have

$$\begin{aligned}
&S_{\psi}^{\vec{b}}(a_j)(x) \\
&= \left[ \int \int_{\Gamma(x)} \left| \int_{B_j} \prod_{i=1}^m (b_i(x) - b_i(z)) \psi_t(y-z) a_j(z) dz \right|^2 \frac{dydt}{t^{n+1}} \right]^{1/2} \\
&\leq \left[ \int \int_{\Gamma(x)} \left( \int_{B_j} \prod_{i=1}^m |b_i(x) - b_i(z)| |\psi_t(y-z) - \psi_t(y-x_0)| |a_j(z)| dz \right)^2 \frac{dydt}{t^{n+1}} \right]^{1/2} \\
&\leq C|x-x_0|^{-(n+\varepsilon)} \int_{B_j} |x_0 - z|^\varepsilon |a_j(z)| \prod_{i=1}^m |b_i(x) - b_i(z)| dz \\
&\leq C|x-x_0|^{-(n+\varepsilon)} \int_{B_j} |x_0 - z|^\varepsilon |a_j(z)| \prod_{i=1}^m |b_i(x) - b_i(y)| dz \\
&\leq C|x-x_0|^{-(n+\varepsilon)} \\
&\times \sum_{i=0}^m \sum_{\sigma \in C_i^m} |(\vec{b}(x) - \vec{b})_\sigma| \int_{B_j} |x_0 - z|^\varepsilon |a_j(z)| |(\vec{b}(y) - \vec{b})_{\sigma^c}| dz \\
&\leq C|x-x_0|^{-(n+\varepsilon)} \sum_{i=0}^m \sum_{\sigma \in C_i^m} |(\vec{b}(x) - \vec{b})_\sigma| 2^{j\varepsilon} \cdot 2^{-j\alpha} \cdot 2^{jn(1-1/q)} \|\vec{b}_{\sigma^c}\|_{BMO} \\
&\leq C|x-x_0|^{-(n+\varepsilon)} \cdot 2^{j(\varepsilon+n(1-1/q)-\alpha)} \sum_{i=0}^m \sum_{\sigma \in C_i^m} |(\vec{b}(x) - \vec{b})_\sigma| \|\vec{b}_{\sigma^c}\|_{BMO};
\end{aligned}$$

So

$$\begin{aligned}
&\|S_{\psi}^{\vec{b}}(a_j)\chi_k\|_{L^q} \leq C 2^{j(\varepsilon+n(1-1/q)-\alpha)} \|\vec{b}_{\sigma^c}\|_{BMO} \\
&\times \left[ \int_{B_k} \left( |x|^{-(n+\varepsilon)} \sum_{i=0}^m \sum_{\sigma \in C_i^m} |(\vec{b}(x) - \vec{b})_\sigma| \right)^q dx \right]^{1/q}
\end{aligned}$$

$$\begin{aligned} &\leq C\|\vec{b}_{\sigma^c}\|_{BMO}2^{j(\varepsilon+n(1-1/q)-\alpha)} \cdot 2^{-k(n+\varepsilon)+kn/q} \\ &\leq C\|\vec{b}\|_{BMO}; \end{aligned}$$

then

$$\begin{aligned} I &= C \left[ \sum_{j=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{k-3} |\lambda_j| \|S_{\psi}^{\vec{b}}(a_j)\chi_k\|_{L^q} \right)^p \right]^{1/p} \\ &\leq C\|\vec{b}\|_{BMO} \\ &\quad \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{k-3} |\lambda_j| 2^{[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]} \right)^p \right]^{1/p} \\ &\leq C\|\vec{b}\|_{BMO} \\ &\times \begin{cases} \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \sum_{j=-\infty}^{k-3} |\lambda_j|^p 2^{[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]p} \right]^{1/p}, \\ 0 < p \leq 1 \\ \left[ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left( \sum_{j=-\infty}^{k-3} |\lambda_j|^p 2^{p[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]/2} \right) \right. \\ \quad \left. \times \left( \sum_{j=-\infty}^{k-3} 2^{p'[j(\varepsilon+n(1-1/q)-\alpha)-k(n+\varepsilon)+kn/q]} \right)^{p/p'} \right]^{1/p}, \\ 1 < p < \infty \end{cases} \\ &\leq C\|\vec{b}\|_{BMO} \\ &\quad \begin{cases} \left[ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \sum_{k=j+3}^{\infty} 2^{(j-k)(\varepsilon+n(1-1/q)-\alpha)p} \right]^{1/p}, \\ 0 < p \leq 1 \\ \left[ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \sum_{k=j+3}^{\infty} 2^{(j-k)(\varepsilon+n(1-1/q)-\alpha)p/2} \right]^{1/p}, \\ 1 < p < \infty \end{cases} \\ &\leq C\|\vec{b}\|_{BMO} \left( \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right)^{1/p} \\ &\leq C\|f\|_{H\dot{K}_{q,\vec{b}}^{\alpha,p}}. \end{aligned}$$

**Remark.** Theorem 2 also hold for nonhomogeneous Herz-type spaces, we omit the details.

## REFERENCES

- [1] J. Alvarez, Continuity properties for linear commutators of Calderón-Zygmund operators, Collect. Math., 49(1998), 17-31.
- [2] R. Coifman, R. Rochberg and G. Weiss, Factorization theorem for Hardy space in several variables, Ann. of Math., 103(1976), 611-635.

- [3] J.Garcia-Cuerva and M.L.Herrero, A Theory of Hardy spaces associated to Herz Spaces, Proc. London Math. Soc., 69(1994), 605-628.
- [4] L.Z.Liu, Weighted weak type  $(H^1, L^1)$  estimates for commutators of Littlewood-Paley operators, Indian J. of Math., 45(1)(2003), 71-78.
- [5] L.Z.Liu, Weighted Block-Hardy spaces estimates for commutators of Littlewood-Paley operators, Southeast Asian Bull. of Math., 27(2004), 833-838.
- [6] L.Z.Liu, Weighted weak type estimates for commutators of Littlewood-Paley operator, Japanese J. of Math., 29(1)(2003), 1-13.
- [7] L.Z.Liu, S.Z.Lu and J.S.Xu, Boundedness for commutators of Littlewood-Paley operators, Adv. in Math.(China), 32(2003), 473-480.
- [8] S.Z.Lu and D.C.Yang, The local versions of  $H^p(R^n)$  spaces at the origin, Studia. Math., 116(1995), 147-158.
- [9] S.Z.Lu and D.C.Yang, The decomposition of the weighted Herz spaces and its applications, Sci. in China(ser.A), 38(1995), 147-158.
- [10] S.Z.Lu and D.C.Yang, The weighted Herz type Hardy spaces and its applications, Sci. in China(ser.A), 38(1995), 662-673.
- [11] S.Z.Lu and D.C.Yang, The continuity of commutators on Herz-type space, Michigan Math. J., 44(1997), 255-281.
- [12] C.Pérez, Endpoint estimate for commutators of singular integral operators, J. Func. Anal., 128(1995), 163-185.
- [13] C.Pérez and R.Trujillo-Gonzalez , Sharp weighted estimates for multilinear commutators, J. London Math. Soc., 65(2002), 672-692.
- [14] E.M.Stein, Harmonic Analysis: real variable methods, orthogonality and oscillatory integrals, Princeton Univ. Press, Princeton NJ., 1993.
- [15] A.Torchinsky, The real variable methods in harmonic analysis, Pure and Applied Math., 123, Academic Press, New York, 1986.

DEPARTMENT OF MATHEMATICS, CHANGSHA UNIVERSITY OF SCIENCE AND TECHNOLOGY, CHANGSHA, 410076, P.R.OF CHINA

*E-mail address:* acrossbow@tom.com

*E-mail address:* lanzheliu@163.com

Received February 26, 2006