

GENERALIZED BOUNDARY VALUE PROBLEMS  
WITH LINEAR GROWTH

VALTER ŠEDA, Bratislava

(Received June 5, 1997)

*Abstract.* It is shown that for a given system of linearly independent linear continuous functionals  $l_i: C^{n-1} \rightarrow \mathbb{R}$ ,  $i = 1, \dots, n$ , the set of all  $n$ -th order linear differential equations such that the Green function for the corresponding generalized boundary value problem (BVP for short) exists is open and dense in the space of all  $n$ -th order linear differential equations. Then the generic properties of the set of all solutions to nonlinear BVP-s are investigated in the case when the nonlinearity in the differential equation has a linear majorant. A periodic BVP is also studied.

*Keywords:* generic properties, periodic boundary value problem

*MSC 1991:* 34B15

## 1. INTRODUCTION

B. Rudolf in [14] has shown that for a given system of linearly independent linear continuous functionals  $l_i: C^n([a, b], \mathbb{R}) \rightarrow \mathbb{R}$ ,  $i = 1, \dots, n$ , there exists a linear differential equation

$$(1) \quad (L(x) \equiv) x^{(n)} + \sum_{k=1}^n p_k(t)x^{(n-k)} = 0, \quad a \leq t \leq b$$

such that the BVP (1),

$$(2) \quad l_i(x) = 0, \quad i = 1, \dots, n$$

has only the trivial solution. This result also holds when the functionals  $l_i$  are given in the space  $C^{n-1}([a, b], \mathbb{R})$ . In this paper we will prove that the set  $S$  of all

differential equations (1) such that (1), (2) has only the trivial solution is open and dense in the space of all  $n$ -th order linear differential equations and we will derive some consequences of that result. Besides the classical existence theorems the generic properties of the set of all solutions to nonlinear BVP-s having a linear majorant will also be studied. The main tool for showing these properties will be a priori estimates like Leray-Schauder estimations. Finally, a special case will be investigated, namely the periodic BVP.

Throughout the paper we will assume that  $n \geq 1, -\infty < a < b < \infty, p_k \in C([a, b], \mathbb{R}), k = 1, \dots, n, l_i: C^{n-1}([a, b], \mathbb{R}) \rightarrow \mathbb{R}$  is a linear continuous functional,  $i = 1, \dots, n$  and the functionals  $l_i, i = 1, \dots, n$ , are linearly independent.

Let  $C^0 = (C([a, b], \mathbb{R}), \|\cdot\|_0)$  be a Banach space with the sup-norm  $\|\cdot\|_0$  and let the topology in  $C^l = C^l([a, b], \mathbb{R})$  be given by the norm  $\|\cdot\|_l$ , whereby

$$\|x\|_l = \max_{k=0, \dots, l} \|x^{(k)}\|_0, \quad l = 1, \dots, n.$$

Further let  $C_n = C^0 \times \dots \times C^0$  ( $n$  times) be the product space with the norm  $\|(x_1, \dots, x_n)\| = \sum_{k=1}^n \|x_k\|_0$ . Then  $C_n$  is a Banach space and the equation (1) can be represented by the  $n$ -tuple  $(p_1, \dots, p_n)$ .

## 2. REGULAR CASE

We will start with the following definition.

**Definition 1.** The BVP (1), (2) will be called regular if and only if it has only the trivial solution.

**Theorem 1.** Let a system of linearly independent linear continuous functionals  $l_i, i = 1, \dots, n$ , be given. Then the set  $S$  of all  $n$ -tuples  $(p_1, \dots, p_n) \in C_n$  such that the BVP (1), (2) is regular is nonempty, open and dense in the space  $C_n$ .

**Proof.** By the Rudolf theorem, [14],  $S \neq \emptyset$ . Suppose that there exists a sequence of nontrivial solutions  $y_m$  to the BVP-s

$$(1_m) \quad (L_m(x) \equiv) x^{(n)} + \sum_{k=1}^n p_{k,m}(t)x^{(n-k)} = 0, \quad a \leq t \leq b$$

$$(2) \quad l_i(x) = 0, \quad i = 1, \dots, n$$

where

$$(3) \quad \|(p_{1,m}, \dots, p_{n,m}) - (p_1, \dots, p_n)\| \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Here  $p_{k,m} \in C^0$ ,  $k = 1, \dots, n$ ,  $m = 1, 2, \dots$ .

The solutions  $y_m$  can be normalized by the condition

$$(4) \quad (y_m(a))^2 + (y'_m(a))^2 + \dots + (y_m^{(n-1)}(a))^2 = 1.$$

By Corollary 4.1, [5], (3) and (4) imply that there is a subsequence  $y_{m_l}$  of  $y_m$  and a solution  $y$  of (1) such that  $\|(y_{m_l}, \dots, y_{m_l}^{(n-1)}) - (y, \dots, y^{(n-1)})\| \rightarrow 0$  as  $l \rightarrow \infty$ . Hence  $y$  is a nontrivial solution of (1), (2) and thus  $C_n \setminus S$  is closed. This implies that  $S$  is open in  $C_n$ .

Now we prove that  $S$  is dense in the space  $C_n$ . Again by the Rudolf theorem, there is a differential equation

$$(5) \quad x^{(n)} + \sum_{k=1}^n r_k(t)x^{(n-k)} = 0$$

such that the BVP (5), (2) is regular. Denote by  $A: D(A) \subset C^{n-1} \rightarrow C$  the operator

$$Ax = \sum_{k=1}^n (p_k(t) - r_k(t))x^{(n-k)}$$

where

$$D(A) = \{x \in C^{n-1}: l_i(x) = 0, i = 1, \dots, n\}.$$

Then the linear operator  $L - A: D(L) \subset C^n \rightarrow C$  which is defined on

$$(6) \quad D(L) = \{x \in C^n: l_i(x) = 0, i = 1, \dots, n\}$$

is onto and one-to-one. By Lemma 4 ([18], p. 512), its inverse mapping  $(L - A)^{-1}: C \rightarrow C^n$  is continuous and as a mapping from  $C$  to  $C^{n-1}$  it is completely continuous. Since the equation  $L(x) = 0$  is equivalent to the equation

$$(7) \quad x = (L - A)^{-1}(-Ax)$$

and the operator  $K = -(L - A)^{-1} \circ A: D(A) \subset C^{n-1} \rightarrow C^{n-1}$  is completely continuous, either (7) has only the trivial solution or 1 is an eigenvalue of  $K$ . In the latter case there is an  $\varepsilon > 0$  such that the equation  $\lambda x = (L - A)^{-1} \circ (-Ax)$  as well as  $Lx = (1 - \frac{1}{\lambda})Ax$  has only the trivial solution for all  $\lambda \in (1 - \varepsilon, 1) \cup (1, 1 + \varepsilon)$ . This means that for these  $\lambda$  the BVP  $L(x) + (\frac{1}{\lambda} - 1)Ax = 0$ , (2) is regular.  $\square$

Consider now a sequence of generalized boundary conditions

$$(2_j) \quad l_i^{(j)}(x) = 0, \quad i = 1, \dots, n,$$

$j = 1, 2, \dots$ , where

$$l_i^{(j)}: C^{n-1} \rightarrow \mathbb{R}$$

is a linear continuous functional,  $i = 1, \dots, n$ ,  $j = 1, 2, \dots$  and for each  $j = 1, 2, \dots$  the functionals  $l_i^{(j)}$ ,  $i = 1, \dots, n$  are linearly independent.

Denote by  $S_j$  the set of all  $n$ -tuples  $(p_1, \dots, p_n) \in C_n$  such that the BVP (1), (2<sub>j</sub>) is regular. On the basis of the Baire theorem ([13], Theorem 2.2), Theorem 1 implies that the set  $\bigcap_{j=1}^{\infty} S_j$  is dense in  $C_n$  and hence the following corollary holds.

**Corollary 1.** *If a sequence of the boundary conditions (2<sub>j</sub>),  $j = 1, 2, \dots$  is given, then the set of all  $n$ -tuples  $(p_1, \dots, p_n) \in C_n$  for which the BVP-s (1), (2<sub>j</sub>),  $j = 1, 2, \dots$  are regular is dense in the space  $C_n$ .*

Let  $\{t_j\}_{j=0}^{\infty} \subset [a, b]$  be an injective sequence of points in  $[a, b]$ . A point  $t_j$  is conjugate to  $t_0$  for the equation

$$(8) \quad x'' + p_1(t)x' + p_2(t)x = 0$$

([8], p. 216) if and only if the BVP (8),

$$x(t_0) = 0, \quad x(t_j) = 0$$

is not regular. Hence, by Corollary 1, the set of all pairs  $(p_1, p_2)$  such that all conjugate points to  $t_0$  in  $[a, b]$  for the equation (8) (when they exist) are different from  $\{t_j\}_{j=1}^{\infty}$  is dense in  $C_2$ .

Now we introduce the notions which are well-known in the coincidence theory developed by J. Mawhin (see e.g. [7]). According to (1), the operator  $L: D(L) \subset C^{n-1} \rightarrow C^0$  is defined by

$$(9) \quad L(x) = x^{(n)} + \sum_{k=1}^n p_k(t)x^{(n-k)}, \quad x \in D(L)$$

where  $D(L)$  is determined by (6). By the Rudolf theorem  $L$  is a Fredholm mapping of index zero. Hence, there exist linear continuous projectors  $P: C^{n-1} \rightarrow C^{n-1}$ ,  $Q: C^0 \rightarrow C^0$  such that

$$R(P) = N(L), \quad N(Q) = R(L)$$

and

$$C^{n-1} = N(L) \oplus N(P), \quad C^0 = R(Q) \oplus R(L)$$

as topological direct sums. Further, the restriction  $L_P$  of  $L$  to  $D(L) \cap N(P)$  is one-to-one and onto  $R(L)$  so that its algebraic inverse  $K_P: R(L) \rightarrow D(L) \cap N(P)$  is well defined.

We shall distinguish two cases: Either (1), (2) is regular or not. When the BVP (1), (2) is regular, then  $P(x) \equiv 0$ ,  $x \in C^{n-1}$ ,  $Q(y) \equiv 0$ ,  $y \in C^0$  and  $C^{n-1} = N(P)$ ,  $C^0 = R(L)$ . In this case  $L_P = L$ ,  $K_P = L^{-1}$ . The operator  $L^{-1}: C^0 \rightarrow D(L) \subset C^{n-1}$  is constructed with help of the Green function  $G = G(t, s)$ ,  $a \leq t, s \leq b$  for the BVP (1), (2) given in [18].  $L^{-1}$  is defined in Lemma 1, ([18], p. 510) by the relation

$$(10) \quad L^{-1}(x)(t) = \int_a^b G(t, s)x(s) ds, \quad a \leq t \leq b, \quad x \in C^0.$$

Further, for each  $k \in \{1, \dots, n-1\}$  (if  $n \geq 2$ ) we have

$$(11) \quad (L^{-1}(x))^{(k)}(t) = \int_a^b \frac{\partial^k G(t, s)}{\partial t^k} x(s) ds, \quad a \leq t \leq b, \quad x \in C^0.$$

By Lemma 3 ([18], p. 512), the function  $\varphi(t) = \int_a^b |G(t, s)| ds$ ,  $a \leq t \leq b$ , is continuous on  $[a, b]$  and hence, by the result in [6], p. 187,

$$(12) \quad \|L^{-1}\| = \max_{a \leq t \leq b} \int_a^b |G(t, s)| ds.$$

A similar relation holds for the norm  $\|L_k^{-1}\|$  of the linear operator standing on the right-hand side of (11). Hence

$$(13) \quad \|L_k^{-1}\| = \max_{a \leq t \leq b} \int_a^b \left| \frac{\partial^k G(t, s)}{\partial t^k} \right| ds.$$

For brevity, in what follows we will write  $\|L_0^{-1}\|$  instead of  $\|L^{-1}\|$ .

Let  $f = f(t, x_1, \dots, x_{p+1}) \in C([a, b] \times \mathbb{R}^{p+1}, \mathbb{R})$ ,  $g \in C([a, b], \mathbb{R})$ , where  $0 \leq p \leq n-1$ . The following existence lemma holds for the nonlinear perturbation of the regular BVP.

**Lemma 1.** *If the BVP (1), (2) is regular and if there exist positive constants  $c_1, \dots, c_{p+1}, d$  such that*

$$(14) \quad \delta = \sum_{k=1}^{p+1} c_k \|L_{k-1}^{-1}\| < 1,$$

$$(15) \quad |f(t, x_1, \dots, x_{p+1})| \leq \sum_{k=1}^{p+1} c_k |x_k| + d, \quad a \leq t \leq b, \quad x_1, \dots, x_{p+1} \in \mathbb{R}$$

and

$$(16) \quad |g(t)| \leq M, \quad a \leq t \leq b,$$

then each possible solution  $x$  of the BVP (2),

$$(17) \quad L(x) = f(t, x, \dots, x^{(p)}) + g(t)$$

satisfies a priori estimates

$$(18) \quad \left( \min_{k=1, \dots, p+1} c_k \right) \|x\|_p \leq \sum_{k=1}^{p+1} c_k \|x^{(k-1)}\|_0 \leq \frac{\delta}{1-\delta} (d + M),$$

and for each  $g \in C^0$  there exists a solution of the BVP (17), (2).

**Proof.** By means of Lemmas 5 and 6 in [18], p. 516, the problem (17), (2) is equivalent to the fixed point problem for the operator

$$(19) \quad T(x)(t) = \int_a^b G(t, s) [f(s, x(s), \dots, x^{(p)}(s)) + g(s)] ds, \quad a \leq t \leq b, \quad x \in C^p.$$

This operator as a mapping from  $C^p$  to  $C^p$  is completely continuous. By the Leray-Schauder principle the existence of a fixed point of  $T$  will be proved if it is shown that the set of all possible solutions of the family of equations

$$(20) \quad x = \lambda T(x), \quad 0 \leq \lambda \leq 1$$

is a priori bounded (in the norm  $\|\cdot\|_p$ ) independently of  $\lambda$ .

Let  $\lambda \in [0, 1]$ ,  $x$  be a possible solution of (20) and let  $k \in \{1, \dots, p+1\}$ . Then

$$x^{(k-1)}(t) = \lambda \int_a^b \frac{\partial^{k-1} G(t, s)}{\partial t^{k-1}} [f(s, x(s), \dots, x^{(p)}(s)) + g(s)] ds, \quad a \leq t \leq b,$$

and by (12), (13), (15), (16) we have

$$\|x^{(k-1)}\|_0 \leq \|L_{(k-1)}^{-1}\| \left( \sum_{j=1}^{p+1} c_j \|x^{(j-1)}\|_0 + d + M \right).$$

Hence

$$\sum_{k=1}^{p+1} c_k \|x^{(k-1)}\|_0 \leq \delta \left( \sum_{k=1}^{p+1} c_k \|x^{(k-1)}\|_0 + d + M \right)$$

and thus (18) is true. □

**Example.** Let  $0 < \eta < 1$ . Consider the Green function for the problem

$$(21) \quad \begin{aligned} x'' &= 0, \\ x(0) &= 0, \quad x(1) - x(\eta) = 0. \end{aligned}$$

Since

$$x(t) = t(1 - \eta)^{-1} \left[ (\eta - 1) \int_0^\eta g(s) \, ds + \int_\eta^1 [sg(s) - g(s)] \, ds \right] + \int_0^t (t - s)g(s) \, ds,$$

$0 \leq t \leq 1$ , is the unique solution of the BVP  $x'' = g(t)$ , (21), the Green function  $G = G(t, s)$  for that problem is determined by

$$G(t, s) = G_1(t, s) + K(t, s) \quad 0 \leq t, s \leq 1$$

where

$$G_1(t, s) = \begin{cases} -t & 0 \leq s \leq \eta \\ -\frac{1-s}{1-\eta}t & \eta \leq s \leq 1 \end{cases},$$

$$K(t, s) = \begin{cases} t - s & 0 \leq s \leq t \leq 1 \\ 0 & 0 \leq t < s \leq 1 \end{cases}$$

and hence  $G(t, s) < 0$  for  $0 < s < 1$ ,  $0 < t \leq 1$ ,  $G(t, 0) = G(t, 1) = 0$ ,  $0 \leq t \leq 1$ ,  $G(0, s) = 0$ ,  $0 \leq s \leq 1$ .

Therefore the solution  $x_0(t) = -\frac{1}{2}(1 + \eta)t + \frac{t^2}{2}$ ,  $0 \leq t \leq 1$  of the problem  $x'' = 1$ , (21) satisfies  $|x_0(t)| = \int_0^1 |G(t, s)| \, ds$ ,  $0 \leq t \leq 1$  and hence

$$(22) \quad \|L_0^{-1}\| = \max_{0 \leq t \leq 1} |x_0(t)| = \frac{1}{8}(1 + \eta)^2.$$

Since  $\frac{\partial G(t, s)}{\partial t} \leq 0$  for  $0 \leq t \leq \eta$ ,  $0 \leq s \leq 1$ , we have

$$\int_0^1 \left| \frac{\partial G(t, s)}{\partial t} \right| \, ds = |x_0'(t)| \leq |x_0'(0)| = \frac{1}{2}(1 + \eta)$$

in  $[0, \eta]$ . A direct calculation gives

$$\int_0^1 \left| \frac{\partial G(t, s)}{\partial t} \right| \, ds = \frac{1}{2(1 - \eta)} [2t^2 + \eta^2 + 1 - 2t(1 + \eta)] \leq \frac{1}{2}(1 - \eta), \quad \eta \leq t \leq 1.$$

Thus

$$(23) \quad \|L_1^{-1}\| = \frac{1}{2}(1 + \eta).$$

By means of (22), (23), Lemma 1 gives an existence statement for the nonlinear BVP (21),

$$(24) \quad x'' = f(t, x, x'),$$

which completes the statements in [4].

**Remark 1.** Estimations of  $\|L_k^{-1}\|$  for a BVP of the third and of the fourth order can be found in [16], Lemma 3 and in [10], Lemma 4, respectively.

The existence and uniqueness of a solution to (17), (2) is guaranteed by

**Theorem 2.** *If the BVP (1), (2) is regular and if there exist constants  $c_1, \dots, c_{p+1}$  such that (14) is true and*

$$(25) \quad \begin{aligned} |f(t, x_1, \dots, x_{p+1}) - f(t, y_1, \dots, y_{p+1})| &\leq \sum_{k=1}^{p+1} c_k |x_k - y_k|, \\ a \leq t \leq b, \quad x_1, \dots, x_{p+1}, y_1, \dots, y_{p+1} &\in \mathbb{R}, \end{aligned}$$

then the BVP (17), (2) has a unique solution.

**Proof.** Consider the space  $C_1^p = C^p([a, b], \mathbb{R})$  provided with the norm  $\|x\|_{p,1} = \sum_{k=1}^{p+1} c_k \|x^{(k-1)}\|_0$ . With respect to inequalities

$$\left( \min_{k=1, \dots, p+1} c_k \right) \|x\|_p \leq \|x\|_{p,1} \leq \left( \sum_{k=1}^{p+1} c_k \right) \|x\|_p$$

the norms  $\|\cdot\|_p, \|\cdot\|_{p,1}$  are equivalent and hence  $C_1^p$  is a Banach space. Consider the operator  $T: C_1^p \rightarrow C_1^p$  given by (19). In view of (25), we have

$$\|T^{(k-1)}(x) - T^{(k-1)}(y)\|_0 \leq \|L_{k-1}^{-1}\| \sum_{k=1}^{p+1} c_k \|x^{(k-1)} - y^{(k-1)}\|_0$$

for each  $k = 1, \dots, p+1$  and  $x, y \in C_1^p$ . Thus

$$\|T(x) - T(y)\|_{p,1} \leq \delta \|x - y\|_{p,1},$$

which means that  $T$  is a strict contraction on  $C_1^p$ . The result follows by the Banach fixed point theorem.  $\square$

**Remark 2.** In the case of homogeneous boundary conditions, Theorem 2 generalizes Theorem 1.1.1 in [1].

Corollary 1 and Lemma 1 imply

**Theorem 3.** *If a sequence of the boundary conditions  $(2_j)$ ,  $j = 1, 2, \dots$ , is given and the function  $f$  satisfies the condition*

$$\lim_{|x_1| + \dots + |x_{p+1}| \rightarrow \infty} \frac{|f(t, x_1, \dots, x_{p+1})|}{|x_1| + \dots + |x_{p+1}|} = 0$$

uniformly in  $t \in [a, b]$ , then there exists a sequence of  $n$ -tuples  $(p_{1,m}, \dots, p_{n,m}) \in C_n$ ,  $m = 1, 2, \dots$ , such that (3) is fulfilled and the BVP (2<sub>j</sub>),

$$(17_m) \quad L_m(x) = f(t, x, \dots, x^{(p)}) + g(t)$$

has a solution  $x_{m,j}$  for each  $m = 1, 2, \dots$  and each  $j = 1, 2, \dots$ .

Another sufficient condition for the existence of a solution to (17), (2) is given in the next lemma.

**Lemma 2.** *Let there exist a sequence of  $n$ -tuples  $(p_{1,m}, \dots, p_{n,m}) \in C_n$ ,  $m = 1, 2, \dots$  such that (3) is satisfied and the BVP (17<sub>m</sub>), (2) has a solution  $x_m$  for each  $m = 1, 2, \dots$ . Let the sequence  $\{x_m\}$  be bounded in  $C^{n-1}$ . Then there exists a subsequence  $\{x_{m_l}\}$  of  $\{x_m\}$  and a solution  $x$  of the problem (17), (2) such that  $x_{m_l} \rightarrow x$  as  $l \rightarrow \infty$  in  $C^n$ .*

*Proof.* By (17<sub>m</sub>), the sequence  $\{x_m\}$  is even bounded in  $C^n$ . Hence, by the Ascoli theorem, there exists a subsequence  $\{x_{m_l}\}$  of the sequence  $\{x_m\}$  and a function  $x \in C^{n-1}$  such that  $x_{m_l} \rightarrow x$  as  $l \rightarrow \infty$  in  $C^{n-1}$ . In view of (3) and (17<sub>m</sub>) the sequence  $x_{m_l}^{(n)}$  is uniformly convergent on  $[a, b]$  and hence  $x \in C^n$  and  $x_{m_l} \rightarrow x$  as  $l \rightarrow \infty$  in  $C^n$ . Thus  $x$  is a solution of (17), (2).  $\square$

Now we prove generic properties of the set of all solutions to (17), (2). To that aim we need the definition of the range of bifurcation  $R_b$  of the BVP (17), (2) (see Definition 4.1 in [20], p.29). For the sake of completeness we will give it here.

First we introduce the Banach space  $X_0 = (D(L), \|\cdot\|_n)$  ([20], p.28). Then the set  $R_b$  of all  $g \in C^0$  with the property that there is a solution  $x$  of the BVP (17), (2) and a sequence  $g_k \rightarrow g$  as  $k \rightarrow \infty$  such that the BVP (17), (2) for  $g = g_k$  has at least two different solutions  $x_k, z_k$  for each  $k$  and  $x_k \rightarrow x, z_k \rightarrow x$  in  $X_0$  for  $k \rightarrow \infty$  is called the range of bifurcation of the BVP (17), (2).

By Lemma 1, Theorems 4.1 and 4.2 in [20], pp.31–32, we get the following theorem.

**Theorem 4.** *If the problem (1), (2) is regular and there exist positive constants  $c_1, \dots, c_{p+1}, d$  such that (14), (15) are true, then the following statements hold:*

1. *For each  $g \in C^0$  the set  $S_g$  of all solutions of the BVP (17), (2) is nonempty and compact.*
2. *If  $C^0 \setminus R_b \neq \emptyset$ , then each component of that set is nonempty, open and hence a region. The number  $n_g$  of solutions of the BVP (17), (2) is finite and constant on each component of the set  $C^0 \setminus R_b$ .*
3. *If  $R_b = \emptyset$ , then the problem (17), (2) has a unique solution for each  $g \in C^0$  and this solution continuously depends on  $g$  as a mapping from  $C^0$  onto  $X_0$ .*

4. If  $\frac{\partial f}{\partial x_i} \in C([a, b] \times \mathbb{R}^{p+1}, \mathbb{R})$ ,  $i = 1, \dots, p+1$ , then the open set  $C^0 \setminus R_b$  is dense in  $C_0$  and hence,  $R_b$  is nowhere dense in  $C^0$ .

**Proof.** Since (1), (2) is regular, the operator  $L$  given by (9) satisfies the assumption (H.1) of Theorem 4.1. Since  $f$  is continuous, (H.2) is satisfied and in the case that  $\frac{\partial f}{\partial x_i}$  are continuous, (H.4) is fulfilled. By the apriori estimates (18) it follows that also (H.3) holds. Then Lemma 1 and Theorems 4.1, 4.2 imply the statements.  $\square$

**Remark 3.** We see that under the assumptions of Theorem 4, uniqueness of the BVP (17), (2) implies correctness of that BVP, that is the existence, uniqueness and continuous dependence of the solution  $x$  of the BVP (17), (2) on  $g$ .

### 3. NON REGULAR CASE

This case is more complicated as the previous one. Now we apply the results of [17] and [20]. Denote by  $F$  the Nemitskij operator  $F: C^p \rightarrow C^0$  which is defined by

$$(26) \quad F(x) = f \circ x, \quad x \in C^p$$

The properties of the operator  $L$  and  $F$  are given by the following lemma.

**Lemma 3.** *The following statements hold:*

- (i) *For each integer  $k$ ,  $0 \leq k \leq n-1$ , the operator  $L: D(L) \subset C^k \rightarrow C^0$  is a linear Fredholm operator of index zero.*
- (ii) *If there exists a continuous linear operator  $A: D(L) \subset C^r \rightarrow C^0$  with  $0 \leq r \leq n-1$  such that  $L - A: D(L) \subset C^r \rightarrow C^0$  is one-to-one, then  $L - A$  is onto, the inverse operator  $(L - A)^{-1}: C^0 \rightarrow D(L) \subset C^{n-1}$  is completely continuous and  $(L - A)^{-1}$  as a mapping from  $C^0$  into  $C^n$  is continuous.*
- (iii) *The operator  $K_P: R(L) \subset C^0 \rightarrow D(L) \cap N(P) \subset C^{n-1}$  is completely continuous.*
- (iv)  *$F + g: C^p \rightarrow C^0$  is continuous and maps bounded sets in  $C^p$  into bounded sets in  $C^0$ .*
- (v)  *$F + g: C^p \rightarrow C^0$  is  $L$ -completely continuous.*

**Proof.** (i) By the Rudolf theorem, there exists a continuous linear operator  $A: C^r \rightarrow C^0$  with  $0 \leq r \leq n-1$  such that  $L - A: D(L) \subset C^r \rightarrow C^0$  is one-to-one. Then by Lemmas 1 and 4, [18], the operator  $L - A: X_0 \rightarrow C^0$  is a homeomorphism

of  $X_0$  onto  $C^0$  and  $A: X_0 \rightarrow C^0$  is a linear completely continuous operator. Nikoľskij theorem ([20], p.21) implies that  $L: X_0 \rightarrow C^0$  is a linear Fredholm operator of index zero. The same is true about  $L: D(L) \subset C^k \rightarrow C^0$ .

(ii), (iii) If  $L - A: D(L) \subset C^r \rightarrow C^0$  is one-to-one, then it is onto, and by Lemma 4, [18],  $(L - A)^{-1}: C^0 \rightarrow C^{n-1}$  is completely continuous and  $(L - A)^{-1}: C^0 \rightarrow C^n$  is continuous and hence, by Remark 1 and Lemma 1 in [17], p.555, the operators  $L: D(L) \subset C^r \rightarrow C^0$  and  $L_P: D(L) \cap N(P) \subset C^r \rightarrow C^{n-1}$  are closed and  $K_P: R(L) \subset C^0 \rightarrow C^r$  is completely continuous. Since  $A$  is continuous also as a mapping from  $C^{n-1}$  to  $C^0$ ,  $K_P: R(L) \subset C^0 \rightarrow C^{n-1}$  is completely continuous, too.

(iv) The statement follows from the continuity of the functions  $f$  and  $g$ .

(v) Let  $E \subset C^p$  be a bounded set. Then by (iii) and (iv) the mappings  $Q \circ (F + g)$ ,  $K_P \circ (I - Q) \circ (F + g)$  are continuous on  $E$  and the sets  $Q \circ (F + g)(E)$  and  $K_P \circ (I - Q) \circ (F + g)(E)$  are relatively compact in  $C^0$  and in  $C^p$ , respectively. This implies the statement.  $\square$

**Remark 4.** By (iii), the statements (iv), (v) also hold for the restriction  $F + g: C^k \rightarrow C^0$ ,  $p < k \leq n - 1$ .

On the basis of Theorem 3 ([17], p.561), the following lemma holds which is analogous to Lemma 1.

**Lemma 4.** *Suppose that the BVP (1), (2) is not regular and the following assumptions hold:*

- (a)  $R(L) \cap N(L) = \{0\}$ ;
- (b) *there exists a continuous linear operator  $A: C^r \rightarrow C^0$  with  $0 \leq r \leq n - 1$  such that  $L - A: D(L) \subset C^r \rightarrow C^0$  is one-to-one;*
- (c) *there exist constants  $c_1, \dots, c_p, d > 0$  such that (15) is true.*  
Let  $d_1 > 0$  and let  $s = \max(p, r)$ .
- (d) *The constant  $c = \sum_{k=1}^{p+1} c_k$  satisfies*

$$(27) \quad c < \frac{1}{\|K_P\|} \frac{d_1}{1 + d_1}$$

where  $\|K_P\|$  is the norm of  $K_P: R(L) \subset C^0 \rightarrow C^s$ .

Let  $\varepsilon = \pm 1$ .

- (e) *There exists an  $R_1 > 0$  with the following property:*

$$\varepsilon F(\bar{x} + \tilde{x}) + \varepsilon g + k\bar{x} \notin R(L)$$

for all  $x = \bar{x} + \tilde{x} \in D(L)$ ,  $\bar{x} \in N(L)$ ,  $\tilde{x} \in N(P)$ ,  $k \in \mathbb{R}$  such that

$$\|\bar{x}\|_s \geq R_1, \quad \|\tilde{x}\|_s \leq d_1 \|\bar{x}\|_s \quad \text{and} \quad k > 0.$$

Then the problem (17), (2) has a solution. Moreover, if  $g$  satisfies (16) and (f) there exists an  $R_3 \geq R_2$  where

$$R_2 = \frac{\|K_P\|(d+M)}{d_1(1 - \|K_P\|c) - \|K_P\|c}$$

such that

$$F(\bar{x} + \tilde{x}) + g \notin R(L)$$

for all  $x = \bar{x} + \tilde{x} \in D(L)$ ,  $\bar{x} \in N(L)$ ,  $\tilde{x} \in N(P)$  satisfying

$$\|\bar{x}\|_s \geq R_3, \quad \|\tilde{x}\|_s \leq d_1\|\bar{x}\|_s,$$

then any solution  $x$  of the BVP (17), (2) satisfies the inequalities

$$(28) \quad \|\bar{x}\|_s < R_3$$

and

$$(29) \quad \|\tilde{x}\|_s \leq \frac{\|K_P\|c}{1 - \|K_P\|c}\|\bar{x}\|_s + \frac{\|K_P\|(d+M)}{1 - \|K_P\|c}.$$

*Proof.* On the basis of Lemma 3, the operator  $L: D(L) \subset C^s \rightarrow C^0$  satisfies the assumptions (L<sub>1</sub>), (L<sub>2</sub>) and (L<sub>3</sub>) in [17], pp. 554–555 with  $X = C^s$ ,  $Z = C^0$ , and by the assumption (a) of this lemma (L<sub>4</sub>) is satisfied, too. Lemma 3 with Remark 4 also implies that  $F + g: C^s \rightarrow C^0$  is continuous, maps bounded sets in  $C^s$  into bounded sets in  $C^0$  and is  $L$ -completely continuous. By virtue of (15), the assumption (F<sub>5</sub>) of Theorem 3 in [17], p. 561, is satisfied. Then (27) together with the assumption (e) imply that also (F<sub>6</sub>) is fulfilled. By the just mentioned Theorem 3 the existence statement follows.

Now we prove the a priori estimates (28) and (29). By Lemma 1 ([17], p. 555), and by (15), (16) any solution  $x = \bar{x} + \tilde{x}$  of (17), (2),  $\bar{x} \in N(L)$ ,  $\tilde{x} \in N(P)$ , satisfies the inequalities

$$\|\tilde{x}\|_s \leq \|K_P\|\|L(x)\|_0 \leq \|K_P\|\|F(x) + g\|_0 \leq \|K_P\|c\|\tilde{x}\|_s + \|K_P\|c\|\bar{x}\|_s + \|K_P\|(d+M)$$

and hence (29) is true. Since (27) is equivalent to  $\|K_P\|c/(1 - \|K_P\|c) < d_1$ , the right-hand side of (29) is less than or equal to  $d_1\|\bar{x}\|_s$  if and only if

$$\|\bar{x}\|_s \geq R_2.$$

Hence for  $\|\bar{x}\|_s \geq R_2$  we have

$$(30) \quad \|\tilde{x}\|_s \leq d_1\|\bar{x}\|_s.$$

By the assumption (f) the solution  $x$  of  $L(x) = F(x) + g$  cannot satisfy  $\|\bar{x}\|_s \geq R_3$ , (30) and thus (28) and (29) are true.  $\square$

By virtue of Lemma 4 the proof of the following theorem is similar to that of Theorem 4.

**Theorem 5.** *If the problem (1), (2) is not regular, the assumptions (a)–(d) of Lemma 4 hold, further for each  $g \in C^0$  the assumption (e) of that lemma is satisfied and for each  $M > 0$  and each  $g \in C^0$  satisfying (16) the assumption (f) of Lemma 4 holds, then all statements of Theorem 4 hold.*

Consider the BVP (2),

$$(17') \quad L(x) = f(t, x) + g(t) + h(t, x, x', \dots, x^{(p)})$$

where  $f \in C([a, b] \times \mathbb{R}, \mathbb{R})$ ,  $g \in C([a, b])$ ,  $h \in C([a, b] \times \mathbb{R}^{p+1}, \mathbb{R})$ ,  $0 \leq p \leq n - 1$ . We show that under simple assumptions on  $N(L)$ ,  $R(L)$  Theorem 5 implies the following theorem.

**Theorem 6.** *Assume that the following conditions are satisfied:*

(i)

$$\begin{aligned} N(L) &= \{x \in D(L) : x \text{ is a constant on } [a, b]\}, \\ R(L) &= \{y \in C^0 : \int_a^b y(x) dx = 0\}; \end{aligned}$$

(ii) *there exists a continuous linear operator  $A: C^r \rightarrow C^0$  with  $0 \leq r \leq n - 1$  such that  $L - A: D(L) \subset C^r \rightarrow C^0$  is one-to-one;*

(iii) *there exist constants  $c, d, \delta > 0$  such that*

$$|f(t, x_1)| \leq c|x_1| + d, \quad |h(t, x_1, \dots, x_{p+1})| \leq \delta \text{ for } a \leq t \leq b, x_1, \dots, x_{p+1} \in \mathbb{R};$$

(iv)

$$(31) \quad 2c\|K_P\| < 1$$

where  $\|K_P\|$  is the norm of  $K_P: R(L) \subset C^0 \rightarrow C^s$ ,  $s = \max(p, r)$ ;

(v) for  $\varepsilon = 1$  or  $\varepsilon = -1$

$$(32) \quad \lim_{x \rightarrow \infty} \varepsilon f(t, x) = \infty, \quad \lim_{x \rightarrow -\infty} \varepsilon f(t, x) = -\infty \text{ uniformly in } t \in [a, b].$$

Then all statements of Theorem 4 hold for the BVP (17'), (2).

**Proof.** We shall show that all assumptions of Theorem 5 concerning the BVP (17'), (2) hold and by this theorem Theorem 6 follows. The assumptions (i)–(iii)

imply that the problem (1), (2) is not regular and the assumptions (a)–(c) of Lemma 4 are satisfied where instead of  $d$  we have  $d + \delta$ . In view of (31) there exists a  $d_1$  such that

$$(33) \quad 0 < d_1 < 1$$

and (27) is true. Hence the assumption (d) of Lemma 4 is also fulfilled.

Now with respect to (i) we can choose the projector  $P(x) = \frac{1}{b-a} \int_a^b x(t) dt$ ,  $x \in C^s$ . Then  $P(x) = \bar{x}$  for each  $x \in D(L)$  and  $x(t) = \bar{x} + \tilde{x}(t)$ ,  $a \leq t \leq b$ . On the basis of (33), the condition  $\|\tilde{x}\|_s \leq d_1 \|\bar{x}\|_s$  implies that  $|\tilde{x}(t)| \leq d_1 |\bar{x}|$ ,  $a \leq t \leq b$  and hence

$$(34) \quad x(t) \geq (1-d_1)\bar{x} \text{ for } \bar{x} > 0 \text{ and } x(t) \leq (1-d_1)\bar{x} \text{ for } \bar{x} < 0, \quad a \leq t \leq b, \quad x \in D(L).$$

Suppose further that  $g$  satisfies (16). Since  $F: C^p \rightarrow C^0$  is now determined by the relation

$$(26') \quad F(x) = f \circ x + h \circ x$$

and  $\varepsilon F(\bar{x} + \tilde{x}) + \varepsilon g + k\bar{x} \in R(L)$  if and only if

$$\int_a^b \varepsilon [f(t, \bar{x} + \tilde{x}(t)) + g(t) + h(t, \bar{x} + \tilde{x}(t), \dots, \tilde{x}^{(p)}(t))] dt + k\bar{x}(b-a) = 0,$$

on the basis of (34) we get that both conditions (e) and (f) of Lemma 4 will be satisfied if for all sufficiently great  $|\bar{x}|$  and  $k \geq 0$  we have

$$\text{sign} [\varepsilon f(t, \bar{x} + \tilde{x}(t)) + \varepsilon g + \varepsilon h(t, x(t), \dots, x^{(p)}(t)) + k\bar{x}] = \text{sign } \bar{x}.$$

This follows by the boundedness of  $g, h$  and (32). The proof is complete.  $\square$

If  $L(x) = x^{(n)}$  and the conditions (2) are of the form

$$(35) \quad x^{(i)}(a) - x^{(i)}(b) = 0, \quad i = 0, \dots, n-1,$$

then the condition (i) is satisfied and  $A = cI: C^0 \rightarrow C^0$  where  $c \neq 0$  is sufficiently small and  $I$  is the identity in  $C^0$ . Hence  $r = 0$  and thus  $s = p$  in conditions (ii) and (iv), respectively.

**Corollary 2.** *If the conditions (iii), (iv) (with  $s = p$ ) and (v) of Theorem 6 are satisfied, then all statements of Theorem 4 hold for the BVP (35),*

$$(36) \quad x^{(n)} = f(t, x) + g(t) + h(t, x, \dots, x^{(p)}).$$

**Remark 5.** The operator  $K_P$  for certain periodic BVP-s is constructed in [9], [11].

Consider a special case of (36), (35), namely the BVP (35),

$$(37) \quad x^{(n)} = f(x) + g(t) + h(t, x)$$

where  $f \in C(\mathbb{R}, \mathbb{R})$ . Similarly as in [2], the results will depend on the fact whether  $n$  is odd or even. We will denote the scalar product and the norm in  $L^2([a, b], \mathbb{R})$  by  $(\cdot, \cdot)$  and  $\|\cdot\|_{L^2}$ , respectively.

**Lemma 5.** *Suppose that  $n = 2m + 1$ ,  $n \geq 3$ ,  $\varepsilon = 1$  or  $\varepsilon = -1$ ,  $f$  satisfies the condition*

$$(32') \quad \lim_{x \rightarrow \infty} \varepsilon f(x) = \infty, \quad \lim_{x \rightarrow -\infty} \varepsilon f(x) = -\infty,$$

$g$  fulfils (16) and  $h$  satisfies

$$(38) \quad |h(t, x_1)| \leq \delta \text{ for } a \leq t \leq b, \quad x_1 \in \mathbb{R}$$

with a  $\delta > 0$ . Then the following statements hold:

- (1) *There exists a constant  $R > 0$  such that each possible solution  $x$  of the BVP (37), (35) where  $x(t) = \bar{x} + \tilde{x}(t)$ ,  $a \leq t \leq b$ ,  $\bar{x} = \frac{1}{b-a} \int_a^b x(t) dt$ , satisfies the inequalities*

$$(39) \quad |\bar{x}| \leq R,$$

$$(40) \quad \|\tilde{x}\|_0 \leq \frac{1}{3} \left( \frac{b-a}{2\pi} \right)^{n-2} \left( \frac{b-a}{2} \right)^2 (M + \delta).$$

- (2) *For each  $c_1 > 0$  sufficiently small there exists an  $R_1 > 0$  such that all possible solutions  $x(t) = \bar{x} + \tilde{x}(t)$ ,  $a \leq t \leq b$ , of (35),*

$$(41) \quad x^{(n)} - (1 - \mu)\varepsilon c_1 x = \mu[f(x) + h(t, x)], \quad 0 < \mu < 1$$

*satisfy the inequalities*

$$(39') \quad |\bar{x}| \leq R_1$$

$$(40') \quad \|\tilde{x}\|_0 \leq \frac{1}{3} \left( \frac{b-a}{2\pi} \right)^{n-2} \left( \frac{b-a}{2} \right)^2 \delta.$$

**P r o o f.** 1. If  $x$  is a possible solution of (37), (35), then, similarly as in Lemma 1, [2], we get

$$(-1)^{\frac{n-1}{2}} \|x^{(\frac{n+1}{2})}\|_{L^2}^2 = (x^{(n)}, x') = (f(x(\cdot)), x') + (g, x') + (h(\cdot, x(\cdot)), x')$$

and in view of (35), (16), (38) we have

$$(42) \quad \|x^{(\frac{n+1}{2})}\|_{L^2}^2 \leq (M + \delta)(b - a)\|x'\|_0.$$

By Sobolev and Wirtinger inequalities ([12], pp. 216–217),

$$(43) \quad \|x'\|_0 \leq 3^{-1/2} \frac{(b-a)^{1/2}}{2} \left(\frac{b-a}{2\pi}\right)^{\frac{n-3}{2}} \|x^{(\frac{n+1}{2})}\|_{L^2}.$$

(42) and (43) imply that

$$\|x^{(\frac{n+1}{2})}\|_{L^2} \leq \frac{b-a}{2} \left(\frac{b-a}{3}\right)^{1/2} \left(\frac{b-a}{2\pi}\right)^{\frac{n-3}{2}} (M + \delta)$$

and thus

$$\|x'\|_{L^2} \leq \frac{b-a}{2} \left(\frac{b-a}{3}\right)^{1/2} \left(\frac{b-a}{2\pi}\right)^{n-2} (M + \delta),$$

which implies (40).

If (39) were not true, there would exist solutions  $x_k(t) = \bar{x}_k + \tilde{x}_k(t)$ ,  $a \leq t \leq b$ ,  $k = 1, 2, \dots$  of (37), (35) such that either  $\lim_{k \rightarrow \infty} \bar{x}_k = \infty$  or  $\lim_{k \rightarrow \infty} \bar{x}_k = -\infty$ . Only the first case will be considered. Then in view of (16), (38), (40) and the first condition in (38')  $f(x_k(t)) + g(t) + h(t, x_k(t))$  would be of constant sign for all sufficiently great  $k$  and hence,  $x_k^{(n-1)}(b) - x_k^{(n-1)}(a) \neq 0$ . This contradiction with (35) proves (39). We remark that the contradiction is also attained in the case when  $x_k$  are solutions of  $x^{(n)} = f(x) + g_k(t) + h(t, x, \dots, x^{(p)})$  and all  $g_k$  satisfy  $|g_k(t)| \leq M$ ,  $a \leq t \leq b$ .

2. If we start with (41) instead of (37) and proceed in the same way as above, we come to the inequality

$$\|x^{(\frac{n+1}{2})}\|_{L^2}^2 \leq \delta(b-a)\|x'\|_0$$

which now replaces (42). This inequality leads to (40').

If there existed solutions  $x_k(t) = \bar{x}_k + \tilde{x}_k(t)$ ,  $a \leq t \leq b$ ,  $k = 1, 2, \dots$  of  $x^{(n)} = (1 - \mu_k)\varepsilon c_1 x + \mu_k[f(x) + h(t, x)]$  with  $\lim_{k \rightarrow \infty} \bar{x}_k = \infty$ , then with respect to (38), (40') and the first condition in (32'), the functions  $(1 - \mu_k)\varepsilon c_1 x_k(t) + \mu_k[f(x_k(t)) + h(t, x_k(t))]$  would be of constant sign for all sufficiently great  $k$ . This again contradicts (35) and thus (39') is true. Similarly we proceed in the case when  $\lim_{k \rightarrow \infty} \bar{x}_k = -\infty$ .  $\square$

**Lemma 6.** Suppose that  $n = 2m$ ,  $n \geq 2$ ,  $\varepsilon = 1$  or  $\varepsilon = -1$ ,  $f$  satisfies (32') and the following condition:

There exists  $\beta$ ,  $0 \leq \beta < \left(\frac{2\pi}{b-a}\right)^n$ , such that

$$(44) \quad (-1)^{n/2}(f(v) - f(w))(v - w) \leq \beta(v - w)^2, \quad v, w \in \mathbb{R},$$

$g$  fulfils (16) and  $h$  satisfies (38). Then the following statements hold:

- (1) There exists a constant  $R > 0$  such that each possible solution  $x$  of the BVP (37), (35) where  $x(t) = \bar{x} + \tilde{x}(t)$ ,  $a \leq t \leq b$ ,  $\bar{x} = \frac{1}{b-a} \int_a^b x(t) dt$ , satisfies the inequalities

$$(45) \quad |\bar{x}| \leq R,$$

$$(46) \quad \|\tilde{x}\|_0 \leq \left(1 - \beta \left(\frac{b-a}{2\pi}\right)^n\right)^{-1} \frac{1}{3} \left(\frac{b-a}{2}\right)^2 \left(\frac{b-a}{2\pi}\right)^{n-2} (M + \delta).$$

- (2) For each  $c_1$ ,  $0 < c_1 < \beta$ ,  $c_1$  sufficiently small, there exists an  $R_1 > 0$  such that all possible solutions  $x(t) = \bar{x} + \tilde{x}(t)$ ,  $a \leq t \leq b$  of (41), (35) satisfy the inequalities

$$(45') \quad |\bar{x}| \leq R_1,$$

$$(46') \quad \|\tilde{x}\|_0 \leq \left(1 - \beta \left(\frac{b-a}{2\pi}\right)^n\right)^{-1} \frac{1}{3} \left(\frac{b-a}{2}\right)^2 \left(\frac{b-a}{2\pi}\right)^{n-2} \delta.$$

**Proof.** 1. If  $x$  is a possible solution of the BVP (37), (35), then, similarly as in the proof of Lemma 2, [2], we get that

$$\begin{aligned} \|x^{(\frac{n}{2})}\|_{L^2}^2 &= (-1)^{\frac{n}{2}}(x^{(n)}, \tilde{x}) \\ &= (-1)^{\frac{n}{2}}(f(x(\cdot)) - f(\bar{x}), x - \bar{x}) + (-1)^{\frac{n}{2}}(g, \tilde{x}) + (-1)^{\frac{n}{2}}(h(\cdot, x(\cdot)), \tilde{x}). \end{aligned}$$

Then by (44), (16), (38)

$$(47) \quad \|x^{(\frac{n}{2})}\|_{L^2}^2 \leq \beta \|\tilde{x}\|_{L^2}^2 + (b-a)(M + \delta) \|\tilde{x}\|_0.$$

Again Sobolev and Wirtinger inequalities imply that

$$(48) \quad \|\tilde{x}\|_{L^2} \leq \left(\frac{b-a}{2\pi}\right)^{\frac{n}{2}} \|x^{(\frac{n}{2})}\|_{L^2},$$

$$(49) \quad \|\tilde{x}\|_0 \leq \frac{1}{2} \left(\frac{b-a}{3}\right)^{\frac{1}{2}} \left(\frac{b-a}{2\pi}\right)^{\frac{n-2}{2}} \|x^{(\frac{n}{2})}\|_{L^2}.$$

From (47), (48) and (49) we get

$$\|x^{(\frac{n}{2})}\|_{L^2} \leq \left(1 - \beta \left(\frac{b-a}{2\pi}\right)^n\right)^{-1} \frac{b-a}{2} \left(\frac{b-a}{3}\right)^{\frac{1}{2}} \left(\frac{b-a}{2\pi}\right)^{\frac{n-2}{2}} (M + \delta)$$

and further

$$\|x'\|_{L^2} \leq \left(1 - \beta \left(\frac{b-a}{2\pi}\right)^n\right)^{-1} \frac{b-a}{2} \left(\frac{b-a}{3}\right)^{\frac{1}{2}} \left(\frac{b-a}{2\pi}\right)^{n-2} (M + \delta),$$

which implies (46).

If (45) were not true, then similarly as in the proof of statement 1 in Lemma 5, the existence of solutions  $x_k(t) = \bar{x}_k + \tilde{x}_k(t)$ ,  $a \leq t \leq b$ ,  $k = 1, 2, \dots$  of (37), (35) with the property  $\lim_{k \rightarrow \infty} \bar{x}_k = \infty$  or  $\lim_{k \rightarrow \infty} \bar{x}_k = -\infty$  in view of (46), (16), (38) would imply that  $f(x_k(t)) + g(t) + h(t, x_k(t))$  is of constant sign for all sufficiently great  $k$  and this would lead to a contradiction with (35). Thus (45) is proved.

2. Since  $0 < c_1 < \beta$  and  $(x, \tilde{x}) = (\bar{x}, \tilde{x})$ , each solution  $x(t) = \bar{x} + \tilde{x}(t)$ ,  $a \leq t \leq b$  of (41), (35) satisfies the inequality

$$\begin{aligned} \|x^{(\frac{n}{2})}\|_{L^2}^2 &\leq [(1 - \mu)c_1 + \mu\beta] \|\tilde{x}\|_{L^2}^2 + (b-a)\delta \|\tilde{x}\|_0 \\ &\leq \beta \|\tilde{x}\|_{L^2}^2 + (b-a)\delta \|\tilde{x}\|_0 \end{aligned}$$

which replaces (47). Therefore (46) with  $M = 0$  implies (46'). The inequality (45') can be proved in the same way as (39') has been proved.  $\square$

**Remark 6.** It is clear that the condition (44) is equivalent to the following condition: If  $n = 4m$  ( $n = 4m + 2$ ),  $m \geq 1$ , then the function  $F(x) = f(x) - \beta x$  ( $F(x) = f(x) + \beta x$ ) is nonincreasing (nondecreasing) in  $\mathbb{R}$ .

**Theorem 7.** Suppose that  $n \geq 2$ ,  $\varepsilon = 1$  or  $\varepsilon = -1$ ,  $f$  satisfies (32'),  $h$  fulfils (38) and when  $n = 2m$ , there exists  $\beta$ ,  $0 \leq \beta < \left(\frac{2\pi}{b-a}\right)^n$  such that (44) is satisfied. Then the statements 1-3 of Theorem 4 hold for the BVP (37), (35). Moreover, if  $f \in C^1(\mathbb{R}, \mathbb{R})$  and  $\frac{\partial h}{\partial x} \in C([a, b] \times \mathbb{R}, \mathbb{R})$ , then also statement 4 of Theorem 4 holds for that BVP.

**Proof.** Proceeding in a similar way as in the proof of Theorem 4 we see that the assumptions (H.1), (H.2) and in the case that  $f'$ ,  $\frac{\partial h}{\partial x}$ , are continuous, also (H.4) of Theorems 4.1 and 4.2, [20], are satisfied. Since the BVP  $x^{(n)} - \varepsilon c_1 x = 0$ , (35), is regular for all sufficiently small  $c_1 > 0$ , the a priori estimates given in Lemmas 5 and 6 imply that also (H.3) and (H.5) are fulfilled. Then the result follows by Theorems 4.1 and 4.2 as well as by Corollary 4.2 in [20].  $\square$

By Remark 6, Theorem 3 in [19] and Theorem 5.2 in [20], Theorem 7 implies the following corollary.

**Corollary 3.** *Suppose that  $n \geq 2$ ,  $\varepsilon = 1$  or  $\varepsilon = -1$ ,  $f$  satisfies (32'),  $h$  fulfils (38) with a positive constant  $\delta$  and that the following conditions are true:*

- (a) *If  $n = 2m + 1$ , then the function  $f(\cdot) + h(t, \cdot)$  is either nonincreasing in  $\mathbb{R}$  or nondecreasing in  $\mathbb{R}$  for every  $t \in [a, b]$ .*
- (b) *If  $n = 4m$ , then the function  $f(\cdot) + h(t, \cdot)$  is nonincreasing in  $\mathbb{R}$  for every  $t \in [a, b]$  and there exists a  $\beta$ ,  $0 \leq \beta < (\frac{2\pi}{b-a})^n$  such that the function  $f(x) - \beta x$  of the variable  $x$  is nonincreasing in  $\mathbb{R}$ .*
- (c) *If  $n = 4m - 2$ , then the function  $f(\cdot) + h(t, \cdot)$  is nondecreasing in  $\mathbb{R}$  for every  $t \in [a, b]$  and there exists a  $\beta$ ,  $0 \leq \beta < (\frac{2\pi}{b-a})^n$  such that the function  $f(x) + \beta x$  of the variable  $x$  is nondecreasing in  $\mathbb{R}$ . Then there exists a closed set  $R_b \subset C^0$  such that for each  $g \in C^0 \setminus R_b$  the BVP (37), (35) has a unique solution, for each  $g \in R_b$  the set  $S_g$  of all solutions of that BVP is convex and  $R_b = \overline{R}_2$ , where  $R_2 \subset R_b$  is the set of all  $g \in C^0$  for which the BVP (37), (35) has infinitely many solutions.*

**Remark 7.** We see that under the assumptions of Corollary 3 the following alternative holds: Either the BVP (37), (35) has a unique solution or it has infinitely many solutions, more precisely a nontrivial convex compact set of solutions.

#### References

- [1] *S. R. Bernfeld, V. Lakshmikantham: An Introduction to Nonlinear Boundary Value Problems. Academic Press, Inc., New York, 1974.*
- [2] *S.H. Ding, J. Mawhin: A multiplicity result for periodic solutions of higher order ordinary differential equations. Differential Integral Equations 1 (1988), 31–39.*
- [3] *J.-P. Gossez, P. Omari: Periodic solutions of a second order ordinary differential equation: A necessary and sufficient condition for nonresonance. J. Differential Equations 94 (1991), 67–82.*
- [4] *Ch. P. Gupta: A note on a second order three-point boundary value problem. J. Math. Anal. Appl. 186 (1994), 277–281.*
- [5] *Ph. Hartman: Ordinary Differential Equations. Wiley-Interscience, New York, 1969.*
- [6] *L. V. Kantorovich, G. P. Akilov: Functional Analysis. Nauka, Moscow, 1977. (In Russian.)*
- [7] *J. Mawhin: Topological Degree Methods in Nonlinear Boundary Value Problems. CBMS Regional Conf. Ser. in Math. 40, American Math. Soc., Providence, 1979.*
- [8] *F. Neuman: Global Properties of Linear Ordinary Differential Equations. Academia, Praha, 1991.*
- [9] *L. Pinda: A remark on the existence of small solutions to a fourth order boundary value problem with large nonlinearity. Math. Slovaca 43 (1993), 149–170.*
- [10] *L. Pinda: On a fourth order periodic boundary value problem. Arch. Math. (Brno) 30 (1994), 1–8.*

- [11] *E. Pinda*: Landesman-Lazer type problems at an eigenvalue of odd multiplicity. Arch. Math. (Brno) *30* (1994), 73–84.
- [12] *N. Rouche, J. Mawhin*: Équations différentielles ordinaires, T. 2: Stabilité et solutions périodiques. Masson et Cie, Paris, 1973.
- [13] *W. Rudin*: Functional Analysis. Mc. Graw-Hill Book Co., New York, 1973.
- [14] *B. Rudolf*: The generalized boundary value problem is a Fredholm mapping of index zero. Arch. Math. (Brno) *31* (1995), 55–58.
- [15] *B. Rudolf*: A multiplicity result for a periodic boundary value problem. (Preprint).
- [16] *V. Šeda*: On the three-point boundary-value problem for a nonlinear third order ordinary differential equation. Arch. Math. 2, Scripta Fac. Sci., Nat. UJEP Brunensis *8* (1972), 85–98.
- [17] *V. Šeda*: Some remarks to coincidence theory. Czechoslovak Math. J. *38* (1988), 554–572.
- [18] *V. Šeda*: Quasilinear and approximate quasilinear method for generalized boundary value problems. WSSIA *1* (1992), 507–529.
- [19] *V. Šeda, J. J. Nieto, M. Gera*: Periodic boundary value problems for nonlinear higher order ordinary differential equations. Appl. Math. Comp. *48* (1992), 71–82.
- [20] *V. Šeda*: Fredholm mappings and the generalized boundary value problem. Differential Integral Equations *8* (1995), 19–40.

*Author's address: Valter Šeda*, Department of Mathematical Analysis, Comenius University, Mlynská dolina, 842 15 Bratislava, Slovakia, e-mail: [seda@fmph.uniba.sk](mailto:seda@fmph.uniba.sk).