

THE DIRECTED DISTANCE DIMENSION OF ORIENTED GRAPHS

GARY CHARTRAND, MICHAEL RAINES, PING ZHANG, Kalamazoo

(Received January 26, 1998)

Abstract. For a vertex v of a connected oriented graph D and an ordered set $W = \{w_1, w_2, \dots, w_k\}$ of vertices of D , the (directed distance) representation of v with respect to W is the ordered k -tuple $r(v \mid W) = (d(v, w_1), d(v, w_2), \dots, d(v, w_k))$, where $d(v, w_i)$ is the directed distance from v to w_i . The set W is a resolving set for D if every two distinct vertices of D have distinct representations. The minimum cardinality of a resolving set for D is the (directed distance) dimension $\dim(D)$ of D . The dimension of a connected oriented graph need not be defined. Those oriented graphs with dimension 1 are characterized. We discuss the problem of determining the largest dimension of an oriented graph with a fixed order. It is shown that if the outdegree of every vertex of a connected oriented graph D of order n is at least 2 and $\dim(D)$ is defined, then $\dim(D) \leq n - 3$ and this bound is sharp.

Keywords: oriented graphs, directed distance, resolving sets, dimension

MSC 2000: 05C12, 05C20

1. INTRODUCTION

For an oriented graph D of order n , an ordered set $W = \{w_1, w_2, \dots, w_k\}$ of vertices of D , and a vertex v of D , the k -vector (ordered k -tuple)


$$r(v \mid W) = (d(v, w_1), d(v, w_2), \dots, d(v, w_k))$$

is referred to as the (*directed distance*) *representation of v with respect to W* , where $d(x, y)$ denotes the directed distance from x to y , that is, the length of a shortest directed $x - y$ path in D . Since directed $x - y$ paths need not exist in D , even if D is connected (its underlying graph is connected), the vector $r(v \mid W)$ need not exist as well. If $r(v \mid W)$ exists for every vertex v of D , then the set W is called a *resolving set* for D if every two distinct vertices of D have distinct representations. A resolving set of minimum cardinality is called a *basis* for D and this cardinality is

the (*directed distance*) *dimension* $\dim(D)$ of D . Of course, not every oriented graph has a dimension. An oriented graph of dimension k is also called *k-dimensional*.

To determine whether an ordered set $W = \{w_1, w_2, \dots, w_k\}$ of vertices in an oriented graph D is a resolving set, we need only show that the representations of the vertices of $V(D) - W$ are distinct since $r(w_i | W)$ is the only representation whose i th coordinate is 0.

The directed distance dimension of an oriented graph is a natural analogue of the metric dimension of a graph that was introduced independently by Harary and Melter [2] and Slater [3], [4]. This concept was also investigated in [1] as a result of studying a problem in pharmaceutical chemistry.



D

Figure 1. An oriented graph D with dimension 2

In the oriented graph D of Figure 1, let $W_1 = \{u, v\}$. The five representations of the vertices of D with respect to W_1 are $r(u | W_1) = (0, 2)$, $r(v | W_1) = (1, 0)$, $r(w | W_1) = (2, 1)$, $r(x | W_1) = (2, 1)$, and $r(y | W_1) = (1, 3)$. Since x and w have the same representation, W_1 is not a resolving set for D .

The five representations of the vertices of D with respect to $W_2 = \{u, v, w\}$ are

$$\begin{aligned} r(u | W_2) &= (0, 2, 2), & r(v | W_2) &= (1, 0, 3), & r(w | W_2) &= (2, 1, 0), \\ r(x | W_2) &= (2, 1, 1), & r(y | W_2) &= (1, 3, 3) \end{aligned}$$

Since these five 3-vectors are distinct, W_2 is a resolving set for D . However, W_2 is not a basis for D . To see this, let $W_3 = \{x, y\}$. Then $r(u | W_3) = (1, 3)$, $r(v | W_3) = (2, 1)$, $r(w | W_3) = (3, 1)$, $r(x | W_3) = (0, 2)$, and $r(y | W_3) = (2, 0)$, which are distinct as well. So W_3 is a resolving set for D . Since there is no 1-element resolving set for D , it follows that W_3 is a basis and $\dim(D) = 2$.

Now let T be the tournament shown in Figure 2. Table 1 gives all 2-element choices for W and shows that for each such choice, there exist two equal 2-vectors, thus showing that $\dim(T) \geq 3$. However, $\dim(T) = 3$ since $\{v_1, v_3, v_6\}$ is a basis for T . Figure 3 shows an oriented graph D containing T as an induced subdigraph. The set $W = \{x, y\}$ is a basis of D , so $\dim(D) = 2$. Hence we have the possibly

unexpected property that the 3-dimensional tournament T is an induced subdigraph of the 2-dimensional oriented graph D .




Figure 2. The tournament T




Figure 3. The digraph D

There is a fundamental question here—one whose answer is not known to us, but one which deserves further study. What is a necessary and sufficient condition for the dimension of a digraph D to be defined? Certainly, if D is strong, then $\dim(D)$ is defined. Also, if D is connected and contains a vertex such that $D - v$ is strong, then $\dim(D)$ is defined. This last statement follows because if $\text{od } v > 0$, then $V(D) - \{v\}$ is a resolving set; while if $\text{id } v > 0$, then $V(D)$ is a resolving set. There are numerous other sufficient conditions for $\dim(D)$ to be defined.

W	equivalent vectors
$\{v_1, v_2\}$	$r(v_5 W) = r(v_7 W) = (1, 2)$
$\{v_1, v_3\}$	$r(v_6 W) = r(v_7 W) = (1, 1)$
$\{v_1, v_4\}$	$r(v_5 W) = r(v_7 W) = (1, 2)$
$\{v_1, v_5\}$	$r(v_6 W) = r(v_7 W) = (1, 2)$
$\{v_1, v_6\}$	$r(v_2 W) = r(v_3 W) = (2, 2)$
$\{v_1, v_7\}$	$r(v_5 W) = r(v_6 W) = (1, 1)$
$\{v_2, v_3\}$	$r(v_1 W) = r(v_6 W) = (1, 1)$
$\{v_2, v_4\}$	$r(v_5 W) = r(v_7 W) = (2, 2)$
$\{v_2, v_5\}$	$r(v_1 W) = r(v_5 W) = (1, 2)$
$\{v_2, v_6\}$	$r(v_4 W) = r(v_5 W) = (2, 1)$
$\{v_2, v_7\}$	$r(v_4 W) = r(v_5 W) = (2, 1)$
$\{v_3, v_4\}$	$r(v_1 W) = r(v_2 W) = (1, 1)$
$\{v_3, v_5\}$	$r(v_6 W) = r(v_7 W) = (1, 2)$
$\{v_3, v_6\}$	$r(v_1 W) = r(v_2 W) = (1, 2)$
$\{v_3, v_7\}$	$r(v_2 W) = r(v_6 W) = (1, 1)$
$\{v_4, v_5\}$	$r(v_2 W) = r(v_3 W) = (1, 1)$
$\{v_4, v_6\}$	$r(v_1 W) = r(v_2 W) = (1, 2)$
$\{v_4, v_7\}$	$r(v_1 W) = r(v_3 W) = (1, 2)$
$\{v_5, v_6\}$	$r(v_2 W) = r(v_3 W) = (1, 2)$
$\{v_5, v_7\}$	$r(v_2 W) = r(v_4 W) = (1, 1)$
$\{v_6, v_7\}$	$r(v_4 W) = r(v_5 W) = (1, 1)$

Table 1.

2. 1-DIMENSIONAL ORIENTED GRAPHS

In this section we characterize those oriented graphs having dimension 1. We also describe some properties of bases for 1-dimensional oriented graphs.

Theorem 2.1. *Let D be a nontrivial oriented graph of order n . Then $\dim(D) = 1$ if and only if there exists a vertex v in D such that*

- (i) D contains a hamiltonian path P with terminal vertex v such that $\text{id}_D v = 1$;
and
- (ii) if the hamiltonian path P in (i) is of the form

$$v_{n-1}, v_{n-2}, \dots, v_1, v,$$

then, for each pair i, j of integers with $1 \leq i < j \leq n - 1$, the digraph $D - E(P)$ contains no arc of the form (v_j, v_i) .

PROOF. Assume that $\dim(D) = 1$. Let $W = \{v\}$, $v \in V(D)$, be a basis of D . Then the distance $d(u, v)$ from u to v is defined for each vertex u in D and the set $\{d(u, v); u \in V(D)\} = \{0, 1, \dots, n-1\}$. Thus, we may assume that $V(D) = \{v, v_1, v_2, \dots, v_{n-1}\}$ where $d(v_i, v) = i$ ($1 \leq i \leq n-1$). Clearly, $\text{id } v = 1$. Since $d(v_{n-1}, v) = n-1$, there exists a hamiltonian path in D , namely $P: v_{n-1}, v_{n-2}, \dots, v_1, v$, so (i) holds. Furthermore, if there exists a pair i, j of integers ($1 \leq i < j \leq n-1$) such that the arc (v_j, v_i) is in $D - E(P)$, then $j \neq i+1$ and $d(v_j, v) = d(v_{i+1}, v)$ (shown in Figure 4). This contradicts the fact that $\{d(u, v); u \in V(D)\}$ consists of n distinct integers, so (ii) holds.




Figure 4.

Conversely, assume that there is a vertex v in D such that (i) and (ii) hold. We show that $W = \{v\}$ is a resolving set of D . Since $d(u, v)$ is defined for each $u \in V(D)$, it suffices to show that the set $\{d(v_i, v); 1 \leq i \leq n-1\}$ consists of $n-1$ distinct integers. Suppose that this is not the case. Then there exist integers i, j ($1 \leq i < j \leq n-1$) such that $d(v_j, v) = d(v_i, v) = \ell$. Let P_1 be a $v_i - v$ path and P_2 a $v_j - v$ path in D such that P_1 and P_2 have the same length ℓ . Since $\text{id } v = 1$, there exists a vertex $v_k \neq v$ in D that belongs to both P_1 and P_2 . Assume that v_k is the vertex with largest index k such that the path $v_k, v_{k-1}, \dots, v_1, v$ is on both P_1 and P_2 (see Figure 5).




Figure 5.

Let $(v_{k_1}, v_k) \in E(P_1)$ and $(v_{k_2}, v_k) \in E(P_2)$ where $(v_{k_1}, v_k) \neq (v_{k_2}, v_k)$. Clearly, $k_1 > k$ and $k_2 > k$. It follows that at least one of these arcs is in $D - E(P)$, but this is a contradiction to (ii). \square

We now present some facts concerning bases in 1-dimensional oriented graphs.

Theorem 2.2. *Let D be a digraph of order n with $\dim(D) = 1$. Furthermore, let v_1 and v_2 be distinct vertices of D with $d(v_1, v_2) = 2$ such that both $\{v_1\}$ and*

$\{v_2\}$ are bases of D . If v is a vertex of D such that $(v_1, v), (v, v_2) \in E(D)$, then $\{v\}$ is also a basis of D .

Proof. To show that $\{v\}$ is a basis of D , we show that for each $u \in V(D)$, the distance $d(u, v)$ is defined and the set $\{d(u, v); u \in V(D)\}$ consists of n distinct integers.

First notice that $\text{id } v = 1$, for otherwise there exist distinct vertices x and y of D such that $d(x, v) = d(y, v) = 1$. Since $\text{id } v_2 = 1$, by Theorem 2.1, we have

$$d(x, v_2) = d(y, v_2) = d(x, v) + 1 = 2$$

This contradicts the fact that $\{v_2\}$ is a basis of D .

Furthermore, suppose that there exist vertices u, w in D such that $d(u, v) = d(w, v)$. Since $\text{id } v = 1$, each $u - v$ path contains the arc (v_1, v) as its terminal arc, as does each $w - v$ path, so

$$d(u, v_1) = d(w, v_1) = d(u, v) - 1$$

Again, this contradicts the fact that $\{v_1\}$ is a basis of D . □

We now have an immediate consequence of Theorem 2.2.

Corollary 2.3. *If D is a 1-dimensional oriented graph of order $n \geq 3$ such that $\{v\}$ is a basis of D for every vertex v in D , then D is a directed cycle.*

Proof. Let $V(D) = \{v_1, v_2, \dots, v_n\}$. By Theorem 2.2, $\text{id } v = 1$ for every vertex v of D . Moreover, D contains a hamiltonian path P . We can assume that

$$P: v_n, v_{n-1}, \dots, v_2, v_1$$

Next, we show that D contains the cycle

$$C_n: v_n, v_{n-1}, \dots, v_2, v_1, v_n$$

Since $\text{id } v_n = 1$, there exists a unique vertex v such that $(v, v_n) \in E(D)$. If $v \neq v_1$, then $(v_i, v_n) \in E(D)$ for some i ($2 \leq i \leq n-1$). Since $\{v_n\}$ is a basis of D , there exists a hamiltonian path in D with terminal vertex v_n . However, since every vertex has indegree 1, the only possible path in D with v_n as its terminal vertex is

$$P': v_{n-1}, v_{n-2}, \dots, v_{i+1}, v_i, v_n$$

Since P' has length $n-i$, it is not a hamiltonian path. This contradicts the fact that $\{v_n\}$ is a basis. So D contains the cycle C_n . Furthermore, since $\text{id } v = 1$, D cannot contain any arc except those in C_n . So $D = C_n$. □

We can improve Corollary 2.3 slightly.

Corollary 2.4. *If D is a 1-dimensional oriented graph of order $n \geq 3$ such that*

$$|\{v; \{v\} \text{ is a basis of } D\}| \geq n - 1$$

then D is a directed cycle.

Proof. Let $V(D) = \{v, v_1, v_2, \dots, v_{n-1}\}$. Without loss of generality, we assume that $\{v_i\}$ is a basis of D for $1 \leq i \leq n - 1$. By Corollary 2.3, it suffices to show that $\{v\}$ is a basis as well.

We claim that $\text{od } v > 0$. Suppose that this is not the case. Then for each vertex $u (\neq v)$, the distance $d(v, u)$ is not defined, which contradicts the fact that $\{u\}$ is a basis of D . Hence, there is a vertex $x (\neq v)$ such that $(v, x) \in E(D)$. Since $\{x\}$ is also a basis of D , then by Theorem 2.1(i), D contains a hamiltonian path with terminal vertex x and $\text{id } x = 1$. This implies that there exists a vertex y distinct from x and v such that $(y, v) \in E(D)$. It follows that $d(y, x) = 2$ and by Theorem 2.2, $\{v\}$ is also a basis of D . \square

The bound in Corollary 2.4 cannot be improved in general. For example, consider the oriented graph D of order n in Figure 6. Since $\{v_i\}$ is a basis for D for $1 \leq i \leq n - 2$, $\dim(D) = 1$. However, neither $\{v_{n-1}\}$ nor $\{v_n\}$ is a basis D . So $|\{v; \{v\} \text{ is a basis of } D\}| = n - 2$ and D is not a directed cycle.




Figure 6. An oriented graph with $(n - 2)$ 1-element bases

There is only one 1-dimensional oriented tree of every order.

Theorem 2.5. *For every oriented tree T , $\dim(T) = 1$ or $\dim(T)$ is undefined. Furthermore, if $\dim(T) = 1$, then T is a directed hamiltonian path.*

PROOF. There are certainly oriented trees whose dimension is undefined, for example, any orientation of a star $K_{1,t}$, where $t \geq 3$. Now let T be an oriented tree whose dimension is defined. Since T contains no cycles, for every pair x, y of vertices, whenever $d(x, y)$ is defined, $d(y, x)$ is undefined. Thus $\dim(T) = 1$.

If $\dim(T) = 1$, then, by Theorem 2.1, T contains a hamiltonian path P and so $T = P$. \square

3. ON ORIENTED GRAPHS WITH LARGE DIMENSION

We have characterized those oriented graphs with dimension 1. But how large can the dimension of an oriented graph of order n be? In this section, we describe upper bounds for the dimension of a connected oriented graph in terms of lower bounds for the outdegrees of its vertices. The outdegree of every vertex in the oriented graph D of Figure 7 is 2, yet $\dim(D)$ is undefined. Such examples exist regardless of the outdegrees.




Figure 7. The oriented graph D

Theorem 3.1. *If D is a connected oriented graph of order $n \geq 3$ with $\text{od } v \geq 1$ for all $v \in V(D)$ such that $\dim(D)$ is defined, then $\dim(D) \leq n - 2$.*

PROOF. Let D be an oriented graph satisfying the hypothesis of the theorem. Certainly $\dim(D) \leq n - 1$. Assume, to the contrary, that $\dim(D) = n - 1$. Let $W = \{v_1, v_2, \dots, v_{n-1}\}$ be a basis for D and let $V(D) - W = \{x\}$. Since $\text{od } x \geq 1$, assume, without loss of generality, that x is adjacent to v_1 . Also, since $\text{od } v_1 \geq 1$, we may assume that v_1 is adjacent to v_2 . Since $\dim(D) = n - 1$, $r(v_i \mid W - \{v_i\}) = r(x \mid W - \{v_i\})$ for $1 \leq i \leq n - 1$. Since x is adjacent to v_1 , it follows that v_2 is adjacent to v_1 , but this contradicts the fact that D is an oriented graph. \square

We now describe a class of oriented graphs. For $k \geq 2$, let D_k be an oriented graph with vertex set

$$V(D_k) = \{u, v, w_1, w_2, \dots, w_k\}$$

and let $E(D_k)$ consist of the arc (u, v) and the arcs (v, w_j) and (w_j, u) for $1 \leq j \leq k$. The oriented graph D_k is shown in Figure 8. Then D_k has order $n = k + 2$ and $\text{od } v \geq 1$ for all $v \in V(D_k)$. We claim that $\dim(D_k) = n - 3$.




Figure 8. The oriented graph D_k with minimum outdegree 1

First we show that $\dim(D_k) \leq n - 3$. Let $W = \{w_2, w_3, \dots, w_k\}$, where then $|W| = n - 3$. The distances $d(u, w_2) = 2$, $d(v, w_2) = 1$, and $d(w_1, w_2) = 3$ show that W is a resolving set for D_k and so $\dim(D_k) \leq n - 3$. On the other hand, at least $k - 1$ of the vertices w_1, w_2, \dots, w_k must belong to every resolving set of D_k since the distance from any two of these vertices to every other vertex of D_k is the same. Hence $\dim(D_k) \geq n - 3$ and so $\dim(D) = n - 3$. Of course, this does not show that sharpness of the bound in Theorem 3.1, except that if D_1 is the directed 3-cycle, then $\dim(D_1) = 1 = n - 2$.

We can, however, improve the bound in Theorem 3.1 if we require that the out-degree of every vertex is at least 2.

Theorem 3.2. *If D is a connected oriented graph of order $n \geq 5$ with $\text{od } v \geq 2$ for all $v \in V(D)$ such that $\dim(D)$ is defined, then $\dim(D) \leq n - 3$.*

Proof. Suppose, to the contrary, that D contains a basis \mathcal{B} of cardinality $n - 2$. Let $\mathcal{B} = \{v_1, v_2, \dots, v_{n-2}\}$, and $V(D) - \mathcal{B} = \{x, y\}$. For each i ($1 \leq i \leq n - 2$), $\mathcal{B} - \{v_i\}$ is not a resolving set. Hence for each such i , some two of the three vertices x, y, v_i have the same representations with respect to $\mathcal{B} - \{v_i\}$. We consider two cases.

Case 1: For some i ($1 \leq i \leq n - 2$), x and y have the same representations with respect to $\mathcal{B} - \{v_i\}$. Assume, without loss of generality, that x and y have the same representations with respect to $W = \mathcal{B} - \{v_{n-2}\}$. Then x and y have the same out-neighbors in W . Since x and y have distinct representations with respect to \mathcal{B} , exactly one of x and y is adjacent to v_{n-2} ; for if neither x nor y is adjacent to v_{n-2} , then $d(x, v_{n-2}) = d(y, v_{n-2})$. Therefore, we may assume that y is adjacent to v_{n-2} .

Let $W' = \{v_1, v_2, \dots, v_{n-4}, v_{n-2}\}$. Two of x, y , and v_{n-3} have the same representations with respect to W' . However, y is adjacent to v_{n-2} and x is not, so x and y do not have the same representations with respect to W' . Thus there are two possibilities.

Subcase 1.1: $r(x | W') = r(v_{n-3} | W')$. We claim that x is adjacent to at most one of v_1, v_2, \dots, v_{n-2} . Suppose that this is not the case. Then we can assume without loss of generality that x is adjacent to v_1 and v_2 . Then $r(v_1 | \mathcal{B} - \{v_1\}) = r(x | \mathcal{B} - \{v_1\})$ or $r(v_1 | \mathcal{B} - \{v_1\}) = r(y | \mathcal{B} - \{v_1\})$. Similarly, $r(v_2 | \mathcal{B} - \{v_2\}) = r(x | \mathcal{B} - \{v_2\})$ or $r(v_2 | \mathcal{B} - \{v_2\}) = r(y | \mathcal{B} - \{v_2\})$. Since the out-neighbors of y in W are the same as the out-neighbors of x in W , we have that v_2 is an out-neighbor of v_1 and that v_1 is an out-neighbor of v_2 . Since D is an oriented graph, this is impossible, so, as claimed, x is adjacent to at most one of v_1, v_2, \dots, v_{n-2} . Now, since $\text{od } x \geq 2$, it follows that x is adjacent to y and exactly one vertex from v_1, v_2, \dots, v_{n-2} , say v_1 . However, since for $1 \leq i \leq n-3$, $r(v_i | \mathcal{B} - \{v_i\}) = r(x | \mathcal{B} - \{v_i\})$ or $r(v_i | \mathcal{B} - \{v_i\}) = r(y | \mathcal{B} - \{v_i\})$, it follows that v_1 is an out-neighbor of every vertex in the set $\{x, y, v_2, v_3, \dots, v_{n-3}\}$, so $\text{od } v_1 \leq 1$, which contradicts the assumption that every vertex in D has out-degree at least 2.

Subcase 1.2: $r(y | W') = r(v_{n-3} | W')$. We first suppose that x is adjacent to some vertex in W' , say v_1 . Because of the assumptions in Case 1 and Subcase 1.2, it follows that y and v_{n-3} are also adjacent to v_1 . However, since for $2 \leq i \leq n-3$, $r(v_i | \mathcal{B} - \{v_i\}) = r(x | \mathcal{B} - \{v_i\})$ or $r(v_i | \mathcal{B} - \{v_i\}) = r(y | \mathcal{B} - \{v_i\})$, it follows that v_1 is an out-neighbor of every vertex in the set $\{x, y, v_2, v_3, \dots, v_{n-3}, v_{n-2}\}$, so $\text{od } v_1 = 0$, which is a contradiction. Therefore, x is not adjacent to any of $v_1, v_2, \dots, v_{n-4}, v_{n-2}$. Thus, since $\text{od } x \geq 2$, it follows that x must be adjacent to both y and v_{n-3} . But y is adjacent to v_{n-3} as well, because x and y have the same representations with respect to W . Since x is not adjacent to any of v_1, v_2, \dots, v_{n-4} , it follows that y is not adjacent to any of v_1, v_2, \dots, v_{n-4} . Now $r(y | W') = r(v_{n-3} | W')$, so it follows that v_{n-3} is not adjacent to any of v_1, v_2, \dots, v_{n-4} . All of this implies that $\text{od } v_{n-3} = 1$, which is a contradiction.

Case 2: For every i ($1 \leq i \leq n-2$), x and y have distinct representations with respect to $\mathcal{B} - \{v_i\}$. We next prove that every vertex of \mathcal{B} is an out-neighbor of x or y but at most one vertex of \mathcal{B} is an out-neighbor of both x and y . To prove this, we first show that among the out-neighbors y_1, y_2, \dots, y_k of y in \mathcal{B} , at most one y_i has the same representation as y with respect to $\mathcal{B} - \{y_i\}$. Suppose that this is not the case. Then we may assume that $r(y_1 | \mathcal{B} - \{y_1\}) = r(y | \mathcal{B} - \{y_1\})$ and that $r(y_2 | \mathcal{B} - \{y_2\}) = r(y | \mathcal{B} - \{y_2\})$. The first equality tells us that y_2 is an out-neighbor of y_1 and the second equality tells us that y_1 is an out-neighbor of y_2 , contradicting the fact that D is an oriented graph. Similarly, among the out-neighbors x_1, x_2, \dots, x_ℓ of x in \mathcal{B} , at most one x_j has the same representation as x with respect to $\mathcal{B} - \{x_j\}$.

Next, we show that for each i ($1 \leq i \leq n-2$), at least one of x and y is adjacent to v_i . This follows from the fact that if neither x nor y is adjacent to v_i , then no other

vertex v_j from $\mathcal{B} - \{v_i\}$ can be adjacent to v_i since $r(v_j | \mathcal{B} - \{v_j\}) = r(x | \mathcal{B} - \{v_j\})$ or $r(v_j | \mathcal{B} - \{v_j\}) = r(y | \mathcal{B} - \{v_j\})$. Thus $\text{id } v_i = 0$, which is impossible since $d(z, v_i)$ must be defined for all $z \in V(D)$. Finally, x and y are simultaneously adjacent to at most one vertex v_i ($1 \leq i \leq n-2$), for if v_a and v_b are distinct out-neighbors of both x and y , then v_a and v_b are out-neighbors of each other, which is impossible.

This creates a natural partition of the vertices of \mathcal{B} into either two or three subsets, depending on whether there exists a vertex to which x and y are simultaneously adjacent. We now consider these two subcases.

Subcase 2.1: There exists a unique common out-neighbor of x and y .

We assume, without loss of generality, that v_{n-2} is an out-neighbor of both x and y . Furthermore, we can assume, without loss of generality, that the set $X = \{v_1, v_2, \dots, v_k\}$ consists of the out-neighbors of x and not y , and that the set $Y = \{v_{k+1}, v_{k+2}, \dots, v_{n-3}\}$ consists of the out-neighbors of y and not x . We further assume, without loss of generality, that the representations of y and v_{n-2} with respect to $\mathcal{B} - \{v_{n-2}\}$ are the same. Therefore, there is no vertex in $v_j \in Y$ for which the representations of y and v_j with respect to $\mathcal{B} - \{v_j\}$ are the same. Therefore, for every $v_j \in Y$, the representations of x and v_j with respect to $\mathcal{B} - \{v_j\}$ are the same.

Since x is adjacent to every vertex in X , every vertex in Y is adjacent to every vertex in $X \cup \{v_{n-2}\}$. Now, there is at most one $v_i \in X$ for which the representations of x and v_i are the same with respect to $\mathcal{B} - \{v_i\}$. Therefore, if $|X| \geq 2$, there exists at least one vertex $v_i \in X$ for which the representations of y and v_i with respect to $\mathcal{B} - \{v_i\}$ are the same. Hence, such a vertex v_i is adjacent to every vertex in Y , but this implies that D is not an oriented graph since for any $v_j \in Y$, there is an arc from v_i to v_j and an arc from v_j to v_i . Therefore, $|X| \leq 1$. But if $|X| = 1$, then v_1 is the only vertex that could possibly be an out-neighbor of v_{n-2} . This contradicts the assumption that the out-degree of every vertex in D is at least 2, so $|X| = 0$. We have already seen that every vertex in $Y \cup \{x\}$ is adjacent to vertex v_{n-2} , so even if $|X| = 0$, we have that $\text{od } v_{n-2} = 0$, which cannot occur.

Subcase 2.2: No vertex is a common out-neighbor of x and y .

We assume, without loss of generality, that the set $X = \{v_1, v_2, \dots, v_k\}$ consists of the out-neighbors of x and not y , and that the set $Y = \{v_{k+1}, v_{k+2}, \dots, v_{n-2}\}$ consists of the out-neighbors of y and not x . Recall that there is at most one $v_i \in X$ such that the representations of v_i and x with respect to $\mathcal{B} - \{v_i\}$ are equal and at most one $v_j \in Y$ such that the representations of v_j and y with respect to $\mathcal{B} - \{v_j\}$ are equal. This produces three possibilities to consider.

Subcase 2.2.1: For every $v_i \in X$ and $v_j \in Y$, the representations of v_i and y with respect to $\mathcal{B} - \{v_i\}$ are the same and the representations of v_j and x with respect to

$\mathcal{B} - \{v_j\}$ are the same. Then every vertex in Y is adjacent to every vertex in X , and every vertex in X is adjacent to every vertex in Y . This contradicts the fact that D is an oriented graph as long as X and Y are both nonempty. However, if X or Y is empty, then $\text{od } x \leq 1$ or $\text{od } y \leq 1$, respectively, which is a contradiction.

Subcase 2.2.2: There is exactly one $v_i \in X$ for which the representations of v_i and x with respect to $\mathcal{B} - \{v_i\}$ are equal and there is no $v_j \in Y$ for which v_j and y have the same representations with respect to $\mathcal{B} - \{v_j\}$. (Note that this subcase is symmetric to the case when there is exactly one $v_j \in Y$ for which the representations of v_j and y with respect to $\mathcal{B} - \{v_j\}$ are equal and for which there is no $v_i \in X$ such that v_i and x have the same representations with respect to $\mathcal{B} - \{v_i\}$.) Now every vertex in Y has the same out-neighbors as x , namely the vertices in the set X . So if $Y \neq \emptyset$, then every vertex in Y is adjacent to every vertex in X . Furthermore, every vertex in $X - \{v_i\}$ has the same out-neighbors as y . So if $|X| \geq 2$, then there is at least one vertex in X which is adjacent to every vertex in Y . But this produces a contradiction since D is an oriented graph. Note that if $Y = \emptyset$, then y is adjacent to at most one vertex, namely x , and this is a contradiction.

Assume now that $|X| \leq 1$ (so $|Y| \geq 2$). If $|X| = 1$, then $v_i = v_1$ and since every vertex in Y is adjacent to v_i , the vertex v_i is adjacent to no vertex except possibly y . Hence, $\text{od } v_i \leq 1$, which is a contradiction. If $X = \emptyset$, then x has no out-neighbors except possibly for y , but this contradicts the assumption that the out-degree of x is at least 2.

Subcase 2.2.3: There exists exactly one $v_i \in X$ for which the representations of v_i and x with respect to $\mathcal{B} - \{v_i\}$ are the same and exactly one $v_j \in Y$ for which the representations of v_j and y with respect to $\mathcal{B} - \{v_j\}$ are the same. First, suppose that $|X| \geq 2$ and $|Y| \geq 2$. Then there exists at least one vertex $v \in X$ for which the representations of v and y with respect to $\mathcal{B} - \{v\}$ are the same. Therefore, v is adjacent to every vertex in Y . Similarly, there is at least one vertex $w \in Y$ for which the representations of w and x with respect to $\mathcal{B} - \{w\}$ are the same. Therefore, w is adjacent to every vertex in X . However, since $v \in X$ and $w \in Y$, it follows that v is adjacent to w and w is adjacent to v . This contradicts the fact that D is an oriented graph.

Next suppose that $|X| = 1$, that $|Y| \geq 2$, and that $X = \{v_1\}$. Then the out-neighbors of x are y and v_1 . Furthermore, v_1 is an out-neighbor of every vertex in $Y - \{v_j\}$. The only possible out-neighbors of v_1 are y and v_j . However, if v_i is adjacent to v_j , then x is adjacent to v_j , which contradicts the fact that $v_j \notin X$. Therefore, $\text{od } v_i \leq 1$, contradicting the fact that every vertex in D has out-degree at least 2. The case where $|Y| = 1$ and $|X| \geq 2$ is similar. \square

The sharpness of the bound in Theorem 3.1 is not illustrated by the digraph D_k shown in Figure 8 since the outdegrees of most vertices of D_k are 1. We can, however, show that the upper bound in Theorem 3.2 is sharp. Let F_k be an oriented graph with vertex set

$$V(F_k) = \{u_1, u_2, v_1, v_2, w_1, w_2, \dots, w_k\}$$

and let $E(F_k)$ consist of (1) the arcs (u_i, v_j) for $1 \leq i, j \leq 2$ and (2) the arcs (v_i, w_j) and (w_j, u_i) for $1 \leq i \leq 2$ and $1 \leq j \leq k$. The oriented graph F_k is shown in Figure 9. Then F_k has order $n = k + 4$ and the property that $\text{od } v \geq 2$ for all $v \in V(F_k)$. We claim that $\dim(F_k) = n - 3$.




Figure 9. The oriented graph F_k with minimum outdegree 2

First we show that $\dim(F_k) \leq n - 3$. Let $W = \{u_1, v_1, w_2, w_3, \dots, w_k\}$, where then $|W| = n - 3$. The distances $d(u_2, w_2) = 2$, $d(v_2, w_2) = 1$, and $d(w_1, w_2) = 3$ show that W is a resolving set for F_k and so $\dim(F_k) \leq n - 3$. Next we show that $\dim(F_k) \geq n - 3$. Let W be a resolving set for F_k . Certainly at least $k - 1$ of the vertices w_1, w_2, \dots, w_k must belong to W since the distance from any two of these vertices to every other vertex of F_k is the same. Moreover, at least one of u_1 and u_2 must belong to W since the distance from u_1 and u_2 every other vertex of F_k is the same. For the same reason, at least one of v_1 and v_2 must belong to W . Hence $\dim(F_k) \geq n - 3$ and so $\dim(F_k) = n - 3$.

No additional restriction on the outdegrees of the vertices of an oriented graph yields an improved bound, however. Let $r \geq 2$ be an integer. In the oriented graph of Figure 8, replace u_1, u_2 by the r vertices u_1, u_2, \dots, u_r and v_1, v_2 by the r vertices v_1, v_2, \dots, v_r and add the appropriate arcs. The resulting oriented graph H_k has $\text{od } v \geq r$ for all $v \in V(H_k)$, but $\dim(H_k) = n - 3$.

References

- [1] *G. Chartrand, L. Eroh, M. Johnson, O. R. Oellermann*: Resolvability in graphs and the metric dimension of a graph. Preprint.
- [2] *F. Harary, R. A. Melter*: On the metric dimension of a graph. *Ars Combin.* 2 (1976), 191–195.
- [3] *P.J. Slater*: Leaves of trees. *Congress. Numer.* 14 (1975), 549–559.
- [4] *P.J. Slater*: Dominating and reference sets in graphs. *J. Math. Phys. Sci.* 22 (1988), 445–455.

Authors' addresses: Gary Chartrand, Michael Raines, Ping Zhang, Department of Mathematics and Statistics Western Michigan University Kalamazoo, MI 49008, USA, e-mail: zhang@math-stat.wmich.edu.