Memoirs on Differential Equations and Mathematical Physics
VOLUME 11, 1997, 89–103

Sergo Kharibegashvili

ON THE SOLVABILITY OF A MULTIDIMENSIONAL VERSION OF THE GOURSAT PROBLEM FOR A SECOND ORDER HYPERBOLIC EQUATION WITH CHARACTERISTIC DEGENERATION

Abstract. A multidimensional version of the Goursat problem is considered for a second order hyperbolic equation with characteristic degeneration. Using the technique of functional spaces with a negative norm the correct formulation of this problem in the Sobolev weighted space is given.

Mathematics Subject Classification. 35L80.

Key words and phrases. Hyperbolic equation with characteristic degeneration, multidimensional version of the Goursat problem, Sobolev weighted space, functional space with a negative norm.

რეზიუმე. განხილულია გურსას ამოცანის მრავალგანზომილებიანი ვერსია მეორე რიგის ჰიპერბოლური განტოლებისათვის მახასიათებელი გადაგვარებით. უარყოფით ნორმიანი ფუნქციონალური სივრცეების ტექნიკის გამოყენებით მოცემულია ამ ამოცანის კორექტული ფორმულირება სობოლევის წონიან სივრცეში.

In the space of variables x_1, x_2, t we shall consider a second order degenerating hyperbolic equation of the form

$$Lu \equiv u_{tt} - u_{x_1x_1} - (|x_2|^m u_{x_2})_{x_2} + a_1 u_{x_1} + a_2 u_{x_2} + a_3 u_t + a_4 u = F, \tag{1}$$

where a_i, $i = 1, \ldots, 4$, F are the given real functions and u is the desired real function, $1 \le m = const < 2$.

Denote by

$$D : \frac{2}{2-m} x_2^{\frac{2-m}{2}} < t < 1 - \frac{2}{2-m} x_2^{\frac{2-m}{2}}, \quad 0 < x_2 < \Big(\frac{2-m}{4}\Big)^{\frac{2}{2-m}}$$

the unbounded domain lying in a half-space $x_2 > 0$ bounded by the characteristic surfaces

$$S_1 : t - \frac{2}{2-m} x_2^{\frac{2-m}{2}} = 0, \quad 0 < x_2 < \Big(\frac{2-m}{4}\Big)^{\frac{2}{2-m}},$$
$$S_2 : t + \frac{2}{2-m} x_2^{\frac{2-m}{2}} = 1, \quad 0 < x_2 < \Big(\frac{2-m}{4}\Big)^{\frac{2}{2-m}}$$

of equation (1) and by the two-dimensional surface $S_0 : x_2 = 0$, $0 < t < 1$ on which this equation has characteristic degeneration. It will be assumed below that in the domain D the coefficients a_i, $i = 1, \ldots, 4$, of equation (1) are the bounded functions from the class $C^2(\overline{D})$.

For equation (1) we shall consider a multidimensional version of the Goursat problem formulated as follows: in the domain D find a solution $u(x_1, x_2, t)$ of equation (1) satisfying the boundary condition

$$u \mid_{S_1} = 0. \tag{2}$$

In a similar manner we formulate the problem for the equation

$$L^* v \equiv v_{tt} - v_{x_1x_1} - (|x_2|^m v_{x_2})_{x_2} - (a_1 v)_{x_1} - \\ - (a_2 v)_{x_2} - (a_3 v)_t + a_4 v = F \tag{3}$$

in the domain D using the boundary condition

$$v \mid_{S_2} = 0, \tag{4}$$

where L^* is the formal conjugate operator of L.

Similar problems, in which, along with condition (2), it is required that the condition $u \mid_{S_0} = 0$ or $\frac{\partial u}{\partial n} \mid_{S_0} = 0$ be fulfilled on the section S_0 of the boundary ∂D of the domain D, are investigated in [1–6] for $m = 0$ when equation (1) is not the degenerating one and has, in its principal part, a wave operator. As will be shown below, by virtue of the degeneration character of equation (1), where $1 \le m < 2$, we can get rid of the fulfillment of any boundary condition on the section S_0 of the boundary ∂D of the domain D, since problem (1), (2) will turn out to be correctly formulated. In the case of a second order hyperbolic equation with noncharacteristic degeneration of the form

$$u_{tt} - |x_2|^m u_{x_1x_1} - u_{x_2x_2} + a_1 u_{x_1} + a_2 u_{x_2} + a_3 u_t + a_4 u = F$$

a multidimensional variant of the first Darboux problem is studied in [7]. Other variants of the multidimensional Goursat and Darboux problems are treated in [8–10].

Denote by E and E^* the classes of functions from the Sobolev space $W_2^2(D)$ satisfying the boundary condition (2) or (4), respectively. Let $W_+(W_+^*)$ be the Hilbert space with weight obtained by the closure of the space $E(E^*)$ with respect to the norm

$$\|u\|_{1,+}^2 = \int\limits_D (u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2 + u^2) dD.$$

Remark 1. Since $m \geq 1$, by virtue of the familiar embedding theorems for Sobolev weighted spaces [11] the class of functions $E_0(E_0^*)$ belonging to the space $C^\infty(\overline{D})$, having the bounded carriers (i.e., diam supp $u < +\infty$), satisfying the boundary condition (2) ((4)) and vanishing in some neighborhood (each function has its own neighborhood) of the surface S_0, is a dense subspace of the weighed space $W_+(W_+^*)$. Therefore, below it will be sometimes convenient for us to use, instead of the spaces E and E^*, the spaces E_0 and E_0^*.

Denote by $W_-(W_-^*)$ the space with negative norm constructed with respect to $L_2(D)$ and $W_+(W_+^*)$ [12].

Consider the condition

$$M = \sup_{\overline{D}} \left| x_2^{-\frac{m}{2}} a_2(x_1, x_2, t) \right| < +\infty \tag{5}$$

on the lower coefficient a_2 in equation (1).

The uniqueness theorem for solutions of problem (1), (2) belonging to the Sobolev space $W_2^2(D)$ is provided by

Lemma 1. *Let condition* (5) *be fulfilled. Then for any* $u \in W_2^2(D)$ *satisfying the condition*

$$\int\limits_{S_1} \left[u^2 + x_2^{\frac{m}{2}} u_{x_1}^2 + x_2^{-\frac{m}{2}} \left(\frac{\partial u}{\partial N} \right)^2 \right] ds < +\infty \tag{6}$$

there holds the following a priori estimate

$$\|u\|_{1,+} \leq c\big(\|f\|_{1,*} + \|F\|_{L_2(D)} \big), \tag{7}$$

where the positive constant c *does not depend on* u; $f = u\,|_{S_1}$, $F = Lu$,

$$\|f\|_{1,*}^2 = \int\limits_{S_1} \left[f^2 + x_2^{\frac{m}{2}} f_{x_1}^2 + x_2^{-\frac{m}{2}} \left(\frac{\partial f}{\partial N} \right)^2 \right] ds,$$

$\frac{\partial}{\partial N}\,|_{S_1} = -(1 + x_2^{-m})^{-\frac{1}{2}} \left[\frac{\partial}{\partial t} + x_2^m \frac{\partial}{\partial x_2} \right]$ *is the derivative with respect to the conormal which is the internal differential operator on the characteristic surface* S_1.

Proof. Let $n = (\nu_1, \nu_2, \nu_0)$ be the unit vector of the external normal to ∂D, i.e., $\nu_1 = cos(\widehat{n, x_1})$, $\nu_2 = cos(\widehat{n, x_2})$, $\nu_0 = cos(\widehat{n, t})$. By definition, the derivative with respect to the conormal on the boundary ∂D of the domain D for the operator L is calculated by the formula

$$\frac{\partial}{\partial N} = \nu_0 \frac{\partial}{\partial t} - \nu_1 \frac{\partial}{\partial x_1} - x_2^m \nu_2 \frac{\partial}{\partial x_2}.$$

Applying integration by parts, we have for $u \in W_2^2(D)$ and $\lambda = const > 0$:

$$2\int\limits_D e^{-\lambda t} u_{tt} u_t dD = \int\limits_{\partial D} e^{-\lambda t} u_t^2 \nu_0 ds + \int\limits_D \lambda e^{-\lambda t} u_t^2 dD, \tag{8}$$

$$-2\int\limits_D e^{-\lambda t}\big[u_{x_1x_1}u_t + (x_2^m u_{x_2})_{x_2} u_t\big]dD = -2\int\limits_{\partial D} e^{-\lambda t}(u_{x_1}u_t\nu_1 +$$

$$+x_2^m u_{x_2}u_t\nu_2)ds + 2\int\limits_D e^{-\lambda t}(u_{x_1}u_{x_1t} + x_2^m u_{x_2}u_{x_2t})dD =$$

$$= -2\int\limits_{\partial D} e^{-\lambda t}(u_{x_1}u_t\nu_1 + x_2^m u_{x_2}u_t\nu_2)ds + \int\limits_D e^{-\lambda t}\frac{\partial}{\partial t}(u_{x_1}^2 + x_2^m u_{x_2}^2)dD =$$

$$= -2\int\limits_{\partial D} e^{-\lambda t}(u_{x_1}u_t\nu_1 + x_2^m u_{x_2}u_t\nu_2)ds + \int\limits_{\partial D} e^{-\lambda t}(u_{x_1}^2 + x_2^m u_{x_2}^2)\nu_0 ds +$$

$$+\int\limits_D e^{-\lambda t}\lambda(u_{x_1}^2 + x_2^m u_{x_2}^2)dD. \tag{9}$$

It is easy to verify that

$$\begin{gathered}\nu_0\big|_{S_0} = \nu_1\big|_{S_0} = 0, \quad \frac{\partial u}{\partial N}\Big|_{S_0} = 0,\\ n\big|_{S_1} = \big(0, (1+x_2^{-m})^{-\frac{1}{2}} x_2^{-\frac{m}{2}}, -(1+x_2^{-m})^{-\frac{1}{2}}\big),\\ \nu_0\big|_{S_2} \geq 0, \quad (\nu_0^2 - \nu_1^2 - x_2^m\nu_2^2)\big|_{S_1\cup S_2} = 0.\end{gathered} \tag{10}$$

On multiplying both parts of equation (1) by $2e^{-\lambda t}u_t$, where $F = Lu$, and integrating the resulting expression with respect to the domain D we obtain by virtue of (6) and (8)–(10)

$$2(Lu, e^{-\lambda t}u_t)_{L_2(D)} = \int\limits_{S_1\cup S_2} e^{-\lambda t}\big[(u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2)\nu_0 -$$

$$-2(u_{x_1}u_t\nu_1 + x_2^m u_{x_2}u_t\nu_2)\big]ds + 2\int\limits_D e^{-\lambda t}\big[a_1u_{x_1} + a_2u_{x_2} +$$

$$+a_3u_t + a_4u\big]u_t dD + \int\limits_D e^{-\lambda t}\lambda\big[u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2\big]dD =$$

$$= \int\limits_{S_1\cup S_2} e^{-\lambda t}\nu_0^{-1}\big[(\nu_0 u_{x_1} - \nu_1 u_t)^2 + x_2^m(\nu_0 u_{x_2} - \nu_2 u_t)^2 +$$

$$+(\nu_0^2 - \nu_1^2 - x_2^m\nu_2^2)u_t^2\big]ds + 2\int\limits_D e^{-\lambda t}\big[\lambda(u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2) +$$

$$+2(a_1u_{x_1} + a_2u_{x_2} + a_3u_t + a_4u)u_t\big]dD \geq$$

$$\geq 2\int\limits_D e^{-\lambda t}\big[\lambda(u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2) + 2(a_1u_{x_1} + a_2u_{x_2} + a_3u_t + a_4u)u_t\big]dD -$$

$$-\int\limits_{S_1} e^{-\lambda t}\Big[(1+x_2^{-m})^{-\frac{1}{2}}u_{x_1}^2+(1+x_2^{-m})^{\frac{1}{2}}\Big(\frac{\partial u}{\partial N}\Big)^2\Big]ds\geq$$

$$\geq 2\int\limits_D e^{-\lambda t}\big[\lambda(u_t^2+u_{x_1}^2+x_2^m u_{x_2}^2)+2(a_1u_{x_1}+a_2u_{x_2}+a_3u_t+a_4u)u_t\big]dD-$$

$$-2\int\limits_{S_1}\Big[x_2^{\frac{m}{2}}u_{x_1}^2+x_2^{-\frac{m}{2}}\Big(\frac{\partial u}{\partial N}\Big)^2\Big]ds. \tag{11}$$

In deriving inequality (11), we used the fact that

$$\Big(\frac{\partial u}{\partial N}\Big)^2\Big|_{S_1}=x_2^m(\nu_0u_{x_2}-\nu_2u_t)^2\big|_{S_1}.$$

The structure of the domain D allows one to easily verify the validity of the inequality

$$\int\limits_D u^2dD\leq c_0\Big[\int\limits_{S_1}u^2ds+\int\limits_D u_t^2dD\Big] \tag{12}$$

for some $c_0=const>0$ not depending on $u\in W_2^2(D)$.

By inequality (5) we readily obtain

$$|2a_2u_xu_t|\leq 2M(x_2^{\frac{m}{2}}u_{x_2})u_t\leq M(x_2^mu_{x_2}^2+u_t^2). \tag{13}$$

By virtue of (12) and (13), inequality (11) implies for sufficiently large λ that

$$2(Lu,e^{-\lambda t}u_t)_{L_2(D)}\geq c_1\int\limits_D(u_t^2+u_{x_1}^2+x_2^mu_{x_2}^2+u^2)dD-$$

$$-c_2\int\limits_{S_1}\Big[u^2+x_2^{\frac{m}{2}}u_{x_1}^2+x_2^{-\frac{m}{2}}\Big(\frac{\partial u}{\partial N}\Big)^2\Big]ds, \tag{14}$$

where the positive constants c_1 and c_2 do not depend on u and the constant c_1 can be chosen arbitrarily large depending on λ. Therefore (14) obviously implies estimate (7). ■

Remark 2. Since for the operator L the derivative with respect to the conormal $\frac{\partial}{\partial N}$ is the internal differential operator on the characteristic surfaces of equation (1), by virtue of (2) and (4) we find for the functions $u\in E$ and $v\in E^*$ that

$$\frac{\partial u}{\partial N}\Big|_{S_1}=0,\quad \frac{\partial v}{\partial N}\Big|_{S_2}=0. \tag{15}$$

Lemma 2. *Let condition* (5) *be fulfilled. Then for all* $u\in E$, $v\in E^*$ *we have the inequalities*

$$\|Lu\|_{W_-^*}\leq c_1\|u\|_{W_+}, \tag{16}$$

$$\|L^*v\|_{W_-}\leq c_2\|v\|_{W_+^*}, \tag{17}$$

where the positive constants c_1 *and* c_2 *do not depend on* u *and* v, *respectively,* $\|\cdot\|_{W_+}=\|\cdot\|_{W_+^*}=\|\cdot\|_{1,+}$.

Proof. By the definition of a negative norm for $u \in E$ and by equalities (2), (4), (10), (15) we have

$$\|Lu\|_{W_-^*} = \sup_{v\in W_+^*} \|v\|_{W_+^*}^{-1}(Lu,v)_{L_2(D)} = \sup_{v\in E^*} \|v\|_{W_+^*}^{-1}(Lu,v)_{L_2(D)} =$$

$$= \sup_{v\in E^*} \|v\|_{W_+^*}^{-1} \int\limits_D \big[u_{tt}v - u_{x_1x_1}v - (x_2^m u_{x_2})_{x_2}v + a_1u_{x_1}v + a_2u_{x_2}v +$$

$$+a_3u_tv + a_4uv\big]dD = \sup_{v\in E^*} \|v\|_{W_+^*}^{-1} \int\limits_{\partial D} [u_tv\nu_0 - u_{x_1}v\nu_1 - x_2^m u_{x_2}v\nu_2]ds +$$

$$+ \sup_{v\in E^*} \|v\|_{W_+^*}^{-1} \int\limits_D [-u_tv_t + u_{x_1}v_{x_1} + x_2^m u_{x_2}v_{x_2} + a_1u_{x_1}v + a_2u_{x_2}v +$$

$$+a_3u_tv + a_4uv]dD = \sup_{v\in E^*} \|v\|_{W_+^*}^{-1} \int\limits_{S_1\cup S_2} \frac{\partial u}{\partial N}v\ ds +$$

$$+ \sup_{v\in E^*} \|v\|_{W_+^*}^{-1} \int\limits_D [-u_tv_t + u_{x_1}v_{x_1} + x_2^m u_{x_2}v_{x_2} + a_1u_{x_1}v + a_2u_{x_2}v +$$

$$+a_3u_tv + a_4uv]dD = \sup_{v\in E^*} \|v\|_{W_+^*}^{-1} \int\limits_D [-u_tv_t + u_{x_1}v_{x_1} + x_2^m u_{x_2}v_{x_2} +$$

$$+a_1u_{x_1}v + a_2u_{x_2}v + a_3u_tv + a_4uv]dD. \tag{18}$$

In view of condition (5) and the Schwartz inequality we obtain

$$\Big|\int\limits_D [-u_tv_t + u_{x_1}v_{x_1} + x_2^m u_{x_2}v_{x_2}]dD\Big| \le 3\Big[\int\limits_D (u_t^2 + u_{x_1}^2 +$$

$$+x_2^m u_{x_2}^2)dD\Big]^{\frac12}\Big[\int\limits_D (v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2)dD\Big]^{\frac12} \le 3\|u\|_{W_+}\|v\|_{W_+^*}, \tag{19}$$

$$\Big|\int\limits_D [a_1u_{x_1}v + a_2u_{x_2}v + a_3u_tv + a_4uv]dD\Big| \le$$

$$\le \sup_D |a_1|\ \|u_{x_2}\|_{L_2(D)}\|v\|_{L_2(D)} + M\Big(\int\limits_D x_2^m u_{x_2}^2 dD\Big)^{\frac12}\|v\|_{L_2(D)} +$$

$$+\sup_D |a_3|\ \|u_t\|_{L_2(D)}\|v\|_{L_2(D)} + \sup_D |a_4|\ \|u\|_{L_2(D)}\|v\|_{L_2(D)} \le$$

$$\le \Big(M + \sum_{i=1,i\ne2}^{4} \sup_D |a_i|\Big)\|u\|_{W_+}\|v\|_{W_+^*} = \widetilde{c}\|u\|_{W_+}\ \|v\|_{W_+^*}. \tag{20}$$

From (18)–(20) it follows that

$$\|Lu\|_{W_-^*} \le (3+\widetilde{c})\sup_{v\in E^*} \|v\|_{W_+^*}^{-1}\|u\|_{W_+}\|v\|_{W_+^*} = c_1\|u\|_{W_+},$$

which proves inequality (16). Since the proof of inequality (17) is quite similar to that of inequality (16), Lemma 2 is thereby completely proved. ■

Remark 3. By virtue of inequality (16) ((17)) the operator $L : W_+ \to W_-^*$ ($L : W_+^* \to W_-$) with the dense definition domain of $E(E^*)$ admits a closure which is a continuous operator from the space $W_+(W_+^*)$ into the space $W_-^*(W_-)$. If we denote this closure as previously by $L(L^*)$, it will be defined throughout the Hilbert space $W_+(W_+^*)$.

Lemma 3. *Problems* (1), (2) *and* (3), (4) *are mutually conjugate, i.e., the equality*

$$(Lu, v) = (u, L^*v). \tag{21}$$

holds for any $u \in W_+$ *and* $v \in W_+^*$.

Proof. By Remark 3 it is enough to prove equality (21) when $u \in E$ and $v \in E^*$. In that case it is obvious that $(Lu, v) = (Lu, v)_{L_2(D)}$. therefore we have

$$\begin{aligned}
(Lu, v) = (Lu, v)_{L_2(D)} = \int\limits_{\partial D} [u_t v\nu_0 - u_{x_1} v\nu_1 - x_2^m u_{x_2} v\nu_2] ds + \\
+ \int\limits_{\partial D} [a_1\nu_1 + a_2\nu_2 + a_3\nu_0] uv \, ds + \int\limits_D \big[- u_t v_t + u_{x_1} v_{x_1} + \\
+ x_2^m u_{x_2} v_{x_2} - u(a_1 v)_{x_1} - u(a_2 v)_{x_2} - u(a_3 v)_t + a_4 uv\big] dD = \\
= \int\limits_{\partial D} [u_t v\nu_0 - u_{x_1} v\nu_1 - x_2^m u_{x_2} v\nu_2] ds + \int\limits_{\partial D} [a_1\nu_1 + a_2\nu_2 + \\
+ a_3\nu_0] uv \, ds - \int\limits_{\partial D} [uv_t\nu_0 - uv_{x_1}\nu_1 - x_2^m uv_{x_2}\nu_2] ds + \\
+ \int\limits_D \big[uv_{tt} - uv_{x_1x_1} - u(x_2^m v_{x_2})_{x_2} - u(a_1 v)_{x_1} - \\
- u(a_2 v)_{x_2} - u(a_3 v)_t + a_4 uv\big] dD = \int\limits_{\partial D} \Big[\Big(v \frac{\partial u}{\partial N} - u \frac{\partial v}{\partial N} \Big) + \\
+ (a_1\nu_1 + a_2\nu_2 + a_3\nu_0) uv \Big] ds + (u, L^*v)_{L_2(D)}.
\end{aligned} \tag{22}$$

Since condition (5) implies $a_2\big|_{S_0} = 0$, by virtue of (2), (4), (10) and (15) we readily obtain equality (21) from (22), which proves Lemma 3. ■

Consider the conditions

$$\Omega\big|_{S_1} \le 0, \quad (\lambda\Omega + \Omega_t)\big|_D \le 0, \tag{23}$$

where the second inequality is fulfilled for sufficiently large λ, $\Omega = a_{1x_1} + a_{2x_2} + a_{3t} - a_4$.

Lemma 4. *Let conditions* (5) *and* (23) *be fulfilled. Then for any* $u \in W_+$ *we have the inequality*

$$c\|u\|_{L_2(D)} \le \|Lu\|_{W_-^*} \tag{24}$$

where the positive constant c *does not depend on* u.

Proof. By Remarks 1 and 3 it is enough to show that inequality (24) is fulfilled when $u \in E_0$. If $u \in E_0$ and thus vanishes in some neighborhood of the surface S_0, then one can easily verify that the function

$$v(x_1, x_2, t) = \int_t^{\varphi_2(x_1,x_2)} e^{-\lambda\tau} u(x_1, x_2, \tau) d\tau, \quad \lambda = const > 0,$$

where $t = \varphi_2(x_1, x_2)$ is an equation of the characteristic surface S_2, belongs to the space E_0^* and the equalities

$$v_t(x_1, x_2, t) = -e^{-\lambda t} u(x_1, x_2, t), \quad u(x_1, x_2, t) = -e^{\lambda t} v_t(x_1, x_2, t). \tag{25}$$

are fulfilled.

In view of (10), (15) and (25) we have

$$(Lu, v)_{L_2(D)} = \int_{\partial D} \Big[v \frac{\partial u}{\partial N} + (a_1\nu_1 + a_2\nu_2 + a_3\nu_0)uv \Big] ds +$$

$$+ \int_D [-u_t v_t + u_{x_1} v_{x_1} + x_2^m u_{x_2} v_{x_2} - u a_{1x_1} v - u a_1 v_{x_1} - u a_{2x_2} v -$$

$$- u a_2 v_{x_2} - u a_{3t} v - u a_3 v_t + a_4 uv] dD = \int_D e^{-\lambda t} u_t u \, dD +$$

$$+ \int_D e^{\lambda t} [-v_{x_1 t} v_{x_1} - x_2^m v_{x_2 t} v_{x_2} + a_{1x_1} v_t v + a_1 v_t v_{x_1} + a_{2x_2} v_t v +$$

$$+ a_2 v_t v_{x_2} + a_{3t} v_t v + a_3 v_t^2 - a_4 v_t v] dD. \tag{26}$$

By (2) we obtain similarly to (8) and (9)

$$\int_D e^{-\lambda t} u_t u \, dD = \frac{1}{2} \int_{\partial D} e^{-\lambda t} u^2 \nu_0 \, ds + \frac{1}{2} \int_D e^{-\lambda t} \lambda u^2 \, dD =$$

$$= \frac{1}{2} \int_{S_2} e^{-\lambda t} u^2 \nu_0 \, ds + \frac{1}{2} \int_D e^{\lambda t} \lambda v_t^2 \, dD =$$

$$= \frac{1}{2} \int_{S_2} e^{\lambda t} v_t^2 \nu_0 \, ds + \frac{1}{2} \int_D e^{\lambda t} \lambda v_t^2 \, dD, \tag{27}$$

$$\int_D e^{\lambda t} [-v_{x_1 t} v_{x_1} - x_2^m v_{x_2 t} v_{x_2}] dD = -\frac{1}{2} \int_{\partial D} e^{\lambda t} [v_{x_1}^2 + x_2^m v_{x_2}^2] \nu_0 \, ds +$$

$$+ \frac{1}{2} \int_D e^{\lambda t} \lambda [v_{x_1}^2 + x_2^m v_{x_2}^2] dD. \tag{28}$$

Since $v\big|_{S_2} = 0$, for some α we have $v_t = \alpha\nu_0$, $v_{x_1} = \alpha\nu_1$, $v_{x_2} = \alpha\nu_2$ on S_2. Therefore, recalling that the surface S_2 is characteristic, we obtain

$$(v_t^2 - v_{x_1}^2 - x_2^m v_{x_2}^2)\big|_{S_2} = \alpha^2 (\nu_0^2 - \nu_1^2 - x_2^m \nu_2^2)\big|_{S_2} = 0. \tag{29}$$

By virtue of $\nu_0\big|_{S_0} = 0$, $\nu_0\big|_{S_1} \le 0$, and equalities (4), (29) we find that

$$\frac{1}{2}\int\limits_{S_2} e^{\lambda t} v_t^2 \nu_0 \, ds - \frac{1}{2}\int\limits_{\partial D} e^{\lambda t}[v_{x_1}^2 + x_2^m v_{x_2}^2]\nu_0 \, ds =$$

$$= \frac{1}{2}\int\limits_{S_2} e^{\lambda t} v_t^2 \nu_0 \, ds - \frac{1}{2}\int\limits_{S_1} e^{\lambda t}[v_{x_1}^2 + x_2^m v_{x_2}^2]\nu_0 \, ds -$$

$$-\frac{1}{2}\int\limits_{S_2} e^{\lambda t}[v_{x_1}^2 + x_2^m v_{x_2}^2]\nu_0 \, ds \ge \frac{1}{2}\int\limits_{S_2} e^{\lambda t} v_t^2 \nu_0 \, ds - \frac{1}{2}\int\limits_{S_2} e^{\lambda t}[v_{x_1}^2 +$$

$$+ x_2^m v_{x_2}^2]\nu_0 \, ds = \frac{1}{2}\int\limits_{S_2} e^{\lambda t}[v_t^2 - v_{x_1}^2 - x_2^m v_{x_2}^2]\nu_0 \, ds = 0. \tag{30}$$

Taking into account (27), (28) and (30), we obtain from (26)

$$(Lu, v)_{L_2(D)} = \frac{1}{2}\int\limits_{S_2} e^{\lambda t} v_t^2 \nu_0 \, ds + \frac{1}{2}\int\limits_{D} e^{\lambda t} \lambda v_t^2 \, dD -$$

$$-\frac{1}{2}\int\limits_{\partial D} e^{\lambda t}[v_{x_1}^2 + x_2^m v_{x_2}^2]\nu_0 \, ds + \frac{1}{2}\int\limits_{D} e^{\lambda t}\lambda[v_{x_1}^2 + x_2^m v_{x_2}^2]dD +$$

$$+\int\limits_{D} e^{\lambda t}\big[a_1 v_t v_{x_1} + a_2 v_t v_{x_2} + a_3 v_t^2 + (a_{1x_1} + a_{2x_2} + a_{3t} - a_4) v_t v\big]dD \ge$$

$$\ge \frac{\lambda}{2}\int\limits_{D} e^{\lambda t}[v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2]dD + \int\limits_{D} e^{\lambda t}[a_1 v_t v_{x_1} +$$

$$+ a_2 v_t v_{x_2} + a_3 v_t^2 + (a_{1x_1} + a_{2x_2} + a_{3t} - a_4) v_t v]dD. \tag{31}$$

Using $\nu_0\big|_{S_1} \le 0$ and conditions (4), (10), (23) and performing integration by parts we derive

$$\int\limits_{D} e^{\lambda t}(a_{1x_1} + a_{2x_2} + a_{3t} - a_4) v_t v \, dD = \frac{1}{2}\int\limits_{\partial D} e^{\lambda t}(a_{1x_1} + a_{2x_2} +$$

$$+ a_{3t} - a_4) v^2 \nu_0 ds - \frac{1}{2}\int\limits_{D} e^{\lambda t}\big[\lambda(a_{1x_1} + a_{2x_2} + a_{3t} - a_4) +$$

$$+ (a_{1x_1} + a_{2x_2} + a_{3t} - a_4)_t\big] v^2 dD \ge 0, \tag{32}$$

where λ is a sufficiently large positive number.

With (32) taken into account (31) implies

$$(Lu, v)_{L_2(D)} \ge \frac{\lambda}{2}\int\limits_{D} e^{\lambda t}[v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2]dD +$$

$$+\int\limits_{D} e^{\lambda t}[a_1 v_t v_{x_1} + a_2 v_t v_{x_2} + a_3 v_t^2]dD \ge \frac{\lambda}{2}\int\limits_{D} e^{\lambda t}[v_t^2 + v_{x_1}^2 +$$

$$+x_2^m v_{x_2}^2]dD - \Big| \int\limits_D e^{\lambda t}[a_1 v_t v_{x_1} + a_2 v_t v_{x_2} + a_3 v_t^2]dD\Big|. \tag{33}$$

Assuming

$$\mu = \max\big(\sup_D |a_1|,\ \sup_D |a_3|\big)$$

by condition (5) we find that

$$\Big| \int\limits_D e^{\lambda t}[a_1 v_t v_{x_1} + a_2 v_t v_{x_2} + a_3 v_t^2]dD\Big| \le$$

$$\le \int\limits_D e^{\lambda t}\Big[\frac{\mu}{2}(v_{x_1}^2 + v_t^2) + M\frac{1}{2}(x_2^m v_{x_2}^2 + v_t^2) + \mu v_t^2\Big]dD \le$$

$$\le \Big(\frac{1}{2}M + \frac{3}{2}\mu\Big) \int\limits_D e^{\lambda t}[v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2]dD. \tag{34}$$

By virtue of (34) and (25) inequality (33) implies

$$(Lu, v)_{L_2(D)} \ge \Big[\frac{\lambda}{2} - \Big(\frac{1}{2}M + \frac{3}{2}\mu\Big)\Big] \int\limits_D e^{\lambda t}[v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2]dD \ge$$

$$\ge \sigma\Big[\int\limits_D e^{\lambda t} v_t^2 dD\Big]^{\frac{1}{2}}\Big[\int\limits_D [v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2]dD\Big]^{\frac{1}{2}} =$$

$$= \sigma\Big[\int\limits_D e^{-\lambda t} u^2 dD\Big]^{\frac{1}{2}}\Big[\int\limits_D [v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2]dD\Big]^{\frac{1}{2}} \ge$$

$$\ge \sigma \inf_D e^{-\lambda t}\|u\|_{L_2(D)}\Big[\int\limits_D [v_t^2 + v_{x_1}^2 + x_2^m v_{x_2}^2]dD\Big]^{\frac{1}{2}}, \tag{35}$$

where $\sigma = \Big(\frac{\lambda}{2} - \Big(\frac{1}{2}M + \frac{3}{2}\mu\Big)\Big) > 0$ for sufficiently large λ, and $\inf_D e^{-\lambda t} = const > 0$ by the structure of the domain D.

Since $v\big|_{S_2} = 0$, similarly to (12) one can easily show that the inequality

$$\int\limits_D v^2 dD \le c_0 \int\limits_D v_t^2 dD$$

is valid for some $c_0 = const > 0$ not depending on v. Thus we conclude that, in the space $W_+(W_+^*)$, the norm

$$\|u\|^2_{W_+(W_+^*)} = \int\limits_D (u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2 + u^2)dD$$

is equivalent to the norm

$$\|u\|^2 = \int\limits_D (u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2)dD. \tag{36}$$

Therefore, retaining the previous notation $\|u\|_{W_+(W_+^*)}$ for norm (36), we obtain from (35)

$$(Lu, v)_{L_2(D)} \geq \sigma \inf_D e^{-\lambda t} \|u\|_{L_2(D)} \, \|v\|_{W_+^*}. \tag{37}$$

If now we apply the generalized Schwartz inequality

$$(Lu, v) \leq \|Lu\|_{W_-^*} \, \|v\|_{W_+^*}$$

to the left-hand side of (37), then after reducing by $\|v\|_{W_+^*}$, we obtain inequality (24) where $c = \sigma \inf_D e^{-\lambda t} = const > 0$. Lemma 4 is thereby completely proved. ■

Consider the conditions

$$a_4\big|_{S_2} \geq 0, \quad (\lambda a_4 - a_{4t})\big|_D \geq 0, \tag{38}$$

where the second inequality holds for sufficiently large λ.

Lemma 5. *Let conditions* (5) *and* (38) *be fulfilled. Then for any* $v \in W_+^*$ *the inequality*

$$c\|v\|_{L_2(D)} \leq \|L^* v\|_{W_-} \tag{39}$$

holds for a constant $c = const > 0$ *which does not depend on* $v \in W_+^*$.

Proof. Like in the case of Lemma 4, by Remarks 1 and 3 it is enough to show that inequality (39) is valid for $v \in E_0^*$. Assume that $v \in E_0^*$ and introduce into the consideration the function

$$u(x_1, x_2, t) = \int\limits_{\varphi_1(x_1, x_2)}^{t} e^{\lambda \tau} v(x_1, x_2, \tau) d\tau, \quad \lambda = const > 0,$$

where $t = \varphi_1(x_1, x_2)$ is an equation of the characteristic surface S_1. It is easy to verify that the function $u(x_1, x_2, t)$ belongs to the class E_0 and the following equalities are fulfilled:

$$u_t(x_1, x_2, t) = e^{\lambda t} v(x_1, x_2, t), \quad v(x_1, x_2, t) = e^{-\lambda t} u_t(x_1, x_2, t). \tag{40}$$

From (10), (15) and (40) we have

$$(L^* v, u)_{L_2(D)} = \int\limits_{\partial D} \Big[u \frac{\partial v}{\partial N} - (a_1 \nu_1 + a_2 \nu_2 + a_3 \nu_0) vu \Big] ds +$$

$$+ \int\limits_D [-v_t u_t + v_{x_1} u_{x_1} + x_2^m v_{x_2} u_{x_2} + a_1 v u_{x_1} + a_2 v u_{x_2} + a_3 v u_t +$$

$$+ a_4 uv] dD = - \int\limits_D e^{\lambda t} v_t v \, dD + \int\limits_D e^{-\lambda t} [u_{x_1 t} u_{x_1} + x_2^m u_{x_2 t} u_{x_2}] dD +$$

$$+ \int\limits_D e^{-\lambda t} [a_1 u_{x_1} + a_2 u_{x_2} + a_3 u_t + a_4 u] u_t \, dD. \tag{41}$$

Similarly to (27)–(30), we can prove the equalities

$$-\int\limits_D e^{\lambda t} v_t v \, dD = -\frac{1}{2}\int\limits_{\partial D} e^{\lambda t} v^2 \nu_0 \, ds + \frac{1}{2}\int\limits_D e^{\lambda t}\lambda v^2 \, dD =$$

$$= -\frac{1}{2}\int\limits_{S_1} e^{\lambda t} v^2 \nu_0 \, ds + \frac{1}{2}\int\limits_D e^{-\lambda t}\lambda u_t^2 \, dD =$$

$$= -\frac{1}{2}\int\limits_{S_1} e^{-\lambda t} u_t^2 \nu_0 \, ds + \frac{1}{2}\int\limits_D e^{-\lambda t}\lambda u_t^2 \, dD, \tag{42}$$

$$\int\limits_D e^{-\lambda t}[u_{x_1 t} u_{x_1} + x_2^m u_{x_2 t} u_{x_2}] dD = \frac{1}{2}\int\limits_{\partial D} e^{-\lambda t}[u_{x_1}^2 + x_2^m u_{x_2}^2]\nu_0 \, ds +$$

$$+\frac{1}{2}\int\limits_{\partial D} e^{-\lambda t}\lambda[u_{x_1}^2 + x_2^m u_{x_2}^2] dD, \tag{43}$$

$$(u_t^2 - u_{x_1}^2 - x_2^m u_{x_2}^2)\big|_{S_1} = 0, \tag{44}$$

as well as the inequality

$$-\frac{1}{2}\int\limits_{S_1} e^{-\lambda t} u_t^2 \nu_0 \, ds + \frac{1}{2}\int\limits_{\partial D} e^{-\lambda t}[u_{x_1}^2 + x_2^m u_{x_2}^2]\nu_0 \, ds =$$

$$= -\frac{1}{2}\int\limits_{S_1} e^{-\lambda t} u_t^2 \nu_0 \, ds + \frac{1}{2}\int\limits_{S_1} e^{-\lambda t}[u_{x_1}^2 + x_2^m u_{x_2}^2]\nu_0 \, ds +$$

$$+\frac{1}{2}\int\limits_{S_2} e^{-\lambda t}[u_{x_1}^2 + x_2^m u_{x_2}^2]\nu_0 \, ds \geq$$

$$\geq -\frac{1}{2}\int\limits_{S_1} e^{-\lambda t}[u_t^2 - u_{x_1}^2 - x_2^m u_{x_2}^2]\nu_0 \, ds = 0. \tag{45}$$

In deriving (45), we used the fact that $\nu_0\big|_{S_2} \geq 0$.

By virtue of (42)–(45) equality (41) implies

$$(L^* v, u)_{L_2(D)} \geq \frac{1}{2}\int\limits_D e^{-\lambda t}\lambda[u_t^2 + u_{x_1}^2 + x_2^m u_{x_2}^2] dD +$$

$$+\int\limits_D e^{-\lambda t}[a_1 u_{x_1} + a_2 u_{x_2} + a_3 u_t + a_4 u] u_t \, dD. \tag{46}$$

Using the fact that $\nu_0\big|_{S_2} \geq 0$ and conditions (2), (10), (38) and performing integration by parts, we obtain

$$\int\limits_D e^{-\lambda t} a_4 u u_t dD = \frac{1}{2}\int\limits_{\partial D} e^{-\lambda t} a_4 u^2 \nu_0 ds +$$

$$+\frac{1}{2}\int\limits_D e^{-\lambda t}(\lambda a_4 - a_{4t}) u^2 dD \geq 0. \tag{47}$$

By (47) we find from (46) that

$$(L^*v,u)_{L_2(D)} \geq \frac{1}{2}\int\limits_D e^{-\lambda t}\lambda[u_t^2+u_{x_1}^2+x_2^m u_{x_2}^2]dD+$$

$$+\int\limits_D e^{-\lambda t}[a_1u_{x_1}+a_2u_{x_2}+a_3u_t]u_t\,dD \geq \frac{\lambda}{2}\int\limits_D e^{-\lambda t}[u_t^2+u_{x_1}^2+$$

$$+x_2^m u_{x_2}^2]dD-\Big|\int\limits_D e^{-\lambda t}[a_1u_{x_1}+a_2u_{x_2}+a_3u_t+a_4u]u_t\,dD\Big|.$$

Hence, like in deriving inequality (35), from (33) we obtain

$$(L^*v,u)_{L_2(D)} \geq \Big[\frac{\lambda}{2}-\Big(\frac{1}{2}M+\frac{3}{2}\mu\Big)\Big]\inf_D e^{-\lambda t}\|v\|_{L_2(D)}\ \|u\|_{W_+}. \tag{48}$$

For sufficiently large λ the latter inequality immediately implies (39). This proves Lemma 5. ■

Definition 1. For $F \in L_2(D)$ the function u will be called a strongly generalized solution of problem (1), (2) from the class W_+ provided that $u \in W_+$ and there exists a sequence of functions $u_n \in E_0$ such that $u_n \to u$ in the space W_+ and $Lu_n \to F$ in the space W_-^*, i.e.,

$$\lim_{n\to\infty}\|u_n-u\|_{W_+}=0, \quad \lim_{n\to\infty}\|Lu_n-F\|_{W_-^*}=0.$$

Definition 2. For $F \in W_-^*$ the function u will be called a strongly generalized solution of problem (1), (2) from the class L_2 provided that $u \in L_2(D)$ and there exists a sequence of functions $u_n \in E_0$ such that $u_n \to u$ in the space $L_2(D)$ and $Lu_n \to F$, $n \to \infty$, in the space W_-^*, i.e.,

$$\lim_{n\to\infty}\|u_n-u\|_{L_2(D)}=0. \quad \lim_{n\to\infty}\|Lu_n-F\|_{W_-^*}=0.$$

By the results of [13] Lemmas 2–5 give rise to the following theorems.

Theorem 1. *Let conditions* (5), (23) *and* (38) *be fulfilled. Then for any* $F \in W_-^*$ *there exists a unique strongly generalized solution* u *of problem* (1), (2) *from the class* L_2, *for which the estimate*

$$\|u\|_{L_2(D)} \leq c\|F\|_{W_-^*} \tag{49}$$

where the positive constant c *does not depend on* F, *is valid.*

Theorem 2. *Let conditions* (5), (23) *and* (38) *be fulfilled. Then for any* $F \in L_2(D)$ *there exists a unique strongly generalized solution* u *of problem* (1), (2) *from the class* W_+, *for which estimate* (49) *holds.*

Similar results hold for problem (3), (4) as well.

References

1. J. Hadamard, Lectures on Cauchy's problem in partial differential equations. *New Haven: Yale University Press*, 1933.

2. J. Tolen, Problém de Cauchy sur la deux hypersurfaces caracteristiques sécantes. *C.R. Acad. Sci. Paris Sér.* A-B291(1980), No. 1, 49–52.

3. S. Kharibegashvili, On a characteristic problem for the wave equation. *Proc. I. Vekua Inst. Appl. Math., Tbilisi University Press* **47**(1922), 76–82.

4. S. Kharibegashvili, On a spatial problem of Darboux type for a second order hypebolic equation. *Georgian Math. J.* **2**(1995), No. 3, 299–311.

5. S. Kharibegashvili, On the solvability of a spatial problem of Darboux type for the wave equation. *Georgian Math. J.* **2**(1995), No. 4, 385–394.

6. S. Kharibegashvili, On the solvability of a noncharacteristic spatial problem of Darboux type for the wave equation. *Georgian Math. J.* **3**(1996), No. 1, 53–68.

7. S. Kharibegashvili, On the solvability of a multidimensional version of the first Darboux problem for a modal second order degenerating hyperbolic equation. *Georgian Math. J.* **4**(1997), No. 4, 341–354.

8. A. V. Bitsadze, On mixed type equations on three-dimensional domains (Russian). *Dokl. Akad. Nauk SSSR* **143**(1962), No. 5, 1017–1019.

9. A. M. Nakhushev, A multidimensional analogy of the Darboux problem for hyperbolic equations (Russian). *Dokl. Akad. Nauk SSSR* **194**(1970), No. 1, 31–34.

10. T. Sh. Kalmenov, On multidimensional regular boundary value problems for the wave equation (Russian). *Izv. Akad. Nauk Kazakh. SSR. Ser. Fiz.-Mat.* (1982), No. 3, 18–25.

11. H. Triebel, Interpolation theory, function spaces, differential operators. *VEB Deutscher Verlag der Wissenshchaften, Berlin,* 1978.

12. Yu. Berezanski. Expansion with respect to the eigenfunctions of self-conjugate operators (Russian). *Naukova Dumka, Kiev,* 1965.

13. I. I. Lyashko, V. P. Didenko, O. Ye. Tsitritski. Filtration of noises (Russian). *Naukova Dumka, Kiev*, 1979.

(Received 27.12.1994)

Author's address:
Department of Theoretical Mechanics (4)
Georgian Technical University
77, M. Kostava St., Tbilisi 380075
Georgia