

## MULTI POINT BOUNDARY VALUE PROBLEMS FOR SECOND ORDER DIFFERENTIAL INCLUSIONS

M. Benchora and S. K. Ntouyas

**Abstract.** In this paper we investigate the existence of solutions on a compact interval to a multi-point boundary value problem for a class of second order differential inclusions. We shall rely on a fixed point theorem for condensing maps due to Martelli.

### 1. Introduction

Let  $a_i, b_j \in \mathbf{R}$ , with all of the  $a_i$ 's, (respectively,  $b_j$ 's), having the same sign,  $\xi_i, \zeta_j \in (0, 1)$ ,  $i = 1, 2, \dots, m-2$ ,  $j = 1, 2, \dots, n-2$ ,  $0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < 1$ ,  $0 < \zeta_1 < \zeta_2 < \dots < \zeta_{n-2} < 1$ . The main purpose of this paper is to get results on the solvability of the following boundary value problems (BVPs for short) for second order differential inclusions of the forms

$$\begin{cases} y''(t) \in F(t, y(t)), & t \in (0, 1) \\ y(0) = \sum_{i=1}^{m-2} a_i y'(\xi_i), & y(1) = \sum_{j=1}^{n-2} b_j y(\zeta_j) \end{cases} \quad (\text{A})$$

$$\begin{cases} y''(t) \in F(t, y(t)), & t \in (0, 1) \\ y(0) = \sum_{i=1}^{m-2} a_i y'(\xi_i), & y'(1) = \sum_{j=1}^{n-2} b_j y'(\zeta_j) \end{cases} \quad (\text{B})$$

$$\begin{cases} y''(t) \in F(t, y(t)), & t \in (0, 1) \\ y(0) = \sum_{i=1}^{m-2} a_i y(\xi_i), & y(1) = \sum_{j=1}^{n-2} b_j y(\zeta_j) \end{cases} \quad (\text{C})$$

and

$$\begin{cases} y''(t) \in F(t, y(t)), & t \in (0, 1) \\ y(0) = \sum_{i=1}^{m-2} a_i y(\xi_i), & y'(1) = \sum_{j=1}^{n-2} b_j y'(\zeta_j) \end{cases} \quad (\text{D})$$

where  $F: J \times \mathbf{R} \rightarrow 2^{\mathbf{R}}$  is a multivalued map with compact convex values.

*AMS Subject Classification:* 34 A 60, 34 B 10, 34 B 15

*Keywords and phrases:* Multi point boundary value problems, differential inclusion, existence, fixed point

The study of multi-point boundary value problems for second order ordinary differential equations was initiated by Il'in and Moiseev in [12, 13] motivated by the work of Bitsadze and Samarskii on nonlocal elliptic boundary value problems, [2, 3, 4].

Existence of solutions on compact intervals for multi-point boundary value problems for second order differential equations was given by Gupta in [6], Gupta et al in [7–10]. However, to our knowledge, this type of problems has not been studied for the multivalued case.

It is well known (c.f. [12]) that if a function  $y \in C^1$  satisfies one of the boundary conditions stated above and  $a_i, b_j, i = 1, 2, \dots, m-2, j = 1, 2, \dots, n-2$  are as above, then there exist  $\eta \in [\xi_1, \xi_{m-2}], \tau \in [\zeta_1, \zeta_{n-2}]$  such that

$$y(0) = \alpha y'(\eta), \quad y(1) = \beta y(\tau)$$

$$y(0) = \alpha y'(\eta), \quad y'(1) = \beta y'(\tau)$$

$$y(0) = \alpha y(\eta), \quad y(1) = \beta y(\tau)$$

$$y(0) = \alpha y(\eta), \quad y'(1) = \beta y'(\tau)$$

respectively with  $\alpha = \sum_{i=1}^{m-2} a_i, \beta = \sum_{j=1}^{n-2} b_j$ . Hence the multi-point BVPs (A)–(D) can be reduced to a corresponding four-point BVP. The method of proof for the existence of a solution for a four-point BVP and for a multi-point BVP (A)–(D) is the same.

In order not to hide the main ideas behind general and technically complicated statements, we restrict our discussion to the following four-point BVP

$$y'' \in F(t, y), \quad t \in J = [0, 1] \tag{1.1}$$

$$y(0) = y'(\eta), \quad y(1) = y(\tau) \tag{1.2}$$

where  $F: J \times \mathbf{R} \rightarrow 2^{\mathbf{R}}$  is a multivalued map with compact convex values and  $\eta, \tau \in (0, 1)$ . This is a special case of the BVP (A) when  $\alpha = \beta = 1$ . All the other four-point BVP and the general multi-point BVP are examined in a similar way, with obvious modifications.

The method we are going to use is to reduce the existence of solutions to problem (1.1)–(1.2) to the search for fixed points of a suitable multivalued map on the Banach space  $C(J, \mathbf{R})$ . In order to prove the existence of fixed points, we shall rely on a fixed point theorem for condensing maps due to Martelli [15].

## 2. Preliminaries

In this section, we introduce notations, definitions, and preliminary facts from multivalued analysis which are used throughout this paper.

Let  $(X, \|\cdot\|)$  be a Banach space. A multivalued map  $G: X \rightarrow 2^X$  is convex (closed) valued if  $G(x)$  is convex (closed) for all  $x \in X$ .  $G$  is bounded on bounded sets if  $G(B) = \bigcup_{x \in B} G(x)$  is bounded in  $X$  for any bounded set  $B$  of  $X$  (i.e.  $\sup_{x \in B} \{\sup\{\|y\| : y \in G(x)\}\} < \infty$ ).

$G$  is called upper semicontinuous (u.s.c.) on  $X$  if for each  $x_* \in X$  the set  $G(x_*)$  is a nonempty, closed subset of  $X$ , and if for each open set  $B$  of  $X$  containing  $G(x_*)$ , there exists an open neighbourhood  $V$  of  $x_*$  such that  $G(V) \subseteq B$ .

$G$  is said to be completely continuous if  $G(B)$  is relatively compact for every bounded subset  $B \subseteq X$ .

If the multivalued map  $G$  is completely continuous with nonempty compact values, then  $G$  is u.s.c. if and only if  $G$  has a closed graph (i.e.  $x_n \rightarrow x_*$ ,  $y_n \rightarrow y_*$ ,  $y_n \in G(x_n)$  imply  $y_* \in G(x_*)$ ).

$G$  has a fixed point if there is  $x \in X$  such that  $x \in G(x)$ .

In the following  $CC(X)$  denotes the set of all nonempty compact and convex subsets of  $X$ .

A multivalued map  $G: J \rightarrow CC(E)$  is said to be measurable if for each  $x \in E$  the function  $Y: J \rightarrow \mathbf{R}$  defined by

$$Y(t) = d(x, G(t)) = \inf\{|x - z| : z \in G(t)\}$$

is measurable.

DEFINITION 2.1. A multivalued map  $F: J \times \mathbf{R} \rightarrow 2^{\mathbf{R}}$  is said to be an  $L^1$ -Carathéodory map if

- (i)  $t \mapsto F(t, y)$  is measurable for each  $y \in \mathbf{R}$ ;
- (ii)  $y \mapsto F(t, y)$  is upper semicontinuous for almost all  $t \in J$ ;
- (iii) for each  $k > 0$ , there exists  $h_k \in L^1(J, \mathbf{R}_+)$  such that

$$\|F(t, y)\| = \sup\{|v| : v \in F(t, y)\} \leq h_k(t)$$

for all  $|y| \leq k$  and for almost all  $t \in J$ .

An upper semi-continuous map  $G: X \rightarrow 2^X$  is said to be condensing if for any subset  $B \subseteq X$  with  $\alpha(B) \neq 0$ , we have  $\alpha(G(B)) < \alpha(B)$ , where  $\alpha$  denotes the Kuratowski measure of noncompactness. For properties of the Kuratowski measure, we refer to Banas and Goebel [1].

We remark that a completely continuous multivalued map is the easiest example of a condensing map. For more details on multivalued maps see the books of Deimling [5] and Hu and Papageorgiou [11].

We will need the following hypotheses:

- (H1)  $F: J \times \mathbf{R} \rightarrow CC(\mathbf{R})$  is an  $L^1$ -Carathéodory multivalued map.
- (H2) There exists a function  $H \in L^1(J, \mathbf{R}_+)$  such that

$$\|F(t, y)\| := \sup\{|v| : v \in F(t, y)\} \leq H(t) \text{ for almost all } t \in J \text{ and all } y \in \mathbf{R}.$$

DEFINITION 2.2. A function  $y: J \rightarrow \mathbf{R}$  is called a solution for the BVP (1.1)–(1.2) if  $y$  and its first derivative are absolutely continuous and  $y''$  (which exists almost everywhere) satisfies the differential inclusion (1.1) a.e. on  $J$  and the condition (1.2).

Our considerations are based on the following lemmas.

LEMMA 2.3. [14] *Let  $I$  be a compact real interval and  $X$  be a Banach space. If  $F$  is a multivalued map satisfying (H1) and  $\Gamma$  is a linear continuous mapping from  $L^1(I, X)$  to  $C(I, X)$ , then the operator*

$$\Gamma \circ S_F: C(I, X) \longrightarrow CC(C(I, X)), \quad y \longmapsto (\Gamma \circ S_F)(y) := \Gamma(S_{F,y})$$

*is a closed graph operator in  $C(I, X) \times C(I, X)$ .*

LEMMA 2.4. [15] *Let  $X$  be a Banach space and  $N: X \longrightarrow CC(X)$  be a u.s.c. condensing map. If the set*

$$\Omega := \{y \in X : \lambda y \in Ny \text{ for some } \lambda > 1\}$$

*is bounded, then  $N$  has a fixed point.*

### 3. Main Result

Now, we are able to state and prove our main theorem.

THEOREM 3.1. *Assume that Hypotheses (H1)–(H2) hold. Then the BVP (1.1)–(1.2) has at least one solution on  $J$ .*

*Proof.* Let  $C(J, \mathbf{R})$  be the Banach space provided with the norm

$$\|y\|_\infty := \sup\{|y(t)| : t \in J\}, \text{ for } y \in C(J, \mathbf{R}).$$

Transform the problem (1.1)–(1.2) into a fixed point problem. Consider the multivalued map,  $N: C(J, \mathbf{R}) \longrightarrow 2^{C(J, \mathbf{R})}$  defined by:

$$Ny = \left\{ h \in C(J, \mathbf{R}) : h(t) = \int_0^t (t-s)g(s) ds + \int_0^\eta g(s) ds \right. \\ \left. + \frac{1+t}{1-\tau} \left[ \int_0^\tau (\tau-s)g(s) ds - \int_0^1 (1-s)g(s) ds \right] \right\}$$

where

$$g \in S_{F,y} = \left\{ g \in L^1(J, \mathbf{R}) : g(t) \in F(t, y(t)) \text{ for a.e. in } J \right\}.$$

REMARK 3.2. (i) It is clear that the fixed points of  $N$  are solutions to (1.1)–(1.2).

(ii) For each  $y \in C(J, \mathbf{R})$  the set  $S_{F,y}$  is nonempty (see Lasota and Opial [14]).

We shall show that  $N$  satisfies the assumptions of Lemma 2.4. The proof will be given in several steps.

*Step 1.*  $Ny$  is convex for each  $y \in C(J, \mathbf{R})$ .

Indeed, if  $h_1, h_2$  belong to  $Ny$ , then there exist  $g_1, g_2 \in S_{F,y}$  such that for each  $t \in J$  we have

$$h(t) = \int_0^t (t-s)g_i(s) ds + \int_0^\eta g_i(s) ds \\ + \frac{1+t}{1-\tau} \left[ \int_0^\tau (\tau-s)g_i(s) ds - \int_0^1 (1-s)g_i(s) ds \right], \quad i = 1, 2.$$

Let  $0 \leq \alpha \leq 1$ . Then for each  $t \in J$  we have

$$\begin{aligned} (\alpha h_1 + (1 - \alpha)h_2)(t) &= \int_0^t (t - s)\{\alpha g_1(s) + (1 - \alpha)g_2(s)\} ds \\ &\quad + \int_0^\eta \{\alpha g_1(s) + (1 - \alpha)g_2(s)\} ds \\ &\quad + \frac{1+t}{1-\tau} \left[ \int_0^\tau (\tau - s)\{\alpha g_1(s) + (1 - \alpha)g_2(s)\} ds \right. \\ &\quad \left. - \int_0^1 (1 - s)\{\alpha g_1(s) + (1 - \alpha)g_2(s)\} ds \right]. \end{aligned}$$

Since  $S_{F,y}$  is convex (because  $F$  has convex values) then

$$\alpha h_1 + (1 - \alpha)h_2 \in Ny.$$

*Step 2.  $N$  is bounded on bounded sets of  $C(J, \mathbf{R})$ .*

Indeed, it is enough to show that there exists a positive constant  $c$  such that for each  $h \in Ny$ ,  $y \in B_r = \{y \in C(J, \mathbf{R}) : \|y\|_\infty \leq r\}$  one has  $\|h\|_\infty \leq c$ .

If  $h \in Ny$ , then there exists  $g \in S_{F,y}$  such that for each  $t \in J$  we have

$$\begin{aligned} h(t) &= \int_0^t (t - s)g(s) ds + \int_0^\eta g(s) ds \\ &\quad + \frac{1+t}{1-\tau} \left[ \int_0^\tau (\tau - s)g(s) ds - \int_0^1 (1 - s)g(s) ds \right], \quad t \in J. \end{aligned}$$

By (H1) we have for each  $t \in J$  that

$$|h(t)| \leq \int_0^t h_r(s) ds + \int_0^\eta h_r(s) ds + \frac{2}{1-\tau} \left[ \int_0^\tau (\tau - s)h_r(s) ds + \int_0^1 (1 - s)h_r(s) ds \right].$$

Then

$$\|h\|_\infty \leq \int_0^1 h_r(s) ds + \int_0^\eta h_r(s) ds + \frac{2}{1-\tau} \left[ \int_0^\tau (\tau - s)h_r(s) ds + \int_0^1 (1 - s)h_r(s) ds \right] = c.$$

*Step 3.  $N$  sends bounded sets of  $C(J, \mathbf{R})$  into equicontinuous sets.*

Let  $t_1, t_2 \in J$ ,  $t_1 < t_2$  and  $B_r$  be a bounded set of  $C(J, \mathbf{R})$ . For each  $y \in B_r$  and  $h \in Ny$ , there exists  $g \in S_{F,y}$  such that

$$\begin{aligned} h(t) &= \int_0^t (t - s)g(s) ds + \int_0^\eta g(s) ds \\ &\quad + \frac{1+t}{1-\tau} \left[ \int_0^\tau (\tau - s)g(s) ds - \int_0^1 (1 - s)g(s) ds \right], \quad t \in J. \end{aligned}$$

Thus we obtain

$$\begin{aligned}
|h(t_2) - h(t_1)| &\leq \int_0^{t_2} (t_2 - s) \|g(s)\| ds + \int_{t_1}^{t_2} (t_1 - s) \|g(s)\| ds \\
&\quad + \frac{t_2 - t_1}{1 - \tau} \left[ \int_0^\tau (\tau - s) \|g(s)\| ds + \int_0^1 (1 - s) \|g(s)\| ds \right] \\
&\leq \int_0^{t_2} (t_2 - s) h_r(s) ds + \int_{t_1}^{t_2} (t_1 - s) h_r(s) ds \\
&\quad + \frac{t_2 - t_1}{1 - \tau} \left[ \int_0^\tau (\tau - s) h_r(s) ds + \int_0^1 (1 - s) h_r(s) ds \right].
\end{aligned}$$

As  $t_2 \rightarrow t_1$  the right-hand side of the above inequality tends to zero.

As a consequence of Step 2, Step 3 together with the Arzela-Ascoli theorem we can conclude that  $N$  is completely continuous.

*Step 4.  $N$  has a closed graph.*

Let  $y_n \rightarrow y_*$ ,  $h_n \in Ny_n$ , and  $h_n \rightarrow h_*$ . We shall prove that  $h_* \in Ny_*$ .

$h_n \in Ny_n$  means that there exists  $g_n \in S_{F, y_n}$  such that

$$\begin{aligned}
h_n(t) &= \int_0^t (t - s) g_n(s) ds + \int_0^\eta g_n(s) ds \\
&\quad + \frac{1 + t}{1 - \tau} \left[ \int_0^\tau (\tau - s) g_n(s) ds - \int_0^1 (1 - s) g_n(s) ds \right], \quad t \in J.
\end{aligned}$$

We must prove that there exists  $g_* \in S_{F, y_*}$  such that

$$\begin{aligned}
h_*(t) &= \int_0^t (t - s) g_*(s) ds + \int_0^\eta g_*(s) ds \\
&\quad + \frac{1 + t}{1 - \tau} \left[ \int_0^\tau (\tau - s) g_*(s) ds - \int_0^1 (1 - s) g_*(s) ds \right], \quad t \in J.
\end{aligned}$$

Now, we consider the linear continuous operator

$$\Gamma: L^1(J, \mathbf{R}) \rightarrow C(J, \mathbf{R})$$

$$\begin{aligned}
g \mapsto \Gamma(g)(t) &= \int_0^t (t - s) g(s) ds + \int_0^\eta g(s) ds \\
&\quad + \frac{1 + t}{1 - \tau} \left[ \int_0^\tau (\tau - s) g(s) ds - \int_0^1 (1 - s) g(s) ds \right], \quad t \in J.
\end{aligned}$$

From Lemma 2.3, it follows that  $\Gamma \circ S_F$  is a closed graph operator.

Moreover from the definition of  $\Gamma$  we have

$$h_n(t) \in \Gamma(S_{F, y_n}).$$

Since  $y_n \rightarrow y_*$ , it follows from Lemma 2.3 that

$$\begin{aligned}
h_*(t) &= \int_0^t (t - s) g_*(s) ds + \int_0^\eta g_*(s) ds \\
&\quad + \frac{1 + t}{1 - \tau} \left[ \int_0^\tau (\tau - s) g_*(s) ds - \int_0^1 (1 - s) g_*(s) ds \right], \quad t \in J
\end{aligned}$$

for some  $g_* \in S_{F, y_*}$ .

Step 5. The set

$$\Omega := \{y \in C(J, \mathbf{R}) : \lambda y \in Ny \text{ for some } \lambda > 1\}$$

is bounded.

Let  $y \in \Omega$ . Then  $\lambda y \in Ny$  for some  $\lambda > 1$ . Thus there exists  $g \in S_{F,y}$  such that

$$\begin{aligned} y(t) &= \lambda^{-1} \int_0^t (t-s)g(s) ds + \lambda^{-1} \int_0^\eta g(s) ds \\ &\quad + \lambda^{-1} \frac{1+t}{1-\tau} \left[ \int_0^\tau (\tau-s)g(s) ds - \int_0^1 (1-s)g(s) ds \right], \quad t \in J. \end{aligned}$$

This implies by (H2) that for each  $t \in J$  we have

$$|y(t)| \leq \int_0^t (t-s)H(s) ds + \int_0^\eta H(s) ds + \frac{2}{1-\tau} \left[ \int_0^\tau (\tau-s)H(s) ds + \int_0^1 (1-s)H(s) ds \right].$$

Thus

$$\begin{aligned} \|y\|_\infty &\leq \int_0^1 (1-s)H(s) ds + \int_0^\eta H(s) ds \\ &\quad + \frac{2}{1-\tau} \left[ \int_0^\tau (\tau-s)H(s) ds + \int_0^1 (1-s)H(s) ds \right] = K. \end{aligned}$$

This shows that  $\Omega$  is bounded.

Set  $X := C(J, \mathbf{R})$ . As a consequence of Lemma 2.4 we deduce that  $N$  has a fixed point which is a solution of (1.1)–(1.2) on  $J$ . ■

#### REFERENCES

- [1] J. Banas and K. Goebel, *Measures of Noncompactness in Banach Spaces*, Marcel-Dekker, New York, 1980.
- [2] A. V. Bitsadze, *On the theory of nonlocal boundary value problem*, Soviet. Math. Dokl. **30** (1) (1984), 8–10.
- [3] A. V. Bitsadze, *On a class of conditionally solvable nonlocal boundary value problems for harmonic functions*, Soviet. Math. Dokl. **31** (1) (1985), 91–94.
- [4] A. V. Bitsadze and A. A. Samarskii, *On some simple generalizations of linear elliptic boundary value problem*, Soviet. Math. Dokl. **10** (2) (1969), 398–400.
- [5] K. Deimling, *Multivalued Differential Equations*, Walter de Gruyter, Berlin-New York, 1992.
- [6] C. P. Gupta, *A generalized multi-point boundary value problem for a second order ordinary differential equation*, Appl. Math. Comput. **89** (1998), 133–146.
- [7] C.P. Gupta, S.K. Ntouyas and P.Ch. Tsamatos, *On an m-point boundary value problem for second order ordinary differential equations*, Nonlinear Anal. **23** (1994), 1427–1436.
- [8] C.P. Gupta, S.K. Ntouyas and P.Ch. Tsamatos, *Existence results for m-point boundary value problems*, Differ. Equ. Dyn. Syst. **2** (1994), 289–298.
- [9] C.P. Gupta, S.K. Ntouyas and P.Ch. Tsamatos, *On the solvability of some m-point boundary value problems*, Appl. Math. **41** (1996), 1–17.
- [10] C.P. Gupta, S.K. Ntouyas and P.Ch. Tsamatos, *Existence results for multi-point boundary value problem for second order ordinary differential equations*, Bull. Greek Math. Soc., to appear.

- [11] Sh. Hu and N. Papageorgiou, *Handbook of Multivalued Analysis, Volume I: Theory*, Kluwer Academic Publishers, Dordrecht, 1997.
- [12] V.A. Il'in and E.I. Moiseev, *Nonlocal boundary value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects*, Differential Equations **23** (7) (1987), 803–810.
- [13] V.A. Il'in and E.I. Moiseev, *Nonlocal boundary value problem of the second kind for a Sturm-Liouville operator*, Differential Equations **23** (8) (1987), 979–987.
- [14] A. Lasota and Z. Opial, *An application of the Kakutani-Ky Fan theorem in the theory of ordinary differential equations*, Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys. **13** (1965), 781–786.
- [15] M. Martelli, *A Rothe's type theorem for noncompact acyclic-valued map*, Boll. Un. Math. Ital. **11** (1975), 3, suppl., 70–76.

(received 14.02.2001, in revised form 30.11.2001)

Département de Mathématiques, Université de Sidi Bel Abbès, BP 89, 22000 Sidi Bel Abbès, Algérie

*E-mail*: benchohra@yahoo.com

Department of Mathematics, University of Ioannina, 451 10 Ioannina, Greece

*E-mail*: sntouyas@cc.uoi.gr